

# Evaluation of Inflammatory Response System (IRS) and Compensatory Immune Response System (CIRS) in Adolescent Major Depression

Nikola Ferencova<sup>1</sup>, Zuzana Visnovcova<sup>1</sup>, Igor Ondrejka<sup>2</sup>, Dana Funakova<sup>2</sup>, Igor Hrtanek<sup>2</sup>, Simona Kelcikova<sup>3</sup>, Ingrid Tonhajzerova<sup>4</sup>

<sup>1</sup>Biomedical Centre Martin, Jessenius Faculty of Medicine in Martin, Comenius University in Bratislava, Martin, Slovak Republic; <sup>2</sup>Psychiatric Clinic, Jessenius Faculty of Medicine in Martin, Comenius University in Bratislava, University Hospital Martin, Martin, Slovak Republic; <sup>3</sup>Department of Midwifery, Jessenius Faculty of Medicine in Martin, Comenius University in Bratislava, Martin, Slovak Republic; <sup>4</sup>Department of Physiology, Jessenius Faculty of Medicine in Martin, Comenius University in Bratislava, Martin, Slovak Republic

Correspondence: Ingrid Tonhajzerova, Department of Physiology, Jessenius Faculty of Medicine in Martin, Comenius University in Bratislava, Mala Hora 4C, Martin, Slovak Republic, Tel +421432633425, Email [ingrid.tonhajzerova@uniba.sk](mailto:ingrid.tonhajzerova@uniba.sk)

**Purpose:** Nowadays, the role of two tightly interconnected systems, the inflammatory response system (IRS) and the compensatory immune response system (CIRS) in depression, is increasingly discussed. Various studies indicate pro-inflammatory activity in adolescent depression; however, there is an almost complete lack of findings about IRS and CIRS balance. Thus, we aimed to assess different IRS and CIRS indices, profiles, and IRS/CIRS ratios in drug-naïve MDD patients at adolescent age, with respect to sex.

**Patients and Methods:** One hundred MDD adolescents (40 boys, average age: 15.4±1.2 yrs.) and 60 controls (28 boys, average age: 15.3±1.5 yrs.) were examined. Evaluated parameters were 1. plasma levels of interleukin (IL)-1 $\alpha$ , IL-1 $\beta$ , IL-2, IL-4, IL-6, IL-8, IL-10, interferon gamma, tumor necrosis factor alpha (TNF- $\alpha$ ), soluble receptor of IL-6 (sIL-6R), soluble receptors of TNF- $\alpha$  (sTNF-R1, sTNF-R2); 2. profiles: IL-6 trans-signaling, M1 macrophage signaling, helper T lymphocytes (Th) 1 profile, regulatory T lymphocytes (Treg)+Th2, allIRS, and allCIRS; 3. IRS vs CIRS activity ratios: TNF- $\alpha$ /TNF-R1, TNF- $\alpha$ /TNF-R2, TNF- $\alpha$ /sTNF-Rs (ie sTNF-R1 +sTNF-R2), Th1/Th2, Th1/Treg, Th1/Th2+Treg, M1/Th2, M1/Treg, M1/Treg+Th2, allIRS/allCIRS.

**Results:** MDD patients showed increased IL-4, IL-10, TNF- $\alpha$ , sIL-6R, Treg+Th2, allIRS, allCIRS, and TNF- $\alpha$ /sTNF-Rs, and decreased Th1/Th2+Treg. MDD females showed increased IL-10 and TNF- $\alpha$  compared to control females. MDD males showed increased IL-4, IL-10, sIL-6R, Treg+Th2, and TNF- $\alpha$ /TNF-R1 compared to control males. Increased sTNF-R1 was found in MDD males compared to MDD females. Positive correlations were found between CDI score and sIL-6R and IL-10 in the total group and between CDI score and IL-10 in adolescent males.

**Conclusion:** Our study for the first time extensively evaluated IRS and CIRS interactions revealing enhanced pro-inflammatory TNF- $\alpha$  signaling and IL-6 trans-signaling in association with increased IL-10- and IL-4-mediated anti-inflammatory activity in first-episode depression at the adolescent age. Moreover, results reflect the sex-specific simultaneous activation of IRS and CIRS pathways in adolescent depression.

**Keywords:** major depressive disorder, adolescent age, comprehensive cytokine analysis, neuroimmune interactions, sex differences

## Introduction

Major depressive disorder (MDD) represents a common mood disorder affecting approximately 300 million people worldwide<sup>1</sup> with increasing incidence, particularly at adolescent age.<sup>2,3</sup> Moreover, MDD is the largest mental health contributor to the disease burden in adolescents.<sup>4</sup> However, the response rates to the psycho- and pharmaco-therapy in adolescent depression are only modest,<sup>5</sup> thus a better understanding of the underlying pathophysiological mechanisms in MDD can contribute to personalized assessment and treatment of depression. Although various risk factors and etiological models in depression have been proposed, in the last few decades, the role of immune dysregulation as one of the most robust mechanisms contributing to MDD pathophysiology has been increasingly discussed.<sup>6,7</sup> The presence

of the inflammatory process in depressive patients can be easily revealed by complete blood cells count reflecting cell blood composition including red blood cells (RBC), RBC distribution width (RDW), platelets, mean platelet volume, and white blood cells (WBC) with differential (eg neutrophils and lymphocytes) as one of the most widespread and frequently used laboratory tests in the clinical setting. More specifically, increased WBC as indicators of the overall immune system activity associated with neutrophilia have been repeatedly reported in adult and adolescent depressive patients compared to controls.<sup>8–11</sup> Further, measures of complete blood cells count can be used for the estimation of the neutrophil-to-lymphocyte ratio (NLR), platelet-to-lymphocyte ratio (PLR), and RDW-to-platelet ratio – other parameters determined as markers of conditions associated with inflammation including depression.<sup>11,12</sup> Specifically, increased NLR and PLR have been the most replicated finding in adult and also adolescent MDD studies.<sup>10–15</sup> In summary, this hematological test can represent a helpful method for identifying the subgroup of depressive patients with “inflamed” depression.


Nowadays, the role of two tightly interconnected systems – inflammatory response system (IRS) and compensatory immune response system (CIRS) – is at the center of scientific interest in the inflammation–depression relationship according to their relevance for the activity of the specific parts of the immune system. However, the precise understanding of the IRS and/or CIRS dysregulation in MDD pathogenesis is just in the beginning.<sup>16,17</sup>

## Inflammatory Response System and Compensatory Immune Response System – Focus on the Adolescent Depression

Various immune cells including macrophages and T lymphocytes, and their effector cytokines seem to play a crucial role in IRS and CIRS pathways.<sup>16</sup> Macrophages can be through the macrophage polarization process discriminated into M1 macrophages whose activation leads to the secretion of pro-inflammatory cytokines such as interleukin (IL)-1 $\beta$ , IL-6, and tumor necrosis factor-alpha (TNF- $\alpha$ ), or into M2 macrophages releasing anti-inflammatory cytokines (eg IL-10).<sup>18</sup> T lymphocytes express on their surface T cell receptor and CD4 (ie helper T lymphocytes) or CD8 (ie cytotoxic T lymphocytes) glycoprotein. The helper T lymphocytes (Th) differentiate into multiple subsets characterized by distinct immune functions (eg Th1 cells release pro-inflammatory IL-2 and interferon-gamma (IFN- $\gamma$ ), Th2 cells release anti-inflammatory IL-4, Th17 release pro-inflammatory IL-17, and T regulatory (Treg) cells release anti-inflammatory IL-10).<sup>19</sup> Additionally, macrophages and T lymphocytes are bi-directionally modulated. While M1 macrophages promote Th1 responses, M2 macrophages drive the differentiation and recruitment of Th2 and Treg lymphocytes. On the other hand, cytokines released by T lymphocytes influence the macrophage polarization; the exposure of naïve monocytes to Th1 cytokines drives the macrophage polarization towards the M1 phenotype, while Th2 and Treg cytokines promote macrophage polarization towards the M2 phenotype.<sup>20,21</sup>

With respect to adult depression, IRS is characterized by mild chronic inflammation caused by M1 macrophages activation leading to increased production of TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and soluble IL-6 receptor (sIL-6R) as well as enhanced Th1 and Th17 responses associated with increased production of IFN- $\gamma$ , IL-2, and IL-17. Many meta-analyses confirmed that adult depression is accompanied by an immune-inflammatory response with increased levels of pro-inflammatory cytokines, acute phase proteins, and other components released by activated IRS.<sup>22–27</sup> On the other hand, MDD-linked CIRS components likely counter-regulating IRS effects involve the increased Th2 and Treg activity resulting in elevated production of anti-inflammatory cytokines IL-4 and IL-10, increased levels of soluble IL-1 receptor antagonist (sIL-1RA), soluble IL-2 receptor (sIL-2R), and soluble TNF receptors (sTNF-R1, sTNF-R2).<sup>16,17,25</sup> The main elements of IRS and CIRS pathways in MDD are presented in [Figure 1](#).

Despite multiple studies evaluating the relationship between adolescent depression and pro-inflammatory/anti-inflammatory cytokines, the precise IRS- and CIRS-related knowledge is severely limited by the inconsistency of individual studies. More specifically, pro-inflammatory IL-2 and IFN- $\gamma$ , as the cytokines showing case–control significant differences in most studies, have been increased,<sup>28,29</sup> decreased<sup>30</sup> or unchanged<sup>31</sup> in adolescent depressive patients compared to healthy controls. Further, pro-inflammatory IL-1 $\beta$ , IL-6, and TNF- $\alpha$  as the most frequently evaluated cytokines in multiple studies<sup>30–38</sup> have shown significant differences between depressive adolescents and controls (ie increased levels of these cytokines in depression) only in few of them.<sup>33,37</sup> Regarding anti-inflammatory



**Figure 1** Simplified schematic diagram of crucial immune cells and mediators involved in inflammatory response system (IRS) and compensatory immune response system (CIRS) pathways in major depression. IRS pathway includes: 1. activated M1 macrophages releasing interleukin (IL)-1 $\beta$ , IL-6 associated with soluble IL-6 receptor (sIL-6R), tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ); 2. activated T helper 1 cells (Th1) releasing IFN- $\gamma$ ; and 3. activated Th17 cells releasing IL-17. CIRS pathway consists of: 1. activated M2 macrophages releasing IL-10; 2. activated Th2 cells releasing IL-4; and 3. activated T regulatory (Treg) cells releasing IL-10. Other markers involved in CIRS pathway include soluble cytokine receptors (sIL-2R, sTNF-R1, sTNF-R2) and soluble IL-1 receptor antagonist (sIL-1RA). Macrophages and T-lymphocytes interplay with each other and can drive their polarization. More specifically, Th cells (Th1, Th2, Th17) drive the polarization of M0 to M1 and/or M2 macrophages. Moreover, Th1 cells bi-directionally interact with M1, while Th2 and Treg cells with M2.

cytokines IL-4 and IL-10, no significant differences in IL-4 and/or IL-10,<sup>28,31,39</sup> increased levels of IL-4 and/or IL-10,<sup>29,33</sup> or decreased levels of IL-10<sup>30</sup> were found in adolescent depression compared to healthy controls. Similarly, regarding meta-analyses, D'Acunto et al<sup>40</sup> reported significantly increased levels of TNF- $\alpha$  with no differences in other included cytokines (ie, IL-1  $\beta$ , IL-4, IL-6, IL-8, IL-10, and IFN- $\gamma$ ) in adolescent MDD patients, while more recent meta-analysis revealed significantly higher levels of IL-1 $\beta$  with no differences in the levels of the other cytokines (IL-2, IL-6, IFN- $\gamma$ , and TNF- $\alpha$ ) in depressive adolescents compared to healthy controls.<sup>41</sup> Moreover, a meta-analysis by Colasanto et al<sup>42</sup> reported positive associations between depression and IL-6 in adolescents, with IL-6 as a predictor of future depression. Lastly, no relationship between IL-1 $\beta$ , IL-2, IL-6, IL-10, IFN- $\gamma$ , and TNF- $\alpha$  and pediatric internalizing disorders was observed in the meta-analysis by Howe and Lynch.<sup>43</sup> The heterogeneous findings in the relation to depression and inflammation can be influenced by various factors including genetics, medication use, and sex. Thus, further research in this area is needed to reveal the exact depression-linked immune alterations.

## Genetic Contribution in the Relationship Between Inflammation and Depression

The substantial heritable link between depression and inflammation<sup>44</sup> has determined a potential genetic substrate for “inflamed” depression. In this context, the most replicated and relevant gene variants involved in both depression and immune activation (according to the review by Barnes et al<sup>45</sup>) include polymorphisms in the genes for interleukin IL-1 $\beta$ ,<sup>46-48</sup> IL-6,<sup>48-50</sup> IL-10,<sup>51,52</sup> TNF- $\alpha$ ,<sup>51,53</sup> CRP,<sup>54,55</sup> monocyte chemoattractant protein-1,<sup>56,57</sup> and phospholipase A2.<sup>58</sup> However, it is important to note that even for the most replicated findings there are discrepant results in the immune effects of the genetic variants and the resulting effects on depression. For example, a polymorphism in the IL-1 $\beta$  promoter at position 511 has been associated with higher depressive symptoms severity whether the polymorphism is associated with increased IL-1 $\beta$  production (allele 511T) or decreased IL-1 $\beta$  production (allele 511C).<sup>59-61</sup> Similar findings have been reported with polymorphisms in TNF- $\alpha$  and CRP promoters.<sup>62</sup> Further, the genome-wide association analyses identified the 44 risk gene variants in MDD with 4 of them (namely *LAC1*, *OLFM4*, *TIAF1*, and *NR4A2*) being associated with immune responses.<sup>63</sup> Besides, a recent weighted gene co-expression network analysis (WGCNA) identified three genes related to immune responses (*IL1RAP*, *UBE2W*, and *UBE2D1*) as hub genes of adolescent

MDD.<sup>64</sup> Further research involving genome-wide association studies, transcriptomics and WGCNA is needed to exactly shed light on the genetic contributions of inflammation-related depression.

## The Effect of Treatment in the Relation Between Depression and Inflammation – the Focus on Antidepressants and Anti-Cytokines

Antidepressant treatment seems to complexly interact with IRS and CIRS pathways; the recent largest meta-analysis has reported significantly decreased levels of IL-6, TNF- $\alpha$ , and IL-10 in adult MDD patients after conventional antidepressant treatment.<sup>25</sup> More specifically, treatment by selective serotonin reuptake inhibitors (SSRIs) led to the reduction in levels of pro-inflammatory cytokines TNF- $\alpha$ , IL-6, and IL-1 $\beta$ <sup>65,66</sup> but also to the reduction in levels of anti-inflammatory cytokines IL-4 and IL-10.<sup>67</sup> Contrary, treatment by serotonin and noradrenaline reuptake inhibitors (SNRIs such as venlafaxine and duloxetine) seems to be associated with an increase in TNF- $\alpha$ <sup>68</sup> and IL-6 levels.<sup>69</sup> This difference between SSRIs and SNRIs can be explained by the known pro-inflammatory effect of noradrenaline.<sup>70</sup> Further, ketamine (a non-competitive N-methyl-D-aspartate glutamate antagonist) infusions as a “novel” antidepressant<sup>71</sup> have reduced the levels of pro-inflammatory cytokines IL-6 and TNF- $\alpha$  in MDD.<sup>72</sup> However, a simultaneous decrease in the levels of pro-inflammatory (IL-1 $\beta$ , IL-6, TNF- $\alpha$ ) as well as anti-inflammatory cytokines (IL-4 and IL-10) in MDD after ketamine treatment has also been reported.<sup>73</sup> Regarding adolescent age, evaluation of the treatment effect on inflammation-related depression using antidepressants (SSRIs) has shown mixed results. Specifically, SSRIs have been repeatedly shown to reduce depressive symptoms and influence (increase or decrease) the levels of at least one cytokine.<sup>33,74,75</sup> However, the findings did not always align with the pro-inflammatory or anti-inflammatory nature of each cytokine. For example, while pro-inflammatory cytokine TNF- $\alpha$  has been reported to decrease post-treatment,<sup>33,75</sup> another pro-inflammatory cytokine IL-2 has been observed to increase.<sup>30</sup>

On the other hand, as currently available monoaminergic antidepressants are ineffective for over a third of MDD patients,<sup>76</sup> targeting inflammation may represent a promising way to develop novel antidepressants.<sup>77</sup> Monoclonal antibodies against pro-inflammatory cytokines as the most intriguing anti-inflammatory treatment for depression allow precise targeting of specific immune pathways and offer a way to personalized medicine. Regarding adult age, the treatment by cytokine inhibitors has been shown to improve depressive symptoms.<sup>7</sup> Specifically, TNF inhibitors such as etanercept or adalimumab and anti-IL-6 antibody sirukumab have been more efficacious compared to placebo in the reduction of depressive symptoms.<sup>78–80</sup> Further, the large meta-analysis by Kappelmann et al<sup>81</sup> reported that anti-cytokine treatments (adalimumab, etanercept and infliximab – TNF inhibitors, and tocilizumab – IL-6 inhibitor) significantly improved depressive symptom severity in patients with chronic inflammatory conditions resistant to antidepressant treatment. Regarding adolescent age, the effect of anti-cytokine agents in MDD patients has not been evaluated yet, however, the effect of the anti-inflammatory agent rosuvastatin influencing the cytokine levels has been estimated in adolescent MDD patients with no efficacy in lowering depressive symptoms or cytokine levels.<sup>82</sup> In summary, the effect of antidepressants on immune dysregulation as well as of anti-cytokines on depressive symptoms remains to be further explored, particularly at the adolescent age.

## Sex Differences in the Relationship Between Depression and Inflammation

Depression is twice as common in females compared to males.<sup>83</sup> This sex gap emerges around age 13 with increasing magnitude across adolescence and persisting through the rest of the lifespan.<sup>84</sup> Moreover, females are also more susceptible to autoimmune disorders.<sup>85</sup> Given mounting evidence that females are at a greater risk for depression and show increased inflammatory activity, it is critical to examine sex differences in the inflammation-depression relationship. However, the findings regarding the role of sex in the relationship between inflammation and depression in adults are mixed. For example, a stronger relationship between inflammatory markers (eg IL-6) and depressive symptoms was found in males compared to females.<sup>86</sup> Conversely, females receiving interferon-alpha showed greater increases in depressive mood compared to males.<sup>87</sup> Besides, no sex differences were found in the relationship between inflammation and depression<sup>88,89</sup> Notably, although already adolescent females are identified as particularly vulnerable to

inflammation-linked depression,<sup>29</sup> to the best of our knowledge, the findings reflecting the role of sex in the relationship between inflammation and first-episode depression prior to pharmacotherapy at the adolescent age are missing.

This study aimed to 1) evaluate IRS and CIRS pathways by assessment of various cytokines and soluble IL receptors in association with cytokine profiles during the acute phase of the first-episode drug-naïve MDD patients at adolescent age, 2) assess IRS/CIRS ratios as there is an almost complete lack of information on IRS versus CIRS activity in adolescent depression, 3) evaluate IRS and CIRS indices in adolescent females (MDD vs controls), in adolescent males (MDD vs controls), and in adolescent MDD females vs adolescent MDD males. To the best of our knowledge, this is the first study comprehensively assessing IRS versus CIRS balance in adolescent depression, with respect to sex.

## Materials and Methods

### Subjects

We have examined 100 MDD adolescent patients (average age: 15.4±1.2 years, BMI: 20.5±0.3 kg/m<sup>2</sup>, 40 males – average age: 15.6±0.2 years, BMI: 20.2±0.4 kg/m<sup>2</sup>; 60 females – average age: 15.4±0.2 years, BMI: 20.6±0.32 kg/m<sup>2</sup>) and 60 controls (average age: 15.3±1.5 years, BMI: 19.9±0.3 kg/m<sup>2</sup>, 28 males – average age: 15.1±0.3 years, BMI: 19.8±0.4 kg/m<sup>2</sup>; 32 females – average age: 15.4±0.3 years, BMI: 19.9±0.4 kg/m<sup>2</sup>). The patients suffering from first-episode MDD were recruited from the inpatients admitted to the Psychiatric Clinic of Jessenius Faculty of Medicine and University Hospital in Martin and were examined prior to the pharmacotherapy during the first days of hospitalization. The diagnosis of MDD without other comorbid psychiatric disorders (eg attention-deficit/hyperactivity disorder, anxiety disorders) was classified by a thorough clinical investigation based on an unstructured diagnostic interview by a staff child/adolescent psychiatrist according to the Diagnostic and Statistical Manual of Mental Disorders (DSM-5<sup>90</sup>). The control participants have never been treated for any mental disorder. Participants included in both groups were free of acute and chronic infections, allergies, endocrine- or immune-related diseases, cancer, and systemic diseases, as determined by self-report or doctor's report. Patients and controls with a history of antidepressant or antipsychotic treatment, immunomodulatory treatment, analgesic/anti-inflammatory use, antibiotic therapy, or substance abuse (alcohol, marijuana, heroin, cocaine, or methamphetamine) were also excluded. All participants filled the Children's Depression Inventory (CDI) to detect the presence and severity of depressive symptoms.<sup>91</sup>

### Blood Analysis

Fasting peripheral venous blood samples (5 mL) were collected into EDTA test tubes by venipuncture in the morning. Consequently, the blood samples were centrifuged for 15 minutes at 2500 rpm and 4°C (refrigerated centrifugation, Hettich Universal 320R, Tuttlingen, Germany), and the obtained plasma was kept frozen at -80°C until the analysis. Selected cytokines (interleukins - IL-1 $\alpha$ , IL-1 $\beta$ , IL-2, IL-4, IL-6, IL-8, IL-10, IFN- $\gamma$ , TNF- $\alpha$  – analytes involved in Cytokine array I), soluble cytokine receptors (sIL-6R, sTNF-R1, sTNF-R2) – analytes involved in Cytokine array IV were analyzed and quantified using a biochip array technology (Evidence Investigator, Randox, United Kingdom). Biochip array technology is a precision multiplex immunoassay-based testing platform allowing for the simultaneous quantitative detection of a wide range of analytes from a single sample. Plasma assays were provided in a separate kit from Randox Laboratories Ltd. (United Kingdom). Each kit (Cytokine array I – CTK I and Cytokine array IV – CTK IV) contains test-biochip cartridges, calibrators, assay diluent, conjugate, wash buffer, and signal reagent (luminol and peroxide, 1:1). Quality Controls (CTK I controls and CTK IV controls) were used for each run validation. There are 3 levels of quality control – low (level 1), medium (level 2), and high (level 3). A nine-point calibration was performed for each assay series as recommended. After the addition of 100  $\mu$ L of plasma samples to the biochip, analytes present in the sample bind to the specific biochip-bound ligands. The degree of binding is determined using a chemiluminescent light source and quantified using a Charge Coupled Device camera and imaging system.

Consequently, z-unit weighted composite scores were calculated from raw cytokines and soluble cytokine receptors data (in logarithmic transformation according to)<sup>92</sup> by using formula:  $Z = (x - M)/SD$ ,<sup>93</sup> where x=proband's raw cytokine or soluble cytokine receptor level, M=mean cytokine or soluble cytokine receptor level in combined group (ie depressive and control group together), and SD=standard deviation of cytokine or soluble cytokine receptor level in combined



group. These standardized data were used to assess indices reflecting individual IRS and/or CIRS activation profiles and ratios,<sup>16,94</sup> namely: IL-6 trans-signaling =  $zIL-6+zsIL-6R$ , M1 signaling =  $zIL-1\beta+zIL-6+zTNF-\alpha$ , Th1 profile =  $zIL-2+zIFN-\gamma$ , Treg+Th2 profile =  $zIL-10+zIL-4$ , allIRS profile (ie indicants of adolescent depression-linked immune-inflammatory activation<sup>41</sup>) =  $zIL-1\beta+zIL-2+zIL-6+zsIL-6R+zTNF-\alpha+zIFN-\gamma$ , allCIRS profile (ie all indicants of immune-regulatory activation) =  $zsTNF-R1+zsTNF-R2+zIL-4+zIL-10$ , TNF- $\alpha$ /sTNF-R1 ratio =  $zTNF-\alpha-zsTNF-R1$ , TNF- $\alpha$ /sTNF-R2 ratio =  $zTNF-\alpha-zsTNF-R2$ , TNF- $\alpha$ /sTNF-Rs ratio =  $zTNF-\alpha-z(zsTNF-R1+zsTNF-R2)$ , Th1/Th2 ratio =  $z(zIFN-\gamma+zIL-2)-zIL-4$ , Th1/Treg ratio =  $z(zIFN-\gamma+zIL-2)-zIL-10$ , Th1/Th2+Treg ratio =  $z(zIFN-\gamma+zIL-2)-z(zIL-4+zIL-10)$ , M1/Th2 ratio =  $z(zIL-1\beta+zIL-6+zTNF-\alpha)-zIL-4$ , M1/Treg ratio =  $z(zIL-1\beta+zIL-6+zTNF-\alpha)-zIL-10$ , M1/Treg+Th2 ratio =  $z(zIL-1\beta+zIL-6+zTNF-\alpha)-z(zIL-10+zIL-4)$ , allIRS/allCIRS ratio =  $z(zIL-1\beta+zIL-2+zIL-6+zsIL-6R+zTNF-\alpha+zIFN-\gamma)-z(zsTNF-R1+zsTNF-R2+zIL-4+zIL-10)$  - a subtraction was performed as these are log-transformed values.<sup>16,94</sup>

## Statistical Analysis

The data were explored and analyzed in Jamovi version 1.2.27 (Sydney, Australia). The Shapiro–Wilk normality test was used for the evaluation of data distributions (Gaussian/non-Gaussian). The analysis of variance (ANOVA) was used to test the effects of group and sex of Gaussian-distributed IRS and CIRS-related data (IL-6 trans-signaling, Th1 profile, allIRS, allCIRS, TNF- $\alpha$ /sTNF-R2, TNF- $\alpha$ /sTNF-Rs, Th1/Treg, M1/Th2, M1/Treg, M1/Treg+Th2, allIRS/allCIRS) and CDI. Next, the statistical test of the main effect was followed by post hoc pairwise comparisons between the groups and sexes, which was corrected by the Bonferroni method used to adjust probability (p) values when performing multiple statistical tests in any context. The non-parametric (Kruskal–Wallis) ANOVA was used for the between-groups and between-sexes comparisons for non-Gaussian distributed IRS and CIRS-related data (IL-1 $\alpha$ , IL-1 $\beta$ , IL-2, IL-4, IL-6, IL-8, IL-10, IFN- $\gamma$ , TNF- $\alpha$ , sIL-6R, sTNF-R1, sTNF-R2, TNF- $\alpha$ /sTNF-R1, Treg+Th2 profile, Th1/Th2, Th1/Treg+Th2, M1 signaling). Gaussian distributed data were expressed as mean  $\pm$  SEM and non-Gaussian distributed data as median (interquartile ranges). Besides, the associations between CDI score and IRS and CIRS indices were analyzed using Spearman's rank-order correlation test. A value of  $p \leq 0.05$  (two-tailed) was considered statistically significant.

## Results

### Evaluation of Selected Cytokines, Soluble Cytokine Receptors, Profiles, and Ratios

The main effect of group evaluated by ANOVA was significant for allIRS profile ( $F_{[4]} = 48.60$ ,  $p=0.045$ ), allCIRS profile ( $F_{[10]}=24.40$ ,  $p=0.002$ ), and TNF/TNFRs ratio ( $F_{[4]} = 9.68$ ,  $p=0.029$ ).

A post hoc pairwise comparison with the Bonferroni adjusted p revealed significantly higher allIRS and allCIRS profiles and TNF/TNFRs ratio in MDD adolescents compared to controls ( $p=0.045$ ,  $p=0.002$ ,  $p=0.029$ , respectively). The main effect of group was without significant changes for IL-6 trans-signaling, Th1 profile, TNF/TNF-R2, M1/Treg, M1/Th2, M1/Treg+Th2, Th1/Treg, and allIRS/allCIRS ratios ( $p=0.304$ ,  $p=0.800$ ,  $p=0.523$ ,  $p=0.710$ ,  $p=0.268$ ,  $p=0.973$ ,  $p=0.051$ , and  $p=0.908$ , respectively).

The non-parametric ANOVA revealed significantly increased IL-4, IL-10, TNF- $\alpha$ , sIL-6R, and Treg+Th2 profile in adolescent MDD patients compared to controls ( $p=0.046$ ,  $p<0.001$ ,  $p=0.002$ ,  $p=0.004$ , and  $p<0.001$ ; respectively), and significantly decreased Th1/Treg+Th2 ratio in MDD adolescents compared to controls ( $p=0.011$ ). No significant between-groups changes were found for IL-1 $\alpha$ , IL-1 $\beta$ , IL-2, IL-6, IL-8, IFN- $\gamma$ , sTNF-R1, sTNF-R2, M1 signaling, TNF- $\alpha$ /sTNF-R1 ratio, and Th1/Th2 ratio ( $p=0.774$ ,  $p=0.237$ ,  $p=0.880$ ,  $p=0.319$ ,  $p=0.087$ ,  $p=0.161$ ,  $p=0.400$ ,  $p=0.065$ ,  $p=0.054$ ,  $p=0.159$ , and  $p=0.496$ , respectively). All results are summarized in Table 1.

### Evaluation of Sex Differences in IRS and CIRS Activity

The main effect of sex evaluated by ANOVA was significant for Th1/Treg ( $F_{[4]}=8.42$ ,  $p=0.035$ ) and allIRS/allCIRS ( $F_{[7]}=51.17$ ,  $p=0.009$ ) ratios. The main effect of sex was without significant changes for IL-6 trans-signaling, Th1 profile, allIRS profile, allCIRS profile, TNF/TNF-Rs ratio, TNF/TNF-R2 ratio, M1/Treg ratio, M1/Th2 ratio, and M1/Treg+Th2

**Table 1** Evaluated Cytokines, Soluble Cytokine Receptors, IRS and CIRS-Related Profiles, and IRS vs CIRS Activity Ratios

Evaluated Indices	IRS, CIRS, IRS vs CIRS Function	Controls			MDD			p-value			
		Total <sup>a</sup> (N=60)	Females <sup>b</sup> (N=32)	Males <sup>c</sup> (N=28)	Total <sup>d</sup> (N=100)	Females <sup>e</sup> (N=60)	Males <sup>f</sup> (N=40)	a vs d	b vs e	c vs f	e vs f
<b>Cytokines</b>											
IL-1 $\alpha$ (pg/mL)	IRS	0.26 (0.19–0.35)	0.27 (0.20–0.35)	0.25 (0.18–0.33)	0.26 (0.19–0.40)	0.25 (0.19–0.33)	0.26 (0.21–0.51)	0.774	0.531	0.278	0.229
IL-1 $\beta$ (pg/mL)	IRS	1.17 (0.87–1.93)	1.39 (0.96–2.12)	1.09 (0.80–1.81)	1.37 (1.00–2.04)	1.41 (1.02–2.17)	1.21 (0.98–1.69)	0.237	0.517	0.428	0.216
IL-2 (pg/mL)	IRS	2.50 (1.50–5.22)	2.88 (1.73–5.26)	2.22 (1.41–3.31)	2.34 (1.60–4.06)	2.60 (1.82–4.06)	1.98 (1.49–3.93)	0.880	0.612	0.945	0.195
IL-4 (pg/mL)	CIRS	2.08 (1.84–2.73)	2.29 (1.87–2.79)	1.98 (1.80–2.41)	2.34 (2.02–2.67)	2.33 (2.01–2.82)	2.36 (2.07–2.63)	<b>0.046</b>	0.354	<b>0.044</b>	0.827
IL-6 (pg/mL)	IRS	0.79 (0.63–1.24)	0.84 (0.69–1.29)	0.69 (0.57–1.14)	0.92 (0.69–1.19)	0.98 (0.76–1.31)	0.88 (0.63–1.07)	0.319	0.594	0.345	0.132
IL-8 (pg/mL)	IRS	3.99 (3.04–5.91)	3.89 (2.83–5.91)	4.05 (3.59–5.58)	3.31 (2.58–5.21)	3.47 (2.51–5.10)	3.22 (2.63–5.38)	0.087	0.319	0.126	0.872
IL-10 (pg/mL)	CIRS	0.77 (0.58–0.99)	0.79 (0.64–0.94)	0.70 (0.54–1.02)	0.95 (0.70–1.44)	0.99 (0.68–1.49)	0.94 (0.73–1.31)	<b>&lt;0.001</b>	<b>0.017</b>	<b>0.007</b>	0.952
IFN- $\gamma$ (pg/mL)	IRS	0.42 (0.28–0.56)	0.46 (0.31–0.66)	0.34 (0.21–0.53)	0.45 (0.30–0.79)	0.44 (0.28–0.82)	0.45 (0.31–0.76)	0.161	0.993	0.052	0.759
TNF- $\alpha$ (pg/mL)	IRS	2.46 (2.00–2.96)	2.29 (1.78–2.83)	2.74 (2.21–3.14)	2.89 (2.29–3.42)	2.84 (2.16–3.19)	3.12 (2.63–3.47)	<b>0.002</b>	<b>0.006</b>	0.074	0.064
<b>Soluble cytokine receptors</b>											
sIL-6R (ng/mL)	IRS	0.91 (0.52–1.42)	0.98 (0.57–1.58)	0.90 (0.46–1.36)	1.33 (0.87–1.80)	1.34 (0.91–1.81)	1.32 (0.65–1.79)	<b>0.004</b>	0.062	<b>0.035</b>	0.616
sTNF-R1 (ng/mL)	CIRS	0.48 (0.39–0.59)	0.42 (0.37–0.55)	0.54 (0.44–0.69)	0.49 (0.44–0.57)	0.48 (0.42–0.55)	0.51 (0.47–0.63)	0.400	0.198	0.956	<b>0.015</b>
sTNF-R2 (ng/mL)	CIRS	0.44 (0.23–0.69)	0.38 (0.22–0.69)	0.48 (0.27–0.69)	0.54 (0.38–0.73)	0.52 (0.37–0.70)	0.59 (0.47–0.77)	0.065	0.178	0.177	0.156
<b>Profiles</b>											
IL-6 trans signaling	IRS	-0.20 $\pm$ 0.23	-0.27 $\pm$ 0.21	-0.10 $\pm$ 0.49	0.23 $\pm$ 0.20	0.43 $\pm$ 0.27	-0.08 $\pm$ 0.28	0.304	0.656	0.998	0.997
MI signaling	IRS	-0.52 (-2.17–1.36)	-0.42 (-1.76–1.59)	-1.80 (-2.32–0.22)	0.06 (-1.52–1.43)	0.24 (-1.35–1.85)	-0.11 (-1.51–1.09)	0.054	0.309	0.086	0.421
Th1 profile	IRS	-0.03 $\pm$ 0.23	0.39 $\pm$ 0.30	-0.47 $\pm$ 0.34	0.04 $\pm$ 0.16	0.07 $\pm$ 0.21	-0.01 $\pm$ 0.24	0.800	0.998	0.997	0.996

(Continued)

Table 1 (Continued).

Evaluated Indices	IRS, CIRS, IRS vs CIRS Function	Controls			MDD			p-value			
		Total <sup>a</sup> (N=60)	Females <sup>b</sup> (N=32)	Males <sup>c</sup> (N=28)	Total <sup>d</sup> (N=100)	Females <sup>e</sup> (N=60)	Males <sup>f</sup> (N=40)	a vs d	b vs e	c vs f	e vs f
Treg+Th2	CIRS	-0.36 (-1.34-0.24)	-0.17 (-1.09-0.53)	-0.74 (-1.74- -0.23)	0.28 (-0.80-1.21)	0.22 (-0.97-1.33)	0.28 (-0.54-0.80)	<b>&lt;0.001</b>	0.074	<b>&lt;0.001</b>	0.549
allIRS	IRS	-1.04±0.46	-0.38±0.56	-1.80±0.74	0.17±0.36	0.43±0.48	-0.21±0.55	<b>0.045</b>	0.996	0.449	0.998
allCIRS	CIRS	-0.57±0.25	-0.56±0.35	-0.58±0.37	0.30±0.15	0.22±0.20	0.41±0.23	<b>0.002</b>	0.230	0.111	0.997
<b>Ratios</b>											
TNF- $\alpha$ /sTNF-Rs	IRS vs CIRS	-0.53±0.24	-0.46±0.33	-0.62±0.34	0.05±0.14	0.12±0.17	-0.05±0.23	<b>0.029</b>	0.598	0.852	0.998
TNF- $\alpha$ /sTNF-R1	IRS vs CIRS	-0.19 (-0.58-0.38)	0.03 (-0.45-0.72)	-0.47 (-0.75- -0.09)	0.04 (-0.48-0.48)	-0.01 (-0.36-0.49)	0.07 (-0.58-0.48)	0.159	0.970	<b>0.047</b>	0.823
TNF- $\alpha$ /sTNF-R2	IRS vs CIRS	0.06±0.17	0.03±0.25	0.09±0.23	0.16±0.09	0.12±0.12	0.23±0.14	0.523	0.979	0.958	0.950
Th1/Th2	IRS vs CIRS	-0.15 (-1.05-0.99)	0.47 (-1.03-1.23)	-0.23 (-1.18-0.68)	-0.36 (-1.23-0.69)	-0.27 (-0.94-0.70)	-0.44 (-1.23-0.67)	0.496	0.369	0.958	0.998
Th1/Treg	IRS vs CIRS	0.28±0.21	0.67±0.32	-0.15±0.25	-0.18±0.13	-0.11±0.17	-0.28±0.22	0.051	0.085	0.996	0.998
Th1/Treg+Th2	IRS vs CIRS	0.10 (-0.73-1.40)	0.57 (-0.77-1.51)	-0.03 (-0.69-0.74)	-0.49 (-0.73-1.40)	-0.48 (-1.25-0.60)	-0.61 (-1.24-0.78)	<b>0.011</b>	0.053	0.076	0.784
M1/Th2	IRS vs CIRS	-0.52±0.31	-0.42±0.37	-0.63±0.52	-0.04±0.21	0.24±0.27	-0.47±0.34	0.268	0.997	0.998	0.729
M1/Treg	IRS vs CIRS	0.03±0.30	0.18±0.35	-0.17±0.53	0.18±0.22	0.30±0.27	-0.01±0.37	0.710	0.995	0.992	0.998
M1/Treg+Th2	IRS vs CIRS	0.11±0.30	0.00±0.32	-0.22±0.48	0.01±0.20	0.09±0.26	-0.28±0.28	0.973	0.998	0.997	0.998
allIRS/allCIRS	IRS vs CIRS	-0.48±0.38	0.04±0.50	-1.13±0.54	-0.35±0.33	0.30±0.43	-1.27±0.47	0.908	0.998	0.998	0.072

**Notes:** Values are expressed as mean±SEM for Gaussian-distributed data and as median (interquartile ranges) for non-Gaussian distributed data. p-value ≤ 0.05 (in bold) is considered statistically significant. <sup>a</sup>Control group; <sup>b</sup>Control adolescent females; <sup>c</sup>Control adolescent males; <sup>d</sup>Depressive group; <sup>e</sup>Depressive adolescent females; <sup>f</sup>Depressive adolescent males.

**Abbreviations:** IRS, inflammatory response system; CIRS, compensatory immune response system; MDD, major depressive disorder; IL, interleukin; IFN- $\gamma$ , interferon-gamma; TNF- $\alpha$ , tumor necrosis factor-alpha; sIL-6R, soluble interleukin-6 receptor; sTNF-R1, soluble tumor necrosis factor receptor 1; sTNF-R2, soluble tumor necrosis factor receptor 2; M1, macrophages M1; Th1, T helper cells 1; Treg, T regulatory cells; Th2, T helper cells 2; allIRS, all indices involved in the adolescent depression-linked inflammatory response system; allCIRS, all indices involved in the compensatory immune response system; sTNF-Rs, soluble tumor necrosis factor receptors (ie sTNF-R1+sTNF-R2).



ratio ( $p=0.617$ ,  $p=0.082$ ,  $p=0.084$ ,  $p=0.769$ ,  $p=0.522$ ,  $p=0.613$ ,  $p=0.381$ ,  $p=0.214$ , and  $p=0.377$ , respectively). The post hoc pairwise comparison revealed no significant changes in any of the above-mentioned indices between adolescent females and males.

The non-parametric ANOVA revealed significantly increased IL-10 and TNF- $\alpha$  in adolescent MDD females compared to control females ( $p=0.017$  and  $p=0.006$ , respectively). On the other hand, adolescent MDD males had significantly increased IL-4, IL-10, sIL-6R, Treg+Th2 profile, and TNF- $\alpha$ /TNF-R1 ratio compared to control males ( $p=0.044$ ,  $p=0.007$ ,  $p=0.035$ ,  $p<0.001$ , and  $p=0.047$ ; respectively). Lastly, significantly increased sTNF-R1 was found in adolescent MDD males compared to adolescent MDD females ( $p=0.015$ ). All results are summarized in [Table 1](#).


## CDI Analysis and Correlation Between Depressive Symptoms and IRS and CIRS Indices

The main effect of the group evaluated by ANOVA was significant for the CDI score ( $F_{[364]} = 16,093.70$ ,  $p<0.001$ ). The main effect of sex evaluated by ANOVA was not significant for CDI score (females vs males =  $16.50\pm 1.34$  vs  $16.10\pm 1.64$ ,  $p=0.300$ ). The post hoc pairwise comparison revealed significantly increased CDI score in adolescent MDD patients compared to controls ( $25.10\pm 0.89$  vs  $3.75\pm 0.45$ ,  $p<0.001$ ), significantly increased CDI score in adolescent MDD females compared to adolescent control females ( $24.30\pm 1.21$  vs  $3.78\pm 0.62$ ,  $p<0.001$ ), and significantly increased CDI score in adolescent MDD males compared to adolescent control males ( $26.50\pm 1.24$  vs  $3.71\pm 0.66$ ,  $p<0.001$ ).

Correlation analysis in the total group (depressive group and control group) revealed significant positive correlations between CDI score and IL-10 and sIL-6R ( $r=0.167$ ,  $p=0.041$ ;  $r=0.163$ ,  $p=0.050$ ; respectively, [Figure 2A](#)). Further, significant positive correlations were found between CDI score and IL-10 ( $r=0.306$ ,  $p=0.017$ ) in adolescent males ([Figure 2B](#)). No significant correlations were found between the CDI score and adolescent females.

## Discussion

This study for the first time such extensively investigated IRS and CIRS indices, profiles, and ratios in depressive adolescents. The main significant results can be summarized as follows: 1) increased levels of TNF- $\alpha$  and sIL-6R associated with increased allIRS profile indicating IRS activation, 2) increased TNF- $\alpha$ /sTNF-Rs ratio reflecting enhanced IRS-linked TNF- $\alpha$  signaling, 3) increased levels of IL-10 and IL-4 associated with increased Treg+Th2 and allCIRS profiles indicating CIRS activation, 4) decreased Th1/Treg+Th2 ratio reflecting enhanced CIRS activity, 5) increased IL-10 and TNF- $\alpha$  in adolescent MDD females compared to adolescent control females, 6) increased IL-4, IL-10, sIL-6R, Treg+Th2 profile, and TNF- $\alpha$ /TNF-R1 ratio in adolescent MDD males compared to adolescent control males, 7) increased sTNF-R1 in adolescent MDD males compared to adolescent MDD females, and 8) positive correlations between CDI score and sIL-6R and IL-10 indicating that more severe depressive symptoms are associated with enhanced



**Figure 2** Correlation analysis. (A) Positive correlations between depressive symptoms and sIL-6R and IL-10 in the total group; (B) Positive correlations between depressive symptoms and IL-10 in adolescent males.

pro-inflammatory IL-6 trans-signaling and anti-inflammatory IL-10 signaling in adolescent depression. Importantly, adolescent males showed positive correlations between CDI score and IL-10, while no correlations between CDI score and evaluated indices were found in adolescent females. Our results indicate discrete abnormalities in both, IRS and CIRS, pathways associated with sex-specific differences in drug-naïve depressive patients already at adolescent age. Several mechanisms are supposed.

## Adolescent Depression-Linked IRS and CIRS (Dys)regulation

Multiple immune cells and immune effector molecules ensure the maintenance of IRS/CIRS functional homeostasis (ie IRS and CIRS represent components of the integrated immune-signaling system with CIRS counter-regulating the IRS effects). While IRS exerting pro-inflammatory effects consists of activated M1 cells (IL-1 + IL-6 + TNF- $\alpha$ ), Th1 cells (IL-2 + IL-12 + IFN- $\gamma$ ), and Th17 cells (IL-17 + IL-6), CIRS exerting anti-inflammatory effects constitutes from activated Th2 cells (IL-4), Treg cells (IL-10 + transforming growth factor beta), and cytokine receptors (eg sIL-1RA, sIL-2R, sTNF-Rs).<sup>17</sup> Disruption of the IRS/CIRS balance (eg M1/M2, Th1/Th2 balance) caused by exogenous adverse conditions such as psychosocial stress can lead to the chronic inflammatory response.<sup>17,20</sup> Dysregulated IRS/CIRS axes seem to play a crucial role in the complex adult MDD immunopathology influencing the onset of depression, number of depressive episodes as well as recovery.<sup>17</sup> Importantly, the depression rate dramatically increases from childhood into young adulthood,<sup>3</sup> thus, the knowledge about the IRS and CIRS interactions and depression at the adolescent age can be crucial.<sup>95</sup> However, the number of IRS and CIRS studies at this age period is limited.

In line with a recent systematic review and meta-analysis of cytokines in children and adolescent depression,<sup>40</sup> our results confirmed increased peripheral levels of TNF- $\alpha$  in depressive adolescents. The cytokine TNF- $\alpha$  plays a key role in the inflammatory response. Increased TNF- $\alpha$  levels may drive differentiation of monocytes, T-cells proliferation, activation of IL-1 and IL-6 production as well as acute phase response.<sup>96</sup> Moreover, besides its pro-inflammatory effects, TNF- $\alpha$  can also promote modifications of neuronal synaptic transmission.<sup>97</sup> These properties are mediated through two specific TNF receptors (TNF-R1 and TNF-R2) expressed on the cell surfaces.<sup>98</sup> The TNF receptors can also be found in the circulation as the soluble TNF receptors (sTNF-R1 and sTNF-R2) predominantly generated by proteolytic cleavage of TNF receptors by the TNF- $\alpha$ -converting enzyme (TACE, also called a disintegrin and metalloprotease 17 (ADAM 17)).<sup>99</sup> Multiple inflammatory signals (eg IL-2, IL-6, IL-10, TNF- $\alpha$  itself) may cause the shedding of TNF receptors. These soluble receptors acting as decoy receptors bind TNF- $\alpha$  and thus attenuate TNF- $\alpha$  signaling resulting in inhibition of TNF- $\alpha$  effects.<sup>100</sup> From this perspective, the meta-analysis by Köhler et al<sup>25</sup> reported elevated sTNF-R2 plasma levels in adult depression. In our study, there was a tendency to an increase in sTNF-R2 plasma levels in adolescent MDD compared to controls. These findings may contribute to the assumption that increased sTNF-R2 levels may counter-regulate enhanced TNF- $\alpha$ -mediated pro-inflammatory effects through the negative regulatory feedback.<sup>16</sup> Importantly, our study for the first time evaluated TNF- $\alpha$ /TNF- $\alpha$  receptors ratios (ie TNF- $\alpha$ /sTNF-R1, TNF- $\alpha$ /sTNF-R2, and TNF- $\alpha$ /sTNF-Rs) in depressive adolescents compared to controls with significantly increased TNF- $\alpha$ /sTNF-Rs indicating dominant pro-inflammatory TNF- $\alpha$  signaling during the acute phase of depression in adolescent drug-naïve patients.

Further, increased TNF- $\alpha$  may enhance the production of another important cytokine, IL-6, characterized by bimodal immune activity mediated through two separate pathways (classical signaling and trans-signaling). In the classical signaling pathway, IL-6 binds to membrane-bound receptor (IL-6R), the complex of IL-6 and IL-6R associates with a second protein (glycoprotein – gp130) inducing dimerization and initiation of intracellular signaling. Similar to TNF receptors, IL-6R can be proteolytically cleaved from the cell membrane by the metalloproteinase ADAM17 to generate a soluble form (sIL-6R).<sup>101</sup> IL-6 bound to sIL-6R forms a complex that can associate with gp130 to initiate signal transduction, this process is known as trans-signaling.<sup>102</sup> While IL-6 classical signaling contributes to anti-inflammatory properties of this cytokine, IL-6 trans-signaling is reported to be pro-inflammatory.<sup>103</sup> It is important to note that IL-6 signaling can be upregulated by increased sIL-6R levels alone, without a change in IL-6 levels.<sup>104</sup> In this way, our results of higher levels of sIL-6R are in accordance with the studies indicating enhanced IL-6 trans-signaling and thus pro-inflammatory activity in adult depression,<sup>105–107</sup> and extend them into adolescent age.

Contrary to pro-inflammatory responses, a predominant CIRS-linked anti-inflammatory effect in depression may occur when naïve T helper (Th0) cells differentiate into Treg and/or Th2 cells producing IL-10 and IL-4. The cytokine

IL-10 signals through a receptor complex (IL-10R alpha and two IL-10R beta subunits) and regulates the activity of monocytes, macrophages, and Th1 cells resulting in the release suppression of pro-inflammatory cytokines such as TNF- $\alpha$ , IFN- $\gamma$ , IL-1 $\beta$ , IL-6.<sup>108,109</sup> Similarly, another cytokine with anti-inflammatory properties, IL-4, signals through two different receptor complexes (a type I receptor expressed on hematopoietic cells or a type II receptor expressed on non-hematopoietic cells). IL-4 mediates the alternative activation of M2 macrophages attenuating the pro-inflammatory activity by the release of transforming growth factor-beta, sIL-1RA, and IL-10.<sup>16,110</sup> Therefore, IL-10 and IL-4 seem to play a crucial role in the control of the inflammatory responses. Our results of increased IL-10 and IL-4 plasma levels are in accordance with the adult MDD studies,<sup>25,111</sup> and extend this finding into the adolescent age. Moreover, our study revealed increased Treg+Th2 and allCIRS profiles in association with decreased Th1/Treg+Th2 ratio confirming enhanced CIRS activity during the acute phase of depression in adolescent patients.

Taken together, abnormalities in the plasma levels of selected cytokines, soluble cytokine receptors in association with altered selected cytokine profiles and ratios provide the evidence for simultaneous signs of IRS and CIRS pathways activation in depression at adolescent age. In addition, contrary to an assumption that during the acute phase of mood disorder in adult patients the IRS is more active compared to CIRS,<sup>112</sup> our study revealed concurrent activation of both, IRS and CIRS confirmed by the unchanged IRS/CIRS ratio during the acute phase of depression in adolescent patients. Further, our study revealed the association between increased severity of depressive symptoms (based on the total CDI score) and IRS as well as CIRS biomarkers (ie increased sIL-6R and IL-10) in depressive adolescents. This finding is in accordance with adult MDD studies (eg<sup>106,113</sup>) and extends this knowledge into the adolescent age.

With regard to sex, this study identified different responses of IRS- and CIRS-related indices between females and males. Specifically, adolescent MDD females had increased levels of TNF- $\alpha$  and IL-10 compared to control females, reflecting enhanced TNF- $\alpha$  signaling counter-balanced by heightened anti-inflammatory IL-10 signaling. This result is partially in accordance with the finding of the positive association between TNF- $\alpha$  and depression in adult MDD females.<sup>114</sup> Similarly, increased TNF- $\alpha$  responses in females induced by a low-dose endotoxin administration were associated with mood disturbances.<sup>115</sup> In the same way, other pro-inflammatory markers such as CRP and IL-6 were associated with an increased risk of depression among females.<sup>9,116–118</sup> On the other hand, adolescent MDD males showed increased IL-4, IL-10, sIL-6R, Treg+Th2 profile, and TNF- $\alpha$ /sTNF-R1 ratio compared to control males, indicating increased pro-inflammatory, but particularly more enhanced anti-inflammatory activity. These findings are partially in accordance with increased pro-inflammatory activity in adult depressive males<sup>119,120</sup> and extend them into adolescent age. In contrast, other studies reported no association between inflammation and depression in adult males.<sup>115,118</sup> Moreover, sTNF-R1 was significantly increased in adolescent MDD males compared to adolescent MDD females and the correlation of IL-10 with the severity of depressive symptoms was found only in adolescent males. Thus, sex-specific differences found in this study could be related to immune variability across the lifespan from childhood through adolescence into adulthood.<sup>121</sup> Specifically for adolescence, this crucial developmental period is associated with a decline in lymphatic tissue accompanied by the reduction of circulating lymphocytes, partially initiated by the puberty-onset release of sex hormones. In this context, estrogen and androgen receptors are expressed on all major components of the innate and adaptive immune systems.<sup>122,123</sup> With respect to sex hormones-related immunomodulation, testosterone appears to attenuate inflammation as it decreases pro-inflammatory cytokines,<sup>124</sup> while estrogen may exert biphasic (ie pro-inflammatory and anti-inflammatory) effects.<sup>125</sup> Altogether, our findings are the first complexly exploring the sex-specific IRS and CIRS association in adolescent depression and can serve as preliminary evidence of the sex-dependent relationship among adolescent depression and cytokines, soluble cytokine receptors, cytokine profiles and IRS/CIRS activity ratios. However, the convergence impact of biological sex, sex hormones, and peripheral immunity on the mood in adolescents remains to be elucidated.

## Communication Between Peripheral Cytokine Activity and the Brain – The Role of Neuroimmune Interactions in Depression

Some peripheral cytokines can cross the blood–brain barrier (BBB) to act on astrocytes, microglia, and neurons throughout the brain contributing to the normal neurophysiological processes (neuronal differentiation, astrocyte

proliferation, etc.).<sup>126</sup> Peripheral cytokines can reach the brain through three major pathways: 1) humoral – cytokines access the brain through leaky regions in BBB such as the circumventricular organs,<sup>127</sup> 2) neural – activation of cytokine receptors on afferent vagal nerve fibers transducing cytokine signals to the brain,<sup>128</sup> and 3) cellular – chemokines released by activated microglia and adhesion molecules expressed in the central nervous system (CNS) can attract activated peripheral cells (monocytes and T cells) to the meninges and brain parenchyma.<sup>129</sup>

Notably, persistent peripheral inflammation can disrupt BBB leading to enhanced BBB permeability.<sup>130</sup> In turn, peripheral pro-inflammatory cytokines can more easily cross BBB to reach the brain leading to the activation of microglia. Microglial activation can then lead to the production of glutamate, reactive oxygen species, nitric oxide, proteases, as well as pro-inflammatory cytokines such as IL-1 $\beta$ , IL-6, and TNF- $\alpha$ . Excess and/or prolonged inflammatory cytokine activity within the brain may result in disturbed neurotransmitter metabolism, neuroendocrine function, neurogenesis/apoptosis, mitochondrial biogenesis, and neural plasticity affecting cognitive, emotional, and behavioral attributes associated with depression.<sup>131–136</sup>

Further, peripheral pro-inflammatory cytokines activate the enzyme indoleamine-2,3-dioxygenase (IDO), an important immunoregulator, breaking down the tryptophan into kynurenine (KYN).<sup>137,138</sup> KYN produced at the periphery can be transported into the brain where it is through the immune-inducible enzyme, kynurenine monooxygenase, presented in microglia, metabolized into neurotoxic KYN metabolites including quinolinic acid. Quinolinic acid, together with glutamate released by activated microglia, activates N-methyl-D-aspartate (NMDA) receptors throughout the brain contributing to depressive symptoms.<sup>139</sup> This pathway can explain why ketamine-linked blockade of NMDA receptors can abrogate depression-like behavior induced by lipopolysaccharide in experimental studies<sup>140</sup> indicating the promising potential of ketamine as a future treatment for depression.<sup>141</sup> Importantly, this pathway does not discard the role of serotonin in inflammation-linked depression. From this view, the inflammation modulates serotonin signaling by increasing the expression of serotonin transporters re-uptaking the synaptic serotonin resulting in the decreased amount of serotonin available at the synapses.<sup>142,143</sup> This inflammation-linked serotonin depletion process and consequently diminished brain serotonin activity has been reported to be associated with MDD.<sup>144,145</sup>

In summary, disturbed interactions between the peripheral immune system and CNS may result in central immune dysregulation and subsequently neurobehavioral MDD-linked abnormalities.<sup>139,145</sup> However, elucidating whether neuroimmune dysregulation is an etiological element leading to depression or a depression's consequence remains a chicken-or-egg dilemma.

Several limitations should be considered in our study. Firstly, other important IRS (eg IL-17, acute phase proteins) and CIRS (eg sIL-2R, sIL-1RA, IL-6 classical signaling, haptoglobin) indices were not evaluated in this study. Secondly, the smoking status as a factor influencing the relationship between inflammation and depression was not exactly monitored. Further, all participants came from the same place and were of the same ethnicity, thus our findings might be limited in the general applicability. Lastly, we have compared MDD adolescents and controls to identify potential biomarkers helpful for MDD diagnosing; however, future studies should explore whether our potential biomarkers could be also used to distinguish MDD adolescents from adolescent patients suffering from other mental disorders.

## Conclusion

Adolescent depression is characterized by enhanced pro-inflammatory TNF- $\alpha$  signaling and IL-6 trans-signaling associated with increased IL-10- and IL-4-mediated anti-inflammatory activity. Importantly, the association between IRS and CIRS pathways and adolescent depression seems to be sex-specific. Our findings can contribute to a clearer understanding of IRS and CIRS interactions in the complex immunopathogenesis of depression already at the adolescent age. Since IRS and CIRS can influence the treatment outcomes, further research is needed to identify precise IRS and CIRS biomarkers that could serve as a new therapeutic target for MDD adolescents, also considering the sex.

## Abbreviations

BBB, blood–brain barrier; CDI, Children Depression Inventory; CIRS, compensatory immune response system; CNS, central nervous system; CRP, C-reactive protein; DSM-5, Diagnostic and Statistical Manual of Mental Disorders, fifth edition; gp130, glycoprotein 130; IDO, indoleamine-2,3-dioxygenase; IFN- $\gamma$ , interferon-gamma; IL, interleukin; IRS,

inflammatory response system; KYN, kynurenine; M1, macrophages M1; M2, macrophages M2; MDD, major depressive disorder; NMDA, N-methyl-D-aspartate; NLR, neutrophil-to-lymphocyte ratio; PLR, platelet-to-lymphocyte ratio; RBC, red blood cells; RDW, red blood cell distribution width; sIL-1RA, soluble IL-1 receptor antagonist; sIL-2R, soluble IL-2 receptor; sIL-6R, soluble IL-6 receptor; SNRIs, serotonin and noradrenaline reuptake inhibitors; SSRIs, selective serotonin reuptake inhibitors; sTNF-R1, soluble tumor necrosis factor receptor 1; sTNF-R2, soluble tumor necrosis factor receptor 2; sTNF-Rs, soluble tumor necrosis factor receptors (ie sTNF-R1 + sTNF-R2); TACE/ADAM17, metalloproteinase, TNF- $\alpha$  converting enzyme; Th, helper T lymphocytes; TNF, tumor necrosis factor; TNF- $\alpha$ , tumor necrosis factor-alpha; TNF-R1, tumor necrosis factor receptor 1; TNF-R2, tumor necrosis factor receptor 2; Treg, regulatory T lymphocytes; WGCNA, weighted gene co-expression network analysis WGCNA; WBC, white blood cells.

## Ethics Approval and Consent to Participate

The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Ethics Committee of Jessenius Faculty of Medicine in Martin, Comenius University in Bratislava (protocol code EK23/2021 of 29 June 2021). Informed consent was obtained from all subjects and their legal guardians involved in the study.

## Acknowledgments

The authors thank Mgr. Jana Kohutova for technical support during RANDOX analysis. The authors also thank all participants involved in the study.

## Author Contributions

All authors made a significant contribution to this work whether in the conception, study design, execution, or data acquisition, analysis and interpretation. All authors critically reviewed the article, gave final approval of the version to be published, agreed on the journal to which the article has been submitted; and agreed to be accountable for the contents of the article.

## Funding

This research was funded by the Scientific Grant Agency of the Ministry of Education, Science, Research and Sport of the Slovak Republic under grant VEGA 1/0190/20.

## Disclosure

The authors declare no conflicts of interest in this work.

## References

1. Institute of Health Metrics and Evaluation. Global Health Data Exchange (GHDx). Available from: <http://ghdx.healthdata.org/gbd-results-tool?params=gbd-api-2019-permalink/d780dffbe8a381b25e1416884959e88b>. Accessed August 11, 2022.
2. Weinberger AH, Gbedemah M, Martinez AM, Nash D, Galea S, Goodwin RD. Trends in depression prevalence in the USA from 2005 to 2015: widening disparities in vulnerable groups. *Psychol Med*. 2018;48(8):1308–1315. doi:10.1017/S0033291717002781
3. Mojtabai R, Olfson M, Han B. National trends in the prevalence and treatment of depression in adolescents and young adults. *Pediatrics*. 2016;138(6):e20161878. doi:10.1542/peds.2016-1878
4. World Health Organization. Adolescent mental health. Available from: <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health>. Accessed September 28, 2022.
5. Zhou X, Teng T, Zhang Y, et al. Comparative efficacy and acceptability of antidepressants, psychotherapies, and their combination for acute treatment of children and adolescents with depressive disorder: a systematic review and network meta-analysis. *The Lancet Psychiatry*. 2020;7(7):581–601. doi:10.1016/S2215-0366(20)30137-1
6. Miller AH, Raison CL. Immune system contributions to the pathophysiology of depression. *Focus*. 2008;6(1):36–45. doi:10.1176/foc.6.1.foc36
7. Beurel E, Toups M, Nemeroff CB. The bidirectional relationship of depression and inflammation: double trouble. *Neuron*. 2020;107(2):234–256. doi:10.1016/j.neuron.2020.06.002
8. Maes M, Van Der Planken M, Stevens WJ, et al. Leukocytosis, monocytosis and neutrophilia: hallmarks of severe depression. *J Psychiatr Res*. 1992;26(2):125–134. doi:10.1016/0022-3956(92)90004-8
9. Köhler-Forsberg O, He W, Chang Y, Atlas SJ, Meigs JB, Nierenberg AA. White blood cell count at first depression diagnosis as predictor for risk of subsequent hospitalization with depression. *Neurol Psychiatry Brain Res*. 2017;26:1–6. doi:10.1016/j.npbr.2017.10.002



10. Puangsri P, Ninla-Aesong P. Potential usefulness of complete blood count parameters and inflammatory ratios as simple biomarkers of depression and suicide risk in drug-naive, adolescents with major depressive disorder: simple peripheral markers for inflammation in drug-naive, adolescents with major depressive disorder. *Psychiatry Res.* 2021;305:114216. doi:10.1016/j.psychres.2021.114216
11. Uçar HN, Eray Ş, Murat D. Simple peripheral markers for inflammation in adolescents with major depressive disorder. *Psych Clin Psychopharmacol.* 2018;28:254–260. doi:10.1080/24750573.2018.1423769
12. Mazza MG, Lucchi S, Tringali AGM, Rossetti A, Botti ER, Clerici M. Neutrophil/lymphocyte ratio and platelet/lymphocyte ratio in mood disorders: a meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry.* 2018;84:229–236. doi:10.1016/j.pnpbp.2018.03.012
13. Demircan F, Gözel N, Kılınç F, Ulu R, Atmaca M. The impact of red blood cell distribution width and neutrophil/lymphocyte ratio on the diagnosis of major depressive disorder. *Neurol Ther.* 2016;5(1):27–33. doi:10.1007/s40120-015-0039-8
14. Demir S, Atli A, Bulut M, et al. Neutrophil–lymphocyte ratio in patients with major depressive disorder undergoing no pharmacological therapy. *Neuropsychiatr Dis Treat.* 2015;11:2253–2258. doi:10.2147/NDT.S89470
15. Özyurt G, Binici NC. Increased neutrophil-lymphocyte ratios in depressive adolescents is correlated with the severity of depression. *Psychiatry Res.* 2018;268:426–431. doi:10.1016/j.psychres.2018.08.007
16. Maes M, Carvalho AF. The Compensatory Immune-Regulatory Reflex System (CIRS) in depression and bipolar disorder. *Mol Neurobiol.* 2018;55(12):8885–8903. doi:10.1007/s12035-018-1016-x
17. Debnath M, Berk M, Maes M. Translational evidence for the Inflammatory Response System (IRS)/Compensatory Immune Response System (CIRS) and neuroprogression theory of major depression. *Prog Neuropsychopharmacol Biol Psychiatry.* 2021;111:110343. doi:10.1016/j.pnpbp.2021.110343
18. Classen A, Lloberas J, Celada A. Macrophage activation: classical vs. alternative. *Methods Mol Biol.* 2009;531:29–43. doi:10.1007/978-1-59745-396-7\_3
19. Raphael I, Nalawade S, Eagar TN, Forsthuber TG. T cell subsets and their signature cytokines in autoimmune and inflammatory diseases. *Cytokine.* 2015;74(1):5–17. doi:10.1016/j.cyto.2014.09.011
20. Mills CD, Kincaid K, Alt JM, Heilman MJ, Hill AM. M-1/M-2 Macrophages and the Th1/Th2 Paradigm. *J Immunol.* 2000;164(12):6166–6173. doi:10.4049/jimmunol.164.12.6166
21. Tiemessen MM, Jagger AL, Evans HG, Van Herwijnen MJC, John S, Taams LS. CD4+CD25+Foxp3+ regulatory T cells induce alternative activation of human monocytes/macrophages. *Proc Natl Acad Sci U S A.* 2007;104(49):19446–19451. doi:10.1073/pnas.0706832104
22. Haapakoski R, Mathieu J, Ebmeier KP, Alenius H, Kivimäki M. Cumulative meta-analysis of interleukins 6 and 1 $\beta$ , tumour necrosis factor  $\alpha$  and C-reactive protein in patients with major depressive disorder. *Brain Behav Immun.* 2015;49:206–215. doi:10.1016/j.bbi.2015.06.001
23. Howren MB, Lamkin DM, Suls J. Associations of depression with c-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosom Med.* 2009;71(2):171–186. doi:10.1097/PSY.0b013e3181907c1b
24. Dowlati Y, Herrmann N, Swardfager W, et al. A meta-analysis of cytokines in major depression. *Biol Psychiatry.* 2010;67(5):446–457. doi:10.1016/j.biopsych.2009.09.033
25. Köhler CA, Freitas TH, Maes M, et al. Peripheral cytokine and chemokine alterations in depression: a meta-analysis of 82 studies. *Acta Psychiatr Scand.* 2017;135(5):373–387. doi:10.1111/acps.12698
26. Liu Y, Ho RCM, Mak A. Interleukin (IL)-6, tumour necrosis factor alpha (TNF- $\alpha$ ) and soluble interleukin-2 receptors (sIL-2R) are elevated in patients with major depressive disorder: a meta-analysis and meta-regression. *J Affect Disord.* 2012;139(3):230–239. doi:10.1016/j.jad.2011.08.003
27. Valkanova V, Ebmeier KP, Allan CL. CRP, IL-6 and depression: a systematic review and meta-analysis of longitudinal studies. *J Affect Disord.* 2013;150(3):736–744. doi:10.1016/j.jad.2013.06.004
28. Gabbay V, Klein RG, Alonso CM, et al. Immune system dysregulation in adolescent major depressive disorder. *J Affect Disord.* 2009;115(1–2):177–182. doi:10.1016/j.jad.2008.07.022
29. Henje Blom E, Lekander M, Ingvar M, Åsberg M, Mobarrez F, Serlachius E. Pro-inflammatory cytokines are elevated in adolescent females with emotional disorders not treated with SSRIs. *J Affect Disord.* 2012;136(3):716–723. doi:10.1016/j.jad.2011.10.002
30. Lee H, Song M, Lee J, Kim JB, Lee MS. Prospective study on cytokine levels in medication-naïve adolescents with first-episode major depressive disorder. *J Affect Disord.* 2020;266:57–62. doi:10.1016/j.jad.2020.01.125
31. Byrne ML, O'Brien-Simpson NM, Reynolds EC, et al. Acute phase protein and cytokine levels in serum and saliva: a comparison of detectable levels and correlations in a depressed and healthy adolescent sample. *Brain Behav Immun.* 2013;34:164–175. doi:10.1016/j.bbi.2013.08.010
32. Pallavi P, Sagar R, Mehta M, et al. Serum cytokines and anxiety in adolescent depression patients: gender effect. *Psychiatry Res.* 2015;229(1–2):374–380. doi:10.1016/j.psychres.2015.06.036
33. Pérez-Sánchez G, Becerril-Villanueva E, Arreola R, et al. Inflammatory profiles in depressed adolescents treated with fluoxetine: an 8-week follow-up open study. *Mediators Inflamm.* 2018;2018:4074051. doi:10.1155/2018/4074051
34. Ho PS, Yen CH, Chen CY, Huang SY, Liang CS. Changes in cytokine and chemokine expression distinguish dysthymic disorder from major depression and healthy controls. *Psychiatry Res.* 2017;248:20–27. doi:10.1016/j.psychres.2016.12.014
35. Hung YJ, Hsieh CH, Chen YJ, et al. Insulin sensitivity, proinflammatory markers and adiponectin in young males with different subtypes of depressive disorder. *Clin Endocrinol.* 2007;67(5):784–789. doi:10.1111/j.1365-2265.2007.02963.x
36. Miklowitz DJ, Portnoff LC, Armstrong CC, et al. Inflammatory cytokines and nuclear factor-kappa B activation in adolescents with bipolar and major depressive disorders. *Psychiatry Res.* 2016;241:315–322. doi:10.1016/j.psychres.2016.04.120
37. Peters AT, Ren X, Bessette KL, et al. Interplay between pro-inflammatory cytokines, childhood trauma, and executive function in depressed adolescents. *J Psychiatr Res.* 2019;114:1–10. doi:10.1016/j.jpsychires.2019.03.030
38. Su SC, Sun MT, Wen MJ, Lin CJ, Chen YC, Hung YJ. Brain-derived neurotrophic factor, adiponectin, and proinflammatory markers in various subtypes of depression in young men. *Int J Psychiatry Med.* 2011;42(3):211–226. doi:10.2190/PM.42.3.a
39. Belem da Silva CT, de Abreu Costa M, Kapczynski F, de Aguiar BW, Salum GA, Manfro GG. Inflammation and internalizing disorders in adolescents. *Prog Neuropsychopharmacol Biol Psychiatry.* 2017;77:133–137. doi:10.1016/j.pnpbp.2017.03.023
40. D'Acunto G, Nageye F, Zhang J, Masi G, Cortese S. Inflammatory cytokines in children and adolescents with depressive disorders: a systematic review and meta-Analysis. *J Child Adolesc Psychopharmacol.* 2019;29(5):362–369. doi:10.1089/cap.2019.0015



41. Toenders YJ, Laskaris L, Davey CG, et al. Inflammation and depression in young people: a systematic review and proposed inflammatory pathways. *Mol Psychiatry*. 2022;27(1):315–327. doi:10.1038/s41380-021-01306-8
42. Colasanto M, Madigan S, Korczak DJ. Depression and inflammation among children and adolescents: a meta-analysis. *J Affect Disord*. 2020;277:940–948. doi:10.1016/j.jad.2020.09.025
43. Howe AS, Lynch DA. Cytokine alterations in pediatric internalizing disorders: systematic review and exploratory multi-variate meta-analysis. *Brain Behav Immun Heal*. 2022;24. doi:10.1016/j.bbih.2022.100490
44. Su S, Miller AH, Snieder H, et al. Common genetic contributions to depressive symptoms and inflammatory markers in middle-aged men: the twins heart study. *Psychosom Med*. 2009;71(2):152–158. doi:10.1097/PSY.0b013e31819082ef
45. Barnes J, Mondelli V, Pariante CM. Genetic contributions of inflammation to depression. *Neuropsychopharmacology*. 2017;42(1):81–98. doi:10.1038/npp.2016.169
46. Hwang JP, Tsai SJ, Hong CJ, Yang CH, Hsu CD, Liou YJ. Interleukin-1 beta –511C/T genetic polymorphism is associated with age of onset of geriatric depression. *NeuroMolecular Med*. 2009;11(4):322–327. doi:10.1007/s12017-009-8078-x
47. Baune BT, Dannlowski U, Domschke K, et al. The Interleukin 1 Beta (IL1B) gene is associated with failure to achieve remission and impaired emotion processing in major depression. *Biol Psychiatry*. 2010;67(6):543–549. doi:10.1016/j.biopsych.2009.11.004
48. Tartter M, Hammen C, Bower JE, Brennan PA, Cole S. Effects of chronic interpersonal stress exposure on depressive symptoms are moderated by genetic variation at IL6 and IL1β in youth. *Brain Behav Immun*. 2015;46:104–111. doi:10.1016/j.bbi.2015.01.003
49. Udina M, Moreno-España J, Navinés R, et al. Serotonin and interleukin-6: the role of genetic polymorphisms in IFN-induced neuropsychiatric symptoms. *Psychoneuroendocrinology*. 2013;38(9):1803–1813. doi:10.1016/j.psyneuen.2013.03.007
50. Roetker NS, Yonker JA, Lee C, et al. Multigenic interactions and the prediction of depression in the Wisconsin Longitudinal Study. *BMJ Open*. 2012;2(4):e000944. doi:10.1136/bmjopen-2012-000944
51. Clerici M, Arosio B, Mundo E, et al. Cytokine polymorphisms in the pathophysiology of mood disorders. *CNS Spectr*. 2009;14(8):419–425. doi:10.1017/S1092852900020393
52. Holtzman S, Abbey SE, Chan C, Bargman JM, Stewart DE. A genetic predisposition to produce low levels of IL-10 is related to depressive symptoms: a pilot study of patients with end stage renal disease. *Psychosomatics*. 2012;53(2):155–161. doi:10.1016/j.psym.2011.10.001
53. Cerri AP, Arosio B, Viazzoli C, Confalonieri R, Teruzzi F, Annoni G. 308(G/A) TNF-alpha gene polymorphism and risk of depression late in the life. *Arch Gerontol Geriatr*. 2009;49(Suppl 1):29–34. doi:10.1016/j.archger.2009.09.009
54. Halder I, Marsland AL, Cheong J, Muldoon MF, Ferrell RE, Manuck SB. Polymorphisms in the CRP gene moderate an association between depressive symptoms and circulating levels of C-reactive protein. *Brain Behav Immun*. 2010;24(1):160–167. doi:10.1016/j.bbi.2009.09.014
55. Ancelin ML, Farré A, Carrière I, Ritchie K, Chaudieu I, Ryan J. C-reactive protein gene variants: independent association with late-life depression and circulating protein levels. *Transl Psychiatry*. 2015;5(1):499.
56. Pae CU, Yu HS, Kim TS, et al. Monocyte chemoattractant protein-1 (MCP1) promoter –2518 polymorphism may confer a susceptibility to major depressive disorder in the Korean population. *Psychiatry Res*. 2004;127(3):279–281. doi:10.1016/j.psychres.2004.04.004
57. Altamura AC, Mundo E, Cattaneo E, et al. The MCP-1 Gene (SCYA2) and mood disorders: preliminary results of a case-control association study. *Neuroimmunomodulation*. 2010;17(2):126–131. doi:10.1159/000258696
58. Pae CU, Yu HS, Kim JJ, et al. Bcl2 polymorphism of the cytosolic phospholipase A2 gene and mood disorders in the Korean population. *Neuropsychobiology*. 2004;49(4):185–188. doi:10.1159/000077364
59. Fertuzinhos SMM, Oliveira JRM, Nishimura AL, et al. Analysis of IL-1α, IL-1β, and IL-1RA polymorphisms in dysthymia. *J Mol Neurosci*. 2004;22(3):251–255. doi:10.1385/JMN:22:3:251
60. McCulley MC, Day INM, Holmes C. Association between interleukin 1-β promoter (–511) polymorphism and depressive symptoms in Alzheimer’s disease. *Am J Med Genet*. 2004;124B(1):50–53. doi:10.1002/ajmg.b.20086
61. Yu YWY, Chen TJ, Hong CJ, Chen HM, Tsai SJ. Association study of the interleukin-1beta (C-51 IT) genetic polymorphism with major depressive disorder, associated symptomatology, and antidepressant response. *Neuropsychopharmacology*. 2003;28(6):1182–1185. doi:10.1038/sj.npp.1300172
62. Bufalino C, Hepgul N, Aguglia E, Pariante CM. The role of immune genes in the association between depression and inflammation: a review of recent clinical studies. *Brain Behav Immun*. 2013;31:31–47. doi:10.1016/j.bbi.2012.04.009
63. Wray NR, Ripke S, Mattheisen M, et al. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet*. 2018;50(5):668–681. doi:10.1038/s41588-018-0090-3
64. Zhao B, Fan Q, Liu J, Yin A, Wang P, Zhang W. Identification of key modules and genes associated with major depressive disorder in adolescents. *Genes*. 2022;13(3):464. doi:10.3390/genes13030464
65. Hannestad J, Dellagioia N, Bloch M. The effect of antidepressant medication treatment on serum levels of inflammatory cytokines: a meta-analysis. *Neuropsychopharmacology*. 2011;36(12):2452–2459. doi:10.1038/npp.2011.132
66. Wang L, Wang R, Liu L, Qiao D, Baldwin DS, Hou R. Effects of SSRIs on peripheral inflammatory markers in patients with major depressive disorder: a systematic review and meta-analysis. *Brain Behav Immun*. 2019;79:24–38. doi:10.1016/j.bbi.2019.02.021
67. Więdołcha M, Marcinowicz P, Krupa R, et al. Effect of antidepressant treatment on peripheral inflammation markers – a meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry*. 2018;80(Pt C):217–226. doi:10.1016/j.pnpbp.2017.04.026
68. Piletz JE, Halaris A, Iqbal O, et al. Pro-inflammatory biomarkers in depression: treatment with venlafaxine. *World J Biol Psychiatry*. 2009;10(4):313–323. doi:10.3109/15622970802573246
69. Fornaro M, Martino M, Battaglia F, Colicchio S, Perugi G. Increase in IL-6 levels among major depressive disorder patients after a 6-week treatment with duloxetine 60 mg/day: a preliminary observation. *Neuropsychiatr Dis Treat*. 2011;7(1):51–56. doi:10.2147/NDT.S16382
70. Elenkov I, Wilder R, Chrousos G, Vizi E. The sympathetic nerve—an integrative interface between two supersystems: the brain and the immune system. *Pharmacol Rev*. 2000;52(4):595–638.
71. Matveychuk D, Thomas RK, Swainson J, et al. Ketamine as an antidepressant: overview of its mechanisms of action and potential predictive biomarkers. *Ther Adv Psychopharmacol*. 2020;10:2045125320916657. doi:10.1177/2045125320916657
72. Chen MH, Li CT, Lin WC, et al. Rapid inflammation modulation and antidepressant efficacy of a low-dose ketamine infusion in treatment-resistant depression: a randomized, double-blind control study. *Psychiatry Res*. 2018;269:207–211. doi:10.1016/j.psychres.2018.08.078

73. Zhan Y, Zhou Y, Zheng W, et al. Alterations of multiple peripheral inflammatory cytokine levels after repeated ketamine infusions in major depressive disorder. *Transl Psychiatry*. 2020;10(1):1–9. doi:10.1038/s41398-020-00933-z
74. Becerril-Villanueva E, Pérez-Sánchez G, Alvarez-Herrera S, et al. Alterations in the levels of growth factors in adolescents with major depressive disorder: a longitudinal study during the treatment with fluoxetine. *Mediators Inflamm*. 2019;2019:9130868. doi:10.1155/2019/9130868
75. Amitai M, Taler M, Carmel M, et al. The relationship between plasma cytokine levels and response to selective serotonin reuptake inhibitor treatment in children and adolescents with depression and/or anxiety disorders. *J Child Adolesc Psychopharmacol*. 2016;26(8):727–732. doi:10.1089/cap.2015.0147
76. Rush AJ, Trivedi MH, Wisniewski SR, et al. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR\*D report. *Am J Psychiatry*. 2006;163(11):1905–1917. doi:10.1176/ajp.2006.163.11.1905
77. Jha MK, Trivedi MH. Pharmacogenomics and biomarkers of depression. In: Macaluso M, Preskorn S, editors. *Antidepressants. Handbook of Experimental Pharmacology*. Cham: Springer; 2019:101–113.
78. Tying S, Gottlieb A, Papp K, et al. Etanercept and clinical outcomes, fatigue, and depression in psoriasis: double-blind placebo-controlled randomised Phase III trial. *Lancet*. 2006;367(9504):29–35. doi:10.1016/S0140-6736(05)67763-X
79. Menter A, Augustin M, Signorovitch J, et al. The effect of Adalimumab on reducing depression symptoms in patients with moderate to severe psoriasis: a randomized clinical trial. *J Am Acad Dermatol*. 2010;62(5):812–818. doi:10.1016/j.jaad.2009.07.022
80. Sun Y, Wang D, Salvadore G, et al. The effects of interleukin-6 neutralizing antibodies on symptoms of depressed mood and anhedonia in patients with rheumatoid arthritis and multicentric Castleman's disease. *Brain Behav Immun*. 2017;66:156–164. doi:10.1016/j.bbi.2017.06.014
81. Kappelmann N, Lewis G, Dantzer R, Jones PB, Khandaker GM. Antidepressant activity of anti-cytokine treatment: a systematic review and meta-analysis of clinical trials of chronic inflammatory conditions. *Mol Psychiatry*. 2018;23(2):335–343. doi:10.1038/mp.2016.167
82. Berk M, Mohebbi M, Dean OM, et al. Youth Depression Alleviation with Anti-inflammatory Agents (YoDA-A): a randomised clinical trial of rosuvastatin and aspirin. *BMC Med*. 2020;18(1):16. doi:10.1186/s12916-019-1475-6
83. Platt JM, Bates L, Jager J, McLaughlin KA, Keyes KM. Is the US Gender gap in depression changing over time? A meta-regression. *Am J Epidemiol*. 2021;190(7):1190–1206. doi:10.1093/aje/kwab002
84. Hankin BL, Young JF, Abela JRZ, et al. Depression from childhood into late adolescence: influence of gender, development, genetic susceptibility, and peer stress. *J Abnorm Psychol*. 2015;124(4):803–816. doi:10.1037/abn0000089
85. Xing E, Billi AC, Gudjonsson JE. Sex bias and autoimmune diseases. *J Invest Dermatol*. 2022;142(3):857–866. doi:10.1016/j.jid.2021.06.008
86. Gimeno D, Kivimäki M, Brunner EJ, et al. Associations of C-reactive protein and interleukin-6 with cognitive symptoms of depression: 12-Year follow-up of the Whitehall II study. *Psychol Med*. 2009;39(3):413–423. doi:10.1017/S0033291708003723
87. Koskinas J, Merkouraki P, Manesis E, Hadziyannis S. Assessment of depression in patients with chronic viral hepatitis: effect of interferon treatment. *Dig Dis*. 2002;20(3–4):284–288. doi:10.1159/000067682
88. Bremner MA, Beekman ATF, Deeg DJH, et al. Inflammatory markers in late-life depression: results from a population-based study. *J Affect Disord*. 2008;106(3):249–255. doi:10.1016/j.jad.2007.07.002
89. Bonaccorso S, Marino V, Puzella A, et al. Increased depressive ratings in patients with hepatitis C receiving interferon- $\alpha$ -based immunotherapy are related to interferon- $\alpha$ -induced changes in the serotonergic system. *J Clin Psychopharmacol*. 2002;22(1):86–90. doi:10.1097/00004714-200202000-00014
90. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*. 5th ed. Arlington, VA, USA: American Psychiatric Publishing; 2013.
91. Kovacs M. *Children's Depression Inventory. CDI Manual*. Toronto, Ontario, Canada: Multi-Health Systems; 1992.
92. Guimarães PM, Scavuzzi BM, Stadtlober NP, et al. Cytokines in systemic lupus erythematosus: far beyond Th1/Th2 dualism lupus: cytokine profiles. *Immunol Cell Biol*. 2017;95(9):824–831. doi:10.1038/icb.2017.53
93. Andrade C. Z scores, standard scores, and composite test scores explained. *Indian J Psychol Med*. 2021;43(6):555–557. doi:10.1177/02537176211046525
94. Brunoni AR, Supasitthumrong T, Teixeira AL, et al. Differences in the immune-inflammatory profiles of unipolar and bipolar depression. *J Affect Disord*. 2020;262:8–15. doi:10.1016/j.jad.2019.10.037
95. Chiang JJ, Cole SW, Bower JE, et al. Depressive symptoms and immune transcriptional profiles in late adolescents. *Brain Behav Immun*. 2019;80:163–169. doi:10.1016/j.bbi.2019.03.004
96. Pfeffer K. Biological functions of tumor necrosis factor cytokines and their receptors. *Cytokine Growth Factor Rev*. 2003;14(3–4):185–191. doi:10.1016/S1359-6101(03)00022-4
97. Olmos G, Lladó J. Tumor necrosis factor alpha: a link between neuroinflammation and excitotoxicity. *Mediators Inflamm*. 2014;2014:861231. doi:10.1155/2014/861231
98. Gough P, Myles IA. Tumor necrosis factor receptors: pleiotropic signaling complexes and their differential effects. *Front Immunol*. 2020;11:585880. doi:10.3389/fimmu.2020.585880
99. Chemaly M, McGilligan V, Gibson M, et al. Role of tumour necrosis factor alpha converting enzyme (TACE/ADAM17) and associated proteins in coronary artery disease and cardiac events. *Arch Cardiovasc Dis*. 2017;110(12):700–711. doi:10.1016/j.acvd.2017.08.002
100. Sedger LM, McDermott MF. TNF and TNF-receptors: from mediators of cell death and inflammation to therapeutic giants - past, present and future. *Cytokine Growth Factor Rev*. 2014;25(4):453–472. doi:10.1016/j.cytogfr.2014.07.016
101. Gómez MI, Sokol SH, Muir AB, Soong G, Bastien J, Prince AS. Bacterial induction of TNF- $\alpha$  converting enzyme expression and IL-6 receptor  $\alpha$  shedding regulates airway inflammatory signaling. *J Immunol*. 2005;175(3):1930–1936. doi:10.4049/jimmunol.175.3.1930
102. Mackiewicz A, Schooltink H, Heinrich PC, Rose-John S. Complex of soluble human IL-6-receptor/IL-6 up-regulates expression of acute-phase proteins. *J Immunol*. 1992;149(6):2021–2027.
103. Scheller J, Chalaris A, Schmidt-Arras D, Rose-John S. The pro- and anti-inflammatory properties of the cytokine interleukin-6. *Biochim Biophys Acta Mol Cell Res*. 2011;1813(5):878–888. doi:10.1016/j.bbamcr.2011.01.034
104. Hurst SM, Wilkinson TS, McLoughlin RM, et al. IL-6 and its soluble receptor orchestrate a temporal switch in the pattern of leukocyte recruitment seen during acute inflammation. *Immunity*. 2001;14(6):705–714. doi:10.1016/S1074-7613(01)00151-0

105. Maes M, Anderson G, Kubera M, Berk M. Targeting classical IL-6 signalling or IL-6 trans-signalling in depression? *Expert Opin Ther Targets*. 2014;18(5):495–512. doi:10.1517/14728222.2014.888417
106. Sowa-Kućma M, Styczeń K, Siwek M, et al. Lipid peroxidation and immune biomarkers are associated with major depression and its phenotypes, including treatment-resistant depression and melancholia. *Neurotox Res*. 2018;33(2):448–460. doi:10.1007/s12640-017-9835-5
107. Sowa-Kućma M, Styczeń K, Siwek M, et al. Are there differences in lipid peroxidation and immune biomarkers between major depression and bipolar disorder: effects of melancholia, atypical depression, severity of illness, episode number, suicidal ideation and prior suicide attempts. *Prog Neuropsychopharmacol Biol Psychiatry*. 2018;81:372–383. doi:10.1016/j.pnpbp.2017.08.024
108. Mosser DM, Zhang X. Interleukin-10: new perspectives on an old cytokine. *Immunol Rev*. 2008;226(1):205–218. doi:10.1111/j.1600-065X.2008.00706.x
109. Couper KN, Blount DG, Riley EM. IL-10: the master regulator of immunity to infection. *J Immunol*. 2008;180(9):5771–5777. doi:10.4049/jimmunol.180.9.5771
110. Luzina IG, Keegan AD, Heller NM, Rook GAW, Shea-Donohue T, Atamas SP. Regulation of inflammation by interleukin-4: a review of “alternatives”. *J Leukoc Biol*. 2012;92(4):753–764. doi:10.1189/jlb.0412214
111. Oglódek E. Changes in the serum levels of cytokines: IL-1 $\beta$ , IL-4, IL-8 and IL-10 in depression with and without posttraumatic stress disorder. *Brain Sci*. 2022;12(3):387. doi:10.3390/BRAINS112030387
112. Maes M, Rachayon M, Jirakran K, et al. The immune profile of major dysmood disorder: proof of concept and mechanism using the precision nomothetic psychiatry approach. *Cells*. 2022;11(7):1183. doi:10.3390/cells11071183
113. Maes M, Berk M, Goehler L, et al. Depression and sickness behavior are Janus-faced responses to shared inflammatory pathways. *BMC Med*. 2012;10:66. doi:10.1186/1741-7015-10-66
114. Das R, Emon MPZ, Shahriar M, et al. Higher levels of serum IL-1 $\beta$  and TNF- $\alpha$  are associated with an increased probability of major depressive disorder. *Psychiatry Res*. 2021;295:113568. doi:10.1016/j.psychres.2020.113568
115. Moieni M, Irwin MR, Jevtic I, Olmstead R, Breen EC, Eisenberger NI. Sex differences in depressive and socioemotional responses to an inflammatory challenge: implications for sex differences in depression. *Neuropsychopharmacology*. 2015;40(7):1709–1716. doi:10.1038/npp.2015.17
116. Kim J, Kim JH, Chang KA. Sex difference in peripheral inflammatory biomarkers in drug-naïve patients with major depression in young adulthood. *Biomedicines*. 2021;9(7):708. doi:10.3390/biomedicines9070708
117. Zainal NH, Newman MG. Increased inflammation predicts nine-year change in major depressive disorder diagnostic status. *J Abnorm Psychol*. 2021;130(8):829–840. doi:10.1037/abn0000716
118. Birur B, Amrock EM, Shelton RC, Li L. Sex differences in the peripheral immune system in patients with depression. *Front Psychiatry*. 2017;8:16. doi:10.3389/fpsyt.2017.00108
119. Ford DE, Erlinger TP. Depression and C-reactive protein in US adults: data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med*. 2004;164(9):1010–1014. doi:10.1001/archinte.164.9.1010
120. Ramsey JM, Cooper JD, Bot M, et al. Sex differences in serum markers of major depressive disorder in the Netherlands Study of Depression and Anxiety (NESDA). *PLoS One*. 2016;11(5):e0156624. doi:10.1371/journal.pone.0156624
121. Valiathan R, Ashman M, Asthana D. Effects of ageing on the immune system: infants to elderly. *Scand J Immunol*. 2016;83(4):255–266. doi:10.1111/sji.12413
122. West LS. Defining critical windows in the development of the human immune system. *Hum Exp Toxicol*. 2002;21(9–10):499–505. doi:10.1191/0960327102ht288oa
123. Shames RS. Gender differences in the development and function of the immune system. *J Adolesc Health*. 2002;30:59–70. doi:10.1016/S1054-139X(01)00382-2
124. Traish A, Bolanos J, Nair S, Saad F, Morgentaler A. Do androgens modulate the pathophysiological pathways of inflammation? Appraising the contemporary evidence. *J Clin Med*. 2018;7(12):549. doi:10.3390/jcm7120549
125. Straub RH. The complex role of estrogens in inflammation. *Endocr Rev*. 2007;28(5):521–574. doi:10.1210/er.2007-0001
126. Schwarz MJ. Cytokines, neurophysiology, neuropsychology, and psychiatric symptoms. *Dialogues Clin Neurosci*. 2003;5(2):139–153. doi:10.31887/dens.2003.5.2/mschwarz
127. Quan N, Banks WA. Brain-immune communication pathways. *Brain Behav Immun*. 2007;21(6):727–735. doi:10.1016/j.bbi.2007.05.005
128. Watkins LR, Goehler LE, Relton JK, et al. Blockade of interleukin-1 induced hyperthermia by subdiaphragmatic vagotomy: evidence for vagal mediation of immune-brain communication. *Neurosci Lett*. 1995;183(1–2):27–31. doi:10.1016/0304-3940(94)11105-R
129. D’Mello C, Le T, Swain MG. Cerebral microglia recruit monocytes into the brain in response to tumor necrosis factor signaling during peripheral organ inflammation. *J Neurosci*. 2009;29(7):2089–2102. doi:10.1523/JNEUROSCI.3567-08.2009
130. Huang X, Hussain B, Chang J. Peripheral inflammation and blood–brain barrier disruption: effects and mechanisms. *CNS Neurosci Ther*. 2021;27(1):36–47. doi:10.1111/cns.13569
131. Bakunina N, Pariante CM, Zunszain PA. Immune mechanisms linked to depression via oxidative stress and neuroprogression. *Immunology*. 2015;144(3):365–373. doi:10.1111/imm.12443
132. Miller AH, Maletic V, Raison CL. Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biol Psychiatry*. 2009;65(9):732–741. doi:10.1016/j.biopsych.2008.11.029
133. Tonhajzerova I, Sekaninova N, Bona Olexova L, Visnovcova Z. Novel insight into neuroimmune regulatory mechanisms and biomarkers linking major depression and vascular diseases: the dilemma continues. *Int J Mol Sci*. 2020;21(7):2317. doi:10.3390/ijms21072317
134. Anderson G, Maes M. Oxidative/Nitrosative stress and immuno-inflammatory pathways in depression: treatment implications. *Curr Pharm Des*. 2014;20(23):3812–3847. doi:10.2174/13816128113196660738
135. Dantzer R, O’Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci*. 2008;9(1):46–56. doi:10.1038/nrn2297
136. Camacho-Arroyo I, López-Griego L, Morales-Montor J. The role of cytokines in the regulation of neurotransmission. *Neuroimmunomodulation*. 2009;16(1):1–12. doi:10.1159/000179661
137. Capuron L, Ravaut A, Neveu PJ, Miller AH, Maes M, Dantzer R. Association between decreased serum tryptophan concentrations and depressive symptoms in cancer patients undergoing cytokine therapy. *Mol Psychiatry*. 2002;7(5):468–473. doi:10.1038/sj.mp.4000995

138. Munn DH, Mellor AL. Indoleamine 2,3 dioxygenase and metabolic control of immune responses. *Trends Immunol.* 2013;34(3):137–143. doi:10.1016/j.it.2012.10.001
139. Dantzer R. Neuroimmune interactions: from the brain to the immune system and vice versa. *Physiol Rev.* 2018;98(1):477–504. doi:10.1152/physrev.00039.2016
140. Walker AK, Budac DP, Bisulco S, et al. NMDA receptor blockade by ketamine abrogates lipopolysaccharide-induced depressive-like behavior in C57BL/6J mice. *Neuropsychopharmacology.* 2013;38(9):1609–1616. doi:10.1038/npp.2013.71
141. Sattar Y, Wilson J, Khan AM, et al. A review of the mechanism of antagonism of N-methyl-D-aspartate receptor by ketamine in treatment-resistant depression. *Cureus.* 2018;10(5):e2652. doi:10.7759/cureus.2652
142. Baganz NL, Blakely RD. A dialogue between the immune system and brain, spoken in the language of serotonin. *ACS Chem Neurosci.* 2013;4(1):48–63. doi:10.1021/cn300186b
143. Bin ZC, Blakely RD, Hewlett WA. The proinflammatory cytokines interleukin-1beta and tumor necrosis factor-alpha activate serotonin transporters. *Neuropsychopharmacology.* 2006;31(10):2121–2131. doi:10.1038/sj.npp.1301029
144. Cowen PJ, Browning M. What has serotonin to do with depression? *World Psychiatry.* 2015;14(2):158–160. doi:10.1002/wps.20229
145. Müller N, Schwarz MJ. The immune-mediated alteration of serotonin and glutamate: towards an integrated view of depression. *Mol Psychiatry.* 2007;12(11):988–1000. doi:10.1038/sj.mp.4002006

Journal of Inflammation Research

Dovepress

## Publish your work in this journal

The Journal of Inflammation Research is an international, peer-reviewed open-access journal that welcomes laboratory and clinical findings on the molecular basis, cell biology and pharmacology of inflammation including original research, reviews, symposium reports, hypothesis formation and commentaries on: acute/chronic inflammation; mediators of inflammation; cellular processes; molecular mechanisms; pharmacology and novel anti-inflammatory drugs; clinical conditions involving inflammation. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>