

# Systematic review of endoscopic ultrasound-guided biliary drainage versus percutaneous transhepatic biliary drainage

Authors: Zeinab Hassan<sup>A</sup> and Eyad Gadour<sup>B</sup>

## Introduction

Endoscopic ultrasound-guided biliary drainage (EUS-BD) is a novel technique that allows biliary drainage by echoendoscopy and fluoroscopy using a stent from the biliary tree to the gastrointestinal tract. Percutaneous transhepatic cholangiography biliary drainage (PTBD) is a diagnostic and therapeutic procedure that involves inserting a needle into the biliary tree, followed by the immediate insertion of a catheter. This study examined the technical aspects and outcomes of these different approaches to biliary drainage.

## Materials and methods

We compared the technical aspects and outcomes of two different approaches to biliary drainage: EUS-BD and PTBD. Different databases (including PubMed, Embase, ClinicalTrials.gov, the Cochrane library, Scopus and Google Scholar) were searched according to the PRISMA guidelines to obtain studies comparing PTBD and EUS-BD.

## Results

Among the six studies that fulfilled the inclusion criteria, PTBD patients underwent significantly more reinterventions (4.9 vs 1.3), experienced more post-procedure pain (4.1 vs 1.9) and experienced more late adverse events (53.8% vs 6.6%) than EUS-BD patients (Table 1).<sup>1-6</sup> The EUS-BD group had a higher success rate of biliary drainage (92% vs 46%;  $p > 0.05$ ) and a lower rate of adverse events (20% vs 46%;  $p = 0.05$ ) than PTBD group. There was a significant reduction in total bilirubin in both groups (from 16.4  $\mu\text{mol/L}$  to 3.3  $\mu\text{mol/L}$  for EUS-BD and 17.2  $\mu\text{mol/L}$  to 3.8  $\mu\text{mol/L}$  for PTBD;  $p = 0.002$ ) at the 7-day follow-up. There were no significant differences observed for complication rates between PTBD and EUS-BD (3.3 vs 3.8, respectively). PTBD was associated with a higher adverse event rate than EUS-BD in all procedures, including reinterventions (80.4% vs 15.7%, respectively) and a higher index procedure (39.2% vs 18.2%, respectively).

## Conclusion

The findings of this systematic review revealed that EUS-BD is linked with a higher rate of effective biliary drainage and a more manageable procedure-related adverse event profile than PTBD. EUS-BD could become a first-line biliary drainage treatment instead of

Authors: <sup>A</sup>Stockport NHS Foundation Trust, Stockport, UK; <sup>B</sup>University Hospitals of Morecambe Bay NHS Foundation Trust, Kendal, UK

**Table 1. Rates of clinical and technical success in the included studies**

Study	Technical success		Clinical success	
	EGBD, event/total cases, n	PTBD, event/total cases, n	EGBD, event/total cases, n	PTBD, event/total cases, n
Artifon <i>et al</i> <sup>1</sup>	13/13	12/12	13/13	12/12
Bapaye <i>et al</i> <sup>2</sup>	23/25	26/26	23/25	26/26
Khashab <i>et al</i> <sup>3</sup>	19/22	51/51	19/19	47/51
Giovannini <i>et al</i> <sup>4</sup>	19/20	17/17	18/19	17/17
Jung <i>et al</i> <sup>5</sup>	32/34	31/32	28/32	27/31
Sharaiha <i>et al</i> <sup>6</sup>	43/47	12/13	27/43	3/12

EGBD = endoscopic ultrasound-guided choledochoduodenostomy; PTBD = percutaneous transhepatic biliary drainage.

endoscopic retrograde cholangiopancreatography if the outcomes of clinical studies are positive and technologies are simplified. Prospective, randomised controlled studies are required to clarify these issues. ■

## References

- Artifon EL, Aparicio D, Paione JB *et al*. Biliary drainage in patients with unresectable, malignant obstruction where ERCP fails: endoscopic ultrasonography-guided choledochoduodenostomy vs percutaneous drainage. *J Clin Gastroenterol* 2012;46:768–74.
- Bapaye A, Dubale N, Aher A. Comparison of endosonography-guided vs. percutaneous biliary stenting when papilla is inaccessible for ERCP. *United European Gastroenterol J* 2013;1:285–93.
- Khashab MA, Valeshabad AK, Afghani E *et al*. A comparative evaluation of EUS-guided biliary drainage and percutaneous drainage in patients with distal malignant biliary obstruction and failed ERCP. *Dig Dis Sci* 2015;60:557–65.
- Giovannini M. Learning in therapeutic EUS. *Endosc Ultrasound* 2021;10:317–8.
- Jung Y, Lee J, Cho JY *et al*. Comparison of efficacy and safety between endoscopic submucosal dissection and transanal endoscopic microsurgery for the treatment of rectal tumor. *Saudi J Gastroenterol* 2018;24:115–21.
- Sharaiha RZ, Kumta NA, Desai AP *et al*. Endoscopic ultrasound-guided biliary drainage vs percutaneous transhepatic biliary drainage: predictors of successful outcome in patients who fail endoscopic retrograde cholangiopancreatography. *Surg Endosc* 2016;30:5500–5.