


RESEARCH

Open Access



# Whole-cell tumor vaccines desialylated to uncover tumor antigenic Gal/GalNAc epitopes elicit anti-tumor immunity

Jianmei Huang<sup>1</sup>, Meiyong Li<sup>2</sup>, Bingjie Mei<sup>1</sup>, Junyang Li<sup>1</sup>, Yi Zhu<sup>3</sup>, Qiaoshan Guo<sup>2</sup>, Jianming Huang<sup>2\*</sup> and Guonan Zhang<sup>1,4\*</sup> 

## Abstract

**Background:** Aberrant sialoglycans on the surface of tumor cells shield potential tumor antigen epitopes, escape recognition, and suppress activation of immunocytes.  $\alpha$ 2,3/ $\alpha$ 2,6Gal- and  $\alpha$ 2,6GalNAc (Gal/GalNAc)-linked sialic acid residues of sialoglycans could affect macrophage galactose-type lectins (MGL) mediated-antigen uptake and presentation and promote sialic acid-binding immunoglobulin-like lectins (Siglecs) mediated-immunosuppression. Desialylating sialoglycans on tumor cells could present tumor antigens with Gal/GalNAc residues and overcome glyco-immune checkpoints. Thus, we explored whether vaccination with desialylated whole-cell tumor vaccines (DWCTVs) triggers anti-tumor immunity in ovarian cancer (OC).

**Methods:** Sialic acid (Sia) and Gal/GalNAc residues on OC A2780, OVCAR3, and ID8 cells treated with  $\alpha$ 2-3 neuraminidase ( $\alpha$ 2-3NA) and  $\alpha$ 2-6NA, and Sigec-9 or Siglec-E and MGL on DCs pulsed with desialylated OC cells were identified using flow cytometry (FCM); RT-qPCR determined IFNG expression of T cells, TRBV was sequenced using Sanger sequencing and cytotoxicity of  $\alpha$  $\beta$  T cells was measured with LDH assay; Anti-tumor immunity in vivo was validated via vaccination with desialylated whole-cell ID8 vaccine (ID8 DWCTVs).

**Results:** Gal/GalNAc but not Sia residues were significantly increased in the desialylated OC cells.  $\alpha$ 2-3NA-modified DWCTV increased MGL but decreased Siglec-9 or Siglec E expression on DCs. MGL<sup>bright</sup>/Siglec-9<sup>dim</sup> DCs significantly up-regulated IFNG expression and CD4/CD8 ratio of T cells and diversified the TCR repertoire of  $\alpha$  $\beta$  T-cells that showed enhanced cytotoxic activity. Vaccination with  $\alpha$ 2-3NA-modified ID8 DWCTVs increased MGL<sup>bright</sup>/Siglec-E<sup>dim</sup> DCs in draining lymph nodes, limited tumor growth, and extended survival in tumor-challenged mice.

**Conclusion:** Desialylated tumor cell vaccine could promote anti-tumor immunity and provide a strategy for OC immunotherapy in a clinical setting.

**Keywords:** Sialoglycans, Gal/GalNAc, Siglec-9, MGL, Tumor vaccine, TCR-V $\beta$  repertoire

## Background

Cell surface glycans on membrane-bound proteins are essential regulators of the immune system that regulate immune cell development and function, modulate immune receptor interactions, and serve as ligands for glycan-binding proteins (lectins) expressed by immune cells [1, 2]. Terminal sialic acid residues are abundant in glycan chains of glycoproteins and glycolipids on the

\*Correspondence: wesleyhuangcn2002@163.com; zhanggn@hotmail.com

<sup>1</sup> School of Medicine, University of Electronic Science and Technology of China, Chengdu, China

<sup>2</sup> Biochemistry and Molecular Biology, Sichuan Cancer Institute, Chengdu, China

Full list of author information is available at the end of the article



surface of all live cells forming an outer layer of the cell known initially as glycocalyx [3]. Moreover, they mediate several of the regulatory functions of glycans in the immune system. Sialic acid-containing glycans (sialoglycans) are involved in numerous molecular interactions at the cell surface and have several direct and indirect functions in the immune system [4, 5].

Tumor antigens produced by somatic DNA mutations due to the inherent genetic instability are glycosidically hypersialylated. Aberrant glycosylation is a common feature of many cancers. Alteration of cell surface sialylation is seen with transformation to a tumor cell. Altered glycosylation sites in epithelial cancer cells may play an essential role in tumor progression, as they may affect tumor cell migration and antigen presentation by antigen-presenting cells (APCs) [6–8]. Altered sialylation has long been associated with the malignancy of carcinoma. High expression of sialic acids has been proposed to protect cancer cells from recognition and eradication by the immune system [9].

It has been demonstrated that sialic acids linked to galactose (Gal) and *N*-acetylgalactosamine (GalNAc) via  $\alpha$ -2,3- and  $\alpha$ -2,6-linkages significantly increased in cancer compared to normal tissues [10, 11]. They allow tumor cells to become 'invisible' to avoid immune recognition by DCs as APCs and attack by anti-tumor T cells. The macrophage galactose-type lectin (MGL) on DCs has been shown to induce immunosuppressive responses upon recognizing aberrant sialylation on cancer cells [12, 13]. MGL is the lectin that exclusively binds terminal Gal/GalNAc epitopes of tumor-associated glycan [14, 15]. It is an antigen-uptake receptor for the internalization of Gal/GalNAc carrying immunogens delivered into MHC class I and II compartments, thus improving DC performance to facilitate MHC-restricted antigen presentation to T cells and, most importantly, increasing antigen-specific CD8 + T-cell activation [16–19].

Sialoglycans are recognized by sialic acid-specific receptors on immune cells, such as sialic acid-binding immunoglobulin-like lectins (Siglecs). The human Siglec family consists of 15 members that are broadly expressed by the majority of immune cells. They can be categorized as structurally conserved Siglecs (Siglec-1, -2, -4, and -15) and CD33-related Siglecs (CD33 or Siglec-3, Siglec-5, -6, -7, -8, -9, -10, -11, -12, -14, and -16). As numerous human malignancies express ligands for both Siglec-7 and Siglec-9, the CD33-related Siglec-7 and Siglec-9 are of special importance in the context of tumor immunotherapy [20, 21]. Siglec-7 and Siglec-9 are reported to influence NK cell-dependent tumor immunosurveillance [20]. Siglec-7 binds to 2,8-sialyl residues with the highest affinity, while Siglec-9 preferentially binds to 2,6-sialyl and 2,3-sialyl residues [22]. The sialoglycan-Siglec

interaction suppresses an immune response as sialic acids are considered self-associated molecular patterns [20, 23–25]. The surface glycan profile of human tumor cells was dominated by  $\alpha$ -2,3- and  $\alpha$ -2,6-linked sialic acid-capped complex *N*-glycans and bi-antennary *N*-glycans. Sialic acids can trigger immune inhibitory Siglecs on DCs, affecting antigen presentation to CD4+ and CD8+ T cells [26]. Gal/GalNAc-linked sialic acids of sialoglycans are known to interact with Siglec 9, an inhibitory immune checkpoint expressed broadly on immune cells, including DCs, which can impede T cell-mediated anti-tumor responses through immunoreceptor tyrosine-based inhibitory motifs (ITIM) [27–31].

Poor presentation of tumor antigens to the immune system remains a major obstacle to effective anti-tumor vaccine therapy [32]. Aberrant sialylation of tumor antigens enhances immune evasion and reduces the efficacy of several types of available immunotherapies, including tumor vaccine therapy [9, 33].

These findings suggest that removing  $\alpha$ -2,3Gal- and  $\alpha$ -2,6GalNAc-linked sialic acid residues from sialoglycans on the surface of tumor cells could improve tumor antigen presentation by DCs, block Siglec-sialic acid interactions, and boost anti-tumor immune responses. Thus, targeted desialylation of sialoglycan on tumor cells can be used to design novel anti-tumor vaccines [29, 34–36]. Here, we selectively removed the  $\alpha$ -2,3- and  $\alpha$ -2,6-linked sialic acids on OC cells by  $\alpha$ 2-3 and  $\alpha$ 2-6 neuraminidases, respectively, to generate desialylated whole-cell tumor vaccines (DWCTVs), resulting in the novel delivery approach for tumor antigens with Gal/GalNAc epitopes. Increased Gal/GalNAc epitopes accompanied the decrease of sialic acids on OC cells. DWCTVs could mature and activate DCs, reduce Siglec-9 and elevate MGL on DCs. These Siglec-9<sup>dim</sup>/MGL<sup>bright</sup> DCs diversified the TCR-V $\beta$  repertoires of T cells, resulting in enhanced cytotoxicity against the parental OC cells in vitro. Vaccination with DWCTVs could delay tumor formation, limit tumor growth, and prolong the survival of mice inoculated with tumor cells. The "win-win" effects of DWCTVs might be their blocking of Siglec-sialic acid interaction and increasing MGL-Gal/GalNAc interaction between immune cells and OC cells. DWCTVs in our study provided a promising therapeutic option to render tumors permissive to immune attack to reduce or prevent tumor recurrence.

## Methods

### Cell lines and animals

Human OC cell lines (OVCAR-3 and A2780) and the mouse OC cell line (ID8) were purchased from the Committee on Type Culture Collection of the Chinese Academy of Sciences. OVCAR-3 cells were maintained in

McCoy's 5A (Sigma), containing 1% human insulin and 15% fetal calf serum (FCS). A2780 and ID8 cells were cultured with RPMI-1640 (Gibco) and DMEM (Gibco), respectively, supplemented with 10% FCS. Penicillin (100 IU mL<sup>-1</sup>) and gentamicin (40 IU mL<sup>-1</sup>) were added to the mediums for these three cells. All cells were cultured in a humidified incubator at 37 °C with 5% CO<sub>2</sub>.

Specific pathogen-free, 6–8-week-old female C57BL/6 littermate mice (20 ± 2 g) (Dossy Experimental Animals Co. LTD Chengdu, China) were used for tumorigenicity studies. Animal studies were carried out following protocols approved by the Ethics Committee of Sichuan Cancer Hospital (SCCHEC-04-2019-004).

#### Preparation and characterization of DWCTVs

The cells (1 × 10<sup>6</sup> cells/mL) were pre-treated with 100 ng mL<sup>-1</sup> of mitomycin C (R&D, 3258/10) in serum-free medium for 3 h at 37 °C, washed and resuspended, and then desialylated with 100 mU mL<sup>-1</sup> of α-2-3 and α-2-6 neuraminidases (NAs) from *Clostridium perfringens* (Sigma, N2876) and *Arthrobacter ureafaciens* (Roche, 10269611001) in the same medium for 30 min at 37 °C, respectively. The terminal α-2,3/α-2,6 sialic acid and Gal/GalNAc residues on the resultant cells were identified using flow cytometry (FCM) with Biotinylated MALII/DyLight 488 streptavidin (B-1265-1/SA-5488-1), Fluorescein labeled SNA (FL-1301), and Cy5 labeled PNA (CL-1075-1).

#### DC cultures and maturation

Peripheral blood mononuclear cells (PBMCs) were isolated from the healthy adult female volunteer by Ficoll-Hypaque (1.077 g mL<sup>-1</sup>) gradient centrifugation and plated in a complete RPMI-1640 medium in culture plates for 4 h at 37 °C. The adherent monocytes were cultured in a complete RPMI-1640 medium with 500 U mL<sup>-1</sup> interleukin 4 (IL 4) (R&D, 204-IL-010) and 150 ng mL<sup>-1</sup> granulocyte–macrophage colony-stimulating factor (GM-CSF) (R&D, 215-GM-010) for 7 consecutive days. After then, the immature DCs (imDCs) were harvested, washed, and pulsed with an equal number of WCTV cells in the same medium containing 250 U mL<sup>-1</sup> IL 4 and 75 ng mL<sup>-1</sup> GM-CSF for 3 days. The mature DCs (mDCs) were then analyzed by flow cytometry for DC-surface maturation and activation markers (CD86, CD83), Siglec-9, and MGL expression.

#### T cell stimulation with the mDCs

The nonadherent cells of PBMCs were harvested, washed twice with PBS, and then incubated with the mDCs at a 10:1 ratio of cell number overnight. The stimulated lymphocytes were harvested to detect *IFNG* expression by RT-qPCR and CD4/CD8 ratio by FCM. Then these

lymphocytes were expanded in the same medium with 500 U mL<sup>-1</sup> IL-2 (Solarbio, GMP-11848-HNAE) for 5 days. The TCR-Vβ sequences and cytotoxicity of the resultant T cells were identified by Sanger sequencing and LDH assay, respectively.

#### Real-time quantitative PCR for *IFNG* expression of human T cells

Total RNA was extracted from human T cells using TRIzol reagent (Invitrogen, 15596026). A total of 2 μg RNA was reverse transcribed using Eastep RT Master Mix Kit (Promega, LS2052), and 1 μL cDNA was used for qPCR using Eastep qPCR Master Mix Kit (Promega, LS2062), and the primer pairs for human *IFNG*: forward: 5'-TCG GTAAC TGA CT TGA ATG TCCA-3', reverse: 5'-TCG CTTCCCTGTTT TAGCTGC-3' and the primer pairs for human *GAPDH*: forward 5'-CAAGGTCATCCATGACAACTTTG-3, reverse 5'-TCGCTTCCCTGTTT TAGCTGC-3'

#### FCM assay

All lectins and antibodies were obtained directly from Vector Labs or BD Pharmingen, R&D, Biolegend, Abcam, and Servicebio. According to the manufacturers' protocols, cells were incubated with antibodies for 30 min, and analysis was performed on a BD FACS Canto II flow cytometer.

#### TCR-Vβ repertoire sequencing

TCR beta-chain variable (*TRBV*) gene of human T cells was PCR-sequenced. Briefly, to PCR-amplify and clone the sequences of TCR β-chains, 5'RACE was performed with 1.0 μg of total RNA from human T cells co-incubated with mDCs using RLM-RACE Kit (Thermo, AM1700) and primers specific for the gene (inner primer 5'-TTCTGATGGCTCAAACAC-3' and outer primer 5'-GGTCCACTCGTCATTCTCCGA-3'). Then, 4 μL of the recovered and purified DNA fragments larger than 400 bp were ligated into 1 μL pEASY T3 cloning vector (TRANS, CT301). The cloned DNA was sequenced using Sanger sequencing and analyzed using the IMG T/V-QUEST online tool (IMG T-the international ImMunoGeneTics information system, <http://www.imgt.org>) [37, 38]. The entire full length of TRBV is defined as a cluster containing all three complementarity determining regions (CDRs).

#### Cytotoxicity assay

Cytotoxicity of T cells was measured by the LDH assay kit (Promega, CytoTox 96®-G1780). Briefly, T cells and OC cells were mixed by the ratio of effect and target at 1:1, 5:1, and 10:1 in 1% FCS RPMI 1640 medium in a 96-well V-bottom for 4 h at 37 °C. The subsequent operations

were performed according to the manufacturer's procedure. The percentage of cytotoxicity is calculated with the following formula, % cytotoxicity = (experimental-effector spontaneous-target spontaneous)/(target maximum-target spontaneous) × 100.

### IL-2 and IFN $\gamma$ ELISA

Mice were vaccinated with whole-cell ID8 vaccines, and the serum was frozen at  $-80^{\circ}\text{C}$ . IL-2 and IFN $\gamma$  in serum were quantified using mouse IL-2 and IFN $\gamma$  ELISA plates (R&D VAL607 and VAL602). Plates were analyzed with an Infinite Tecan M200 PRO microplate reader (Switzerland).

### Mouse T cell subsets and DCs assay

Single-cell suspensions from tumor-draining lymph nodes of the vaccinated mice with whole-cell ID8 vaccines were prepared. FCM analysis of T cell subsets and DCs were performed with antibodies specific for CD3e (BD, 553061), CD4 (BD, 553051), CD8a (BD, 552877), and antibodies specific for Siglec E (R&D, FAB5806A), MGL1/2 (R&D, FAB4297P), CD11c (Biolegend, 117324), CD86 (Biolegend, 105116).

### Fluorescent microscopy

Mice ID8 tumor tissue sections with antigen repair were immunostained with primary antibodies against murine CD8 and CD39, and FITC and Cy3 fluorescein-conjugated secondary antibodies against primary antibodies for imaging with fluorescent microscopy and analysis of the intensity of staining with Image J (Fiji) software.

### Tumorigenicity studies

Animals were randomly grouped, and all animals were included in the data analysis except those who unexpectedly died. Female mice were inoculated subcutaneously (s.c.) once a week for three consecutive weeks at one side axilla with  $2 \times 10^5$  whole-cell ID8 vaccines prepared with ID8 tumor tissue as stated above or PBS before subcutaneous challenge with  $1 \times 10^6$  parental ID8 cells in 100  $\mu\text{L}$  PBS at the opposite axilla. Tumor volume at the opposite axilla was determined using calipers. Mice were sacrificed when tumor diameter exceeded 1 cm. The tumor growth and survival outcome were used to evaluate the efficiency of desialylated whole-cell ID8 vaccines in inducing specific anti-tumor immunity.

### Statistical analysis

All statistical analyses were performed using GraphPad Prism 8. Statistical analysis was carried out using a paired t-test, unpaired t-test, and the log-rank (Mantel-Cox) test. A P-value of less than 0.05 was considered statistically significant.

## Results

### Characterization of DWCTVs with Gal/GalNAc epitopes

There are four hydroxyl groups on Gal and GalNAc that may form glycosidic bonds (Fig. 1a). 1-hydroxyl of GalNAc is attached to the hydroxyl group of the protein serine (Ser) or threonine (Thr) residues for initiation of glycosylation, and 3- and 6-hydroxyls of GalNAc could be linked with sialic acids to terminate glycosylation. To uncover Gal/GalNAc epitopes on OC cells, we desialylated sialic acids on intact cells by hydrolysis with  $\alpha$ 2-3NA and  $\alpha$ 2-6NA following mitomycin C to generate DWCTVs (Fig. 1b; Additional file 1: Fig. S1, Additional file 2: Fig. S2).

FCM showed that there was a significant decrease of  $\alpha$ -2,3Sia and  $\alpha$ -2,6 Sia residues and an increase of Gal/GalNAc epitopes on the surface of DWCTVs compared to the corresponding controls ( $P < 0.0001$ ) (Fig. 1c–e; Additional file 3: Fig. S3).

### Maturation and activation of DCs induced by DWCTVs in vitro


To explore the impact of  $\alpha$ -2,3Sia and  $\alpha$ -2,6Sia shedding on DC performance, we co-cultured immature DCs (imDCs) with DWCTVs (Fig. 2a). As shown in Fig. 2b–d, DCs (CD14-imDCs) from monocytes of the healthy female were pulsed with DWCTVs (Additional file 4: Fig. S4) derived from A2780 but not OVCAR3 cells exhibited that the expression of differentiation marker CD83, co-stimulatory molecule CD86, and MGL ( $P < 0.05$ ) were remarkably increased, but Siglec-9 was decreased ( $P < 0.01$ ), as compared to the corresponding controls. Interestingly, the phenotypic alterations of DCs stimulated with  $\alpha$ 2-3NA modified DWCTVs were more obvious than those with  $\alpha$ 2-6NA modified ones, suggesting that the removal of  $\alpha$ -2,3-linked sialic acids on the terminal galactosyl of GalNAc may play an important role for activation and performance of DCs.

### DCs loaded with DWCTVs activate T cells in vitro

To test whether DCs loaded with DWCTVs could activate T cells, the CD4/CD8 ratio and *IFNG* expression of T cells were detected following exposure of PBLs to DCs loaded with DWCTVs (Fig. 3a). The results showed that CD4/CD8 ratio and *IFNG* expression in CD3+ T cells were markedly increased, as compared to the controls ( $P < 0.05$ ), and that DCs loaded with  $\alpha$ 2-3NA modified DWCTVs could prime activation of T cells more effectively than  $\alpha$ 2-6NA modified DWCTVs (Fig. 3b–e).

### DCs loaded with DWCTVs change TCR-V $\beta$ repertoire

To prove the immunogenicity of DCs loaded with DWCTVs, we examined whether these DCs modulate




the TCR-V $\beta$  repertoire diversity of  $\alpha\beta$  T cells. As shown in Table 1, *TRBV* gene sequencing identified no difference in *TRBV* sequences between the control T cells and T cells co-incubated with imDCs. The *TRBV* sequences of T cells co-incubated with DCs pulsed with primary whole-cell tumor vaccines derived from A2780 (9.52% of CDR1/CDR2 AA sequences) and OVCAR3 cells (85.71% of CDR1/CDR2/CDR3 AA sequences) were partially identical to that of the control T cells. Interestingly, there was a great diversity in *TRBV* sequences between T cells stimulated with DCs loaded with DWCTVs and the control T cells. The T cells stimulated with DCs loaded with DWCTV derived from A2780 and OVCAR3 cells fell into 12 and 6 distinct *TRBV* sequences, respectively (Table 1). These demonstrated that the immunogenicity of DCs conferred by DWCTVs could modulate the TCR-V $\beta$  repertoire diversity of  $\alpha\beta$  T cells.

### DWCTVs enhance DC-mediated cytotoxicity of T cells *in vitro*

To further prove the anti-tumor activity of  $\alpha\beta$  T cells with the diversified TCR-V $\beta$  repertoires by the DCs loaded with DWCTVs, we test the cytotoxicity of the activated  $\alpha\beta$  T cells on A2780 and OVCAR3 cells. The LDH assay showed that the killing efficiencies of these T cells significantly improved as compared to the corresponding controls ( $P < 0.05$ ) (Fig. 3f, g). Together, our findings showed that DCs pulsed with DWCTVs not only induced the activation of T cells with high-affinity tumor antigen-specific TCRs but also enhanced their killing activity against parent tumor cells.


### Activation of immune cells by DWCTVs *in vivo*

To verify whether DWCTVs can elicit activation of immune cells *in vivo*, female mice were vaccinated with the inactivated DWCTVs derived from ID8 cells to test



the plasma IFN $\gamma$  and IL-2, T cell subtypes, and MGL and Siglec E of DCs in draining lymph nodes (DLNs) (Fig. 4a). Our results showed that the plasma levels of IFN $\gamma$  and IL-2 reached the peak, and the CD4/CD8 ratio of T cells in DLNs increased from  $1.25 \pm 0.05$

to  $1.37 \pm 0.10$  ( $P < 0.05$ ) (Fig. 4d) around 14 days after the vaccination. Meanwhile, vaccination with  $\alpha 2-3NA$  modified ID8 DWCTV induced an increased expression of MGL, not Siglec E of CD11c+ /CD86+ DCs, while vaccination with  $\alpha 2-6NA$  modified ID8 DWCTV



induced an increased expression of both MGL and Siglec E; however, inoculation with primary ID8 WCTVs induced an increased expression of Siglec E but not MGL (Fig. 4e, f). These findings suggest that vaccination with DWCTVs rather than primary WCTV could evidently elicit immune activation in vivo.

### Vaccination with DWCTVs limits tumor growth and prolongs survival of mice bearing tumor

To determine whether ID8 DWCTVs elicit anti-tumor immunity in vivo, tumorigenicity of parental ID8 cells, tumor growth, and survival of mice bearing ID8 tumor were evaluated after vaccination with ID8 DWCTVs once a week for 3 weeks in challenging

**Table 1** The TCR-V $\beta$  repertoires of  $\alpha\beta$  T cells

Spec	V_gene	J_gene	D_gene	CDR1	CDR2	CDR3	Freq (%)
Control	TRBV30	TRBJ2-5	TRBD2	GTSNPN	SVGIG	AWRTGAGQY	100.0
imDC	TRBV30	TRBJ2-5	TRBD2	GTSNPN	SVGIG	AWRTGAGQY	100.0
Unmodified	TRBV30	TRBJ1-5	TRBD1	GTSNPN	SVGIG	AWSHGNQPQH	4.76
(A2780)	TRBV30	TRVJ2-2	TRBD2	GTSNPN	SVGIG	AWSNWRGGPGELF	4.76
	TRBV5-1	TRBJ2-7	TRBD1	SGHRS	YFSETQ	ASSLRLGDEQY	23.81
	TRBV6-6	TRBJ1-1	TRBD2	MNHNY	SVGAGI	ASSYEGGPGTEAF	14.29
	TRBV20-1	TRBJ1-4	TRBD1	DFQATT	SNEGSKA	SASGLLEELKF	52.38
$\alpha$ 2-3 NA	TRBV24-1	TRBJ1-2	-	KGHDR	SFDVKD	ATSYT	5.00
(A2780)	TRBV20-1	TRBJ2-3	TRBD1	SLARSH	SNEGSKA	SANTGSTDTQY	5.00
	TRBV7-2	TRBJ2-3	TRBD2	SGHTA	FQGNSA	ASSSRSLAGAEDTQY	5.00
	TRBV28	TRBJ2-7	TRBD1	MDHEN	SYDVKM	ASSHRYEQY	5.00
	TRBV10-1	TRBJ2-7	TRBD2	CVCW	SYGVQD	ASCSGSDPILRAV	10.00
	TRBV11-3	TRBJ2-5	TRBD2	SGHNT	YENEEA	ASSPQLAGVFKTQY	10.00
	TRBV28	TRBJ1-2	TRBD1	MDHEN	SYDVKM	ASTAGSNYGYT	10.00
$\alpha$ 2-6 NA	TRBV2	TRBJ1-5	TRBD2	SNHLY	FYNNEI	ASSAPLSNQPQH	50.00
(A2780)	TRBV29-1	TRBJ2-2	TRBD1	SQVTM	ANQGSEA	SAVLRGTGELF	14.29
	TRBV6-6	TRBJ2-1	TRBD2	MNHNY	SVGAGI	ASRKAEGIYNEQF	19.04
	TRBV20-1	TRBJ1-5	TRBD1	DFQATT	SNEGSKA	SARVQANQPQH	23.81
	TRBV7-8	TRBJ1-5	TRBD1	SGHVS	FQNEAQ	ASSAGWGPQH	42.86
Unmodified	TRBV27	TRBJ1-1	TRBD2	MNHEY	SMNVEV	ASSFQRGTEAF	14.29
(OVCAR3)	TRBV30	TRBJ2-5	TRBD2	GTSNPN	SVGIG	AWRTGAGQY	85.71
$\alpha$ 2-3 NA	TRBV6-2	TRBJ2-1	TRBD1	MNHEY	SVGEGT	ASSPTSPRTPEQF	50.00
(OVCAR3)	TRBV5-6	TRBJ2-5	TRBD2	SGHDT	YEEEE	ASAPGGMHEQY	50.00
$\alpha$ 2-6 NA	TRBV3-1	TRBJ2-6	TRBD1	LGHDT	YNNKEL	ASSLLGYSGANVLT	9.52
(OVCAR3)	TRBV21-1	TRBJ1-5	TRBD1	TISF*	SQNEEL	ASSKIRTVGAISPS	14.29
	TRBV29-1	TRBJ2-6	TRBD2	SQVTM	ANQGSEA	SAAASGANVLT	23.81
	TRBV2	TRBJ2-1	TRBD1	SNHLY	FYNNEI	ASSDQSQGQGEF	52.38

TCR  $\beta$ -chains of control T cells and T cells stimulated by imDC and NA modified/unmodified A2780/OVCAR3 cells antigen-loaded DCs were determined by TRBV gene sequencing. Each row represents one clone, and its frequency (Freq.) in its group is indicated to the right. \*Stop codon, Spec: specimen

mouse tumor models (Fig. 5a). Vaccinations with  $\alpha$ 2-3NA and  $\alpha$ 2-6NA modified ID8 DWCTVs significantly delayed tumor formation time ( $27.60 \pm 2.79$  days and  $9.00 \pm 3.16$  days, respectively) compared to the parental ID8 cells and primary ID8 WCTV controls ( $5.00 \pm 1.00$  days and  $6.20 \pm 1.48$  days, respectively, both  $P < 0.05$ ) (Fig. 5b; Additional file 5: Fig. S5), also clearly limited the tumor growth (Fig. 5c). Interestingly, vaccination with  $\alpha$ 2-3NA modified ID8 DWCTVs had more significant effects than  $\alpha$ 2-6NA modified ID8 DWCTV in postponing tumor progression (Fig. 5c). Furthermore, vaccination with  $\alpha$ 2-3NA, not  $\alpha$ 2-6NA modified ID8 DWCTV notably prolonged overall survival of mice bearing ID8 tumor (median OS 119 days,  $n = 5$ ) compared to the primary WCTV control (median OS 90 days,  $n = 5$ ) (Fig. 5d) ( $P < 0.05$ ), suggesting that  $\alpha$ 2-3NA modified DWCTV could prime the anti-tumor immunity against OC.


Moreover, CD8+ T cells and MGL not Siglec E of DCs from DLNs significantly increased in vaccinated

mice inoculated with parental OC cells for 7 days (Fig. 5e–g), suggesting that ID8 DWCTVs could be able to induce the proliferation and activation of memory CD8+ cells. Immunofluorescence histochemistry (IFHC) revealed that vaccination with  $\alpha$ 2-3NA modified ID8 DWCTV significantly increased effector CD8+ T cells and decreased CD8+ /CD39+ exhausted T cells in tumor-infiltrating lymphocytes (TILs) of ID8 tumor tissues compared to the corresponding controls (Fig. 5h–j), suggesting that DWCTVs could also contribute to the anti-tumor immune effect on the local tumor. These might indicate that DWCTVs could activate immune cells in the draining lymph node, and these immune cells circulate to tumor tissues and exert anti-tumor effects (Fig. 6).

## Discussion

Tumor-derived sialoglycans can target different aspects of the immune system to promote evasion responses [39]. Tumor cells mask their specific antigens by glycan






modification which is one of the molecular recognition patterns of immune evasion [34]. Moreover, the tumor-derived sialoglycans serve as potent immunomodulatory, forming an invisibility cloak shielding tumor cells from immune recognition [34, 39, 40]. The dense layer of sialoglycans on tumor cell surfaces avoids the normal occurrence of immunological synapses between cancer and immune cells [9, 41]. Such reduced recognition is believed to be enhanced by hypersialylation of tumor ligands for CLRs expressed by immune cells [42].

The aberrant sialylation forms complex glycoalyx lattices on the tumor cell surface, reducing tumor

immunogenicity [39, 43, 44]. However, the underlying Gal/GalNAc epitopes interact with MGL on DCs to trigger an immune response [12, 45]. Here, DWCTVs could expose tumor antigenic Gal/GalNAc epitopes and reduce sialic acids, conferring them with immunostimulatory potential. The 1-hydroxyl of GalNAc is attached to the hydroxyl group of the protein serine (Ser) or threonine (Thr) residues for initiation of glycosylation by *N*-acetyl-galactosaminyltransferase (ppGalNAc-T) [46], and 3- and 6-hydroxyls of GalNAc could be linked with sialic acids to terminate glycosylation by  $\alpha$ -2,3-sialyltransferase (ST3GAL II) and  $\alpha$ -2,6-sialyltransferase (ST6GAL I) [47].


(See figure on next page.)

**Fig. 5** Effects of DWCTVs on tumorigenicity studies. **a** The timeline of the in vivo tumor challenge experiment. **b** The tumor formation time after tumor challenge. **c** The tumor growth curve of tumor-bearing mice. **d** The survival curves of tumor-challenged mice. **e, f** Quantification of Siglec E and MGL expression on DCs located in DLNs at the 7th d after tumor injection. **g** CD4/CD8 ratio of T cells in DLNs at the 7th d after tumor inoculation. **h** Representative dual immunofluorescence images for TILs in tumor tissues after mice were sacrificed at 50th d. Red, CD8; Green, CD39; Scale bar 20  $\mu$ m. **i** The Mean gray value of CD8 pixel for the tumor section. **j** The Mean gray value of CD8 and CD39 double-positive pixel for the tumor section. The control mice represent normal mice without vaccination and tumor injection. Unmodified,  $\alpha$ 2-3NA and  $\alpha$ 2-6NA represent the vaccination with unmodified WCTV,  $\alpha$ 2-3NA modified DWCTV, and  $\alpha$ 2-6NA modified DWCTV, respectively. Each plot of **e-g** represents the mean of three technical replicates of a mouse. Each plot of **i, j** represents the average of 3 fields of view per tumor section. The statistical test of (**e-g, i, j**) used is unpaired t-tests, while it is paired t-test for (**c**). The survival curve of (**d**) is analyzed using the log-rank test. \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ ; \*\*\*\* $P < 0.0001$



Tn (GalNAcα1-O-Ser/Thr), T (Galβ1-3GalNAcα1-O-Ser/Thr), and STn (NeuAcα2-6GalNAcα1-O-Ser/Thr) antigens are widely expressed in several types of tumors [48]. Especially, the expression of STn on normal cells is

limited but is abundant on OC, which makes STn a relatively specific tumor-associated antigen for vaccination [48]. ST3GAL II transfers sialic acid to the 3-position of the distal GalNAc leading to tumor progression in OC



[49–51]. Recently, the immunogenicity of GalNAc has paved its application in the targeted delivery of drugs [52, 53]. Our study's approach for DWCTVs production was desialylation of OC cells to expose Gal/GalNAc epitopes, which might have great potential to confer specific immunity to OC patients.

The glycan structures of tumor-associated sialoglycans on the surface of tumor cells could affect the interaction with MGL expressed on immature monocyte-derived DCs to orchestrate distinct immune responses. Aberrantly sialylated structures decorating cell surfaces on cancer cells have been shown to induce immunosuppressive responses of MGL on DCs [54]. In contrast, MGL-expressing APCs can take up Tn-derived peptide structures for antigen presentation and induction of T cell responses to elicit anti-tumor immunity [17, 55, 56]. Sialic acids on the termini of neighboring oligosaccharides significantly limit the peptide region recognized by APCs. The 3- and 4-hydroxyl groups of Gal/GalNAc are essential for  $\text{Ca}^{2+}$  binding required for MGL binding. The 2-acetamido group (NHAc) promotes the MGL binding, while the sialylation at 3-hydroxyl of GalNAc might interfere with MGL binding [16, 56], and exposure of the 3-hydroxyl group of GalNAc with  $\alpha$ 2-3NA treatment could enhance the affinity of binding MGL. The significantly increased DWCTV-DC interactions could result from the removal of steric hindrance formed by sialic acids and/or generation of (specific) interaction sites

between uncapped galactosyl residues of GalNAc-Ser/Thr that could interact with MGL on the DCs [4].

Sialoglycans interact with Siglec-9 colocalized with the TCR-CD3 complex to inhibit TCR-mediated cell activation via recruitment of SHP-1 by ITIM phosphorylation to reduce ZAP 70 phosphorylation [57]. In addition, sialoglycans as alternative ligands of CD28 bind to CD80 on APCs, attenuating co-stimulatory signals of antigen-mediated activation of T cells [54]. Previous studies have found that aberrant sialic acid-Siglec interactions are associated with reduced anti-tumor immunity, which could be reversed by desialylation of tumor cells [29, 30]. Siglec-9 is an immunosuppressive sialic acid binding receptor and is involved in immunoregulation through ligation with glycoconjugates with terminal sialic acids on cancer cells. Studies showed that the expression of Siglec-9 was up-regulated in the development of monocytes into immature DCs and was decreased in mature DCs. Because Siglec-9 binds to both  $\alpha$ 2,3- and  $\alpha$ 2,6-sialic acid-linked glycoconjugates or sialoglycans, removal of terminal sialic acids from sialoglycans on the surface of cancer cells by  $\alpha$ 2-3 and  $\alpha$ 2-6NA or blocking of ligation with Siglec-9 by anti-Siglec-9 mAb could ameliorate maturation and activation of DCs [31, 58]. Furthermore, CD86, expressed on DCs, is an activating ligand of co-stimulatory receptor CD28 on T cells and is required for naïve T cells to differentiate into functional effector cells [59]. Thus, by removing sialic acid ligands on the surface

of tumor cells, CD28 is better able to engage CD80, accounting for increased co-stimulation of T cells [54]. As shown in our study, DWCTVs could expose tumor antigenic Gal/GalNAc epitopes masked by aberrant sialoglycans and facilitate Gal/GalNAc-MGL-mediated internalization and presentation of tumor antigens by mature DCs for activation of T cells.

The interaction of CD86<sup>high</sup> DCs with T cells could induce the changes in T cell phenotypes and up-regulation of *IFNG* expression by CD8+ cytotoxic T cells (CTLs) in our study, indicating that DCs loaded with DWCTVs achieve immunogenic property and is capable of initiating T cell activation and proliferation. It is known that the two-signal model of T cell activation needs both peptide-MHC-TCR interaction and co-stimulatory CD28-CD80/CD86 interaction. The mature DCs with higher CD80/CD86 could induce co-stimulatory signals of antigen-mediated activation of naïve T cells and a higher proportion of CD8+ CTLs [54].

As we know, conventional tumor vaccines target tumor-associated antigens (TAAs) overexpressed in cancers that are also expressed in normal tissues and can potentially induce central and peripheral tolerance responses, which results in low vaccination efficiency [60]. Moreover, the vaccines generated by intact tumor cells or tumor cell lysates also usually fail to elicit an effective immune response. Thus, DWCTV modified with  $\alpha$ 2-3NA could enhance the immunogenicity of DCs and provide co-stimulatory signals to activate T cells [61, 62]. CD8+ T cells are the critical effector cells that contribute to the anti-tumor immune response. They comprise various T-cell clones with diverse antigen-specific TCRs. Thus, elucidating the overall anti-tumor responses of diverse T-cell clones is an emerging challenge in tumor immunology. As we know, the lack of cellular immune response to tumor cells results from the poor presentation of antigens by DCs to CD8+ T cells and the inability of tumor vaccines to elicit sufficient DC activation to evoke the DC-derived co-stimulatory signals required to initiate effector CD8+ T-cell responses [63]. Fully effective tumor vaccines must elicit a diverse repertoire of CD8+ T-cell responses. The highly variable CDR3 of the TCR  $\beta$  chain is unique to individual T cell clones and can therefore be used to identify the T cell repertoire responses to the immunogenicity of DCs [64]. TCR-V $\beta$  repertoire diversity of  $\alpha\beta$  T cells in our study demonstrates that DWCTVs could confer the immunogenicity of DCs. TCR diversity within antigen-specific T cell repertoires is essential for effective immunity to eliminate tumor cells [64]. The TCR repertoire of T cells is diversified at the initial phase and proliferated into the "best-fit" clonotypes by continuing exposure once they recognize the tumor antigens presented by the DCs. The enhanced

killing activity here might indicate that the DCs could induce TCRs with optimal structural and affinity characteristics. Cancers with low TCR diversity are reported to be unable to recognize and eliminate tumor cells specifically [65]. Anti-tumor immune responses are highly individualized for the intrinsic differences in TCR repertoire contributed to the heterogeneity of anti-tumor immunity against the same tumors [66]. Together, our findings showed that DCs pulsed with DWCTVs induced the activation of T cells with high-affinity tumor antigen-specific TCRs and enhanced their killing activity against parent tumor cells.

Tumor vaccines are known to generate T<sub>H</sub>1-polarised CD4+ T cells to maintain and sustain anti-tumor CD8+ CTL responses. CD4+ T cells secrete IL-2 to directly activate CD8+ CTLs expressing the high-affinity IL-2 receptor and indirectly induce CD8+ CTL responses through the expression of co-stimulatory signals (B7 family ligands and IL-12) by up-regulating CD40 ligand to interact with CD40 on DCs. Tumor vaccines also boost CD4+ T cell-derived secretion of T<sub>H</sub>1-characteristic tumoricidal cytokines (e.g., IFN $\gamma$ ), which have direct anti-tumor activity. CD4+ T cell responses against tumors tend to be against self-derived epitopes [67]. Desialylation of glycan structures of tumor-associated antigens (self) and tumor-specific antigens (non-self) could facilitate MGL-mediated endocytosis and presentation by DCs [56]. The  $\alpha$ 2-3NA modified ID8 DWCTV with low sialic acid residues and high tumor antigenic Gal/GalNAc epitopes in our study induced Siglec E<sup>low</sup>/MGL<sup>high</sup> DCs and the increased CD4/CD8 ratio in DLNs and the increased expression of IFN $\gamma$  and IL-2 in peripheral plasma, indicating that vaccination with DWCTVs could activate both local and systemic immune responses.

Eliminating tumor cells by CD8+ T cells requires overcoming the immunosuppressive tumor environment. CD39 is a marker of exhausted tumor-infiltrating CD8+ T cells and co-expresses with multiple inhibitory receptors, such as PD-1, LAG-3, and Tim-3 [68, 69]. Siglec-9 is also reported to co-express with these inhibitory receptors [70] and potentially attenuate the anti-tumor function of NK cells and T cells by binding sialoglycans in *trans* on OC cells [71]. The anti-tumor effects exhibited in *vivo* might demonstrate that ID8 DWCTVs desialylated to expose tumor antigenic Gal/GalNAc epitopes via hydrolyzing tumor cells with NAs could increase the immunogenicity of tumors [34, 72] and induce T-cell clones with TCR-V $\beta$  repertoire diversity that overcome tumor-suppressive niches and kill tumor cells. Desialylation is a suitable treatment that contributes to the exposure of tumor-associated glycopeptide epitopes, which may be related to the immune recognition of tumor

antigens on the surface of tumor cells [4, 73]. Thus, vaccination with DWCTVs seems to be a favorable way to elicit an anti-tumor immune response. More importantly, T cells within the tumor microenvironment and peripheral are abolished and irreversibly dysfunctional in tumor development, including reduced T cell activation and attenuated antigen-presenting cell responses. T cell-targeted interventions do not rescue T cell dysfunction marked by CD8+CD39+ T cell clonal expansion, while tumor resection is sufficient to revert the systemic immune landscape [74, 75]. Therefore, immunization with tumor vaccines after tumor load reduction- a way to revive systemic immune function- is an ideal strategy to fight residual tumor cells and induce long-lasting anti-tumor immune responses to prevent disease recurrence [76].

## Conclusion

In summary, we report that DWCTVs with Gal/GalNAc epitopes deliver tumor antigens on the surface of tumor cells to the host immune system. Our findings highlight that DWCTVs could overcome sialoglycan-Siglec interaction-mediated immunosuppression and potentiate DC-mediated T cell anti-tumor immunity against OC *in vitro* and *in vivo*. Thus, vaccination with DWCTVs would become an attractive intervention in improving the survival of OC patients after cytoreductive surgery.

## Abbreviations

OC: Ovarian cancer; DC: Dendritic cell; Gal: Galactose; GalNAc: N-Acetylgalactosamine; Sia: Sialic acid; Siglec 9: Sialic acid-binding immunoglobulin-like lectins 9; MGL: Macrophage galactose type C lectin receptors; PD-1: Programmed cell death protein 1; Tim-3: T cell immunoglobulin and mucin-domain containing-3; LAG-3: Lymphocyte-activation gene 3; SHP 1: Src-homology domain 1; ZAP 70: Zeta-chain (TCR) associated protein kinase 70; TCR: T cell receptor; MHC: Major histocompatibility complex; FCM: Flow cytometry; IFN $\gamma$ : Interferon  $\gamma$ ; LDH: Lactate dehydrogenase; RT-qPCR: Real-time quantitative PCR; MAL II: Maackia Amurensis lectin II; PNA: Peanut agglutinin; SNA: Sambucus nigra lectin; PBMC: Peripheral blood mononuclear cell; IL 4: Interleukin 4; GM-CSF: Granulocyte-macrophage colony-stimulating factor; ELISA: Enzyme-linked immunosorbent assay; CD: Cluster of differentiation; AA: Amino acid; DLN: Draining lymph node; TIL: Tumor infiltrating lymphocyte; RNA: Ribonucleic acid.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12967-022-03714-y>.

**Additional file 1: Fig. S1.** Screening the mitomycin C concentration for OC cell treatment. Representative images under phase contrast microscopy of OC cells treated with 0, 50, and 100 ng ml<sup>-1</sup> mitomycin C for 2 h (a) and 3 h (b) and cultured for 24 h. Scale bars represent 100  $\mu$ m.

**Additional file 2: Fig. S2.** Concentration-dependent curves of NA modification on OC cells. The  $\alpha$ -2,3Sia (MAL II),  $\alpha$ -2,6Sia (SNA), and Gal/GalNAc (PNA) epitopes expression on A2780 (a), OVCAR3 (b), and ID8 (c) cells were detected by FCM respectively after  $\alpha$ 2-3/ $\alpha$ 2-6NA modification. Plots

represent three individual experiments; error bars are standard deviations (SD).

**Additional file 3: Fig. S3.** Representative confocal immunofluorescence images of OC cells. (a) and (b) showed the images of unmodified,  $\alpha$ 2-3NA modified, and  $\alpha$ 2-6NA modified ID8 cells stained by  $\alpha$ -2,3Sia (MAL II, green),  $\alpha$ -2,6Sia (SNA, green), and Gal/GalNAc (PNA, pink). Scale bars: 100  $\mu$ m.

**Additional file 4: Fig. S4.** DCs maturation/activation and subsequently lymphocyte activation. The PBMCs (a), immature DCs (b), co-incubation of DCs and tumor cells (c), and co-culture of mature DCs and lymphocytes (d) were observed under a phase-contrast microscope. Scale bars: 25  $\mu$ m.

**Additional file 5: Fig. S5.** Representative picture of HE staining for mouse ID8 OC. The section for tumors of nonvaccinated mice (a) and mice immunized with unmodified whole-cell ID8 vaccine (b),  $\alpha$ 2-3NA modified DWCTV (c), and  $\alpha$ 2-6NA modified DWCTV (d). Scale bars: 100  $\mu$ m.

## Acknowledgements

Not applicable.

## Author contributions

The work presented here was completed in collaboration with all authors. GZ, JH, and JH conceived and designed this study. JH performed all the experiments, and ML was responsible for TCR identification; BM and JL assisted with animal experimentation. YZ and QG carried out the data processing and statistical analysis. JH drafted this manuscript. JH and GZ revised and reviewed this manuscript. All authors read and approved this manuscript.

## Funding

This work was supported by the Sichuan Key Research and Development Project from the Department of Science and Technology of Sichuan Province (2019YFS0405, Meiyong Li; 2019YFS0036, Guonan Zhang).

## Availability of data and materials

All data generated or analysed during this study are included in this published article and its supplementary information files.

## Declarations

### Ethics approval and consent to participate

This study was approved by the Ethics Committee of Sichuan Cancer Hospital (SCCHEC-04-2019-004, Chengdu, China).

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup>School of Medicine, University of Electronic Science and Technology of China, Chengdu, China. <sup>2</sup>Biochemistry and Molecular Biology, Sichuan Cancer Institute, Chengdu, China. <sup>3</sup>Department of Ultrasound, Sichuan Cancer Hospital, School of Medicine, University of Electronic Science and Technology of China, Chengdu, China. <sup>4</sup>Department of Gynecologic Oncology, Sichuan Cancer Hospital, School of Medicine, University of Electronic Science and Technology of China, Chengdu, China.

Received: 30 July 2022 Accepted: 20 October 2022

Published online: 31 October 2022

## References

- Marth JD, Grewal PK. Mammalian glycosylation in immunity. *Nat Rev Immunol.* 2008;8(11):874–87.
- Baum LG, Cobb BA. The direct and indirect effects of glycans on immune function. *Glycobiology.* 2017;27(7):619–24.

3. Ghasempour S, Freeman SA. The glycocalyx and immune evasion in cancer. *FEBS J*. 2021.
4. Varki A, Gagneux P. Multifarious roles of sialic acids in immunity. *Ann N Y Acad Sci*. 2012;1253:16–36.
5. Pearce OMT, Läubli H. Sialic acids in cancer biology and immunity. *Glycobiology*. 2016;26(2):111–28.
6. Hakomori S. Tumor-associated carbohydrate antigens defining tumor malignancy: basis for development of anti-cancer vaccines. *Adv Exp Med Biol*. 2001;491:369–402.
7. van Gisbergen KP, Aarnoudse CA, Meijer GA, Geijtenbeek TBH, van Kooyk Y. Dendritic cells recognize tumor-specific glycosylation of carcinoembryonic antigen on colorectal cancer cells through dendritic cell-specific intercellular adhesion molecule-3-grabbing nonintegrin. *Cancer Res*. 2005;65(13):5935–44.
8. Munkley J, Scott E. Targeting aberrant sialylation to treat cancer. *Medicines (Basel)*. 2019;6(4):104.
9. Bull C, den Brok M, Adema G. Sweet escape: sialic acids in tumor immune evasion. *Biochem Biophys Acta*. 2014;1846(1):238–46.
10. Borzym-Kluczyk M, Radziejewska I, Cechowska-Pasko M. Increased expression of MUC1 and sialyl Lewis antigens in different areas of clear renal cell carcinoma. *Clin Exp Nephrol*. 2015;19(4):732–7.
11. Reis CA, Osorio H, Silva L, Gomes C, David L. Alterations in glycosylation as biomarkers for cancer detection. *J Clin Pathol*. 2010;63(4):322–9.
12. Higashi N, Fujioka K, Denda-Nagai K, Hashimoto S, Nagai S, Sato T, et al. The macrophage C-type lectin specific for galactose/*N*-acetylgalactosamine is an endocytic receptor expressed on monocyte-derived immature dendritic cells. *J Biol Chem*. 2002;277(23):20686–93.
13. Wu AM, Wu JH, Shen F. Interaction of a novel Tn (GalNAc alpha 1→Ser/Thr) glycoprotein with Gal, GalNAc and GlcNAc specific lectins. *Biochem Biophys Res Commun*. 1994;198(1):251–6.
14. van Vliet SJ, van Liempt E, Saeland E, Aarnoudse CA, Appelmek B, Irimura T, et al. Carbohydrate profiling reveals a distinctive role for the C-type lectin MGL in the recognition of helminth parasites and tumor antigens by dendritic cells. *Int Immunol*. 2005;17(5):661–9.
15. Wu AM. Carbohydrate structural units in glycoproteins and polysaccharides as important ligands for Gal and GalNAc reactive lectins. *J Biomed Sci*. 2003;10(6 Pt 2):676–88.
16. Marcelo F, Garcia-Martin F, Matsushita T, Sardinha J, Coelho H, Oude-Vrielink A, et al. Delineating binding modes of Gal/GalNAc and structural elements of the molecular recognition of tumor-associated mucin glycopeptides by the human macrophage galactose-type lectin. *Chemistry*. 2014;20(49):16147–55.
17. Napoletano C, Rughetti A, Agervig Tarp M, Coleman J, Bennett E, Picco G, et al. Tumor-associated Tn-MUC1 glycoform is internalized through the macrophage galactose-type C-type lectin and delivered to the HLA class I and II compartments in dendritic cells. *Cancer Res*. 2007;67(17):8358–67.
18. Napoletano C, Zizzari IG, Rughetti A, Rahimi H, Irimura T, Clausen H, et al. Targeting of macrophage galactose-type C-type lectin (MGL) induces DC signaling and activation. *Eur J Immunol*. 2012;42(4):936–45.
19. van Kooyk Y. C-type lectins on dendritic cells: key modulators for the induction of immune responses. *Biochem Soc Trans*. 2008;36(Pt 6):1478–81.
20. Jandus C, Boligan K, Chijioke O, Liu H, Dahlhaus M, Démoulin T, et al. Interactions between Siglec-7/9 receptors and ligands influence NK cell-dependent tumor immunosurveillance. *J Clin Invest*. 2014;124(4):1810–20.
21. Ibarlucea-Benitez I, Weitzenfeld P, Smith P, Ravetch JV. Siglecs-7/9 function as inhibitory immune checkpoints in vivo and can be targeted to enhance therapeutic antitumor immunity. *Proc Natl Acad Sci USA*. 2021;118(26):e2107424118.
22. Zheng Y, Ma X, Su D, Zhang Y, Yu L, Jiang F, et al. The roles of Siglec7 and Siglec9 on natural killer cells in virus infection and tumour progression. *J Immunol Res*. 2020;2020:6243819.
23. Beatson R, Tajadura-Ortega V, Achkova D, Picco G, Tsourouktsoglou TD, Klausung S, et al. The mucin MUC1 modulates the tumor immunological microenvironment through engagement of the lectin Siglec-9. *Nat Immunol*. 2016;17(11):1273–81.
24. Crocker PR, Paulson JC, Varki A. Siglecs and their roles in the immune system. *Nat Rev Immunol*. 2007;7(4):255–66.
25. Hudak JE, Canham SM, Bertozzi CR. Glycocalyx engineering reveals a Siglec-based mechanism for NK cell immune evasion. *Nat Chem Biol*. 2014;10(1):69–75.
26. Perdicchio M, Ilarregui J, Verstege M, Cornelissen L, Schetters S, Engels S, et al. Sialic acid-modified antigens impose tolerance via inhibition of T-cell proliferation and de novo induction of regulatory T cells. *Proc Natl Acad Sci USA*. 2016;113(12):3329–34.
27. Zhang J, Nicoll G, Jones C, Crocker P. Siglec-9, a novel sialic acid binding member of the immunoglobulin superfamily expressed broadly on human blood leukocytes. *J Biol Chem*. 2000;275(29):22121–6.
28. Läubli H, Varki A. Sialic acid-binding immunoglobulin-like lectins (Siglecs) detect self-associated molecular patterns to regulate immune responses. *Cell Mol Life Sci*. 2020;77(4):593–605.
29. Bärenwaldt A, Läubli H. The sialoglycan-Siglec glyco-immune checkpoint—a target for improving innate and adaptive anti-cancer immunity. *Expert Opin Ther Targets*. 2019;23(10):839.
30. Adams O, Stanczak M, von Gunten S, Laubli H. Targeting sialic acid-Siglec interactions to reverse immune suppression in cancer. *Glycobiology*. 2018;28(9):640–7.
31. Ohta M, Ishida A, Toda M, Akita K, Inoue M, Yamashita K, et al. Immunomodulation of monocyte-derived dendritic cells through ligation of tumor-produced mucins to Siglec-9. *Biochem Biophys Res Commun*. 2010;402(4):663–9.
32. de Charette M, Marabelle A, Houot R. Turning tumour cells into antigen presenting cells: the next step to improve cancer immunotherapy? *Eur J Cancer*. 2016;68:134–47.
33. Bull C, Stoel MA, den Brok MH, Adema GJ. Sialic acids sweeten a tumor's life. *Cancer Res*. 2014;74(12):3199–204.
34. Gray M, Stanczak M, Mantuano N, Xiao H, Pijnenborg J, Malaker S, et al. Targeted glycan degradation potentiates the anticancer immune response in vivo. *Nat Chem Biol*. 2020;16(12):1376–84.
35. Horrevorts SK, Stolk DA, van de Ven R, Hulst M, van Het Hof B, Duinkerken S, et al. Glycan-modified apoptotic melanoma-derived extracellular vesicles as antigen source for anti-tumor vaccination. *Cancers*. 2019;11(9):1266.
36. van Kooyk Y, Unger WW, Fehres CM, Kalay H, Garcia-Vallejo JJ. Glycan-based DC-SIGN targeting vaccines to enhance antigen cross-presentation. *Mol Immunol*. 2013;55(2):143–5.
37. Giudicelli V, Brochet X, Lefranc M. IMGT/V-QUEST: standardized analysis of the immunoglobulin (IG) and T cell receptor (TR) nucleotide sequences. *Cold Spring Harb Protoc*. 2011;2011(6):695–715.
38. Brochet X, Lefranc M, Giudicelli V. IMGT/V-QUEST: the highly customized and integrated system for IG and TR standardized V-J and V-D-J sequence analysis. *Nucleic Acids Res*. 2008;36(Web site issue):W503–8.
39. van de Wall S, Santegoets K, van Houtum E, Bull C, Adema G. Sialoglycans and Siglecs can shape the tumor immune microenvironment. *Trends Immunol*. 2020;41(4):274–85.
40. Sanford B. An alteration in tumor histocompatibility induced by neuraminidase. *Transplantation*. 1967;5(5):1273–9.
41. Xiao H, Woods E, Vukojicic P, Bertozzi C. Precision glycocalyx editing as a strategy for cancer immunotherapy. *Proc Natl Acad Sci USA*. 2016;113(37):10304–9.
42. Büll C, Collado-Camps E, Kers-Rebel E, Heise T, Søndergaard J, Brok M, et al. Metabolic sialic acid blockade lowers the activation threshold of mDCs for TLR stimulation. *Immunol Cell Biol*. 2016;95(4):408–15.
43. Wielgat P, Trofimiuk E, Czarnomysy R, Braszko JJ, Car H. Sialic acids as cellular markers of immunomodulatory action of dexamethasone on glioma cells of different immunogenicity. *Mol Cell Biochem*. 2019;455(1–2):147–57.
44. Varki A. Since there are PAMPs and DAMPs, there must be SAMPs? Glycan “self-associated molecular patterns” dampen innate immunity, but pathogens can mimic them. *Glycobiology*. 2011;21(9):1121–4.
45. Denda-Nagai K, Aida S, Saba K, Suzuki K, Moriyama S, Oo-Puthinan S, et al. Distribution and function of macrophage galactose-type C-type lectin 2 (MGL2/CD301b): efficient uptake and presentation of glycosylated antigens by dendritic cells. *J Biol Chem*. 2010;285(25):19193–204.
46. Coelho H, Rivas MDL, Grosso AS, Diniz A, Soares CO, Francisco RA, et al. Atomic and specificity details of mucin 1 O-glycosylation process by multiple polypeptide GalNAc-transferase isoforms unveiled by NMR and molecular modeling. *JACS Au*. 2022;2(3):631–45.

47. Teppa RE, Petit D, Plechakova O, Cogez V, Harduin-Lepers A. Phylogenetic-derived insights into the evolution of sialylation in eukaryotes: comprehensive analysis of vertebrate beta-galactoside alpha2,3/6-sialyltransferases (ST3Gal and ST6Gal). *Int J Mol Sci.* 2016;17(8).
48. Davidson B, Gottlieb WH, Ben-Baruch G, Kopolovic J, Goldberg I, Nesland JM, et al. Expression of carbohydrate antigens in advanced-stage ovarian carcinomas and their metastases—a clinicopathologic study. *Gynecol Oncol.* 2000;77(1):35–43.
49. Deschuyter M, Leger DY, Verboom A, Chaunavel A, Maftah A, Petit JM. ST3GAL2 knock-down decreases tumoral character of colorectal cancer cells in vitro and in vivo. *Am J Cancer Res.* 2022;12(1):280–302.
50. Aloia A, Petrova E, Tomiuk S, Bissels U, Deas O, Saini M, et al. The sialylglycolipid stage-specific embryonic antigen 4 marks a subpopulation of chemotherapy-resistant breast cancer cells with mesenchymal features. *Breast Cancer Res.* 2015;17(1):146.
51. Toivonen S, Aitio O, Renkonen O. Alpha 2,3-Sialylation of terminal GalNAc beta 1–3Gal determinants by ST3Gal II reveals the multifunctionality of the enzyme. The resulting Neu5Ac alpha 2–3GalNAc linkage is resistant to sialidases from Newcastle disease virus and *Streptococcus pneumoniae*. *J Biol Chem.* 2001;276(40):37141–8.
52. Cui H, Zhu X, Li S, Wang P, Fang J. Liver-targeted delivery of oligonucleotides with *N*-acetylgalactosamine conjugation. *ACS Omega.* 2021;6(25):16259–65.
53. Biessen EAL, Van Berkel TJC. *N*-acetyl galactosamine targeting: paving the way for clinical application of nucleotide medicines in cardiovascular diseases. *Arterioscler Thromb Vasc Biol.* 2021;41(12):2855–65.
54. Edgar LJ, Thompson AJ, Vartabedian VF, Kikuchi C, Woehl JL, Teijaro JR, et al. Sialic acid ligands of CD28 suppress costimulation of T cells. *ACS Cent Sci.* 2021;7(9):1508–15.
55. Singhal A, Fohn M, Hakomori S. Induction of alpha-*N*-acetylgalactosamine-O-serine/threonine (Tn) antigen-mediated cellular immune response for active immunotherapy in mice. *Cancer Res.* 1991;51(5):1406–11.
56. Gabba A, Bogucka A, Luz J, Diniz A, Coelho H, Corzana F, et al. Crystal structure of the carbohydrate recognition domain of the human macrophage galactose C-type lectin bound to GalNAc and the tumor-associated Tn antigen. *Biochemistry.* 2021;60(17):1327–36.
57. van Houtum EJH, Büll C, Cornelissen LAM, Adema GJ. Siglec signaling in the tumor microenvironment. *Front Immunol.* 2021;12: 790317.
58. Angata T, Varki A. Cloning, characterization, and phylogenetic analysis of siglec-9, a new member of the CD33-related group of siglecs. Evidence for co-evolution with sialic acid synthesis pathways. *J Biol Chem.* 2000;275(29):22127–35.
59. Chen L, Flies DB. Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nat Rev Immunol.* 2013;13(4):227–42.
60. Shindo Y, Hazama S, Nagano H. Cancer vaccine focused on neoantigens. *Gan to kagaku ryoho. Cancer Chemothera.* 2019;46(9):1367–71.
61. Keenan BP, Jaffee EM. Whole cell vaccines—past progress and future strategies. *Semin Oncol.* 2012;39(3):276–86.
62. Korbelik M. Optimization of whole tumor cell vaccines by interaction with phagocytic receptors. *Vaccines.* 2021;9(8).
63. Jhunjhunwala SA-O, Hammer CA-O, Delamarre LA-O. Antigen presentation in cancer: insights into tumour immunogenicity and immune evasion. *Nat Rev Cancer.* 2021;21(5):298–312.
64. Turner SJ, La Gruta NL, Kedzierska K, Thomas PG, Doherty PC. Functional implications of T cell receptor diversity. *Curr Opin Immunol.* 2009;21(3):286–90.
65. Aran A, Garrigos L, Curigliano G, Cortes J, Marti M. Evaluation of the TCR repertoire as a predictive and prognostic biomarker in cancer: diversity or clonality? *Cancers.* 2022;14(7).
66. Woolaver RA, Wang X, Krinsky AL, Waschke BC, Chen SMY, Popolizio V, et al. Differences in TCR repertoire and T cell activation underlie the divergent outcomes of antitumor immune responses in tumor-eradicating versus tumor-progressing hosts. *J Immunother Cancer.* 2021;9(1).
67. Tay RE, Richardson EK, Toh HC. Revisiting the role of CD4(+) T cells in cancer immunotherapy—new insights into old paradigms. *Cancer Gene Ther.* 2021;28(1–2):5–17.
68. Sade-Feldman M, Yizhak K, Bjorgaard SL, Ray JP, de Boer CG, Jenkins RW, et al. Defining T cell states associated with response to checkpoint immunotherapy in melanoma. *Cell.* 2018;175(4):998–1013.
69. Canale FP, Ramello MC, Nunez N, Araujo Furlan CL, Bossio SN, Gorosito Serran M, et al. CD39 Expression defines cell exhaustion in tumor-infiltrating CD8(+) T cells. *Cancer Res.* 2018;78(1):115–28.
70. Stanczak MA, Siddiqui SS, Trefny MP, Thommen DS, Boligan KF, von Gunten S, et al. Self-associated molecular patterns mediate cancer immune evasion by engaging Siglecs on T cells. *J Clin Investig.* 2018;128(11):4912–23.
71. Belisle JAHS, Jennifer GAA, et al. Identification of Siglec-9 as the receptor for MUC16 on human NK cells, B cells, and monocytes. *Mol Cancer.* 2010;9:1–14.
72. Zhou Z, Wang X, Jiang L, Li D, Qian R. Sialidase-conjugated, “NanoNiche” for efficient immune checkpoint blockade therapy. *ACS Appl Bio Mater.* 2021;4(7):5735–41.
73. Büll C, Boltje T, Balneger N, Weischer S, Wassink M, van Gemst J, et al. Sialic acid blockade suppresses tumor growth by enhancing T cell-mediated tumor immunity. *Can Res.* 2018;78(13):3574–88.
74. Allen B, Hiam K, Burnett C, Venida A, DeBarge R, TenVooren I, et al. Systemic dysfunction and plasticity of the immune macroenvironment in cancer models. *Nat Med.* 2020;26(7):1125–34.
75. Yost K, Satpathy A, Wells D, Qi Y, Wang C, Kageyama R, et al. Clonal replacement of tumor-specific T cells following PD-1 blockade. *Nat Med.* 2019;25(8):1251–9.
76. Maringer Y, Walz JS. Tumor vaccination-strategies and time points. *Internist (Berl).* 2021;62(9):991–7.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

