

Identification and Functional Characterization of the *Neisseria gonorrhoeae* *lbpB* Gene Product

GOUR D. BISWAS,^{1*} JAMES E. ANDERSON,¹ CHING-JU CHEN,¹ CYNTHIA NAU CORNELISSEN,²
AND P. FREDERICK SPARLING^{1,3}

Department of Medicine¹ and Department of Microbiology and Immunology,³ School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, and Department of Microbiology and Immunology, Virginia Commonwealth University, Richmond, Virginia 23298²

Received 10 July 1998/Returned for modification 18 August 1998/Accepted 20 October 1998

We cloned *lbpB*, encoding a predicted 80-kDa lipoprotein, upstream of *lbpA*. A nonpolar mutant (*LbpB*⁻ *LbpA*⁺) had normal lactoferrin (LF) binding and grew normally with LF as an iron source, whereas *LbpB*⁻ *LbpA*⁻ and *LbpB*⁺ *LbpA*⁻ strains had reduced binding of LF and did not grow with LF as an iron source. *LbpB* bound LF directly in an affinity purification, suggesting that *LbpB* might play a still-uncharacterized role in the LF iron utilization.

Most organisms use inorganic iron (Fe) as a cofactor in carrying out a variety of metabolic functions (34). In an oxidized environment, ferric iron is highly insoluble and relatively unavailable. Mammals solubilize and sequester extracellular Fe by binding to the glycoproteins transferrin (TF) in serum (34) and lactoferrin (LF) on mucosal surfaces (22). Since there is essentially no free Fe in humans, pathogens have evolved mechanisms to scavenge Fe from host proteins. Many bacteria synthesize and secrete siderophores, low-molecular-weight compounds which bind and transport Fe into the cell through specific siderophore receptors (16). Production of siderophores facilitates infection in several bacterial pathogens (26). *Neisseria gonorrhoeae* and several other gram-negative mucosal pathogens including *Neisseria meningitidis* do not produce siderophores (10, 23) but produce specific receptors for each of the glycoproteins TF and LF, as well as hemoglobin (5, 7, 10, 15, 21).

A receptor for LF, *LbpA*, was identified and the corresponding *lbpA* gene was cloned and characterized in *N. gonorrhoeae* (3) and *N. meningitidis* (29). Recently, a second LF receptor protein, *LbpB*, was identified in *N. meningitidis* (6, 20, 28). Here we report the cloning of *lbpB* from *N. gonorrhoeae* and characterization of the role of *LbpB* in gonococcal LF binding and Fe acquisition.

Cloning of *lbpB*. We previously cloned the entire *lbpA* gene on four overlapping clones (3). pUNCH127 (Fig. 1) included 1,246 bp of DNA upstream of the 5' end of the *lbpA* open reading frame (ORF). To test if the region upstream of *lbpA* was also involved in LF utilization, we inactivated it by inserting (31) an Ω cassette (32) into an *AvaI* site in pUNCH127, creating pUNCH130. Sequence analysis of pUNCH130 confirmed that the Ω fragment was inserted at the *AvaI* site. pUNCH130 DNA was introduced into wild-type gonococcal strain FA19 by transformation (4), resulting in a mutant strain, FA6839. The presence of the Ω fragment in the *AvaI* site in FA6839 was confirmed by Southern analysis with the use of an *lbpB*-specific oligonucleotide probe (positions 2451 to 2470 in the *lbpB* sequence in GenBank) or the 2-kb Ω fragment (32). The mutant FA6839 had a phenotype similar to that of *lbpA*

mutant FA6775 (3), including loss of ability to (i) bind LF in a solid-phase assay, (ii) take up Fe from LF, (iii) utilize LF as an Fe source, and (iv) express *LbpA*. These results suggest that insertion of Ω upstream of *lbpA* had a polar effect on expression of *LbpA*.

DNA upstream of *lbpA* was cloned in two chromosome walking steps involving the use of oligonucleotide primers L58 and L62 (Fig. 1). Southern hybridization analysis showed that L58 hybridized to a 2.3-kb *RsaI* fragment (data not shown). We cloned the 2.3-kb *RsaI* fragment into the *SmaI* site of pUNCH615 (32), creating pUNCH142. Hybridization analysis with probe L62 identified a 1.1-kb *Sau3A* fragment of FA19 DNA. We ligated the 1.1-kb *Sau3A* fragment into the *BamHI* site of pMCL210 (27) harboring *lacZ*, allowing rapid color screening for inserts. Screening by colony hybridization produced two clones, pUNCH192 and pUNCH193, that differed in size. The clone pUNCH192 in *Escherichia coli* produced colonies on X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside) and IPTG (isopropyl- β -D-thiogalactopyranoside) plates that had three phenotypes: slightly blue, white, and blue. Plasmid DNA from a colony belonging to each class was sequenced. Plasmid DNA from a slightly blue colony designated pUNCH192SB (Fig. 1) contained an 1,138-bp *Sau3AI* insert with a complete 5' end of an apparent 2,189-bp-long ORF (Fig. 1). In contrast, the corresponding plasmid isolated from a white colony was 328 bp shorter than pUNCH192SB. The missing DNA included 21 bp at the 5' end of the 2,189-bp ORF. Plasmid DNA from the blue colony contained primarily vector DNA. The clone pUNCH193 produced stable white colonies on similar media and was stable on subculture. Interestingly, pUNCH193 lacked 20 bp of DNA that included 5'-GTC GAA TCA ACG CCG ACC GC-3' at positions 31 to 51 from the ATG codon start site of the 2,189-bp ORF in pUNCH192SB. To verify that the DNA in pUNCH192SB represented wild-type FA19, we PCR amplified and directly sequenced FA19 DNA flanking the deleted 20-bp region. A comparison of the sequences of FA19 DNA in the amplified product and the corresponding DNA in pUNCH192SB showed complete identity, confirming that the clone pUNCH192SB contained wild-type FA19 DNA. Thus, the set of three overlapping clones, pUNCH127, pUNCH142, and pUNCH192SB, represents 2,493 bp of contiguous FA19 DNA upstream of *lbpA*.

Nucleotide sequence analysis. An examination of the sequences in clones pUNCH127, pUNCH142, and pUNCH

* Corresponding author. Mailing address: Department of Medicine, University of North Carolina at Chapel Hill, CB# 7030, Chapel Hill, NC 27599-7005. Phone: (919) 966-3661. Fax: (919) 966-6714. E-mail: gdbis@med.unc.edu.




FIG. 1. Physical map of the *N. gonorrhoeae* chromosome showing genetic organization of the *lbpBA* operon region. The top line represents the restriction map of the cloned FA19 chromosome fragment. The organization of *lbpBA* genes including the direction of transcription as indicated by arrows is shown in the next line. Plasmid pUNCH192SB is a 1.1-kb *Sau3AI* fragment. Plasmids pUNCH142 and pUNCH127 contain 2.3-kb *RsaI* and 2.5-kb *SspI* fragments, respectively. The solid inverted triangle denotes the position of the *aphA3* insertion in strain FA6965 (pUNCH142), and the open inverted triangle denotes the position of the Ω insertion in strain FA6839 (pUNCH127). L58 and L62 refer to the oligonucleotides used as probes in isolating the clones pUNCH142 and pUNCH192SB, respectively.

192SB revealed a 2,189-bp ORF. The last 4 bp (ATGA) of the 2,189-bp ORF partially overlapped with the ATG initiation codon of the *lbpA* ORF. We designated the 2,189-bp ORF located upstream of *lbpA* as *lbpB*. Further upstream, we found 170 bp that were 56% identical to the 3' end of an *E. coli era* gene that encodes a GTP binding protein (1). Immediately downstream of the *era* homolog there was a potential hairpin loop structure at positions 180 to 212. This structure has a ΔG (25°C) value of -20.8 kcal (33) and probably corresponds to a transcription terminator. In addition, a 10-bp inverted repeat at positions 184 to 208 and another 10-bp sequence at positions 1902 to 1911 matched gonococcal uptake sequence (12, 14).

Translation of the *lbpB* coding region predicted a protein of 728 amino acids with a molecular mass of 80 kDa and a pI value of 4.53. Analysis of the deduced N-terminal amino acid sequence of LbpB revealed an 18-amino-acid signal sequence with the lipoprotein modification consensus sequence LSAC at positions 16 to 19 (17). These data suggested that LbpB was a lipoprotein. Based on fluorographic analysis and palmitic acid labeling, Lewis et al. concluded that LbpB in *N. meningitidis* is a lipoprotein (20). LbpB was 31% identical to TbpB of the same gonococcal strain and 81% identical to the recently reported meningococcal LbpB (6, 28). Comparison of the predicted protein sequence of FA19 LbpB with that of meningococcal LbpB (6, 28) showed very similar features that differentiate each from the neisserial TbpB family. Both gonococcal and meningococcal LbpB proteins contained two stretches of negatively charged residues that were not present in TbpB family members: region one, residues 462 to 527, and region two, residues 689 to 708 (Fig. 2). These regions also exhibit considerable variation between meningococcal and gonococcal LbpB proteins. There was a deletion in region one in meningococcal LbpB relative to the gonococcal LbpB, and in region two, there was a deletion in gonococcal LbpB. Overall, these two charged domains were only about 50% identical, compared to 81% for the whole proteins. Each of the LbpB proteins contained an identical 21-amino-acid C terminus that was absent in all neisserial TbpB family members.

***lbpB* gene product.** To study the function of *lbpB*, we disrupted *lbpB* by inserting a nonpolar *aphA3* cassette (24) into an *HpaI* site in pUNCH142, creating pUNCH195 (Fig. 1; Table 1)

Region 1

```

NG 462 DEEDSGINNGEESDEEEEIAEESSEDEVSDEDDNGEDEDEIVEEEADEAEEI
NM 453 DEEDSDIDNGEESSEDEIGDEEBGTEDAAGDEGSEDEATEENEDGEEDEA

NG 512 EEEEEEEEEPEESPEE 527
NM 503 EEPEEESSAE..... 512

```

Region 2

```

NG 689 .....EDIENEVENEADVGEQLEPE 708
NM 675 EGTENKVEADVVDVVDVVDADADV.EQLKPE 705

```

FIG. 2. Lineup of negatively charged residues in region one and region two of the gonococcal (NG) and meningococcal (NM) LbpB proteins. The numbers indicate the position in the unprocessed protein of the first amino acid listed. Amino acids identical between the two proteins are denoted by vertical lines; colons indicate highly conservative substitutions, and dots indicate less conservative substitutions.

and exchanged this mutant allele into the gonococcal strain FA19 as described above, creating FA6965. Sequence analysis confirmed that the cassette was inserted in the correct orientation and translational frame into pUNCH195. Southern hybridization with the *lbpB*-specific probe L66 (positions 1156 to 1176) or the 0.9-kb *aphA3* fragment (24) confirmed the expected insertion in *lbpB* in FA6965. The *lbpB* gene product was identified with the use of polyclonal, affinity-purified antiserum raised against a synthetic 16-mer peptide, RTRDNGINLS-GNGSTN, from the predicted C terminus of LbpB. We used isogenic strains FA6815, FA6985, and FA6986 (Table 1) lacking TbpB and TbpA in order to avoid potential problems due to production of proteins similar in size and structure to LbpB and LbpA. A Western blot probed with LbpB antibodies reacted with a protein of about 95 kDa in the parent strain FA6815 (Fig. 3A, lane 1). A protein of the same size also was produced by FA6985 (*lbpA::mTn3 Cm*) (Fig. 3A, lane 3) but not FA6986 (*lbpB::aphA3*) (Fig. 3A, lane 4). The latter strain did produce the 103-kDa LbpA, as expected (Fig. 3B, lane 4). FA6839 (*lbpB:: Ω*) produced neither LbpA nor LbpB (Fig. 3, lanes 2). These data showed that the 95-kDa protein was the *lbpB* gene product and also provided experimental evidence that *lbpB* and *lbpA* were part of a single transcriptional unit.

We noticed that the nonpolar mutant FA6986 (*lbpB::aphA3*) expressed reduced amounts of LbpA (Fig. 3B, lane 4) compared to the parent strain FA6815 (*lbpB⁺ lbpA⁺*) (Fig. 3B, lane 1). We confirmed this by comparing the intensities of the LbpA protein bands in linear range of detection in Western blots. We estimated that the reduction was approximately five-fold from the level for the parent strain (data not shown). We concluded that the *aphA3* insertion in *lbpB* in FA6986 caused reduced production of the downstream *lbpA* gene product. The reason for reduced expression of LbpA in the *lbpB::aphA3* mutant is not understood. Insertion of the *aphA3* cassette in *lbpB* clearly did allow transcriptional read-through into *lbpA* but may have attenuated transcription downstream of the insertion.

LF binding. To determine whether LbpB participated in LF binding, we performed dot blot assays of LF binding to whole cells. Bound LF was detected with horseradish peroxidase-conjugated anti-human LF antibody (Accurate Chemical & Scientific Corp., Westbury, N.Y.) (3). The results demonstrated that the LbpB⁻ LbpA⁺ mutant FA6986 bound approximately as much LF as the LbpB⁺ LbpA⁺ parent strain FA6815 (data not shown). In contrast, no binding was detected

TABLE 1. Bacterial strains and plasmids

| Strain or plasmid | Relevant characteristics | Source or reference(s) |
|-------------------------------|--|------------------------|
| <i>N. gonorrhoeae</i> strains | | |
| FA19 | LF ⁺ TF ⁺ | 5, 23 |
| FA1090 | LF ⁻ TF ⁺ | 8 |
| FA6775 | LF ⁻ Cm ^r (<i>lbpA</i> ::mTn3 Cm) | 3 |
| FA6815 | TF ⁻ Str ^r Spc ^r (<i>tbpB</i> :: Ω) | 2 |
| FA6839 | LF ⁻ TF ⁺ Str ^r Spc ^r (<i>lbpB</i> :: Ω); transformant, pUNCH130→FA19 | This study |
| FA6965 | LF ⁺ Kan ^r (<i>lbpB</i> :: <i>aphA3</i>); transformant, pUNCH195→FA19 | This study |
| FA6985 | LF ⁻ (<i>lbpA</i> ::mTn3 Cm) TF ⁻ (<i>tbpB</i> :: Ω), Cm ^r Str ^r Spc ^r ; transformant, FA6775→FA6815 | This study |
| FA6986 | LF ⁺ (<i>lbpB</i> :: <i>aphA3</i>) TF ⁻ (<i>tbpB</i> :: Ω) Kan ^r Str ^r Spc ^r ; transformant, FA6965→FA6815 | This study |
| Plasmids | | |
| pHP45 Ω | Source for the Ω fragment (Str ^r Spc ^r) | 32 |
| pMCL210 | Cloning vector, low copy; <i>lacZ</i> α Cm ^r | 27 |
| pUC18K | Source for the <i>aphA3</i> cassette (Kan ^r) | 24 |
| pUNCH127 | Ap ^r Tet ^r ; 2.5-kb FA19 <i>lbpAB SspI</i> fragment cloned into <i>SspI</i> site in pBR322 | 3 |
| pUNCH128 | Ap ^r ; FA19 insert in pUNCH127 cloned into <i>BamHI</i> site in pUNCH615 | 3 |
| pUNCH130 | Str ^r Spc ^r ; Ω cassette inserted into <i>AvaI</i> site in pUNCH128 | This study |
| pUNCH142 | Ap ^r ; 2.3-kb FA19 <i>RsaI</i> fragment cloned into <i>BamHI</i> site in pUNCH615 | This study |
| pUNCH192 | Like pUNCH192SB but unstable | |
| pUNCH192SB | Cm ^r ; 1.1-kb FA19 <i>Sau3A</i> fragment cloned into <i>BamHI</i> site in pMCL210 | This study |
| pUNCH193 | Like pUNCH192SB but deleted | This study |
| pUNCH195 | Kan ^r ; <i>aphA3</i> cassette inserted into <i>HpaI</i> site in pUNCH142 | This study |
| pUNCH615 | Ap ^r ; pBluescript II KS(-) with <i>PstI</i> site replaced with <i>MluI</i> | 32 |

for cells lacking LbpA, regardless of whether LbpB was present. Therefore, the solid-phase binding assay failed to detect LF binding by gonococci expressing LbpB alone. LbpB-specific binding was not detected under slightly different conditions (6, 28) for the solid-phase assay (data not shown).

To better quantitate LF binding, we measured the amount of radiolabeled LF bound by living cells in an equilibrium-phase binding assay developed by Cornelissen and Sparling (9). The specific activity of iodinated LF was 2.9×10^5 cpm/ μ g of LF. Approximately 1×10^7 to 5×10^7 CFU of gonococcal cells grown in chelexed defined medium with no added Fe (CDM-0) to induce Fe stress (5) were mixed with 2 to 100 nM 125 I-LF in the presence of 1% bovine serum albumin and 5 μ M

unlabeled LF in individual wells of a Multi-Screen microtiter dish (0.45- μ m-pore-size filter; MAHV N45; Millipore, Cambridge, Mass.). After being allowed to bind for 20 min at room temperature, unbound LF was removed by filtration, followed by five washes with CDM-0. Filters were dried, punched out, and counted. Specific binding was the difference between total binding without cold LF and binding that occurred in the presence of at least a 50-fold-excess concentration of cold LF. All four strains studied exhibited specific, saturable LF binding (Fig. 4). The wild-type strain bound significantly more LF than did each of the other strains. An apparent trend toward an intermediate level of LF binding in FA6965 (LbpB⁻ LbpA⁺) was noted, but the differences were not statistically significant by the *t* test in conjunction with the Bonferroni procedure (19).




FIG. 3. Expression of LbpB and LbpA in *N. gonorrhoeae*. Shown are Western blots containing total membranes prepared from Fe-starved cells probed with affinity-purified LbpB polyclonal antibody (A) and monoclonal LbpA antibody (B) (3). Lanes: 1, parent strain FA6815 (LbpB⁺ LbpA⁺); 2, FA6839 (LbpB⁻ LbpA⁻); 3, FA6985 (LbpB⁺ LbpA⁻); 4, FA6986 (LbpB⁻ LbpA⁺).




FIG. 4. Isotherms for binding of LF to *lbp* mutants. The curves represent the amount of specifically bound LF as a function of LF concentration. Each point represents the mean of four individual experiments. Symbols: \blacklozenge , FA19 (LbpB⁺ LbpA⁺); \blacksquare , FA6965 (LbpB⁻ LbpA⁺); \circ , FA6775 (LbpB⁺ LbpA⁻); \blacktriangle , FA6839 (LbpB⁻ LbpA⁻).

TABLE 2. K_d and copy number estimates for LF receptors expressed by isogenic mutants

| Strain | Phenotype | K_d (nM) | Copy number (M) |
|--------|-------------------------------------|--------------|----------------------------|
| FA19 | LbpA ⁺ LbpB ⁺ | 4.8 (site 1) | 9.3×10^{-12} (N1) |
| | | 550 (site 2) | 6.9×10^{-11} (N2) |
| FA6965 | LbpA ⁺ LbpB ⁻ | 5.0 (site 1) | 2.3×10^{-12} (N1) |
| | | 95 (site 2) | 2.0×10^{-11} (N2) |
| FA6775 | LbpA ⁻ LbpB ⁺ | 45 (site 1) | 1×10^{-11} (N1) |
| FA6839 | LbpA ⁻ LbpB ⁻ | 45 (site 1) | 1×10^{-11} (N1) |

The data obtained in the liquid-phase LF-binding experiments were analyzed with the computer program Receptor Fit Saturation Two-Site (Lundon Software, Inc., Cleveland Heights, Ohio). This program generates K_d and copy number estimates based on the best fit of observed data to progressively more complex models. Table 2 contains the best estimates of K_d and copy numbers generated by this program. Considerable variability was observed for LF binding to all strains; thus, these estimates are best considered comparisons of LF binding to isogenic strains rather than absolute indicators of intrinsic receptor affinity or copy number. Isogenic strains expressing LbpA exhibited complex binding phenomena consistent with the presence of at least two populations of binding sites, with K_d s ranging from 5 to 500 nM. In contrast, the isogenic mutants that did not express LbpA bound LF to a more homogeneous population of receptors, with a single apparent K_d of approximately 45 nM. This was true of both LbpB⁺ and LbpB⁻ strains, suggesting that LbpB played no detectable role in LF binding under the conditions employed in this assay. Somewhat surprisingly, the mutant that did not express either of the two identified LF-binding proteins (LbpB and LbpA) retained the ability to bind LF. This observation suggested that there were uncharacterized LF-binding components on the gonococcal cell surface.

LF affinity isolation of LbpB. To determine whether LbpB could bind LF in the absence of LbpA, total membranes prepared from Fe-starved FA6815 (LbpB⁺ LbpA⁺) and its isogenic derivatives, FA6985 (LbpB⁺ LbpA⁻) and FA6986 (LbpB⁻ LbpA⁺), were subjected to LF-agarose affinity purification (7, 11). Total membrane proteins were dissolved in 2% Zwittergent 3, 14 (Calbiochem) and rocked with a 50% slurry of LF-agarose overnight at 4°C. The unbound material was removed by centrifugation, and the LF-binding protein was eluted in a Laemmli sample buffer for analytical sodium dodecyl sulfate-polyacrylamide gel electrophoresis and Western blotting. Results (Fig. 5) showed an excellent recovery of LbpB in the absence of LbpA, indicating that LbpB was capable of binding LF under these conditions. LbpA also could be isolated by this procedure from total membranes in the absence of LbpB. The amount of LbpA in membrane preparations from the two strains was roughly the same (lanes A and C), but less LbpA was purified from the LbpB⁻ LbpA⁺ strain than from wild type (lanes D and F). This suggested that LbpA alone did not bind to LF as avidly as it did in the presence of LbpB.

LF utilization. To assess the contribution of LbpB in utilization of Fe from LF, we compared growth in CDM supplemented with LF of strains that carried either intact *lbpB* or inactivated *lbpB*. On agar plates or in CDM broth culture (5), the nonpolar *lbpB* mutant FA6965 (LbpB⁻ LbpA⁺) failed to show a growth defect, growing to a level equal to that of wild-type FA19 (data not shown).

⁵⁵Fe-LF uptake. To detect smaller effects of LbpB in LF utilization, we measured the amount of ⁵⁵Fe-LF taken up by




FIG. 5. Western blot of LF-agarose affinity-purified LbpB and LbpA from total membranes of Fe-starved FA6815 (LbpB⁺ LbpA⁺) (lanes A and D), FA6985 (LbpB⁺ LbpA⁻) (lanes B and E), and FA6986 (LbpB⁻ LbpA⁺) (lanes C and F). The blot was probed with antibodies to LbpA and then LbpB. The anti-LbpB antibody used here was raised against an LbpB His-tagged fusion protein. The 66-kDa band present in lanes A and B was recognized by antiserum elicited against full-length recombinant LbpB but not against an LbpB-specific peptide (Fig. 3 and data not shown). This protein band probably represents a breakdown product of holo-LbpB. Lanes A to C, total membranes (TM); lanes D to F, LF-agarose affinity-purified proteins. MW, molecular weight in thousands.

the use of established procedures (2, 5). Results indicated that the nonpolar *lbpB::aphA3* mutant FA6965 (LbpB⁻ LbpA⁺) acquired ⁵⁵Fe from LF at about 60% of wild-type levels ($P < 0.03$) (data not shown). In contrast, LbpA⁻ mutants, FA6775 and FA6839, failed to take up a significant amount of ⁵⁵Fe from LF ($P < 0.01$). We could not determine whether reduced Fe uptake from LF in FA6965 was due to loss of LbpB or to decreased expression of LbpA.

Conclusions. We cloned an *N. gonorrhoeae lbpB* gene, providing genetic evidence for the existence of a second LF receptor protein in the gonococcus, similar to that described recently for meningococci (6, 20, 28). Despite difficulties encountered in isolating the 5' end of the gonococcal *lbpB* gene, we succeeded in isolating a clone (pUNCH192SB) with an intact 5' end of *lbpB*. Attempts to clone the 5' end of the *lbpB* gene in *N. meningitidis* were unsuccessful (6, 20, 28).

A two-gene *lbpB lbpA* transcriptional unit was implicated by the observations that the 3' end of *lbpB* overlapped the 5' end of *lbpA*, and creation of a polar mutation in *lbpB* resulted in simultaneous loss of expression of the *lbpB* and *lbpA* genes. Recent experiments by Lewis et al. (20) showed that the *lbpB* and *lbpA* genes in meningococci are organized in a single transcriptional unit.

Both the gonococcal and meningococcal LbpB proteins contain two strongly acidic domains, notably absent in TbpB proteins, which are rich in aspartic acid and glutamic acid. This suggests that these regions could be involved in binding of LF, which is highly cationic (34). These regions are quite dissimilar in primary sequence in the known examples in gonococci and meningococci, suggesting that these also could be antigenic domains recognized by the host immune response. Future experiments hopefully will answer whether these domains are involved in binding LF and are immunogenic.

The role of LbpB in mediating Fe acquisition from LF is unclear. The LbpB⁻ mutant (LbpB⁻ LbpA⁺) retained the ability to bind LF, and this was reflected in its capacity to utilize LF as an Fe source. We demonstrated that LbpB, in the

absence of LbpA, could be isolated by LF-agarose affinity, which is the best evidence that LbpB is an LF-binding protein.

We found that the LbpB⁻ LbpA⁺ mutant used LF as an Fe source approximately as well as did the parent, as evidenced by similar growth patterns. The growth patterns of similar meningococcal LbpB⁻ LbpA⁺ mutants varied, but all mutants grew to some extent (28). Differences in growth exhibited by strains in these reports might be explained on the basis of varying levels of LbpA expression in the LbpB⁻ LbpA⁺ strain. For example, negligible growth noted by Lewis et al. might have resulted from a more than 10-fold reduction in LbpA expression (20). Near-normal growth exhibited by the mutant used by Pettersson et al. was correlated with LbpA expression similar to that of LbpB (28). The general conclusion is that LbpA is essential for LF-Fe utilization, whereas LbpB is not essential.

The question of how important LF is to pathogenesis remains. Many pathogens and nonpathogens can survive on the mucosal surface without being able to bind LF or to use LF as a source of Fe. For example, all *Haemophilus influenzae* strains are LF⁻ (18, 30), and about one-half of gonococci are LF⁻ (3, 25). Indeed, gonococcal strain FA1090 is LF⁻ due to a large *lbpBA* deletion (references 13 and 20 and data not shown), yet it causes urethritis readily in male human volunteers (8). Future work is required to clarify the role of LF in bacterial pathogenesis and the roles of LbpB, in particular in utilization of LF.

Nucleotide sequence accession number. The GenBank accession number for the 2,189-bp ORF is AF072890.

This work was supported by Public Health Service grants AI31496 and AI26837 from the National Institute of Allergy and Infectious Diseases.

We are grateful to Chris Elkins for his assistance with preparation of antipeptide sera and affinity isolation procedures. We also thank Chris Thomas for his expert computer assistance. We acknowledge the generosity of P. K. Sen for advice on statistical analysis and Rishikesh Chakravorty for analyzing the LF binding data. We are thankful to Bill Shafer for providing plasmid pUC18K and the UNC-CH Automated DNA Sequencing Facility for DNA sequencing.

REFERENCES

- Ahnn, J., P. E. March, H. E. Takiff, and M. Inouye. 1986. A GTP binding protein of *Escherichia coli* has homology to yeast RAS proteins. *Proc. Natl. Acad. Sci. USA* **83**:8849–8853.
- Anderson, J. E., P. F. Sparling, and C. N. Cornelissen. 1994. Gonococcal transferrin-binding protein 2 facilitates but is not essential for transferrin utilization. *J. Bacteriol.* **176**:3162–3170.
- Biswas, G. D., and P. F. Sparling. 1995. Characterization of *lbpA*, the structural gene for a lactoferrin receptor in *Neisseria gonorrhoeae*. *Infect. Immun.* **63**:2958–2967.
- Biswas, G. D., K. L. Burnstein, and P. F. Sparling. 1986. Linearization of donor DNA during plasmid transformation in *Neisseria gonorrhoeae*. *J. Bacteriol.* **168**:756–761.
- Blanton, K. J., G. D. Biswas, J. Tsai, J. Adams, D. W. Dyer, S. M. Davis, G. G. Koch, P. K. Sen, and P. F. Sparling. 1990. Genetic evidence that *Neisseria gonorrhoeae* produces specific receptors for transferrin and lactoferrin. *J. Bacteriol.* **172**:5225–5235.
- Bonnah, R. A., and A. B. Schryvers. 1998. Preparation and characterization of *Neisseria meningitidis* mutants deficient in production of the human lactoferrin binding proteins LbpA and LbpB. *J. Bacteriol.* **180**:3080–3090.
- Chen, C.-J., P. F. Sparling, L. A. Lewis, D. W. Dyer, and C. Elkins. 1996. Identification and purification of a hemoglobin-binding outer membrane protein from *Neisseria gonorrhoeae*. *Infect. Immun.* **64**:5008–5014.
- Cornelissen, C. N., M. Kelley, M. M. Hobbs, J. E. Anderson, and P. F. Sparling. 1998. The transferrin receptor expressed by gonococcal strain FA1090 is required for the experimental infection of human volunteers. *Mol. Microbiol.* **27**:611–616.
- Cornelissen, C. N., and P. F. Sparling. 1996. Binding and surface-exposure characteristics of the gonococcal transferrin receptor are dependent on both transferrin-binding proteins. *J. Bacteriol.* **178**:1437–1444.
- Cornelissen, C. N., and P. F. Sparling. 1994. Iron piracy: acquisition of transferrin-bound iron by bacterial pathogens. *Mol. Microbiol.* **14**:843–850.
- Elkins, C. Unpublished observations.
- Elkins, C., C. E. Thomas, H. S. Seifert, and P. F. Sparling. 1991. Species-specific uptake of DNA by gonococci is mediated by a 10-base-pair sequence. *J. Bacteriol.* **173**:3911–3913.
- Gonococcal Genome Sequence Database. <http://dna1.chem.uoknor.edu/>.
- Goodman, S. D., and J. J. Scocca. 1988. Identification and arrangement of the DNA sequence recognized in specific transformation of *Neisseria gonorrhoeae*. *Proc. Natl. Acad. Sci. USA* **85**:6982–6986.
- Gray-Owen, S., and A. B. Schryvers. 1996. Bacterial transferrin and lactoferrin receptors. *Trends Microbiol.* **5**:185–191.
- Guerinot, M. L. 1994. Microbial iron transport. *Annu. Rev. Microbiol.* **48**:743–772.
- Hayashi, S., and H. C. Wu. 1990. Lipoproteins in bacteria. *J. Bioenerg. Biomembr.* **22**:451–471.
- Herrington, D. A., and P. F. Sparling. 1985. *Haemophilus influenzae* can use human transferrin as a sole source for required iron. *Infect. Immun.* **48**:248–251.
- Hochberg, Y., and A. Tamhane. 1987. Multiple comparison procedures, p. 30 and 363. John Wiley & Sons, Inc., New York, N.Y.
- Lewis, L. A., K. H. Rhode, B. Behrens, E. Gray, S. I. Toth, B. A. Roe, and D. Dyer. 1998. Identification and molecular analysis of *lbpBA*, which encodes the two-component meningococcal lactoferrin receptor. *Infect. Immun.* **66**:3017–3023.
- Lewis, L. A., E. Gray, Y.-P. Wang, B. A. Roe, and D. Dyer. 1997. Molecular characterization of *hpuAB*, the haemoglobin-heptaglobin-utilization operon of *Neisseria meningitidis*. *Mol. Microbiol.* **23**:737–749.
- Masson, P. L., J. F. Heremans, and C. H. Dive. 1966. An iron-binding protein common to many external secretions. *Clin. Chim. Acta* **14**:735–739.
- McKenna, W. R., P. A. Mickelsen, P. F. Sparling, and D. W. Dyer. 1988. Iron uptake from lactoferrin and transferrin by *Neisseria gonorrhoeae*. *Infect. Immun.* **56**:785–791.
- Menard, R., P. J. Sansonetti, and C. Parsot. 1993. Nonpolar mutagenesis of the *ipa* genes defines IpaB, IpaC, and IpaD as effectors of *Shigella flexneri* entry into epithelial cells. *J. Bacteriol.* **175**:5899–5906.
- Mickelsen, P. A., E. Blackman, and P. F. Sparling. 1982. Ability of *Neisseria gonorrhoeae*, *Neisseria meningitidis*, and commensal *Neisseria* species to obtain iron from lactoferrin. *Infect. Immun.* **35**:915–920.
- Mietzner, T. A., and S. A. Morse. 1994. The role of iron binding proteins in the survival of pathogenic bacteria. *Annu. Rev. Nutr.* **14**:471–493.
- Nakano, Y., Y. Yoshida, Y. Yamashita, and T. Koga. 1995. Construction of a series of pACYC-derived plasmid vectors. *Gene* **162**:157–158.
- Pettersson, A., T. Prinz, A. Umar, J. Biezen, and J. Tommassen. 1998. Molecular characterization of *lbpB*, the second lactoferrin-binding protein of *Neisseria meningitidis*. *Mol. Microbiol.* **27**:599–610.
- Pettersson, A., A. Maas, and J. Tommassen. 1994. Identification of the *iroA* gene product of *Neisseria meningitidis* as a lactoferrin receptor. *J. Bacteriol.* **176**:1764–1766.
- Pidcock, K. A., J. A. Wooten, B. A. Daley, and T. L. Stull. 1988. Iron acquisition by *Haemophilus influenzae*. *Infect. Immun.* **56**:721–725.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Thomas, C. E., and P. F. Sparling. 1994. Identification and cloning of a *fur* homologue from *Neisseria meningitidis*. *Mol. Microbiol.* **11**:725–737.
- Tinoco, I., Jr., P. N. Borer, B. Dengler, M. D. Levine, O. C. Uhlenbeck, D. M. Crothers, and J. Gralla. 1973. Improved estimation of secondary structure in ribonucleic acids. *Nature (London) New Biol.* **246**:40–41.
- Welch, S. 1992. The chemistry and biology of iron, p. 41–59. *In* S. Welch (ed.), *Transferrin: the iron carrier*. CRC Press, Inc., Boca Raton, Fla.