



Published in final edited form as:

Brain Behav Immun. 2018 October ; 73: 1–2. doi:10.1016/j.bbi.2018.06.008.

Interleukin (IL)-6: a good kid hanging out with bad friends (and why sauna is good for health)

Charles L. Raison^{1,2,3}, Jennifer M. Knight⁴, Carmine Pariante⁵

¹School of Human Ecology, University of Wisconsin-Madison, Madison, WI, USA

²School of Medicine and Public Health, University of Wisconsin-Madison, Madison, WI, USA

³Usona Institute, Fitchburg, WI, USA

⁴Departments of Psychiatry, Medicine, and Microbiology & immunology; Medical College of Wisconsin, Milwaukee, WI, USA

⁵Stress, Psychiatry and Immunology Lab, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

As highlighted in a recent review by Del Giudice & Gangestad in *Brain, Behavior & Immunity* (Del Giudice and Gangestad, 2018), increasing data suggest that the cytokine interleukin (IL)-6, universally maligned for its associations with bad mental and physical health, may be like the proverbial good kid who lands in jail merely because he was in the backseat of the car when his bad (cytokine) friends decided to commit a crime. If IL-6, like the good kid with bad buddies, is mostly guilty by association, is a major revision in in order on how we understand the relationship between IL-6 and disease risk and persistence?

Guilty as Charged?

If a kid is in the backseat for the robbery of 10 convenience stores, one has to suspect that he is more than a passive bystander. By this criterion, things don't look good for IL-6, because the cytokine is on the crime scene for all manner of illness.

In psychiatry, every disease has been associated with increased IL-6 levels, from major depressive disorder (MDD) and bipolar disorder, to schizophrenia, panic disorder and posttraumatic stress disorder (PTSD). Beyond mental health, increased IL-6 predicts the future development of cognitive decline, diabetes mellitus, and cardiovascular events (e.g., myocardial infarction). And for those already burdened with a medical illness, increased circulating IL-6 is a bellwether for death, whether the medical condition be dialysis-dependent renal failure, stroke, cardiovascular disease or cancer.

Corresponding Author: Charles L. Raison, MD, 1300 Linden Drive, School of Human Ecology, University of Wisconsin-Madison, Madison, WI 53706, raison@wisc.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

One might try to procure a “get out of jail free” card for IL-6 by invoking the old saw about association not proving causality, were it not for the fact that the pharmacological blockade of IL-6 is successfully used for the treatment of several immune-related diseases in which the cytokine has been implicated as a causal mechanism. Moreover, in patients with two of these diseases (rheumatoid arthritis and Castleman’s disease) treatment with an IL-6 antagonist has been reported to produce an antidepressant response that is over and above improvement in the underlying autoimmune condition itself (Sun et al., 2017).

Is there hope for the kid?

Yet, the story is more complicated than this.

Because of its bifurcated signaling system, IL-6 not only has disease-promoting inflammatory effects, but also provides negative feedback on the release of other proinflammatory cytokines (such as Tumor Necrosis Factor (TNF)-alpha and IL-1-beta) and induces the release of IL-10, the body’s primary anti-inflammatory cytokine. Specifically, when IL-6 docks with a soluble form of its receptor, this complex can interact directly with membrane-bound glycoprotein 130, a protein found on every cell of the body. This process, known as “trans-signaling”, accounts for the inflammatory actions of IL-6. On the other hand, when IL-6 interacts with its membrane-bound receptor, which only exists on a small handful of cell types (hepatocytes, monocytes/macrophages and some lymphocytes), it produces primarily antiinflammatory and regenerative effects, via what has been termed the “classical signaling pathway”.

What happen if you block both? Work from our group has yielded the startling finding that treating patients about to undergo allogeneic hematopoietic cell transplantation with tocilizumab, an IL-6 receptor antagonist that inhibits both trans and classical signaling, increases—rather than decreases—depression and anxiety a month later (Knight et al., 2018). And in a neurotoxic cytokine storm syndrome traditionally blamed on both IL-1 and IL-6, blocking classic and trans IL-6 signaling provides no protection, whereas blocking IL-1 abrogates the syndrome and associated neurotoxicity (Norelli et al., 2018). So, perhaps IL-6 is mostly good, at least in some circumstances? If so, what happens when it shows up alone, without the bad friends around?

Perhaps not such a bad kid after all?

Acute exercise stimulates muscle cells to pump out IL-6 in dizzying amounts, up to 100 times normal resting levels (Pedersen and Febbraio, 2008) – *without* activating TNF or IL-1-beta. And, activated in this way, IL-6 induces a host of biological changes that are generally considered anti-inflammatory, including the induction of the potent anti-inflammatory molecules, IL-10 and IL-1 receptor antagonist (IL-1ra).

It is not only physical exercise.

Hyperthermia, when administered to patients with depression, produces a large increase in circulating IL-6 – again, *without* activating TNF or IL-1-beta (Raison, 2017). And how do people have historically induced hyperthermia? Yes, sauna: recently shown to produce

a range of health benefits, including reduced cardiovascular death, dementia and all-cause mortality (Heinonen and Laukkanen, 2018).

Fasting has significant anti-inflammatory effects in humans and has been shown to induce IL-6 production, which is essential for the health-enhancing metabolic effects of caloric restriction (Wueest et al., 2014). Sleep deprivation and sunlight exposure also acutely increase IL-6.

What do all these interventions (physical exercise, hyperthermia, fasting, sleep deprivation and sunlight exposure) have in common? They have antidepressant or mood-elevating properties. In fact, ketamine, which produces a rapid and profound antidepressant effect, also acutely increases circulating levels of IL-6 (Park et al., 2016). Not bad for a bad kid!

But even these results should be bracketed with caution, given data that inflammation itself, which lowers mood when acute or chronic in normal volunteers (Raison et al., 2005), may actually have antidepressant properties when administered as an acute stimulus in the context of depression or chronic stress (Bauer et al., 1995)

Conclusions

Despite the title, our goal in this Viewpoint has not been to establish any definitive conclusions regarding the degree to which IL-6 is a good kid, a bad kid, or an opportunistic kid who behaves differently depending on who he is hanging out with.

Rather, we have sought to highlight the fact that IL-6 plays remarkably complex, and likely contradictory, roles in a wide array of biological processes of great relevance to mental and physical health. While discouraging to those of us who like parsimony, this complexity should encourage a sense of active curiosity regarding how IL-6 might best be corralled into the service of medicine.

Meanwhile, if you think you know IL-6, think again. And if you can, do your thinking in a sauna.

Financial Disclosures:

Dr. Raison discloses that in the last twelve months he has served as a consultant to Usona Institute, Emory Healthcare, Alkermes and Novartis. This publication was supported in part by the National Center for Advancing Translational Sciences, National Institutes of Health, through Grant Numbers UL1TR001436 and KL2TR001438. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH. Professor Pariante is supported by the UK National Institute for Health Research (NIHR) Biomedical Research Centre at the South London and Maudsley NHS Foundation Trust and King's College London; and by the UK Medical Research Council, grants MR/L014815/1, MR/J002739/1 and MR/N029488/1; he has received research funding from pharmaceutical companies interested in the development of new antidepressants targeting immune mechanisms, such as Johnson & Johnson and Eleusis Ltd, but this amounts to less than 10% of his overall research funding. Moreover, he has received research funding from the Medical Research Council (UK) and the Wellcome Trust for research on depression and inflammation as part of two large consortia that also include Johnson & Johnson, GSK, Pfizer and Lundbeck.

References

Bauer J, Hohagen F, Gimmel E, Bruns F, Lis S, Krieger S, Ambach W, Guthmann A, Grunze H, Fritsch-Montero R, 1995. Induction of cytokine synthesis and fever suppresses REM sleep

- and improves mood in patients with major depression. *Biol. Psychiatry* 38, 611–621. [PubMed: 8573663]
- Del Giudice M, Gangestad SW, 2018. Rethinking IL-6 and CRP: Why they are more than inflammatory biomarkers, and why it matters. *Brain Behav. Immun.* 70, 61–75 10.1016/j.bbi.2018.02.013 [PubMed: 29499302]
- Heinonen I, Laukkanen JA, 2018. Effects of heat and cold on health, with special reference to Finnish sauna bathing. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 314, R629–R638. 10.1152/ajpregu.00115.2017 [PubMed: 29351426]
- Knight J, Costanzo ES, Singh S, Yin Z, Szabo A, Hillard C, Rizzo JD, D'Souza A, Coe CL, Drobycki WR, 2018. Pre-Transplant Tocilizumab is Associated with More Severe Depression, Anxiety, Pain, and Sleep Following Allogeneic Hematopoietic Cell Transplantation. *Biology of Blood and Marrow Transplantation* 24, S260–S261. 10.1016/j.bbmt.2017.12.192
- Norelli M, Camisa B, Barbiera G, Falcone L, Purevdorj A, Genua M, Sanvito F, Ponzoni M, Doglioni Ca., Cristofori P, Traversari C, Bordignon C, Ciceri F, Ostuni R, Bonini C, Casucci M, Bondanza A. 2018. Monocyte-derived IL-1 and IL-6 are differentially required for cytokine-release syndrome and neurotoxicity due to CAR T cells. *Nature*; epub.
- Park M, Newman LE, Gold PW, Luckenbaugh DA, Yuan P, Machado-Vieira R, Zarate CA, 2016. Change in cytokine levels is not associated with rapid antidepressant response to ketamine in treatment-resistant depression. *Journal of psychiatric research* 84, 113–118. 10.1016/j.jpsychires.2016.09.025 [PubMed: 27718369]
- Pedersen BK, Febbraio MA, 2008. Muscle as an endocrine organ: focus on muscle-derived interleukin-6. *Physiol. Rev.* 88, 1379–1406. 10.1152/physrev.90100.2007 [PubMed: 18923185]
- Raison C, 2017. 419. Inflammation in Treatment Resistant Depression: Challenges and Opportunities. *Biological Psychiatry* 81, S171. 10.1016/j.biopsych.2017.02.903
- Raison CL, Demetrashvili M, Capuron L, Miller AH, 2005. Neuropsychiatric adverse effects of interferon-alpha: recognition and management. *CNS Drugs* 19, 105–123. [PubMed: 15697325]
- Sun Y, Wang D, Salvatore G, Hsu B, Curran M, Casper C, Vermeulen J, Kent JM, Singh J, Drevets WC, Wittenberg GM, Chen G, 2017. The effects of interleukin-6 neutralizing antibodies on symptoms of depressed mood and anhedonia in patients with rheumatoid arthritis and multicentric Castleman's disease. *Brain Behav. Immun.* 66, 156–164. 10.1016/j.bbi.2017.06.014 [PubMed: 28676350]
- Wueest S, Item F, Boyle CN, Jirkof P, Cesarovic N, Ellingsgaard H, Böni-Schnetzler M, Timper K, Arras M, Donath MY, Lutz TA, Schoenle EJ, Konrad D, 2014. Interleukin-6 contributes to early fasting-induced free fatty acid mobilization in mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 306, R861–867. 10.1152/ajpregu.00533.2013 [PubMed: 24694381]