

## Review Article

# Recent Advances in the Synthesis of Xanthines: A Short Review

Anandi Kapri,<sup>1</sup> Nitin Gupta,<sup>2</sup> and Sumitra Nain <sup>1</sup>

<sup>1</sup>Department of Pharmacy, Banasthali Vidyapith, Banasthali, Rajasthan, India

<sup>2</sup>Agilent Technologies Pvt. Ltd., 181/46, Industrial Area, Phase-1, Chandigarh, India

Correspondence should be addressed to Sumitra Nain; [nainsumitra@gmail.com](mailto:nainsumitra@gmail.com)

Received 30 April 2022; Revised 31 August 2022; Accepted 26 September 2022; Published 8 November 2022

Academic Editor: Giacomo Garibotto

Copyright © 2022 Anandi Kapri et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Xanthine and its derivatives are considered a pharmacologically potential moiety that manifests immense biological activities. Owing to this much diversity in the biological field, this scaffold has fascinated the attention of many researchers around the globe to scrutinize its basic structure chemically as well as biologically. In recent years, xanthine derivatives have been used therapeutically in different pathological conditions due to their presence in day-to-day life. Herein, we review the recent progress in the synthesis of xanthine and its derivatives. Some of the widely used synthetic strategies such as (a) Traube's synthesis, (b) one-pot synthesis, (c) xanthine-annealed synthesis, and (d) miscellaneous synthesis were compiled in this review paper. The results obtained from this review paper highlight the significance of various xanthine derivatives as possible leads to the development of new drugs. The data compiled in this review paper could help the medicinal chemist in designing new active compounds from the modification of the already existing compounds in the search for novel drug leads. This report concludes that the various synthetic procedures exemplified in this review paper may serve as a support system for the designing of new molecules with a xanthine scaffold. Thus, we hope that this molecule may serve as the prototype in order to find out more active xanthine derivatives.

## 1. Introduction

Xanthine or 3,7-dihydropurine-2,6-dione (see Figure 1), a unique heterocycle is a purine base containing nitrogen as a central atom and composed of a pyrimidine ring fused with an imidazole ring.

It is an essential core element of diverse natural products because their structural fragments are found in various natural and synthetic medicinally active compounds [1]. The versatility of the xanthine moiety displays that it is the essential part of several medicinal agents and some of its derivatives have shown innumerable physiological and pharmacological activities viz. respiratory tract, heart, smooth muscle cells, CNS (central nervous system), kidney, and stomach [2]. In brief, the objective of such studies is to expose the drug-like properties of xanthine and its derivatives in order to build prospects for harnessing the full potential of this scaffold.

Xanthine scaffold has fascinated the attention of researchers in health sciences due to its remarkable properties either chemical or physical [3]. Over the last two decades, the

compound and its derivatives have gained considerable interest [4]. The xanthine scaffold can also act as a basic framework for numerous pharmacologically active scaffolds [5]. Several patent applications were also filed for xanthine derivatives as mentioned in (see Table 1), which displays the therapeutic effectiveness of this scaffold.

Owing to the importance of xanthine moiety in medicinal chemistry and its broad range of biological activity [22]. This review article primarily focuses on the updated knowledge of synthetic methods used to access xanthine scaffolds. In the present work, we have compiled the recent literature that belongs to the synthetic strategies of these derivatives.

## 2. Search Strategy

The data has been compiled from the year 2010 to 13 July 2022. We performed an electronic search to find out the existing literature on xanthine derivatives. For this purpose, the compiled data has been searched from different search engines and databases such as Science Direct, Google




FIGURE 1: Structure of xanthine.

Scholar, Cochrane, PubMed, Scopus, and Scientific information databases. More than ten months were completely used to compile data for this manuscript. The key terms used during the search were “xanthine,” “xanthine derivatives” “synthesis of xanthines” and “targets of xanthine derivatives.” One hundred twenty-two papers were screened and then inclusion and exclusion criteria were applied to prepare this manuscript. Out of these papers, forty-four papers were used to prepare this manuscript. The last search was conducted on 15. March 2022.

The results obtained from this review paper highlight the significance of various xanthine derivatives as possible leads to the development of new drugs. The reactions covered in this review fall under the following categories: (a) Traube’s synthesis, (b) one-pot synthesis, (c) xanthine-annealed synthesis, and (d) miscellaneous synthesis.

### 3. Synthesis Methods

**3.1. Traube’s Method.** Traube’s synthesis is the oldest and most used synthesis method for xanthine analogs. The method consists of the preparation of 5,6-diaminouracil (**17**) from urea or substituted urea. Several xanthine derivatives were synthesized by using uracil (**15**) as a precursor. Two methods were employed for the synthesis. According to method A-the nitrosation reaction (**16**) took place in presence of sodium nitrite and acetic acid followed by reduction (**17**) in presence of sodium dithionate and cyclisation to form substituted xanthine in presence of formic acid and sodium formate. In method B, the nitrosation reaction took place in presence of sodium nitrite, dimethyl formate, and hydrochloric acid followed by reduction and ring closure in presence of triethylorthoformate to form substituted xanthine (**18**). The main disadvantage of this method is that the overall process is tedious because of the ring closure synthetic mechanism (Scheme 1). Some recently used examples of this synthesis method are discussed in this section.

The preparation of 1-,3-,7-,8-substituted xanthines was done by using 3-substituted-6-aminouracil (**19**) as the substrate. This substrate was further alkylated (**20**) with alkyl iodide, followed by nitrosation (**21**) with sodium nitrite and reduction lead to 3-substituted 5,6-diaminouracil (**22-24**), followed by subsequent condensation with corresponding carboxylic acid lead to the corresponding xanthines (**25, 26**) [23] (Scheme 2). The derivatives were evaluated for the A1, A2A, A2B, and A3 subtypes of adenosine receptors. A number of derivatives have shown moderate to high affinity for human A1 and A2B adenosine receptors and lower affinity for A2A and A3 adenosine receptors.

An efficient method was reported for the synthesis of 8-substituted xanthines. The synthesis was performed by using 5, 6-diamino-1, 3-dimethyluracil (**27**) as starting material. The starting material was synthesized by condensing N, N-dimethyl urea, cyanoacetic acid, and acetic anhydride, and subsequent nitrosation and reduction afforded 5, 6-diamino-1, 3-dimethyluracil. Treatment of this compound with a substituted aldehyde (**28**) forms benzylidene derivative (**29**), which on oxidative cyclisation, by refluxing with thionyl chloride afforded the desired xanthine derivative (**30**) [24] (Scheme 3).

A series of 8-substituted xanthines by using 1,3-dimethyl and 1, 3-diethyl 5, 6-diaminouracils (**31**) as starting compounds was reported. The uracil was treated with desired carboxylic acid (**32**) to yield intermediate amides. Furthermore, the amide was reacted with NaOH (sodium hydroxide) to yield 1-, 3-, 7-trisubstituted xanthines (**33**). The obtained compound was further treated with an excess of iodomethane or iodoethane to yield 1-, 3-, 7-, 8-substituted xanthines (**34**) [25] (Scheme 4).

Novel 8-(p-substituted) xanthine derivatives were designed, synthesized, and reported. The synthesis was done by using 5, 6-diamino-1, 3-dimethyluracil (**35**). The treatment of uracil with aldehydes (**36**) yields the corresponding Schiff base (**37**), and further ring closure afforded the desired xanthine derivatives (**38**) [26] (Scheme 5).

The synthesis of 1, 3, 7, 8-substituted xanthine derivatives was reported. The synthesis was done by reacting 1,3-dialkyl-5, 6-diaminouracil (**39**) with that of substituted phenoxy acetic acid (**40**) and EDAC (1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide). In the next step, the obtained intermediate amide was treated with aq. NaOH to yield the desired xanthine analog (**41, 42**) [27] (Scheme 6).

1-, 3-, 8-substituted xanthines were synthesized from 1,3-dipropyl-5, 6-diaminouracil (**43**). The uracil was reacted with corresponding carboxylic acid (**44**) in presence of EDCI [1-ethyl-3-(3-dimethylaminopropyl) carbodiimide] to yield the benzylidene derivative (**45**), and further ring closure was achieved with NaOH to give 8-substituted xanthines (**46-49**) [28] (Scheme 7).

A new reaction for the synthesis of tetrahydropyrazino [2,1f] purinediones was reported. The 5,6-diaminouracil (**50**) was treated with glycolic acid at 100°C, ring closure was achieved with NaOH solution to give 8-hydroxymethylxanthine (**51**). Subsequent alkylation (**52**) at position-7 in the presence of DIPEA (N, N-diisopropylethylamine). In the end, the obtained compound (**53**) was treated with various amines (**54**) resulting in xanthine derivatives (**55**) [29] (Scheme 8).

8-(2-nitroaryl) xanthines were synthesized from dimethyl urea (**56**), cyanoacetic acid, and acetic anhydride. The nitrosation (**57**) and reduction of 6-aminouracil (**58**) was achieved resulting in 5, 6-diaminouracil (**59**), which on further treatment with a corresponding aldehyde (**60**) yielded benzylidene derivative (**61**). These derivatives were further cyclized by refluxing in thionyl chloride to give the required xanthines (**62**). All the synthesized compounds were further biologically evaluated for adenosine receptor subtypes [30] (Scheme 9).

TABLE 1: Patent filed for xanthine derivatives in last twenty years (2002–2020).






S. No.	Patent application number	Year of filing	Applicant name	Patent office	Compound	Category
1	AU 200521 958-B2 [6]	2005	Boehringer Ingelheim International GmbH	Australian	 (1) 8-(3-Amino-piperidin-1-yl)-xanthine	DPP-IV inhibitor
2	CA 02873524 [7]	2007	Boehringer Ingelheim International GmbH	Canadian	1-[(4-methyl-quinazolin-2-yl)methyl]-3-7-(2-butyn-1-yl)-8-(3-(R)-aminopiperidin-1-yl) xanthine	DPP-IV inhibitor
3	EP 2058311A2 [8]	2003	Boehringer Ingelheim International GmbH and Co.	European patent office	 (2) 8-{3-amino-piperidin-1-yl}-xanthine	DPP-IV inhibitor
4	DE 6020050 00 986T2 [9]	2005	Loreal Paris	Deutsches patent under Markenamt	—	Obesity
5	EP 1368349 B1 [10]	2002	Boehringer Ingelheim International GmbH and Co.	European	 (3) Xanthine derivative	DPP-IV
6	EP 1515972 B1 [11]	2003	F.Hoffman-La Roche AG 4070 Basel (CH)	European	 (4) Amide substituted xanthine derivative	Type-II diabetes
7	EP 1599477 B1 [12]	2004	F.Hoffman-La Roche AG 4070 Basel (CH)	European	 (5) Sulphonamide substituted xanthine Derivative	PEPCK inhibitors

TABLE 1: Continued.











S. No.	Patent application number	Year of filing	Applicant name	Patent office	Compound	Category
8	EP 1689748 B1 [13]	2004	Boehringer Ingelheim International GmbH	European	 <p>(6)</p> <p>8-(piperazine-1-yl) and 8-(1,4 diazepam-1-yl)-xanthine</p>	Diabetes
9	ES 2401128 T3 [14]	2006	GlaxoSmithKline LLC	Spain	 <p>(7)</p> <p>Xanthine derivative</p>	Agonist of HM74 A
10	US 7696212 B2 [15]	2010	Boehringer Ingelheim International GmbH and Co. KG	United States	 <p>(8)</p> <p>Xanthine derivative</p>	DPP-IV inhibitor
11	US 7838529 B2 [16]	2010	Boehringer Ingelheim International GmbH, Ingelheimam Rhein (DE)	United States	 <p>(9)</p> <p>Xanthine derivative</p>	Type-2 diabetes mellitus, antiobesity.
12	US 7879864 B2 [17]	2011	Sanofi-Aventis Deutschland GmbH, Frankfurt am main (DE)	United States	 <p>(10)</p> <p>8-amino alkoxy- xanthine</p>	DPP-IV inhibitor
13	US 9221821 B2 [18]	2015	Forest Laboratories Holdings Limited, Hamilton (BM)	United States	 <p>(11)</p> <p>1,3-Substituted aminouracils and xanthine derivatives</p>	-

TABLE 1: Continued.


S. No.	Patent application number	Year of filing	Applicant name	Patent office	Compound	Category
14	US 10202383 B2 [19]	2019	Boehringer Ingelheim International GmbH, Ingelheim am Rhein (DE)	United States	 (12) 8-{3-Amino-piperidin-1-yl}-xanthines	DPP-IV
15	US 10214530 B2 [20]	2019	Max-Delbruck-Centrum Fur Molekulare Medizin, Berlin (De), Forschungs-Verbund-Berlin-E. V., Berlin (De)	United States	 (13) Xanthine derivative	Serotonin based disease
16	US 201111 8464 [21]	2009	Ing-Jun Chen Linya District (TW)	United States	 (14) Halogenated xanthine derivative	Anticancer




SCHEME 1: Synthetic procedure for substituted xanthines by Traube's method.

A novel series of 1-,3-,7-triethyl substituted xanthine derivatives were reported. The 5,6-diamino-1,3-dialkyluracil (**63**) was used as the starting material. The starting material was treated with commercially available carboxylic acid (**64**)


to give 1,3-dialkyl-6-amino-5-carboxamidouracil (**65**) intermediate, subsequent cyclisation of this derivative yield corresponding xanthine derivatives (**66,67**) [31] (Scheme 10).



SCHEME 2: Synthesis of 1-,3-,8-trisubstituted 1H-purine-2,6(3H,9H)-diones.




SCHEME 3: Synthesis of 8-cyclopentylxyphenyl xanthine derivatives.




Reagents and conditions: i) EDAC, dioxane:  $\text{H}_2\text{O}$  (1:1); ii) Na OH (aq), reflux  
 (iii)  $\text{CH}_3\text{I}$  or  $\text{C}_2\text{H}_5\text{I}$ ,  $\text{K}_2\text{CO}_3$ , DMF, ethanol or acetone.


SCHEME 4: Synthetic pathway to xanthine derivatives.



SCHEME 5: Synthesis of 8-(p-substituted) xanthine derivatives.



SCHEME 6: Synthesis of 8-(phenoxyethyl)-xanthine analogs.



Ar/Het = -Phenyl, Isoxazole, Pyrazole

(i)EDCl, MeOH, r.t.; (ii) 10% NaOH, MeOH, reflux; (iii) SEM-Cl, K<sub>2</sub>CO<sub>3</sub>, DMF, r.t.; (iv) H<sub>2</sub>, 10% Pd/C, MeOH:DCM (2:1), r.t.; (v) K<sub>2</sub>CO<sub>3</sub>, acetone, 50°C; (vi) 2N HCl, ethanol, 80°C


SCHEME 7: Synthetic scheme of 1,3,8-substituted xanthines.

8-phenyl-1,3-dimethylxanthine derivatives were synthesized and reported. In this synthesis 5,6-diaminouracil (**68**) was used as the key intermediate. The intermediate was treated with a different substituted aldehyde (**69**) and methanol: acetic acid (4:1) results in the formation of benzylidene derivative (**70**), which further gets cyclized to form xanthine carboxylate ester (**71**), which on treatment with substituted amines at 80–100°C to yield (**72-75**) [32] (Scheme 11).

The fast, efficient synthesis of 6-Amino-5-carboxamidouracils as the predecessor for the 8-substituted xanthines was reported. In procedure A: the 5,6-diaminouracils (**76**) were condensed with aldehydes, results in imine formation (**77**), following oxidative cyclisation resulted in 8-substituted xanthine derivatives (**78**), which is the most common route of xanthine synthesis. In Procedure B, the


uracil (**76**) derivative was treated with carboxylic acid (**80**) and EDAC-HCl (1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (**79**) which lead to the formation of xanthine derivative (**78**). In Procedure-C, the 5,6-diaminouracils (**76**) were made to react with carboxylic acid (**81**) but prior to this reaction the activation of the carboxylic acid to form carboxylic acid chloride (**82**) was done, further leading to the formation of 6-amino-5-carboxamidouracil derivative (**77**) and then subsequent ring closure yield 8-substituted xanthines (**73**) [33] (Scheme 12).

The synthesis of 1,3-disubstituted-8-styrylxanthines under chemo and regioselective conditions was reported. The synthesis was achieved by using 6-aminouracil (**83**) as the starting material. The 6-aminouracil was alkylated at 3-position to form 6-amino-3-propyluracil (**84**), which undergo nitrosation (**85**) and reduction to yield 5,6-diamino-3-



(a) i. glycolic acid (1.2 eq.), 1 h, 100°C, neat, ii. H<sub>2</sub>O, NaOH, pH 12–13, 100°C, 4 h;  
 (b) 1, 2-dibromoethane (6 eq.), DMF, DIPEA, 70°C, 16 h; (c) PBr<sub>3</sub> (4eq.), CH<sub>2</sub>Cl<sub>2</sub>, 0°C to rt, 1 h;  
 (d) amine (2 eq.), dimethoxyethane, DIPEA, rt, 16 h.

SCHEME 8: Schematic pathway of 8-substituted purinediones.




Reagents and reaction conditions: (i) cyanoacetic acid, acetic anhydride, reflux, sodium hydroxide;  
 (ii) glacial acetic acid, sodium nitrite, icebath; (iii) concentrated NH<sub>4</sub>OH, sodium dithionite.  
 (iv) MeOH/CH<sub>3</sub>COOH, room temperature, 18 h; (v) SOCl<sub>2</sub>, reflux, 30–40 min; NH<sub>4</sub>OH

SCHEME 9: Synthesis of 8-substituted xanthines.


propyl uracil (**86**). The obtained compound was condensed with EDCI and cinnamic acid to form 6-amino-3-propyl-5-styrylcarboxamide (**87,88**) and finally, cyclisation was achieved by using alkali to form the resultant xanthine derivative (**89**). All the synthesized compounds have shown good activity for both A1 and A2A adenosine receptors [34] (Scheme 13).

**3.2. Xanthine-Annealed Synthesis.** In xanthine-annealed synthesis, the upgradation of one more ring starting from the bicyclic xanthine scaffold produces several biologically important compounds with modified physiochemical properties. Though, xanthine-annealed synthesis is tedious but not widely used for the synthesis of xanthine derivatives. In this context, the synthesis of tetrahydropyrimido [2,1-f]






SCHEME 10: Synthesis of 8-(p-substituted) xanthines.




SCHEME 11: Systematic study for the synthesis of xanthine carboxylate amides.

purinediones by using a convergent approach was reported. The commercially available amines (**91**) were treated with 8-bromo-7-(3-chloropropyl)-1,3-dialkylxanthine (**90**) in presence of a base in DMF (dimethyl formamide) to form the substituted xanthine derivatives (**92**) [35] (Scheme 14).

Another contribution to this study is based on the synthesis of 8-Benzyl-substituted tetrahydropyrazino [2,1-f] purinediones was reported. The 1,3-dimethyl-8-hydroxymethylxanthine (**93**) was used as the starting material in this study. Primarily, position-7 of xanthine was alkylated




SCHEME 12: Synthesis of 8-substituted xanthine derivatives.




Reagents-i. HMDS, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, n-Pr-I, ii. NaNO<sub>2</sub>, AcOH, iii. Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>, NH<sub>4</sub>OH, iv. C<sub>9</sub>H<sub>8</sub>O<sub>2</sub>, CH<sub>3</sub>OH, v. R-X, DMF, K<sub>2</sub>CO<sub>3</sub>, vi. KOH, CH<sub>3</sub>OH

SCHEME 13: 1,3-disubstituted 8-styrylxanthines.




SCHEME 14: Synthesis of tetrahydropyrimido [2,1-f] purinediones using the convergent approach.




i) 1, 2 dibromoethane, DMF, 80°C, 6 h; ii)  $\text{PBr}_3$ ,  $\text{CH}_2\text{Cl}_2$ , RT, 1 h; iii) dimethoxyethane, N, Ndiisopropylethylamine, RT, 16 h

SCHEME 15: Synthetic pathway towards 8-substituted-purine-2,4(1H,3H)-diones.



Where X = Cl, Br and n = 1, 2, 3


SCHEME 16: Synthesis of tricyclic xanthine derivatives.




SCHEME 17: Synthesis of N8-substituted 1,3-dimethyltetrahydropyrazino [2,1-f] purinediones.

(94) with 1,2 dibromoethane. The hydroxy group present at the 8-position was then converted into the corresponding bromide to form the resulting purinedione derivative (95) which was further treated with different substituted benzylamines to afford the tetrahydropyrazino derivative (96) [36] (Scheme 15).


Another series based on xanthine annealed synthesis was reported. The N-9 benzyl substituted purinediones were synthesized from theophylline (97), which was oxidatively brominated to give the compound (98). Subsequently, 8-bromotheophylline has been alkylated at the N-7 position to obtain an 8-bromo-7-alkyltheophylline derivative (99).



SCHEME 18: One-pot synthesis of xanthines.



SCHEME 19: One-pot three-component reaction-alkyl bromide, thiourea, and 8-bromocaffeine.



SCHEME 20: One-pot method for the synthesis of xanthine core via BDMS accelerated condensation.

Finally, condensation with appropriate amine leads to tricyclic xanthines (**100**) [37] (Scheme 16).

Moreover, 8-Substituted 1,3-dimethyltetrahydropyrazino [2,1-f] purinedione derivatives were reported as multitargeted drugs. Initially, the compound (**101**) was converted to the corresponding bromide to get compound (**102**). Furthermore, the compound was treated with appropriate amine (**103**) to afford tetrahydropyrazino derivatives (**104**) [38] (Scheme 17).


**3.3. One-Pot Method.** According to the emerging importance of xanthine and its derivatives. The development of a novel method for access to xanthine scaffold in less time, without using toxic chemicals, and improvement in yield is still in great demand. In this study, numerous functionalized xanthines were synthesized with better yield, without the use of toxic reagents, and in lower time. Based on the aforementioned information, a one-pot synthesis of xanthine was reported. The 5,6-diaminouracil (**105**) was chosen as the substrate. The substrate was refluxed with acetic anhydride

in acetic acid to obtain compound (**106**) and heating of substrate with malononitrile gave compound (**107**) in good yield [39] (Scheme 18).

In addition, the synthesis of some 8-alkylmercaptocaffeine derivatives has been reported via a one-pot three-component reaction. The treatment of thiourea (**109**) with alkylbromide (**108**) and 8-bromocaffeine (**110**) yielded 8-alkylmercaptocaffeine derivative (**111**) in excellent yield [40] (Scheme 19).


Another one-pot synthesis of 8-xanthine derivatives was reported. The synthesis was done by treatment of 5,6-diaminouracil (**112**) with a simple aldehyde to form a xanthine derivative (**113,114**) through an imine intermediate (**115**). Furthermore, cyclisation of the intermediate yield xanthine derivative (**116**). All the synthesized compounds were observed for A2A adenosine receptor antagonist [41] (Scheme 20).

**3.4. Miscellaneous Synthesis.** Xanthines can also be used as a starting material for the large-scale manufacture of xanthine derivatives. Mainly the alteration at 1-, 3-, 7- and 8- in the



Reagents and conditions: (i)  $\text{Br}_2$ , AcONa, AcOH, r.t.–60°C, 2 h; (ii) 1-Bromo-2-butyne, DIEA, DMF, 80°C, 6 h; (iii) Propargylbromide,  $\text{Cs}_2\text{CO}_3$ , DMF, rt, 1 h; (iv) Morpholine,  $\text{K}_2\text{CO}_3$ , DMF, 75°C, 4 h; (v)  $\text{ArN}_3$ , CuI, THF, rt, 6–8 h

SCHEME 21: Synthesis of 1,2,3-triazole-based xanthenes.




i) chloroacetic acid, chloroacetamide or propyl chloroacetate,  $\text{NaHCO}_3$  and DMF, reflux. ii)  $\text{NaOH}$ ,  $\text{H}_2\text{O}$ , reflux, iii) propanol, dioxane,  $\text{H}_2\text{SO}_4$ , reflux.

SCHEME 22: Synthesis of 3-benzyl-8-propylxanthinyl-7-acetic acid, its ester, and amide.

xanthine scaffold results in derivatives with numerous pharmacological activities. To discover numerous roles of xanthenes, the examination of various methods used for the synthesis became an emerging topic of research. In this context, a novel series of 1,2,3-triazole-based xanthenes were designed and synthesized. The compound (117) was treated

with bromine to form an intermediate (118). The intermediate was then allowed to react with 1-bromo-2-butyne to form compound (119), subsequent interaction of this compound with propargyl bromide gave another purine intermediate (120) which on treatment with morpholine gave morpholinodione intermediate (121). Lastly, the



SCHEME 23: Schematic synthesis of xanthine derivatives.

TABLE 2: Reported potent compounds of xanthines and their targets.

S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
1	1,3,8- and 1,3,7,8-substituted xanthines derivatives [23]	A2B and A1 adenosine receptors	<p style="text-align: center;">Compound 14q</p>
2	8(cyclopentyloxy) phenylxanthines derivatives [24]	A1 and A2 adenosine receptors	<p style="text-align: center;">Compound 6a</p> <p style="text-align: center;">Compound 12</p>

TABLE 2: Continued.







S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
3	8-(phoxymethyl) xanthine and 8-(3-phenylpropyl) xanthine derivatives [25]	A2A adenosine receptor antagonistic properties	 <p>Compound 6f</p>
4	8-(p-substituted-phenyl/benzyl) xanthines derivatives [26]	A2A adenosine receptor	
5	1,3-diethyl-7-methyl-8-(phoxymethyl)-xanthine and 1,3,7-trimethyl-8-(phoxymethyl)-xanthine derivatives [27]	A1 and A2A adenosine receptor antagonists	 <p>Compound 4f</p>  <p>Compound 4b</p>
6	8-(1-prop-2-ynyl-1H-pyrazol-4-yl)-xanthine derivatives [28]	A2B adenosine receptor antagonists	 <p>Compound 36</p>  <p>Compound 46</p>

TABLE 2: Continued.







S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
7	Tetrahydropyrazino-annulated theophylline derivatives [29]	Multi-targeted drugs with adenosine receptor (A1, A2A) and MAO-B antagonistic activity.	 <p>(15d)</p>
			 <p>(20e)</p>
			 <p>(20h)</p>
			 <p>(16a)</p>
			 <p>(16b)</p>
8	8-(2-nitroaryl) xanthines [30]	Human A2A adenosine receptor	 <p>Compound 9e</p>



TABLE 2: Continued.







S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
9	8-(3-phenylpropyl) xanthines 8-(2-phenylethyl) xanthines and 8-(phenoxyethyl) xanthines [31]	Adenosine A1 receptors	<div style="text-align: center;">  <p>Compound 3d</p>  <p>Compound 4d</p>  <p>Compound 5d</p> </div>
10	Carboxylate amides of 8-phenyl-1,3-dimethylxanthine [32]	Adenosine A2A receptors	<div style="text-align: center;">  <p>Compound 13b</p>  <p>Compound 16a</p>  <p>Compound 16d</p> </div>

TABLE 2: Continued.






S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
11	6-amino-5-carboxamidouracils [33]	—	
12	1,3-substituted 8-styrylxanthines [34]	A1 and A2A adenosine receptors antagonist	 <p data-bbox="1182 676 1214 697">11a</p>  <p data-bbox="1182 981 1214 1002">11d</p>
13	Tetrahydropyrimido[2,1-f] purinediones derivatives [35]	A2B adenosine receptor antagonists	 <p data-bbox="1007 1208 1134 1229">Compound 19f</p>  <p data-bbox="1007 1427 1134 1449">Compound 19g</p>

TABLE 2: Continued.







S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
14	8-benzyltetrahydropyrazino [2,1-f] purinediones derivatives [36]	Dual-target-directed A1/A2A adenosine receptor antagonists	 <p>Compound 72</p>  <p>Compound 36</p>
15	1,3-dialkyl-substituted tetrahydropyrimido [1,2-f]purine-2,4-diones [37]	Human A2A adenosine receptor antagonists	 <p>Compound 9u</p>  <p>Compound 10d</p>
16	8-benzyl-substituted tetrahydropyrazino [2,1-f] purinediones [38]	Dual A1/A2A adenosine receptor antagonists	 <p>Compound 41</p>  <p>Compound 57</p>

TABLE 2: Continued.





S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
17	Xanthine derivatives [39]	Antimicrobial and antioxidant activities	 <p>(9)</p>
17	Xanthine derivatives [39]	Antimicrobial and antioxidant activities	 <p>(10)</p>
18	8-alkylmercaptocaffeine derivatives [40]	—	 <p>(12a-q)</p>
19	8-substituted xanthine derivatives [41]	A2A adenosine receptor antagonists	

TABLE 2: Continued.









S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
20	1,2,3-triazole-based xanthine derivatives [42]	Dipeptidyl peptidase-4 inhibitors	<div style="text-align: center;">  <p>(7b)</p>  <p>(7e)</p>  <p>(7g)</p>  <p>(6e)</p> </div>
21	3-benzyl-8-propylxanthinyl-7-acetic acid [43]	—	<div style="text-align: center;">  </div>

TABLE 2: Continued.

S. No.	Substituted xanthine derivative	Disease target	Potent compounds reported
22	1,3,8-substituted tetrahydropyrazino [2,1-f] purinediones [44]	A1 and A2A adenosine receptors and MAO-B	 <p>Compound PSB-18339</p>  <p>Compound 20h</p>  <p>Compound 20e</p>

addition of the 1,3-dipolar cycle of terminal alkyne gave the corresponding 1,2,3-triazoles (**122,123**) in good yield [42] (Scheme 21).

3-benzyl-8-propylxanthinyl-7-acetic acid and its derivatives were designed, synthesized, and reported. The compound (**124**) was taken as a lead compound. The lead compound was treated with chloroacetic acid, chloroacetamide, or propyl chloroacetate to form the target xanthine derivatives (**125-127**) [43] (Scheme 22).

Theophylline (**128**), 8-bromotheophylline (**129**), and theobromine (**130**) were reacted with various 2/3-chloro-N-phenylacetamides or their propanamide analogs to obtain the resultant xanthine derivatives (**131-133**). The synthesized compounds were biologically assessed for in vitro bronchodilator activity [44] (Scheme 23).

#### 4. Reported Potent Compounds of Xanthines and Their Targets

Natural and synthetic compounds consisting of xanthine scaffold showed a variety of pharmacological activities. A large number of biologically active compounds were obtained by incorporating different substituents at different places in this ring [22]. A number of reported potent compounds of xanthines along with the therapeutic disease target are discussed (Table 2).

#### 5. Conclusion

Our prime aim in writing this review paper is that the content presented in this review paper will be beneficial to the field and will provide great help to those researchers working on this scaffold. This review article provides a summary overview of the synthesis of xanthine structures. The development of novel, selective and efficient methods for the formation of xanthine ring starting from commercially available substrates is a pivotal target in the current organic synthesis. In this regard, the synthesis of xanthine derivatives by Traube's method is the most used approach. However, other approaches were also found attractive to researchers. The various synthetic procedures exemplified in this review paper may serve as a support system for the designing of new molecules with xanthine scaffold. We hope that the data compiled in this review paper could help the medicinal chemist in designing new active compounds from the modification of the already existing compounds in the search for novel drug leads.

#### Abbreviations

DMF:	Dimethylformamide
SAR:	Structural activity relationship
$A_1$ , $A_{2A}$ , $A_{2B}$ ,	Adenosine receptor subtypes
$A_3$ :	

EtOH:	Ethanol
H <sub>2</sub> O:	Water
HCl:	Hydrochloric acid
NaNO <sub>2</sub> :	Sodium nitrite
AcOH:	Acetic acid
HCl:	Hydrochloric acid
HCOOH:	Formic acid
HCOONa:	Sodium formate
NaOH:	Sodium hydroxide
HC(OEt) <sub>3</sub> :	Triethylorthoformate
CH <sub>3</sub> I:	Methyl iodide
K <sub>2</sub> CO <sub>3</sub> :	Potassium carbonate
MeOH:	Methanol
COOH:	Carboxylic acid
SOCl <sub>2</sub> :	Thionyl chloride
RT:	Room temperature
DIPEA:	Diisopropylethylamine
NaH:	Sodium hydride
KI:	Potassium iodide
C <sub>2</sub> H <sub>5</sub> I:	Ethyl iodide
CH <sub>3</sub> COOH:	Acetic acid
HBr:	Hydrogen bromide
NaClO <sub>3</sub> :	Sodium chlorate
TEBA:	Benzyltriethylammonium chloride
NEt <sub>3</sub> :	Triethylamine
μM:	Micromolar
CH <sub>2</sub> Cl <sub>2</sub> :	Dichloromethane
Ac <sub>2</sub> O:	Acetic anhydride
CH <sub>2</sub> (CN) <sub>2</sub> :	Malononitrile
AcONa:	Acetic anhydride-sodium acetate
DIEA:	Diisopropylethylamine
Cs <sub>2</sub> CO <sub>3</sub> :	Cesium carbonate
CuI:	Copper iodide
THF:	Tetrahydrofuran
NaHCO <sub>3</sub> :	Sodium hydrogen carbonate
H <sub>2</sub> SO <sub>4</sub> :	Sulfuric acid
EDC HCl:	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EDCI:	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide.

## Conflicts of Interest

The authors declare that they have no conflicts of interest that could appear to influence the work reported in this review article.

## Acknowledgments

The authors are grateful to the Honorable Vice-Chancellor, Banasthali Vidyapith, Rajasthan, for providing essential facilities.

## References

- [1] N. Singh, A. K. Shreshtha, M. S. Thakur, and S. Patra, "Xanthine scaffold: scope and potential in drug development," *Heliyon*, vol. 4, no. 10, Article ID e00829, 2018.
- [2] S. Mourya, R. Bodla, R. Taurean, and A. Sharma, "Simultaneous estimation of xanthine alkaloids (theophylline, theobromine and caffeine) by high-performance liquid chromatography," *International Journal of Drug Regulatory Affairs*, vol. 7, no. 2, pp. 35–41, 2019.
- [3] H. Ashihara, K. Mizuno, T. Yokota, and A. Crozier, "Xanthine alkaloids: occurrence, biosynthesis, and function in plants," *Progress in the Chemistry of Organic Natural Products*, vol. 105, pp. 1–88, 2017.
- [4] E. Szymańska, J. Mazurkiewicz, and K. Kieć-Kononowicz, "Methods for the synthesis of xanthine-derived polycyclic fused systems," *Heterocyclic Compound*, vol. 19, no. 5, pp. 297–310, 2013.
- [5] S. Perviz, H. Khan, and A. Pervaiz, "Plant alkaloids as an emerging therapeutic alternative for the treatment of depression," *Frontiers in Pharmacology*, vol. 7, no. 28, pp. 28–37, 2016.
- [6] F. Himmelsbach, E. Langkopf, M. Eckhardt, M. Tadayyon, and L. Thomas, "8-[3-amino-piperidin-1-yl]-xanthine, the production thereof and the use in the form of a dpp inhibitor," WO2005085246A1, 2005.
- [7] P. Sieger and W. Pfrengle, "Salt forms of 1-[(4-methylquinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8(3-(r)-amino-piperidin-1-yl)-xanthine," CA2745037A1, 2020.
- [8] F. Himmelsbach, E. Langkopf, M. Eckhardt et al., "8-[3-amino-piperidin-1-yl]-xanthines, their production and utilisation as medicine," EP2058311-A2, 2009.
- [9] A. Moreno and B. Piot, "Slimming composition containing a xanthine-based base, a dimethicone copolypol and a polyurethane powder," DE602005000986, 2008.
- [10] F. Himmelsbach, M. Mark, M. Eckhardt, E. Langkopf, R. Maier, and R. Lotz, "Xanthine derivative, production and use thereof as a medicament," EP1368349B1, 2007.
- [11] P. W. Dunten, L. H. Foley, N. J. S. Huby, S. L. Pietranico-Cole, and W. Yun, "Amide substituted xanthine derivatives with gluconeogenesis modulating activity," EP1515972B1, 2009.
- [12] L. H. Foley, N. J. S. Huby, S. L. Pietranico-Cole, W. Yun, and P. W. Dunten, "Sulfonamide substituted xanthine derivatives for use as pepck inhibitors," EP1599477B1, 2005.
- [13] F. Himmelsbach, E. Langkopf, M. Eckhardt, M. Tadayyon, and L. Thomas, "Novel 8-(Piperazine-1-yl)- and 8-(1,4-Diazepan-1-yl)-xanthine, the production and use thereof in the form of a drug," EP1689748B1, 2008.
- [14] R. J. D. Hatley, A. M. Mason, and I. L. Pinto, "Xanthine derivatives as selective agonists of HM74A," ES2401128T3, 2013.
- [15] F. Himmelsbach, E. Langkopf, M. Eckhardt et al., "Xanthine derivatives, the preparation thereof and their use as pharmaceutical compositions," US7696212B2, 2010.
- [16] F. Himmelsbach, E. Langkopf, M. Eckhardt, M. Mark, R. Maier, and R. Lotz, "Xanthine derivatives, their preparation and their use in pharmaceutical compositions," US7838529B2, 2010.
- [17] K. Schoenafinger, G. Jaehne, E. Defossa, C. Buning, G. Tschank, and U. Werner, "Substituted 8-aminoalkoxy-xanthine compounds and compositions useful in the treatment of hyperlipidemia and diabetes," US7879864B2, 2011.
- [18] N. Diep and Y. B. Kalyan, "Methods for the synthesis of 1,3-substituted aminouracils and other xanthine-related compounds," US9221821B2, 2015.
- [19] F. Himmelsbach, E. Langkopf, M. Eckhardt, and R. Maier, "8-[3-amino-piperidin-1-yl]-xanthines, the preparation thereof and their use as pharmaceutical compositions," US10202383B2, 2019.

- [20] M. Bader, E. Specker, S. Matthes et al., "Xanthine derivatives, their use as a medicament, and pharmaceutical preparations comprising the same," US10214530B2, 2019.
- [21] C. Ing-Jun, "Halogenated xanthine derivatives and precursors thereof for anti-cancer and anti-metastasis activity and preparing method thereof," US2011118464, 2019.
- [22] A. Kapri, S. Pant, N. Gupta, and S. Nain, "Recent advances in the biological significance of xanthine and its derivatives: a review," *Pharmaceutical Chemistry Journal*, vol. 56, no. 4, pp. 461–474, 2022.
- [23] J. E. Rodríguez-Borges, X. García-Mera, M. C. Balo et al., "Synthesis and pharmacological evaluation of novel 1, 3, 8- and 1, 3, 7, 8-substituted xanthines as adenosine receptor antagonists," *Bioorganic & Medicinal Chemistry*, vol. 18, no. 5, pp. 2001–2009, 2010.
- [24] R. Bansal, G. Kumar, D. Gandhi, R. Yadav, L. Young, and A. Harvey, "Synthesis of 8-(cyclopentyloxy)phenyl substituted xanthine derivatives as adenosine A2A ligands," *Arzneimittelforschung*, vol. 60, no. 3, pp. 131–136, 2011.
- [25] M. M. Van der Walt, G. Terre'Blanche, A. Petzer, A. C. U. Lourens, and J. P. Petzer, "The adenosine A2A antagonistic properties of selected C8-substituted xanthines," *Bioorganic Chemistry*, vol. 49, pp. 49–58, 2013.
- [26] R. Yadav, R. Bansal, S. Kachler, and K. N. Klotz, "Novel 8-(p-substituted-phenyl/benzyl)xanthines with selectivity for the A2A adenosine receptor possess bronchospasmolytic activity," *European Journal of Medicinal Chemistry*, vol. 75, pp. 327–335, 2014.
- [27] R. Harmse, M. M. van der Walt, J. P. Petzer, and G. Terre'Blanche, "Discovery of 1, 3-diethyl-7-methyl-8-(phenoxyethyl)-xanthine derivatives as novel adenosine A1 and A2A receptor antagonists," *Bioorganic & Medicinal Chemistry Letters*, vol. 26, no. 24, pp. 5951–5955, 2016.
- [28] S. Basu, D. A. Barawkar, V. Ramdas et al., "Design and synthesis of novel xanthine derivatives as potent and selective A2B adenosine receptor antagonists for the treatment of chronic inflammatory airway diseases," *European Journal of Medicinal Chemistry*, vol. 134, pp. 218–229, 2017.
- [29] P. Koch, A. Brunschweiler, V. Namasivayam et al., "Probing substituents in the 1- and 3-position: tetrahydropyrazino-annelated water-soluble xanthine derivatives as multi-target drugs with potent adenosine receptor antagonistic activity," *Frontiers of Chemistry*, vol. 6, no. 206, pp. 206–228, 2018.
- [30] R. Bansal, G. Kumar, S. Rohilla et al., "Synthesis and evaluation of a new series of 8-(2-Nitroaryl) xanthines as adenosine receptor ligands," *Drug Development Research*, vol. 77, no. 5, pp. 241–250, 2016.
- [31] M. M. Van der Walt and G. Terre' Blanche, "1, 3, 7-Triethyl-substituted xanthines-possess nanomolar affinity for the adenosine A1 receptor," *Bioorganic & Medicinal Chemistry*, vol. 23, no. 20, pp. 6641–6649, 2015.
- [32] R. Yadav, R. Bansal, S. Rohilla, S. Kachler, and K.-N. Klotz, "Synthesis and pharmacological characterization of novel xanthine carboxylate amides as A2A adenosine receptor ligands exhibiting bronchospasmolytic activity," *Bioorganic Chemistry*, vol. 65, pp. 26–37, 2016.
- [33] D. Marx, L. M. Wingen, G. Schnakenburg, C. E. Müller, and M. S. Scholz, "Fast, efficient, and versatile synthesis of 6-amino-5-carboxamidouracils as precursors for 8-substituted xanthines," *Frontiers of Chemistry*, vol. 7, no. 56, pp. 56–15, 2019.
- [34] I. D. L. P. d. León, M. d. C. Parra-Cid, A. Muñoz-Zurita et al., "Motor Effects of 1, 3-Disubstituted 8-Styrylxanthines as Adenosine-Receptor Antagonists in Rats," *Pharmacology & Pharmacy*, vol. 4, no. 3, pp. 303–311, 2013.
- [35] P. Koch, R. Akkari, A. Brunschweiler et al., "1, 3-Dialkyl-substituted tetrahydropyrimido[1, 2-f] purine-2, 4-diones as multiple target drugs for the potential treatment of neurodegenerative diseases," *Bioorganic & Medicinal Chemistry*, vol. 21, no. 23, pp. 7435–7452, 2013.
- [36] A. Brunschweiler, P. Koch, M. Schlenk et al., "8-Benzyltetrahydropyrazino[2, 1-f] purinediones: water-soluble tricyclic xanthine derivatives as multitarget drugs for neurodegenerative diseases," *ChemMedChem*, vol. 9, no. 8, pp. 1704–1724, 2014.
- [37] M. Załuski, J. Schabikowski, M. Schlenk et al., "Novel multi-target directed ligands based on annelated xanthine scaffold with aromatic substituents acting on adenosine receptor and monoamine oxidase B. Synthesis, in vitro and in silico studies," *Bioorganic & Medicinal Chemistry*, vol. 27, no. 7, pp. 1195–1210, 2019.
- [38] A. Brunschweiler, P. Koch, M. Schlenk et al., "8-Substituted 1, 3-dimethyltetrahydropyrazino[2, 1-f]purinediones: water-soluble adenosine receptor antagonists and monoamine oxidase B inhibitors," *Bioorganic & Medicinal Chemistry*, vol. 24, no. 21, pp. 5462–5480, 2016.
- [39] S. A. El-Kalyoubi, E. A. Fayed, and A. S. Abdel-Razek, "One pot synthesis, antimicrobial and antioxidant activities of fused uracils: pyrimidodiazepines, lumazines, triazolouracil and xanthines," *Chemistry Central Journal*, vol. 11, no. 1, pp. 66–13, 2017.
- [40] M. N. S. Rad and S. Maghsoudi, "Two-step three-component process for one-pot synthesis of 8-alkylmercaptocaffeine derivatives," *RSC Advances*, vol. 6, no. 74, pp. 70335–70342, 2016.
- [41] P. LaBeaume, M. Dong, M. Sitkovsky et al., "An efficient route to xanthine based A2A adenosine receptor antagonists and functional derivatives," *Organic and Biomolecular Chemistry*, vol. 8, no. 18, pp. 4155–4157, 2010.
- [42] S. Narsimha, K. S. Battula, M. Ravinder, Y. N. Reddy, and V. R. Nagavelli, "Design, synthesis and biological evaluation of novel 1, 2, 3-triazole-based xanthine derivatives as DPP-4 inhibitors," *Journal of Chemical Sciences*, vol. 132, no. 1, pp. 59–9, 2020.
- [43] E. K. Mikhail'chenko, K. V. Aleksandrova, S. Levich, and A. Korzhova, "Synthesis and physical-chemical properties of functional derivatives of 3-benzyl-8-propylxanthinyl-7-acetic acid," *Current Issues in Pharmacy and Medical Sciences*, vol. 1, no. 23, pp. 14–19, 2017.
- [44] A. R. Mohamed, H. H. Georgey, R. F. George, W. I. El-Eraky, D. O. Saleh, and N. M. Abdel Gawad, "Identification of some novel xanthine-based derivatives with bronchodilator activity," *Future Medicinal Chemistry*, vol. 9, no. 15, pp. 1731–1747, 2017.