

# Cytomegalovirus Seropositivity in Older Adults Changes the T Cell Repertoire but Does Not Prevent Antibody or Cellular Responses to SARS-CoV-2 Vaccination

Jessica A. Breznik,<sup>\*,†,‡,§</sup> Angela Huynh,<sup>‡</sup> Ali Zhang,<sup>\*,†,¶</sup> Lucas Bilaver,<sup>\*,†,‡</sup> Hina Bhakta,<sup>‡</sup> Hannah D. Stacey,<sup>\*,†,¶</sup> Jann C. Ang,<sup>\*,†,¶</sup> Jonathan L. Bramson,<sup>\*,†,‡,¶</sup> Ishac Nazy,<sup>‡,||</sup> Matthew S. Miller,<sup>\*,†,¶</sup> Judah Denburg,<sup>‡</sup> Andrew P. Costa,<sup>‡,#,\*\*</sup> Dawn M. E. Bowdish,<sup>\*,†,‡,††</sup> and other members of the COVID-in-LTC Investigator Group<sup>1</sup>

Chronic infection with human CMV may contribute to poor vaccine efficacy in older adults. We assessed the effects of CMV serostatus on Ab quantity and quality, as well as cellular memory recall responses, after two and three SARS-CoV-2 mRNA vaccine doses, in older adults in assisted living facilities. CMV serostatus did not affect anti-Spike and anti-receptor-binding domain IgG Ab levels, nor neutralization capacity against wild-type or  $\beta$  variants of SARS-CoV-2 several months after vaccination. CMV seropositivity altered T cell expression of senescence-associated markers and increased effector memory re-expressing CD45RA T cell numbers, as has been previously reported; however, this did not impact Spike-specific CD4<sup>+</sup> T cell memory recall responses. CMV-seropositive individuals did not have a higher incidence of COVID-19, although prior infection influenced humoral immunity. Therefore, CMV seropositivity may alter T cell composition but does not impede the durability of humoral protection or cellular memory responses after SARS-CoV-2 mRNA vaccination in older adults. *The Journal of Immunology*, 2022, 209: 1892–1905.

**A**ging is associated with an increased frequency of viral respiratory infections and postinfection sequelae (1), as well as reduced efficacy and longevity of protective immunity after vaccination (2). Early in the COVID-19 pandemic, age was identified to be the most significant factor contributing to morbidity and mortality (3), and it was unclear how effective vaccines against the novel SARS-CoV-2 virus would be in older adults. In this population, vaccines that target a memory response (e.g., herpes zoster) are often more immunogenic and effective at preventing illness than vaccines that target viral Ags that are antigenically distant from previous circulating strains (e.g., some seasonal influenza vaccines) (4–7). SARS-CoV-2 mRNA vaccines have been shown to be protective in older adults (8–10), as they are effective at generating cellular and Ab-mediated immunity (11–14). However, in older adults, immune responses after vaccination are generally quite heterogeneous, and immunity may wane faster than in younger adults (11, 13).

In older adults, both immunosenescence and inflammation are thought to influence the immunogenicity of vaccines and longevity of protective immune responses after vaccination (15, 16). Many

studies have also implicated human CMV as a significant contributor to age-associated immune dysfunction. CMV is a common and persistent  $\beta$ -herpesvirus, found in ~60–90% of adults worldwide, and seropositivity increases with age (17, 18). CMV infection is typically asymptomatic in immunocompetent individuals, but its accompanying chronic immune activation fundamentally alters immune cell composition and function. There is a consensus that CMV seropositivity has a long-term impact on the maturation and composition of immune cells, including increased numbers and prevalence of CD8<sup>+</sup> T cells, with expansion of CMV-specific effector and memory cells at the expense of naive T cells (19, 20). Age-associated immunosenescence likewise contributes to similar changes within the T cell repertoire, reducing naive T cells and increasing memory T cell populations, which have impaired proliferation, differentiation, and effector functions (21, 22). A dysfunctional T cell repertoire may also have significant effects on B cell proliferation, differentiation, and maturation (23). Accordingly, CMV seropositivity has been implicated as an exacerbating factor in age-associated immune remodeling and interindividual immune diversity (24–27), as well as a modifying factor that

\*McMaster Immunology Research Centre, McMaster University, Hamilton, Ontario, Canada; <sup>†</sup>Michael G. DeGroot Institute for Infectious Disease Research, McMaster University, Hamilton, Ontario, Canada; <sup>‡</sup>Department of Medicine, Michael G. DeGroot School of Medicine, McMaster University, Hamilton, Ontario, Canada; <sup>§</sup>McMaster Institute for Research on Aging, McMaster University, Hamilton, Ontario, Canada; <sup>¶</sup>Department of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, Ontario, Canada; <sup>||</sup>McMaster Centre for Transfusion Research, McMaster University, Hamilton, Ontario, Canada; <sup>#</sup>Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada; <sup>\*\*</sup>Centre for Integrated Care, St. Joseph's Health System, Hamilton, Ontario, Canada; and <sup>††</sup>Firestone Institute for Respiratory Health, St. Joseph's Healthcare, Hamilton, Ontario, Canada

<sup>1</sup>All authors and their affiliations appear at the end of this article.

ORCID: 0000-0002-5434-9158 (J.A.B.); 0000-0002-1977-9339 (L.B.); 0000-0002-5695-4343 (I.N.); 0000-0003-3426-5069 (M.S.M.); 0000-0001-5398-0063 (J.D.); 0000-0001-9212-5641 (A.P.C.); 0000-0001-6823-2957 (D.M.E.B.).

Received for publication May 20, 2022. Accepted for publication September 3, 2022.

This work was supported by a grant from the Canadian COVID-19 Immunity Task Force and Public Health Agency of Canada awarded to A.P.C. and D.M.E.B. A.P.C. is the Schlegel Chair in Clinical Epidemiology and Aging. D.M.E.B. is the Canada Research

Chair in Aging and Immunity. Funding support for this work was provided by grants from the Canadian Institutes of Health Research, the Ontario Research Foundation COVID-19 Rapid Research Fund, and by the Canadian COVID-19 Immunity Task Force awarded to I.N. M.S.M. is supported, in part, by an Ontario Early Researcher Award and a Canada Research Chair in Viral Pandemics.

Address correspondence and reprint requests to Dr. Dawn M.E. Bowdish, Michael G. DeGroot Institute for Infectious Disease Research, McMaster Immunology Research Centre, McMaster University, MDCL 4020, 1280 Main Street West, Hamilton, ON L8S 4L8, Canada. E-mail address: bowdish@mcmaster.ca

The online version of this article contains supplemental material.

Abbreviations used in this article: AIM, activation-induced marker; CM, central memory; EM, effector memory; EMRA, EM re-expressing CD45RA; HA, hemagglutinin; MNT<sub>50</sub>, microneutralization titer at 50%; N, naive; RBD, receptor-binding domain; TD, terminally differentiated.

This article is distributed under The American Association of Immunologists, Inc., [Reuse Terms and Conditions for Author Choice articles](#).

Copyright © 2022 by The American Association of Immunologists, Inc. 0022-1767/22/\$37.50

may compromise infection outcomes and quality and longevity of immune protection after vaccination (28).

Although there is a paucity of data to date, CMV reactivation in older adults has been suggested to contribute to more severe COVID-19 (29–33), as CMV seropositivity has been associated with impaired Ab production and cellular memory recall responses (34–36). However, CMV seropositivity has also been reported to enhance cellular and Ab immune responses to unrelated bacteria or viruses, through diversification of CD8<sup>+</sup> TCRs and augmentation of basal inflammation (37–40). CMV seropositivity has, in addition, been associated with a reduced humoral response to inactivated split influenza virus vaccines (41–44) and viral vector-based Ebola vaccines (45). However, a recent meta-analysis reported that there was insufficient evidence that CMV-seropositive individuals have decreased Ab production after influenza vaccination (46). Recent data also show that CMV seropositivity in young adults does not affect Ab or cellular responses after vaccination with the adenovirus-based vector vaccine ChAdOx1 nCoV-19 (47). These conflicting reports may suggest context-, time-, and age-dependent effects of chronic CMV infection on immune function. SARS-CoV-2 mRNA vaccines have been widely deployed in Canada, particularly in older adults. Whether CMV seropositivity impacts SARS-CoV-2 mRNA vaccine efficacy and durability of immunity is not yet known. In this study, we investigated effects of CMV serostatus on vaccine-associated humoral protection and cellular memory recall responses several months after two and three doses of SARS-CoV-2 mRNA vaccines in older adults. We found that CMV serostatus does not impede Ab or cellular responses to SARS-CoV-2 vaccination in older adults.

## Materials and Methods

### Participant recruitment and blood collection

Participants in the COVID in Long-Term Care Study (<https://covidinlnc.ca>) were recruited from assisted living facilities (17 nursing homes and 8 retirement homes) in Ontario, Canada, between March and December 2021. All protocols were approved by the Hamilton Integrated Research Ethics Board and other site-specific research ethics boards, and informed consent was obtained. Venous blood was drawn in anti-coagulant-free vacutainers for isolation of serum, as per standard protocols (48). Venous blood was drawn in heparin-coated vacutainers for immunophenotyping and T cell activation assays. Blood was collected at least 7 d after two and/or three mRNA vaccine doses. Participants received two doses of Moderna Spikevax (100 µg; mRNA-1273) or Pfizer Comirnaty (30 µg; BNT162b2) as per the recommended schedules, and a third mRNA vaccine dose in Fall 2021 at least 6 mo from the last dose, as per Province of Ontario guidelines (49). For this study, humoral and cellular data were retrospectively assessed in the context of CMV serostatus from a cohort of 186 participants, 65 y of age and older. Blood was drawn after two and three mRNA vaccine doses from 47 cohort participants. Participant cohort demographics are summarized in Table I.

### Determination of CMV serostatus

CMV seropositivity was determined by ELISA with a human anti-CMV IgG ELISA kit (CMV) (no. ab108639, Abcam) as per the manufacturer's instructions. Serum samples from a participant's first blood draw were diluted 1:40 and assessed in duplicate. Samples with a CMV IgG index above or equal to the positive standard were classified as CMV seropositive, whereas samples with a CMV IgG index less than the positive standard were classified as CMV seronegative.

### Whole-blood immunophenotyping

Circulating immune cell populations were quantitated in whole blood using fluorophore-conjugated mAbs by multicolor flow cytometry with a CytoFLEX LX (four lasers, Beckman Coulter), as per standard protocols (48, 50, 51). CountBright absolute counting beads (no. C36950, Invitrogen/Life Technologies) were used to determine absolute cell counts. Data were analyzed with FlowJo v10.8.1 (Tree Star), following a previously published gating strategy to identify T cell populations (50). Five main subsets of human CD8<sup>+</sup> and CD4<sup>+</sup> T cells (naive [N], central memory [CM], effector memory [EM], EM re-expressing CD45RA [EMRA], and terminally differentiated [TD]) were

identified by their expression of CD45RA, CCR7, CD28, and/or CD57. CD8<sub>N</sub> and CD4<sub>N</sub> were classified as CD45RA<sup>+</sup>CCR7<sup>+</sup>, CD8<sub>CM</sub> and CD4<sub>CM</sub> as CD45RA<sup>+</sup>CCR7<sup>+</sup>, CD8<sub>EM</sub> and CD4<sub>EM</sub> as CD45RA<sup>+</sup>CCR7<sup>-</sup>, CD8<sub>EMRA</sub> and CD4<sub>EMRA</sub> as CD45RA<sup>+</sup>CCR7<sup>-</sup>, and CD8<sub>TD</sub> and CD4<sub>TD</sub> were classified as CD45RA<sup>+</sup>CCR7<sup>-</sup>CD28<sup>-</sup>CD57<sup>-</sup>, as per standard protocols (52).

### Assessment of T cell memory responses to SARS-CoV-2 Spike by an activation-induced marker assay

Ag-specific T cell recall responses were evaluated by an activation-induced marker (AIM) assay as per established protocols (50). Each participant sample was stimulated with a Spike glycoprotein SARS-CoV-2 peptide pool (1 µg/ml) containing overlapping peptides of the complete immunodominant sequence domain (no. 130-126-701, Miltenyi Biotec) as well as influenza hemagglutinin (HA) peptides (4 µl of 0.12 µg/µl HA; AgriFlu, Afluria Tetra inactivated influenza vaccine 2020–2021 season, Seqirus, Maidenhead, U.K.). A negative medium control (unstimulated) and positive stimulation control (polyclonal stimulation with CytoStim at 0.5 µl/well; no. 130-092-173, Miltenyi Biotec) were included with each sample. For each of these four conditions, 100 µl of heparinized venous blood was incubated with an equal volume of IMDM with GlutaMAX supplement (no. 31980030, Invitrogen/Life Technologies) for 44 h in 96-well flat bottom plates at 37°C. Samples were stained with fluorophore-conjugated mAbs and assessed with a CytoFLEX LX (4 lasers, Beckman Coulter, Brea, CA) as previously described (50). Data were analyzed with FlowJo, following a previously published gating strategy to identify AIM<sup>+</sup> T cells (50). Activated T cells (AIM positive) were identified by their coexpression of CD25 and CD134 (OX40) on CD4<sup>+</sup> T cells (53, 54) and coexpression of CD69 and CD137 (4-1BB) on CD8<sup>+</sup> T cells (55). Samples with a T cell count of at least 20 events and ≥2-fold above the unstimulated sample (negative control; i.e., stimulation index ≥ 2) were defined as AIM-positive. Expression of CXCR3 (CD183; Brilliant Violet 421, no. 353716, BioLegend), CCR4 (CD194; Brilliant Violet 605, no. 359418, BioLegend), and CCR6 (CD196; Brilliant Violet 785, no. 353422, BioLegend) was used to identify Th1 (CXCR3<sup>+</sup>CCR6<sup>-</sup>CCR4<sup>-</sup>), Th2 (CXCR3<sup>-</sup>CCR4<sup>+</sup>CCR6<sup>-</sup>), and Th17 (CXCR3<sup>-</sup>CCR4<sup>+</sup>CCR6<sup>+</sup>) AIM<sup>+</sup> CD4<sup>+</sup> T cell subsets.

### Measurements of anti-SARS-CoV-2 Abs and neutralizing capacity

Serum anti-SARS-CoV-2 Spike protein and receptor-binding domain (RBD) IgG, IgA, and IgM Abs were measured by a validated ELISA as previously described (50, 56), with assay cutoff 3 SD above the mean of a pre-COVID-19 population from the same geographic region. Ab neutralization capacity was assessed by cell culture assays with Vero E6 (ATCC CRL-1586) cells and live SARS-CoV-2, with data reported as geometric microneutralization titer at 50% (MNT<sub>50</sub>), which ranged from below detection (MNT<sub>50</sub> = 5; 1:10 dilution) to MNT<sub>50</sub> = 1280 (56). Ab neutralization was measured against the ancestral strain of SARS-CoV-2 and the β variant of concern (B.1.351). The β variant was obtained through BEI Resources (National Institute of Allergy and Infectious Diseases, National Institutes of Health: SARS-related coronavirus 2, isolate hCoV-19/South Africa/KRISP-K005325/2020, NR-54009, contributed by Alex Sigal and Tulio de Oliveira).

### Determination of prior SARS-CoV-2 infection

Due to demand for and limits on availability of testing, participants were not consistently tested for COVID-19 by a nasopharyngeal swab PCR assay, even when they were symptomatic. Serological testing for anti-nucleocapsid SARS-CoV-2 IgG and IgA Abs was performed on all collected samples by the ELISA described above (50, 56), using assay wells coated with 2 µg/ml nucleocapsid Ag (Jackson ImmunoResearch Laboratories). Participants were identified to have had COVID-19 when they were seropositive for IgG or IgA anti-nucleocapsid Abs and had a documented positive nasopharyngeal PCR test prior to any blood collection, as summarized in Table I.

### Statistical analysis

Statistical analyses were conducted using GraphPad Prism version 9 (GraphPad Software, San Diego, CA). Two-group comparisons of dose and CMV seropositivity or prior COVID-19 and CMV seropositivity were assessed by two-way ANOVA. Differences between CMV-seropositive and CMV-seronegative group Ab levels, Ab neutralization capacity, T cell immunophenotype, and T cell memory recall responses were assessed by a Student *t* test with Welch's correction or a Mann-Whitney *U* test, according to data normality. The *p* values are reported as two-tailed, and *p* values <0.05 were considered significant.

## Results

### Participant demographics

Serum anti-CMV IgG Abs were measured by ELISA, and 69.4% ( $n = 129/186$ ) of participants were found to be CMV seropositive. Age and sex distribution were similar between seropositive (median  $85 \pm 6.8$  y, 67.6% female) and seronegative (median  $86 \pm 7.7$  y, 69.8% female) participants (Table I). Blood samples were collected at a median of 179.8 d (CMV seronegative) and 175.4 d (CMV seropositive) after two doses of Moderna Spikevax (100  $\mu$ g; mRNA-1273) or Pfizer Comirnaty (30  $\mu$ g; BNT162b2) administered as per the manufacturer-recommended schedules. In Ontario, Canada, third dose vaccinations were recommended for older adults in congregate living beginning in August 2021 when they were  $>6$  mo after their second vaccinations (57). Participants received third doses in August–September 2021, and blood samples were collected at a median of 82.0 d (CMV seronegative) or 83.7 d (CMV seropositive) after third doses. Participants were classified as having had a previous SARS-CoV-2 infection when they had a documented positive PCR test and/or were seropositive for IgG or IgA nucleocapsid Abs. A positive nasopharyngeal PCR test and/or serum anti-nucleocapsid IgG or IgA Abs were reported in 37.8% of CMV-seronegative participants and 34.4% of CMV-seropositive participants.

### CMV seropositivity does not impede anti-SARS-CoV-2 Ab production or neutralization in older adults

To assess the effects of CMV serostatus on Ab responses after SARS-CoV-2 vaccination, serum anti-SARS-CoV-2 Spike and RBD IgG, IgA, and IgM Ab levels were measured by ELISA (Fig. 1, Table II). The number of responders (i.e., individuals with Abs above the threshold limit of detection) significantly increased between post-second and post-third dose measurements of anti-Spike IgG, IgA, and IgM Abs, as well as anti-RBD IgG Abs, but not anti-RBD IgA or IgM Abs (Table II). For example, 5–7 mo after second dose

vaccinations, anti-Spike IgG Abs were detected in 88.7% of participants, and anti-RBD IgG Abs were detected in 63.5% of participants. Approximately 3 mo after the third vaccine dose, 97.3 and 94.6% of participants had detectable anti-Spike IgG and anti-RBD IgG, respectively. CMV seropositivity did not impact the frequency of responders for anti-Spike and anti-RBD IgG, IgA, and IgM Abs. Accordingly, two-group analyses showed a main effect of vaccine dose, but not CMV serostatus, on serum anti-Spike IgG (Fig. 1A) and anti-RBD IgG (Fig. 1D) Abs, anti-Spike IgA (Fig. 1B) and anti-RBD IgA (Fig. 1E) Abs, as well as anti-Spike IgM (Fig. 1C) Abs. There were no significant main effects of vaccine dose or CMV serostatus on anti-RBD IgM (Fig. 1F) Abs. Therefore, CMV seropositivity does not impede maintenance of antiviral Abs several months after two or three doses of SARS-CoV-2 mRNA vaccines.

To examine potential effects of CMV serostatus on Ab function, serum Ab neutralization capacity was assessed by MNT<sub>50</sub> assays against live ancestral (wild-type) and  $\beta$  variant SARS-CoV-2 (Fig. 1G, 1H). Vaccines were designed against the wild-type virus, whereas the  $\beta$  variant contains mutations that confer increased transmissibility and immune evasion (58). Neutralization of ancestral and  $\beta$  variant SARS-CoV-2 ranged from below the detection limit to MNT<sub>50</sub> = 1280, although mean neutralization was consistently higher against the ancestral virus compared with the  $\beta$  variant after two and three vaccine doses. Neutralization capacity was similar between CMV-seropositive and CMV-seronegative individuals against both ancestral and  $\beta$  variant SARS-CoV-2, although there was a main effect of dose on neutralization capacity. In particular, significant increases in serum Ab neutralization were observed against the  $\beta$  variant between second and third dose vaccinations. Anti-Spike and anti-RBD IgG levels moderately correlated with Ab neutralization capacity, irrespective of CMV serostatus (Table III). As well, modest correlations were generally observed between ancestral SARS-CoV-2 neutralization capacity and anti-Spike and anti-RBD

Table I. Participant demographics


	CMV Seronegative ( $n = 74$ )	CMV Seropositive ( $n = 159$ )	Statistical Assessment
Age (y), median $\pm$ SD (range)	$85 \pm 6.8$ (65–93)	$86 \pm 7.7$ (65–101)	$p = 0.1265^a$
Sex (frequency)	67.6% Female ( $n = 50$ )	69.8% Female ( $n = 111$ )	$p = 0.7618^b$
Sample size postdose 2	$n = 51$	$n = 108$	—
Second dose vaccine combination (frequency)	47.1% Moderna-Moderna ( $n = 24$ ) 52.9% Pfizer-Pfizer ( $n = 27$ )	35.8% Moderna-Moderna ( $n = 39$ ) 60.6% Pfizer-Pfizer ( $n = 66$ ) 2.75% Pfizer-Moderna ( $n = 3$ )	$p = 0.2968^b$
Days between dose 1 and dose 2 (mean $\pm$ SD)	$32.2 \pm 20.8$	$30.4 \pm 17.9$	$p = 0.6697^a$
Days since second dose to blood collection (mean $\pm$ SD)	$179.8 \pm 50.1$	$175.4 \pm 64.1$	$p = 0.9041^a$
Sample size postdose 3 <sup>c</sup>	$n = 23$ (17 repeated)	$n = 51$ (30 repeated)	—
Third dose vaccine combination (frequency)	65.2% Moderna-Moderna-Moderna ( $n = 15$ ) 34.8% Pfizer-Pfizer-Pfizer ( $n = 8$ )	56.9% Moderna-Moderna-Moderna ( $n = 29$ ) 43.1% Pfizer-Pfizer-Pfizer ( $n = 22$ )	$p = 0.6115^b$
Days between dose 2 and dose 3 (mean $\pm$ SD)	$203.9 \pm 21.0$	$211.7 \pm 12.1$	$p = 0.2689^a$
Days since third dose to blood collection (mean $\pm$ SD)	$82.0 \pm 17.3$	$83.7 \pm 11.5$	$p = 0.7597^a$
Prior COVID-19 <sup>d</sup>	37.8% ( $n = 28$ )	34.4% ( $n = 55$ )	$p = 0.6609^a$
Only positive for nasopharyngeal PCR test	28.6% ( $n = 8$ )	36.4% ( $n = 20$ )	$p = 0.6244^b$
Only positive for anti- nucleocapsid Abs	39.3% ( $n = 11$ )	40.0% ( $n = 22$ )	$p > 0.9999^b$
Positive for PCR test and anti-nucleocapsid Abs	32.1% ( $n = 9$ )	23.6% ( $n = 13$ )	$p = 0.7977^b$

<sup>a</sup>Student  $t$  test with Welch's correction or Mann-Whitney  $U$  test by normality.

<sup>b</sup>Fisher's exact test. Comparisons for vaccines postdose 2 were calculated for Moderna-Moderna versus Pfizer-Pfizer.

<sup>c</sup>Forty-seven participants had blood drawn postdose 2 and postdose 3.

<sup>d</sup>Participants were considered positive for prior COVID-19 infection when they had a recorded positive nasopharyngeal PCR test at a date prior to blood collection and/or seropositivity for anti-nucleocapsid IgG or IgA Abs at time of blood collection for Ab and cellular assays.



**FIGURE 1.** Abs and neutralization capacity by CMV serostatus after two and three COVID-19 vaccines in older adults. Serum SARS-CoV-2 anti-Spike and anti-RBD Abs were detected by ELISA, and Ab neutralization capacity was assessed by MNT<sub>50</sub> with live SARS-CoV-2 virus, postdose 2 (PD2), and postdose 3 (PD3) vaccination in CMV-seronegative (–ve) and CMV-seropositive (+ve) individuals. **(A–C)** Anti-Spike Abs: IgG (A), IgA (B), and IgM (C). **(D–F)** Anti-RBD Abs: IgG (D), IgA (E), and IgM (F). **(G and H)** Ab neutralization capacity was assessed against ancestral (G) and β variant (H) SARS-CoV-2. CMV-PD2, *n* = 51; CMV-PD3, *n* = 23; CMV+PD2, *n* = 108; CMV+PD3, *n* = 51. Dotted lines indicate the threshold of detection. Each data point indicates an individual participant, with the center line at the median. Associations between CMV serostatus and vaccine dose were assessed by two-way ANOVA, with a Tukey’s test post hoc analysis of significant main effects. \**p* < 0.05, \*\**p* < 0.01, \*\*\**p* < 0.001, \*\*\*\**p* < 0.0001.

IgA Abs in both CMV-seropositive and CMV-seronegative participants. Therefore, CMV seropositivity does not compromise maintenance of vaccine-elicited Ab neutralization of SARS-CoV-2 several months after two or three vaccine doses.

As serum was collected from some participants after both second and third vaccine doses, these paired data were also assessed independently (Supplemental Fig. 1). Consistent with our pooled participant data, there were main effects of vaccine dose but not CMV

Table II. Responders to vaccination by CMV serostatus: Abs

Ab	Vaccine Doses	Total Responder Frequency			Responder Frequency by CMV Seropositivity				
		<i>n</i>	Frequency (%)	<i>p</i> (Fisher’s Exact Test)	Seronegative		Seropositive		<i>p</i> (Fisher’s Exact Test)
					<i>n</i>	Frequency (%)	<i>n</i>	Frequency (%)	
anti-Spike IgG	2	141/159	88.7	0.0415	47/51	92.2	94/108	87.0	0.4287
	3	72/74	97.3		21/23	91.3	51/51	100	
anti-Spike IgA	2	98/159	61.6	<0.0001	33/51	64.7	65/108	60.2	0.6053
	3	71/74	95.9		21/23	91.3	50/51	98.0	
anti-Spike IgM	2	20/159	12.6	<0.0001	8/51	15.7	12/108	11.1	0.4476
	3	28/74	37.8		8/23	34.8	20/51	39.2	
anti-RBD IgG	2	101/159	63.5	<0.0001	34/51	66.7	67/108	62.0	0.6011
	3	70/74	94.6		21/23	91.3	49/51	96.1	
anti-RBD IgA	2	35/159	22.0	0.0760	12/51	23.5	23/108	21.3	0.8380
	3	25/74	33.8		8/23	13.0	17/51	33.3	
anti-RBD IgM	2	6/159	3.77	>0.9999	3/51	5.88	3/108	2.78	0.3863
	3	3/74	4.05		1/23	4.35	2/51	3.92	

Table III. Associations of Ab and neutralization responses after vaccination by CMV serostatus

	Ab	MNT <sub>50</sub> Wild-Type		MNT <sub>50</sub> $\beta$ Variant	
		CMV-ve	CMV+ve	CMV-ve	CMV+ve
Postdose 2	IgG Spike	****	****	****	****
		$r = 0.8367$	$r = 0.8316$	$r = 0.6963$	$r = 0.7541$
	IgG RBD	****	****	****	****
		$r = 0.7394$	$r = 0.8291$	$r = 0.6848$	$r = 0.6981$
Postdose 2	IgA Spike	****	****	****	****
		$r = 0.6222$	$r = 0.6152$	$r = 0.6402$	$r = 0.5254$
	IgA RBD	**	****	**	****
		$r = 0.4311$	$r = 0.4572$	$r = 0.4813$	$r = 0.3639$
Postdose 3	IgG Spike	***	***	***	***
		$r = 0.6923$	$r = 0.5004$	$r = 0.6545$	$r = 0.5038$
	IgG RBD	****	****	****	****
		$r = 0.8207$	$r = 0.5770$	$r = 0.7475$	$r = 0.6926$
Postdose 3	IgA Spike	**	*	*	NS
		$r = 0.6324$	$r = 0.3228$	$r = 0.4550$	
	IgA RBD	****	*	****	*
		$r = 0.7750$	$r = 0.3081$	$r = 0.7051$	$r = 0.3575$

Spearman's correlation coefficient is reported. CMV-ve, CMV negative; CMV+ve, CMV positive. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$ .

serostatus on anti-Spike and anti-RBD IgG and IgA serum Ab levels. We did observe an interaction between CMV serostatus and vaccine dose on anti-RBD IgM measurements by intraindividual analysis, but most participants had Ab levels below the threshold. Intraindividual analyses also showed that the number of vaccine doses, but not CMV serostatus, had a significant effect on Ab neutralization capacity against wild-type and  $\beta$  variant SARS-CoV-2. Therefore, CMV seropositivity does not significantly impact intraindividual changes in Ab levels or neutralization capacity between two and three doses of SARS-CoV-2 mRNA vaccines.

We next considered postdose 2 and postdose 3 vaccination Ab measurements in the context of prior SARS-CoV-2 infection (Fig. 2). As summarized in Table I, the incidence of COVID-19 (prior to blood collections) was similar between CMV-seronegative and CMV-seropositive participants. We observed a main effect of prior SARS-CoV-2 infection on anti-Spike and anti-RBD IgG, IgA, and IgM Abs after two doses of mRNA vaccines, as well as anti-Spike and anti-RBD IgG and IgA, but not IgM, serum Abs after three vaccine doses. We observed an interaction between prior COVID-19 and CMV serostatus for anti-Spike and anti-RBD IgA Abs postdose 2 and postdose 3, and IgM Abs postdose 2. There was a main effect of CMV serostatus on anti-Spike IgM Abs, but most individuals had levels below the detection threshold. We in addition observed main effects of prior SARS-CoV-2 infection, but not CMV serostatus, on Ab neutralization of ancestral SARS-CoV-2 after two and three vaccine doses, and the  $\beta$  variant after two, but not three, vaccine doses. Collectively, these data indicate that CMV serostatus does not appear to have a major impact on the longevity of circulating anti-Spike or anti-RBD IgG Abs, or total serum Ab neutralization capacity, after SARS-CoV-2 infection or vaccination.

#### *CMV serostatus influences peripheral CD4<sup>+</sup> and CD8<sup>+</sup> T cell immunophenotype in older adults*

To examine the impact of CMV seropositivity on the T cell repertoire, whole-blood CD4<sup>+</sup> and CD8<sup>+</sup> T cell composition was quantitated and the surface expression of CD28 and CD57 was measured by flow cytometry (Fig. 3, Supplemental Fig. 2). Chronic T cell activation and an altered T cell repertoire are characteristics of CMV seropositivity (20, 38, 59). Accordingly, CMV-seropositive individuals had significant changes to their peripheral blood T cell composition. We found no changes in numbers of circulating total leukocytes, total CD4<sup>+</sup> T cells, or CD4<sub>N</sub>, CD4<sub>EM</sub>, CD8<sub>N</sub>, or CD8<sub>EM</sub> T cell populations by CMV serostatus. However, CMV seropositivity

increased numbers of total CD8<sup>+</sup> T cells, as well as CD4<sub>EMRA</sub>, CD4<sub>TD</sub>, CD8<sub>EMRA</sub>, and CD8<sub>TD</sub> T cells, and decreased numbers of CD4<sub>CM</sub> and CD8<sub>CM</sub> T cells.


CD28 is a costimulatory molecule that contributes to TCR Ag-mediated activation of T cells, whereas CD57 is a marker of TD T cells as well as an indicator of immune senescence (60). Repeated T cell activation is associated with upregulation of CD57 and a reduction in CD28 expression (61–63). Consistent with these prior data, comparisons of CD28 and CD57 expression on T cell populations by CMV serostatus in our cohort of older adults (Fig. 3O) revealed increased CD57 expression and reduced CD28 expression on total CD4<sup>+</sup> and CD8<sup>+</sup> T cell populations, as well as more specifically CD4<sub>CM</sub>, CD4<sub>EM</sub>, CD4<sub>EMRA</sub>, CD8<sub>N</sub>, and CD8<sub>EMRA</sub> T cells, in CMV-seropositive individuals. Expression of CD28 was also decreased on CD8<sub>CM</sub> and CD8<sub>EM</sub> cells of CMV-seropositive individuals, although their expression of CD57 was not influenced by CMV serostatus. CMV serostatus did not alter CD57 or CD28 expression on CD4<sub>N</sub> T cells.

As even mild COVID-19 can have lasting effects on immune cell composition (50), we also considered combined effects of prior SARS-CoV-2 infection and CMV serostatus on T cell composition (Fig. 4, Supplemental Fig. 2). Prior SARS-CoV-2 infection was associated with lower total leukocyte counts, and an interaction was observed between CMV serostatus and prior COVID-19 that influenced the numbers of CD8<sub>TD</sub> cells. Otherwise, there were no significant main effects of prior SARS-CoV-2 infection on absolute cell numbers nor the prevalence of the assessed CD4<sup>+</sup> or CD8<sup>+</sup> T cell populations.


In summary, there are significant changes to the relative composition and phenotype of peripheral blood CD8<sup>+</sup> T cell and CD4<sup>+</sup> T cell subsets between CMV-seronegative and CMV-seropositive older adults, irrespective of prior COVID-19. The observed expansion of EMRA and TD T cells, as well as reduced surface expression of the costimulatory molecule CD28 on CD8<sub>N</sub> T cells in particular, may influence vaccine-specific T cell responses.

#### *CMV serostatus influences CD4<sup>+</sup> and CD8<sup>+</sup> T cell SARS-CoV-2 Ag-induced recall responses in older adults*


An AIM assay was used to examine T cell memory responses by stimulation with the SARS-CoV-2 Spike Ag after second and third dose SARS-CoV-2 mRNA vaccinations (Fig. 5, Table IV). SARS-CoV-2 vaccines are unusual in that healthy adults generate strong CD4<sup>+</sup> T cell memory recall responses, but weaker CD8<sup>+</sup> T cell



**FIGURE 2.** Abs and neutralization capacity by CMV serostatus and prior SARS-CoV-2 infection after two and three COVID-19 vaccines in older adults. SARS-CoV-2 anti-Spike and anti-RBD Abs were measured in serum of CMV-seronegative (CMV-) and CMV-seropositive (CMV+) individuals by ELISA, and serum Ab neutralization capacity was assessed by MNT<sub>50</sub> with live SARS-CoV-2 virus. Data are stratified by prior SARS-CoV-2 infection history. (A–F) Post-dose 2: anti-Spike IgG (A), IgA (B), and IgM (C) Abs, and anti-RBD IgG (D), IgA (E), and IgM (F) Abs. (G–L) Postdose 3: anti-Spike IgG (G), IgA (H) and IgM (I) Abs, and anti-RBD IgG (J), IgA (K), and IgM (L) Abs. (M and N) Postdose 2 SARS-CoV-2 neutralization: ancestral (M) and β variant (N). (O and P) Postdose 3 SARS-CoV-2 neutralization: ancestral (O) and β variant (P). PD2: CMV–No, n = 33; CMV–Yes, n = 18; CMV+No, n = 77; CMV+Yes, n = 31. PD3: CMV–No, n = 13; CMV–Yes, n = 10; CMV+No, n = 27; CMV+Yes, n = 24. Dotted lines indicate the threshold of detection. Each data point indicates an individual participant, with the center line at the median. Associations between CMV serostatus and prior COVID-19 were assessed by two-way ANOVA, with a Tukey’s test post hoc analysis of significant main effects and interactions. \**p* < 0.05, \*\**p* < 0.01, \*\*\**p* < 0.001, \*\*\*\**p* < 0.0001.




**FIGURE 3.** Effect of CMV serostatus on the circulating T cell repertoire in older adults. T cell populations in whole blood were assessed by flow cytometry in CMV-seronegative (-ve) and CMV-seropositive (+ve) individuals. **(A)** Relative prevalence of CD4<sup>+</sup> and CD8<sup>+</sup> T cell subsets by CMV serostatus (also see Supplemental Fig. 2). **(B–D)** Absolute cell counts of **(B)** total leukocytes, **(C)** total CD4<sup>+</sup> T cells, and **(D)** CD8<sup>+</sup> T cells. **(E–I)** Absolute cell counts of CD4<sup>+</sup> T cells: **(E)** naive, **(F)** central memory, **(G)** effector memory, **(H)** EMRA, and **(I)** terminally differentiated. **(J–N)** Absolute cell counts of CD8<sup>+</sup> T cells: **(J)** naive, **(K)** central memory, **(L)** effector memory, **(M)** EMRA, and **(N)** terminally differentiated. **(O)** Surface geometric mean expression of CD57 and CD28 on CD4<sup>+</sup> and CD8<sup>+</sup> T cells by CMV serostatus. Blood was assessed either postdose 2 or postdose 3 for each participant. CMV-ve, *n* = 56; CMV+ve, *n* = 128. Each data point in **(B)–(N)** indicates an individual participant, and data are presented as box-and-whisker plots, minimum to maximum, with the center line at the median. The surface marker expression in **(O)** was visualized by concatenating uncompensated events in FlowJo for each participant and indicated T cell population grouped according to CMV serostatus, and then geometric mean fluorescence intensity expression data of each CMV group were overlaid onto the same histogram plot. Associations between T cell subsets and CMV serostatus were assessed by a Student *t* test with Welch's correction or a Mann-Whitney *U* test, according to normality. \**p* < 0.05, \*\**p* < 0.01, \*\*\**p* < 0.001, \*\*\*\**p* < 0.0001.



**FIGURE 4.** Effects of CMV serostatus and prior COVID-19 on the circulating T cell repertoire in older adults. T cell populations in whole blood were assessed by flow cytometry in CMV-seronegative (CMV<sup>-</sup>) and CMV-seropositive (CMV<sup>+</sup>) individuals. Data are stratified by prior SARS-CoV-2 infection history. (A and B) Absolute cell counts of (A) total leukocytes and (B) total T cells. (C–H) Absolute cell counts of CD4<sup>+</sup> T cells: (C) total, (D) naïve, (E) central memory, (F) effector memory, (G) EMRA, and (H) terminally differentiated. (I–N) Absolute cell counts of CD8<sup>+</sup> T cells: (I) total, (J) naïve, (K) central memory, (L) effector memory, (M) EMRA, and (N) terminally differentiated. Blood was assessed either postdose 2 or postdose 3 for each participant. CMV–No,  $n = 35$ ; CMV–Yes,  $n = 21$ ; CMV+No,  $n = 84$ ; CMV+Yes,  $n = 44$ . Each data point indicates an individual participant. Data are presented as box-and-whisker plots, minimum to maximum, with the center line at the median. Associations between CMV serostatus and prior COVID-19 were assessed by two-way ANOVA, with a Tukey’s test post hoc analysis of significant main effects and interactions. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$ .





**FIGURE 5.** CD4<sup>+</sup> and CD8<sup>+</sup> T cell AIM responses to SARS-CoV-2 Spike in older adults. T cell memory responses to SARS-CoV-2 Spike were assessed by AIM assay in CMV-seronegative (CMV<sup>-</sup>) and CMV-seropositive (CMV<sup>+</sup>) individuals postdose 2 (PD2) and postdose 3 (PD3) SARS-CoV-2 vaccination. **(A)** AIM<sup>+</sup>CD8<sup>+</sup> T cells (expressing CD69 and CD137) were measured as a proportion of total CD8<sup>+</sup> T cells. **(B)** AIM<sup>+</sup>CD4<sup>+</sup> T cells (expressing CD25 and OX40) were measured as a proportion of total CD4<sup>+</sup> T cells. **(C–E)** AIM<sup>+</sup>CD4<sup>+</sup> T cell Th1 (C), Th2 (D), and Th17 (E) subsets. Each data point indicates an individual participant. For (A) and (B), data from all individuals are graphed irrespective of whether they meet cutoff requirements for a “positive” result. For (C)–(E), only data from individuals with positive Spike-AIM<sup>+</sup>CD4<sup>+</sup> T cell memory recall responses were graphed (Table IV). CMV–PD2, *n* = 51; CMV–PD3, *n* = 23; CMV+PD2, *n* = 108; CMV+PD3, *n* = 51. Data are presented as box-and-whisker plots, minimum to maximum, with the center line at the median. Associations between CMV serostatus and vaccine dose were assessed by two-way ANOVA, with a Tukey’s test post hoc analysis of significant main effects. \**p* < 0.05, \*\*\**p* < 0.001.

memory responses (64). We also made similar observations in older adults. Most study participants had CD4<sup>+</sup> T cell responses to SARS-CoV-2 Spike (postdose 2, 93.1%; postdose 3, 95.9%), but only 19.5% of participants had Spike-elicited CD8<sup>+</sup> memory T cell responses after two vaccine doses, although this increased to 29.7% of participants after three vaccine doses (Table IV). CMV serostatus did not influence the number of individuals with SARS-CoV-2 Spike-AIM<sup>+</sup>CD4<sup>+</sup> T cells or Spike-AIM<sup>+</sup>CD8<sup>+</sup> T cells after second or third dose vaccinations. Grouped analyses revealed a significant main effect of CMV serostatus on the frequency of Spike-AIM<sup>+</sup>CD8<sup>+</sup> T cells (Fig. 5A). However, despite greater variance of data in CMV-seropositive individuals, post hoc analyses by CMV serostatus were not significant postdose 2 or postdose 3. Intraindividual paired analyses also showed no main effects of vaccine dose or CMV serostatus on activation of Spike-AIM<sup>+</sup>CD8<sup>+</sup> T cells (Supplemental Fig. 3A). Prior COVID-19 did not influence the prevalence of Spike-AIM<sup>+</sup>CD8<sup>+</sup> T cells after two or three vaccine doses, although CMV seropositivity contributed to increased Spike-AIM<sup>+</sup>CD8<sup>+</sup> T cell activation postdose 2 (Fig. 6). Grouped analyses on a population and intraindividual basis showed a main effect of vaccine dose (Fig. 5B, Supplemental Fig. 3B), but the prevalence of Spike-AIM<sup>+</sup>CD4<sup>+</sup> T cells was likewise not different by CMV serostatus (Fig. 5B) or by prior COVID-19 (Fig. 6B, 6G). Therefore, CMV serostatus


contributes to increased CD8<sup>+</sup> T cell, but not CD4<sup>+</sup> T cell, memory recall responses to the SARS-CoV-2 Spike protein several months after second and third dose vaccinations.

CD4<sup>+</sup> memory T cells are comprised of a number of different functional subsets, including Th1, Th2, and Th17 cells (65), which were further characterized. There was a significant effect of CMV serostatus on the frequency of Th1 Spike-AIM<sup>+</sup>CD4<sup>+</sup> T cells, with post hoc analyses showing an increase in CMV-seropositive individuals after two but not three vaccine doses (Fig. 5C, Supplemental Fig. 3C). Th2 and Th17 Spike-AIM<sup>+</sup>CD4<sup>+</sup> T cell frequencies were not influenced by CMV serostatus (Fig. 5D, 5E), although paired analyses by dose and CMV serostatus suggested a significant increase in Th17 responses in CMV-seropositive individuals after third vaccine doses (Supplemental Fig. 3E). Interestingly, when we considered these data in the context of prior SARS-CoV-2 infection, CMV serostatus had a main effect on the prevalence of Spike-AIM<sup>+</sup>CD4<sup>+</sup> Th1 and Th17 T cells, which increased and decreased, respectively, with CMV seropositivity, although only postdose 2 (Fig. 6A, 6E), suggesting dose-dependent modulation of vaccine-associated T cell memory responses by CMV serostatus.

To determine whether the observed contributions of CMV serostatus to increased AIM<sup>+</sup>CD8<sup>+</sup> T cell and AIM<sup>+</sup>CD4<sup>+</sup> Th1 T cell

Table IV. Responders to vaccination by CMV serostatus: SARS-CoV-2 Spike T cell memory recall responses


T Cell Population	Vaccine Doses	Total Responder Frequency				Responder Frequency by CMV Seropositivity				
		<i>n</i>	Frequency (%)	<i>p</i> (Fisher’s Exact Test)	Seronegative		Seropositive		<i>p</i> (Fisher’s Exact Test)	
					<i>n</i>	Frequency (%)	<i>n</i>	Frequency (%)		
Spike-AIM <sup>+</sup> CD8 <sup>+</sup>	2	31/159	19.5	0.0943	6/50	12.0	24/108	22.2	0.1898	
	3	22/74	29.7		3/23	13.0	19/51	37.3	0.0532	
Spike-AIM <sup>+</sup> CD4 <sup>+</sup>	2	148/159	93.1	0.5570	45/50	90.0	102/108	94.4	0.3262	
	3	71/74	95.9		22/23	95.7	49/51	96.1	>0.9999	



**FIGURE 6.** CD4<sup>+</sup> and CD8<sup>+</sup> T cell AIM responses to SARS-CoV-2 Spike after COVID-19 in older adults by CMV serostatus. T cell memory responses to SARS-CoV-2 Spike were assessed by AIM assay in CMV-seronegative (CMV<sup>-</sup>) and CMV-seropositive (CMV<sup>+</sup>) individuals. Data are stratified by prior SARS-CoV-2 infection history. **(A–E)** Postdose 2: (A) AIM<sup>+</sup>CD8<sup>+</sup> T cells (expressing CD69 and CD137) were measured as a proportion of total CD8<sup>+</sup> T cells; (B) AIM<sup>+</sup>CD4<sup>+</sup> T cells (expressing CD25 and OX40) were measured as a proportion of total CD4<sup>+</sup> T cells; AIM<sup>+</sup>CD4<sup>+</sup> T cell Th1 (C), Th2 (D), and Th17 (E) subsets. **(F–I)** Postdose 3: (F) AIM<sup>+</sup>CD8<sup>+</sup> T cells as a proportion of total CD8<sup>+</sup> T cells; (G) AIM<sup>+</sup>CD4<sup>+</sup> T cells as a proportion of total CD4<sup>+</sup> T cells; AIM<sup>+</sup>CD4<sup>+</sup> T cell Th1 (H), Th2 (I), and Th17 (J) subsets. For (C)–(E) and (H)–(J), only data from individuals with positive Spike-AIM<sup>+</sup>CD4<sup>+</sup> T cell memory recall responses were graphed (Table IV). PD2: CMV<sup>-</sup>No,  $n = 33$ ; CMV<sup>-</sup>Yes,  $n = 18$ ; CMV<sup>+</sup>No,  $n = 77$ ; CMV<sup>+</sup>Yes,  $n = 31$ . PD3: CMV<sup>-</sup>No,  $n = 13$ ; CMV<sup>-</sup>Yes,  $n = 10$ ; CMV<sup>+</sup>No,  $n = 27$ ; CMV<sup>+</sup>Yes,  $n = 24$ . Each data point indicates an individual participant. Data are presented as box-and-whisker plots, minimum to maximum, with the center line at the median. Associations between CMV serostatus and prior COVID-19 were assessed by two-way ANOVA, with a Tukey’s test post hoc analysis of significant main effects.

memory responses are consistent across different stimuli, we also examined T cell AIM memory responses after TCR-independent polyclonal stimulation with CytoStim and stimulation with influenza HA Ags (Fig. 7). As we observed for SARS-CoV-2 Spike-activated memory T cells, both CytoStim and HA stimulation resulted in an increased prevalence of AIM<sup>+</sup>CD8<sup>+</sup> T cells in CMV-seropositive individuals (Fig. 7A, 7F), although AIM<sup>+</sup>CD4<sup>+</sup> T cell frequency was not affected by CMV serostatus (Fig. 7B, 7G). Prior COVID-19 did not influence the frequencies of AIM<sup>+</sup> T cells (Supplemental Fig. 4). These data show that the effects of CMV seropositivity

on AIM<sup>+</sup>CD8<sup>+</sup> and AIM<sup>+</sup>CD4<sup>+</sup> T cell frequencies are consistent across different stimuli. CytoStim-stimulated CD4<sup>+</sup> T cells, similar to Spike-stimulated CD4<sup>+</sup> T cells, also showed distinct Th1-skewed AIM<sup>+</sup>CD4<sup>+</sup> T cell responses, although this was not observed after HA stimulation. These data collectively suggest that CMV serostatus alters the T cell response to polyclonal activation (i.e., with CytoStim) and has differential effects on the recall response of memory T cells to specific Ags (i.e., influenza HA or SARS-CoV-2 Spike). However, CMV serostatus does not alter the ability of older adults to generate lasting CD4<sup>+</sup> or CD8<sup>+</sup> T cell memory, nor the incidence of



**FIGURE 7.** CD4<sup>+</sup> and CD8<sup>+</sup> T cell AIM responses to CytoStim and influenza HA in older adults. AIM assays were assessed by flow cytometry analysis after whole-blood stimulation. **(A–E)** CytoStim-induced responses: (A) AIM<sup>+</sup>CD8<sup>+</sup> T cells (expressing CD69 and CD137) as a proportion of total CD8<sup>+</sup> T cells; (B) AIM<sup>+</sup>CD4<sup>+</sup> T cells (expressing CD25 and OX40) as a proportion of total CD4<sup>+</sup> T cells; AIM<sup>+</sup>CD4<sup>+</sup> T cell Th1 (C), Th2 (D), and Th17 (E) subsets. **(F–J)** Influenza HA-induced responses: (F) AIM<sup>+</sup>CD8<sup>+</sup> T cells as a proportion of total CD8<sup>+</sup> T cells; (G) AIM<sup>+</sup>CD4<sup>+</sup> T cells as a proportion of total CD4<sup>+</sup> T cells; AIM<sup>+</sup>CD4<sup>+</sup> T cell Th1 (H), Th2 (I), and Th17 (J) subsets. Each data point indicates an individual participant. Data were pooled from all blood collections, and for participants with two different collection time points, data are only included from postdose 2 assessments. CMV–,  $n = 57$ ; CMV+,  $n = 130$ . For (C)–(E), only data from individuals with >5% CytoStim AIM<sup>+</sup>CD4<sup>+</sup> T cells were graphed. For (H)–(J), only data from individuals with >0% influenza HA AIM<sup>+</sup>CD4<sup>+</sup> T cells were graphed. Data are presented as box-and-whisker plots, minimum to maximum, with the center line at the median. Associations between T cell responses and CMV serostatus were assessed by a Student *t* test with Welch's correction or a Mann–Whitney *U* test, according to normality. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$ .

T cell memory recall activation in response to SARS-CoV-2 Spike protein after mRNA vaccination.

## Discussion

Our data suggest that despite being a significant modifier of peripheral blood T cell composition and phenotype, CMV seropositivity in older adults does not have a negative impact on the longevity of vaccine-elicited Ab quantity and quality, or T cell memory recall responses, several months after second or third doses of SARS-CoV-2 mRNA vaccines. Yet, we found that there were subtle changes in Ab and cellular responses in CMV-seropositive individuals between vaccine doses and in individuals with prior COVID-19. Our study cohort included participants from multiple assisted living facilities, and it did not exclude individuals with particular health conditions (e.g., cancer, diabetes, cardiovascular disease, autoimmune disorders) or prescribed medications (e.g., immune-modulating drugs). Thus, any observed effects of CMV needed to be sufficiently robust to overcome potential effects of those other factors.

Our observations of changes in the peripheral T cell repertoire in CMV-seropositive individuals, and CD4<sup>+</sup> and CD8<sup>+</sup> T cell expression of CD28 and CD57, are consistent with prior publications that reported expansion of exhausted CD4<sub>EMRA</sub> and CD8<sub>EMRA</sub> T cells in CMV-seropositive healthy community-dwelling adults (41, 43). There are conflicting reports as to whether CMV seropositivity is associated with a reduction in naive CD4<sup>+</sup> and CD8<sup>+</sup> T cells. In this investigation, we observed similar numbers of CD4<sub>N</sub> and CD8<sub>N</sub> T cells in seropositive and seronegative individuals, and in particular similar expression of CD57 and CD28 on CD4<sub>N</sub> T cells. Our data therefore suggest that CMV seropositivity does not influence the availability or capacity of circulating naive T cells to respond and

generate memory responses to Ags from novel viruses such as SARS-CoV-2 in older adults.

CMV seropositivity in older adults has been associated with lower frequencies of memory T cells in response to seasonal influenza, although acute infection T cell responses were unchanged (66), and there are conflicting data as to whether CMV seropositivity enhances or impairs influenza virus-specific T cell responses (33, 41, 67, 68). We found that T cell memory recall responses to influenza HA, and moreover SARS-CoV-2 Spike, were similar in CMV-seropositive and CMV-seronegative individuals. Our findings are concordant with observations from a previous study that found CMV serostatus did not alter the ability of older adults to generate memory responses to the (at the time) newly emergent West Nile virus (48). Furthermore, it has been reported that T cell memory responses to the SARS-CoV-2 Spike protein are boosted in convalescent younger adults after vaccination (69, 70). We identified an increase in Spike-specific CD4<sup>+</sup> T cell memory responses between two and three mRNA vaccine doses in our older adult cohort, but we did not observe increased CD8<sup>+</sup> or CD4<sup>+</sup> T cell memory responses in convalescent older adults after vaccination. This may in part be because our analyses of effects of prior COVID-19 were not restricted to a particular time frame after SARS-CoV-2 infection. Importantly, we acknowledge that there may be early effects of CMV seropositivity on the initial generation of Abs or cellular memory that are not apparent at the assessed time points in this study, which were several months since vaccination. Irrespective, these findings suggest that although combined effects of infection and subsequent vaccination may differ by age, older adults can elicit memory T cell responses to infection and vaccination against newly emergent viruses such as SARS-CoV-2.

We observed a distinct Th1 bias after polyclonal T cell activation and in response to the immunodominant regions of the Spike Ag after two doses, but not three doses, of SARS-CoV-2 mRNA vaccines. Th1 skewing of the cellular immune response after influenza virus vaccination has been previously noted in CMV-seropositive infants and young adults and mice (68, 71), as well as in older adults (72). Strong Th1 CD4<sup>+</sup> T cell responses have been associated with lower disease severity in unvaccinated COVID-19 patients (73), and SARS-CoV-2 Spike-elicited CD4<sup>+</sup> T cell memory responses in unvaccinated convalescent individuals have also been identified to be primarily Th1 differentiated (69, 70). However, we did not identify a main effect of prior COVID-19, or an interaction of CMV serostatus and prior COVID-19, on the prevalence of CD4<sup>+</sup> T cell Th1 responses in vaccinated older adults. These data indicate that in older adults CMV seropositivity is associated with Th1-biased CD4<sup>+</sup> T cell responses, which are not further modified by prior SARS-CoV-2 infection. Our observations also suggest that Spike-specific T cell memory recall responses change between two and three vaccine doses in older adults, congruent with observations of changes in memory T cell phenotype between vaccine doses in younger adults (69).

Our data also show that CMV seropositivity does not prevent production of anti-Spike or anti-RBD IgG, IgA, or IgM Abs after SARS-CoV-2 mRNA vaccination, although we did observe interaction effects between CMV seropositivity and prior COVID-19 both postdose 2 and postdose 3 for anti-Spike and anti-RBD IgA Abs, and postdose 2 for IgM Abs. These interactions may be reflective of our limited sample size or differences in time since infection between CMV-seropositive and CMV-seronegative individuals. CMV-seropositive individuals have been reported to have increased B cell proliferation and mutations within the IgH sequences of IgM and IgG, but not IgA, isotypes (74). CMV serostatus could also have a larger effect on maturation of the Ab response via isotype switching, and thus isotype composition after viral infection, which may contribute to our observations, but to our knowledge this has not been extensively explored. Furthermore, our observations are from individuals vaccinated with mRNA vaccines, which were predominately used in older adults in assisted living facilities in Ontario, Canada. The effects of CMV-associated immune dysfunction on the quality and durability of humoral and cellular immune responses after vaccination may be ameliorated with mRNA vaccines, but other vaccine platforms (e.g., inactivated or attenuated whole virus, viral vector, or protein subunit) may have differential outcomes, which should be investigated in future studies.

Early in the pandemic, CMV seropositivity was associated with increased risk of hospitalization in COVID-19 patients (75), and CMV reactivation was later reported to have a higher incidence in patients in intensive care (76). More recently it was reported that unvaccinated individuals with latent CMV, irrespective of anti-CMV Ab levels, age, and sex, are at higher risk of SARS-CoV-2 infection and hospitalization (77). In particular, the exhausted T cells present in CMV-seropositive individuals have been predicted to contribute to more severe COVID-19 pathophysiology (78). Our data do not preclude the possibility that CMV-associated remodeling of innate and adaptive immunity in older adults may contribute to the pathogenesis and severity of SARS-CoV-2 infection. CMV serostatus may in addition impact humoral or cellular protection after vaccination against breakthrough infections with current or emerging variants of concern.

In conclusion, our data show that CMV serostatus alters the T cell repertoire but does not blunt durability of vaccine-elicited cellular memory responses or humoral responses after two and three doses of SARS-CoV-2 mRNA vaccines in older adults in assisted living facilities. Further research is necessary to disentangle the more subtle

effects of CMV serostatus on immunity after vaccination, as well as to assess its role in risk of breakthrough SARS-CoV-2 infections.

## Acknowledgments

Data in this study were collected by the COVID-in-LTC Study Group. We acknowledge administrative and technical assistance from Tara Kajaks, Ahmad Rahim, Megan Hagerman, Braeden Cowbrough, Komal Aryal, Leslie Tan, Sussan Kianpour, Jodie Turner, and Sheneice Joseph of the COVID-in-LTC Study Group at McMaster University. We thank our participants and their families, as well as staff, for support of this study.

## Other Members of the COVID-in-LTC Study Group

Eric D. Brown,<sup>1,2</sup> Gerry Wright,<sup>1,2</sup> David C. Bulir,<sup>3</sup> Mark Loeb,<sup>2,3,4,5</sup> Marek Smieja,<sup>3,4,6</sup> Aaron Jones,<sup>5,7</sup> Parminder Raina,<sup>5,8</sup> Chris Verschoor,<sup>5,9,10</sup> Janet E. McElhaney (posthumous),<sup>9,10</sup> Kevin Brown,<sup>11,12</sup> George A. Heckman,<sup>13,14</sup> John P. Hirdes,<sup>13,14</sup> Michael P. Hillmer,<sup>15,16</sup> Ahmad Von Schlegell,<sup>17,18</sup> Nathan M. Stall,<sup>19,20</sup> Kevin Stinson,<sup>21</sup> and Arthur Sweetman<sup>22</sup>.

<sup>1</sup>Department of Biochemistry & Biomedical Sciences, McMaster University, Hamilton, Ontario, Canada. <sup>2</sup>Michael G. DeGroot Institute for Infectious Disease Research, McMaster University, Hamilton, Ontario, Canada. <sup>3</sup>Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Ontario, Canada. <sup>4</sup>Department of Medicine, Michael G. DeGroot School of Medicine, McMaster University, Hamilton, Ontario, Canada. <sup>5</sup>Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada. <sup>6</sup>St. Joseph's Healthcare Hamilton, Hamilton, Ontario, Canada. <sup>7</sup>Institute for Clinical Evaluative Sciences (ICES), Toronto, Ontario, Canada. <sup>8</sup>McMaster Institute for Research on Aging, McMaster University, Hamilton, Ontario, Canada. <sup>9</sup>Health Science North Research Institute, Sudbury, Ontario, Canada. <sup>10</sup>Northern Ontario School of Medicine, Sudbury, Ontario, Canada. <sup>11</sup>Dalla Lana School of Public Health, University of Toronto, Ontario, Canada. <sup>12</sup>Public Health Ontario, Ontario, Canada. <sup>13</sup>School of Public Health and Health Systems, University of Waterloo, Waterloo, Ontario, Canada. <sup>14</sup>Schlegel Research Institute for Aging, Waterloo, Ontario, Canada. <sup>15</sup>Ontario Ministry of Health, Toronto, Canada. <sup>16</sup>Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Canada. <sup>17</sup>Schlegel Villages, Ontario, Canada. <sup>18</sup>Trillium Health Partners, Ontario, Canada. <sup>19</sup>Department of Medicine, Division of General Internal Medicine and Geriatrics, University of Toronto, Toronto, Ontario, Canada. <sup>20</sup>Sinai Health, Toronto, Ontario, Canada. <sup>21</sup>St. Mary's General Hospital, Kitchener, Ontario, Canada. <sup>22</sup>Department of Economics, McMaster University, Hamilton, Ontario, Canada.

## Disclosures

The authors have no financial conflicts of interest.

## References

- Kline, K. A., and D. M. Bowdish. 2016. Infection in an aging population. *Curr. Opin. Microbiol.* 29: 63–67.
- Lord, J. M. 2013. The effect of ageing of the immune system on vaccination responses. *Hum. Vaccin. Immunother.* 9: 1364–1367.
- Banerjee, A., L. Pasea, S. Harris, A. Gonzalez-Izquierdo, A. Torralbo, L. Shallcross, M. Noursadeghi, D. Pillay, N. Sebire, C. Holmes, et al. 2020. Estimating excess 1-year mortality associated with the COVID-19 pandemic according to underlying conditions and age: a population-based cohort study. *Lancet* 395: 1715–1725.
- Cunningham, A. L., H. Lal, M. Kovac, R. Chlibek, S.-J. Hwang, J. Díez-Domingo, O. Godeaux, M. J. Levin, J. E. McElhaney, J. Puig-Barberà, et al.; ZOE-70 Study Group. 2016. Efficacy of the herpes zoster subunit vaccine in adults 70 years of age or older. *N. Engl. J. Med.* 375: 1019–1032.
- Lal, H., A. L. Cunningham, O. Godeaux, R. Chlibek, J. Díez-Domingo, S. J. Hwang, M. J. Levin, J. E. McElhaney, A. Pöder, J. Puig-Barberà, et al.; ZOE-50 Study Group. 2015. Efficacy of an adjuvanted herpes zoster subunit vaccine in older adults. *N. Engl. J. Med.* 372: 2087–2096.
- Okoli, G. N., F. Racovitan, T. Abdulwahid, C. H. Righolt, and S. M. Mahmud. 2021. Variable seasonal influenza vaccine effectiveness across geographical regions, age groups and levels of vaccine antigenic similarity with circulating virus strains: a systematic review and meta-analysis of the evidence from test-negative design studies after the 2009/10 influenza pandemic. *Vaccine* 39: 1225–1240.

7. Rondy, M., N. El Omeiri, M. G. Thompson, A. Levêque, A. Moren, and S. G. Sullivan. 2017. Effectiveness of influenza vaccines in preventing severe influenza illness among adults: a systematic review and meta-analysis of test-negative design case-control studies. *J. Infect.* 75: 381–394.
8. Brown, K., N. M. Stall, T. Vanniyasingam, S. A. Buchan, N. Daneman, M. P. Hillmer, J. Hopkins, J. Johnstone, A. Maltsev, A. McGeer, et al. Early impact of Ontario's COVID-19 vaccine rollout on long-term care home residents and health care workers. Available at: <https://doi.org/10.47326/ocsat.2021.02.13.1.0>. Accessed: March 19, 2022.
9. Salcher-Konrad, M., S. Smith, and A. Comas-Herrera. 2021. Emerging evidence on effectiveness of COVID-19 vaccines among residents of long-term care facilities. *J. Am. Med. Dir. Assoc.* 22: 1602–1603.
10. Chung, H., S. He, S. Nasreen, M. E. Sundaram, S. A. Buchan, S. E. Wilson, B. Chen, A. Calzavara, D. B. Fell, P. C. Austin, et al.; Canadian Immunization Research Network (CIRN) Provincial Collaborative Network (PCN) Investigators. 2021. Effectiveness of BNT162b2 and mRNA-1273 covid-19 vaccines against symptomatic SARS-CoV-2 infection and severe covid-19 outcomes in Ontario, Canada: test negative design study. *BMJ* 374: n1943.
11. Breznik, J. A., A. Zhang, A. Huynh, M. S. Miller, I. Nazy, D. M. E. Bowdish, and A. P. Costa. 2021. Antibody responses 3–5 months post-vaccination with mRNA-1273 or BNT163b2 in nursing home residents. *J. Am. Med. Dir. Assoc.* 22: 2512–2514.
12. Zhang, A., J. A. Breznik, R. Clare, I. Nazy, M. S. Miller, D. M. E. Bowdish, and A. P. Costa. 2022. Antibody responses to third-dose mRNA vaccines in nursing home and assisted living residents. *J. Am. Med. Dir. Assoc.* 23: 444–446.
13. Brockman, M. A., F. Mwimanzi, H. R. Lapointe, Y. Sang, O. Agafitei, P. K. Cheung, S. Ennis, K. Ng, S. Basra, L. Y. Lim, et al. 2022. Reduced magnitude and durability of humoral immune responses to COVID-19 mRNA vaccines among older adults. *J. Infect. Dis.* 225: 1129–1140.
14. Weng, N. P., and G. Pawelec. 2021. Validation of the effectiveness of SARS-CoV-2 vaccines in older adults in “real-world” settings. *Immun. Ageing* 18: 36.
15. Fulop, T., A. Larbi, G. Dupuis, A. Le Page, E. H. Frost, A. A. Cohen, J. M. Witkowski, and C. Franceschi. 2018. Immunosenescence and inflamm-aging as two sides of the same coin: friends or foes? *Front. Immunol.* 8: 1960.
16. Fulop, T., G. Pawelec, S. Castle, and M. Loeb. 2009. Immunosenescence and vaccination in nursing home residents. *Clin. Infect. Dis.* 48: 443–448.
17. Zuhair, M., G. S. A. Smit, G. Wallis, F. Jabbar, C. Smith, B. Devleeschauwer, and P. Griffiths. 2019. Estimation of the worldwide seroprevalence of cytomegalovirus: a systematic review and meta-analysis. *Rev. Med. Virol.* 29: e2034.
18. Pawelec, G., J. E. McElhaney, A. E. Aiello, and E. Derhovanessian. 2012. The impact of CMV infection on survival in older humans. *Curr. Opin. Immunol.* 24: 507–511.
19. Sylwester, A. W., B. L. Mitchell, J. B. Edgar, C. Taormina, C. Pelte, F. Ruchti, P. R. Sleath, K. H. Grabstein, N. A. Hosken, F. Kern, et al. 2005. Broadly targeted human cytomegalovirus-specific CD4<sup>+</sup> and CD8<sup>+</sup> T cells dominate the memory compartments of exposed subjects. *J. Exp. Med.* 202: 673–685.
20. Chidrawar, S., N. Khan, W. Wei, A. McLarnon, N. Smith, L. Nayak, and P. Moss. 2009. Cytomegalovirus-seropositivity has a profound influence on the magnitude of major lymphoid subsets within healthy individuals. *Clin. Exp. Immunol.* 155: 423–432.
21. Sansoni, P., R. Vescovini, F. Fagnoni, C. Biasini, F. Zanni, L. Zanlari, A. Telera, G. Lucchini, G. Passeri, D. Monti, et al. 2008. The immune system in extreme longevity. *Exp. Gerontol.* 43: 61–65.
22. Goronzy, J. J., and C. M. Weyand. 2017. Successful and maladaptive t cell aging. *Immunity* 46: 364–378.
23. Dörner, T., and A. Radbruch. 2007. Antibodies and B cell memory in viral immunity. *Immunity* 27: 384–392.
24. Yan, Z., H. T. Maecker, P. Brodin, U. C. Nygaard, S. C. Lyu, M. M. Davis, K. C. Nadeau, and S. Andorf. 2021. Aging and CMV discordance are associated with increased immune diversity between monozygotic twins. *Immun. Ageing* 18: 5.
25. Brodin, P., V. Jojic, T. Gao, S. Bhattacharya, C. J. Angel, D. Furman, S. Shen-Orr, C. L. Dekker, G. E. Swan, A. J. Butte, et al. 2015. Variation in the human immune system is largely driven by non-heritable influences. *Cell* 160: 37–47.
26. Mekker, A., V. S. Tchang, L. Haerberli, A. Oxenius, A. Trkola, and U. Karrer. 2012. Immune senescence: relative contributions of age and cytomegalovirus infection. *PLoS Pathog.* 8: e1002850.
27. Olsson, J., A. Wikby, B. Johansson, S. Löfgren, B. O. Nilsson, and F. G. Ferguson. 2000. Age-related change in peripheral blood T-lymphocyte subpopulations and cytomegalovirus infection in the very old: the Swedish longitudinal OCTO immune study. *Mech. Ageing Dev.* 121: 187–201.
28. Crooke, S. N., I. G. Ovsyannikova, G. A. Poland, and R. B. Kennedy. 2019. Immunosenescence and human vaccine immune responses. *Immun. Ageing* 16: 25.
29. Kadambari, S., P. Klenerman, and A. J. Pollard. 2020. Why the elderly appear to be more severely affected by COVID-19: the potential role of immunosenescence and CMV. *Rev. Med. Virol.* 30: e2144.
30. Moss, P. 2020. “The ancient and the new”: is there an interaction between cytomegalovirus and SARS-CoV-2 infection? *Immun. Ageing* 17: 14.
31. Söderberg-Nauclér, C. 2021. Does reactivation of cytomegalovirus contribute to severe COVID-19 disease? *Immun. Ageing* 18: 12.
32. Chen, Y., S. L. Klein, B. T. Garibaldi, H. Li, C. Wu, N. M. Osevala, T. Li, J. B. Margolick, G. Pawelec, and S. X. Leng. 2021. Aging in COVID-19: vulnerability, immunity and intervention. *Ageing Res. Rev.* 65: 101205.
33. Merani, S., G. Pawelec, G. A. Kuchel, and J. E. McElhaney. 2017. Impact of aging and cytomegalovirus on immunological response to influenza vaccination and infection. *Front. Immunol.* 8: 784.
34. Cicin-Sain, L., J. D. Brien, J. L. Uhrlaub, A. Drabig, T. F. Marandu, and J. Nikolich-Zugich. 2012. Cytomegalovirus infection impairs immune responses and accentuates T-cell pool changes observed in mice with aging. *PLoS Pathog.* 8: e1002849.
35. Smithey, M. J., G. Li, V. Venturi, M. P. Davenport, and J. Nikolich-Zugich. 2012. Lifelong persistent viral infection alters the naive T cell pool, impairing CD8 T cell immunity in late life. *J. Immunol.* 189: 5356–5366.
36. Khan, N., A. Hislop, N. Gudgeon, M. Cobbold, R. Khanna, L. Nayak, A. B. Rickinson, and P. A. Moss. 2004. Herpesvirus-specific CD8 T cell immunity in old age: cytomegalovirus impairs the response to a coresident EBV infection. *J. Immunol.* 173: 7481–7489.
37. Barton, E. S., D. W. White, J. S. Cathelyn, K. A. Brett-McClellan, M. Engle, M. S. Diamond, V. L. Miller, and H. W. Virgin IV. 2007. Herpesvirus latency confers symbiotic protection from bacterial infection. *Nature* 447: 326–329.
38. Terrazzini, N., M. Bajwa, S. Vita, D. Thomas, H. Smith, R. Vescovini, P. Sansoni, and F. Kern. 2014. Cytomegalovirus infection modulates the phenotype and functional profile of the T-cell immune response to mycobacterial antigens in older life. *Exp. Gerontol.* 54: 94–100.
39. Pera, A., C. Campos, A. Corona, B. Sanchez-Correa, R. Tarazona, A. Larbi, and R. Solana. 2014. CMV latent infection improves CD8<sup>+</sup> T response to SEB due to expansion of polyfunctional CD57<sup>+</sup> cells in young individuals. [Published erratum appears in 2014 *PLoS One* 9: e96971.] *PLoS One* 9: e88538.
40. Smithey, M. J., V. Venturi, M. P. Davenport, A. S. Buntzman, B. G. Vincent, J. A. Frelinger, and J. Nikolich-Zugich. 2018. Lifelong CMV infection improves immune defense in old mice by broadening the mobilized TCR repertoire against third-party infection. *Proc. Natl. Acad. Sci. USA* 115: E6817–E6825.
41. Derhovanessian, E., H. Theeten, K. Hähnel, P. Van Damme, N. Cools, and G. Pawelec. 2013. Cytomegalovirus-associated accumulation of late-differentiated CD4 T-cells correlates with poor humoral response to influenza vaccination. *Vaccine* 31: 685–690.
42. Saurwein-Teissl, M., T. L. Lung, F. Marx, C. Gschösser, E. Asch, I. Blasko, W. Parson, G. Böck, D. Schönitzer, E. Trannoy, and B. Grubeck-Loebenstein. 2002. Lack of antibody production following immunization in old age: association with CD8<sup>+</sup>CD28<sup>+</sup> T cell clonal expansions and an imbalance in the production of Th1 and Th2 cytokines. *J. Immunol.* 168: 5893–5899.
43. Frasca, D., A. Diaz, M. Romero, A. M. Landin, and B. B. Blomberg. 2015. Cytomegalovirus (CMV) seropositivity decreases B cell responses to the influenza vaccine. *Vaccine* 33: 1433–1439.
44. Trzonkowski, P., J. Myśliwska, E. Szmít, J. Wiekiewicz, K. Lukaszuk, L. B. Brydak, M. Machała, and A. Myśliwski. 2003. Association between cytomegalovirus infection, enhanced proinflammatory response and low level of anti-hemagglutinins during the anti-influenza vaccination—an impact of immunosenescence. *Vaccine* 21: 3826–3836.
45. Bowyer, G., H. Sharpe, N. Venkatraman, P. B. Ndiaye, D. Wade, N. Brenner, A. Mentzer, C. Mair, T. Waterboer, T. Lambie, et al. 2020. Reduced Ebola vaccine responses in CMV<sup>+</sup> young adults is associated with expansion of CD57<sup>+</sup> KLRG1<sup>+</sup> T cells. *J. Exp. Med.* 217: e20200004.
46. van den Berg, S. P. H., K. Warmink, J. A. M. Borghans, M. J. Knol, and D. van Baarle. 2019. Effect of latent cytomegalovirus infection on the antibody response to influenza vaccination: a systematic review and meta-analysis. *Med. Microbiol. Immunol. (Berl.)* 208: 305–321.
47. Sharpe, H. R., N. M. Provine, G. S. Bowyer, P. Moreira Folegatti, S. Belij-Rammerstorfer, A. Flaxman, R. Makinson, A. V. S. Hill, K. J. Ewer, A. J. Pollard, et al. 2022. CMV-associated T cell and NK cell terminal differentiation does not affect immunogenicity of ChAdOx1 vaccination. *JCI Insight* 7: e154187.
48. Verschoor, C. P., V. Kohli, and C. Balion. 2018. A comprehensive assessment of immunophenotyping performed in cryopreserved peripheral whole blood. *Cytometry B Clin. Cytom.* 94: 662–670.
49. Ontario Ministry of Health. 2021. COVID-19 vaccine third dose recommendations. Queen's Printer for Ontario, Toronto. Available at: [https://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/vaccine/COVID-19\\_vaccine\\_third\\_dose\\_recommendations.pdf](https://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/vaccine/COVID-19_vaccine_third_dose_recommendations.pdf). Accessed: December 15, 2021.
50. Kennedy, A. E., L. Cook, J. A. Breznik, B. Cowbrough, J. G. Wallace, A. Huynh, J. W. Smith, K. Son, H. Stacey, J. Ang, et al. 2021. Lasting changes to circulating leukocytes in people with mild SARS-CoV-2 infections. *Viruses* 13: 2239.
51. Loukov, D., S. Karampatos, M. R. Malý, and D. M. E. Bowdish. 2018. Monocyte activation is elevated in women with knee-osteoarthritis and associated with inflammation, BMI and pain. *Osteoarthritis Cartilage* 26: 255–263.
52. Larbi, A., and T. Fulop. 2014. From “truly naive” to “exhausted senescent” T cells: when markers predict functionality. *Cytometry A* 85: 25–35.
53. Zaunders, J. J., M. L. Munier, N. Seddiki, S. Pett, S. Ip, M. Bailey, Y. Xu, K. Brown, W. B. Dyer, M. Kim, et al. 2009. High levels of human antigen-specific CD4<sup>+</sup> T cells in peripheral blood revealed by stimulated coexpression of CD25 and CD134 (OX40). *J. Immunol.* 183: 2827–2836.
54. Seddiki, N., L. Cook, D. C. Hsu, C. Phetsouphanh, K. Brown, Y. Xu, S. J. Kerr, D. A. Cooper, C. M. Munier, S. Pett, et al. 2014. Human antigen-specific CD4<sup>+</sup> CD25<sup>+</sup>CD134<sup>+</sup>CD39<sup>+</sup> T cells are enriched for regulatory T cells and comprise a substantial proportion of recall responses. *Eur. J. Immunol.* 44: 1644–1661.
55. Wolf, M., J. Kuball, W. Y. Ho, H. Nguyen, T. J. Manley, M. Bleakley, and P. D. Greenberg. 2007. Activation-induced expression of CD137 permits detection, isolation, and expansion of the full repertoire of CD8<sup>+</sup> T cells responding to antigen without requiring knowledge of epitope specificities. *Blood* 110: 201–210.
56. Huynh, A., D. M. Arnold, J. W. Smith, J. C. Moore, A. Zhang, Z. Chaglia, B. J. Harvey, H. D. Stacey, J. C. Ang, R. Clare, et al. 2021. Characteristics of Anti-SARS-CoV-2 Antibodies in Recovered COVID-19 Subjects. *Viruses* 13: 697.
57. National Advisory Committee on Immunization. 2021. An Advisory Committee Statement (ACS) National Advisory Committee on Immunization (NACI): guidance on booster COVID-19 vaccine doses in Canada—update December 3, 2021. Public Health Agency of Canada. Available at: <https://www.canada.ca/content/dam/phac-aspc/>

- documents/services/immunization/national-advisory-committee-on-immunization-naci/guidance-booster-covid-19-vaccine-doses/guidance-booster-covid-19-vaccine-doses.pdf Accessed: December 15, 2021.
58. Hirabara, S. M., T. D. A. Serdan, R. Gorjao, L. N. Masi, T. C. Pithon-Curi, D. T. Covas, R. Curi, and E. L. Durigon. 2022. SARS-CoV-2 variants: differences and potential of immune evasion. *Front. Cell. Infect. Microbiol.* 11: 781429.
  59. Klenerman, P., and A. Oxenius. 2016. T cell responses to cytomegalovirus. *Nat. Rev. Immunol.* 16: 367–377.
  60. Strioga, M., V. Pasukoniene, and D. Characiejus. 2011. CD8<sup>+</sup> CD28<sup>-</sup> and CD8<sup>+</sup> CD57<sup>+</sup> T cells and their role in health and disease. *Immunology* 134: 17–32.
  61. Fletcher, J. M., M. Vukmanovic-Stejic, P. J. Dunne, K. E. Birch, J. E. Cook, S. E. Jackson, M. Salmon, M. H. Rustin, and A. N. Akbar. 2005. Cytomegalovirus-specific CD4<sup>+</sup> T cells in healthy carriers are continuously driven to replicative exhaustion. *J. Immunol.* 175: 8218–8225.
  62. Henson, S. M., N. E. Riddell, and A. N. Akbar. 2012. Properties of end-stage human T cells defined by CD45RA re-expression. *Curr. Opin. Immunol.* 24: 476–481.
  63. Brenchley, J. M., N. J. Karandikar, M. R. Betts, D. R. Ambrozak, B. J. Hill, L. E. Crotty, J. P. Casazza, J. Kuruppu, S. A. Migueles, M. Connors, et al. 2003. Expression of CD57 defines replicative senescence and antigen-induced apoptotic death of CD8<sup>+</sup> T cells. *Blood* 101: 2711–2720.
  64. Neidleman, J., X. Luo, M. McGregor, G. Xie, V. Murray, W. C. Greene, S. A. Lee, and N. R. Roan. 2021. mRNA vaccine-induced T cells respond identically to SARS-CoV-2 variants of concern but differ in longevity and homing properties depending on prior infection status. *eLife* 10: e72619.
  65. Crotty, S. 2015. A brief history of T cell help to B cells. *Nat. Rev. Immunol.* 15: 185–189.
  66. van den Berg, S. P. H., J. Lanfermeijer, R. H. J. Jacobi, M. Hendriks, M. Vos, R. van Schuijlenburg, N. M. Nanlohy, J. A. M. Borghans, J. van Beek, D. van Baarle, and J. de Wit. 2021. Latent CMV infection is associated with lower influenza virus-specific memory T-cell frequencies, but not with an impaired T-cell response to acute influenza virus infection. *Front. Immunol.* 12: 663664.
  67. Theeten, H., C. Mathei, K. Peeters, B. Ogunjimi, H. Goossens, M. Ieven, P. Van Damme, and N. Cools. 2016. Cellular interferon gamma and granzyme B responses to cytomegalovirus-pp65 and influenza N1 are positively associated in elderly. *Viral Immunol.* 29: 169–175.
  68. Furman, D., V. Jovic, S. Sharma, S. S. Shen-Orr, C. J. Angel, S. Onengut-Gumuscu, B. A. Kidd, H. T. Maecker, P. Concannon, C. L. Dekker, et al. 2015. Cytomegalovirus infection enhances the immune response to influenza. *Sci. Transl. Med.* 7: 281ra43.
  69. Neidleman, J., X. Luo, J. Frouard, G. Xie, G. Gill, E. S. Stein, M. McGregor, T. Ma, A. F. George, A. Kusters, et al. 2020. SARS-CoV-2-specific T cells exhibit phenotypic features of helper function, lack of terminal differentiation, and high proliferation potential. *Cell Rep. Med.* 1: 100081.
  70. Grifoni, A., D. Weiskopf, S. I. Ramirez, J. Mateus, J. M. Dan, C. R. Moderbacher, S. A. Rawlings, A. Sutherland, L. Premkumar, R. S. Jasti, et al. 2020. Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell* 181: 1489–1501.e15.
  71. Miles, D. J., M. van der Sande, D. Jeffries, S. Kaye, J. Ismaili, O. Ojuola, M. Sanneh, E. S. Touray, P. Waight, S. Rowland-Jones, et al. 2007. Cytomegalovirus infection in Gambian infants leads to profound CD8 T-cell differentiation. *J. Virol.* 81: 5766–5776.
  72. Felismino, E. S., J. M. B. Santos, M. Rossi, C. A. F. Santos, E. L. Durigon, D. B. L. Oliveira, L. M. Thomazelli, F. R. Monteiro, A. Sperandio, J. S. Apostólico, et al. 2021. Better response to influenza virus vaccination in physically trained older adults is associated with reductions of cytomegalovirus-specific immunoglobulins as well as improvements in the inflammatory and CD8<sup>+</sup> T-cell profiles. *Front. Immunol.* 12: 713763.
  73. Rydzynski Moderbacher, C., S. I. Ramirez, J. M. Dan, A. Grifoni, K. M. Hastie, D. Weiskopf, S. Belanger, R. K. Abbott, C. Kim, J. Choi, et al. 2020. Antigen-specific adaptive immunity to SARS-CoV-2 in acute COVID-19 and associations with age and disease severity. *Cell* 183: 996–1012.e19.
  74. Wang, C., Y. Liu, L. T. Xu, K. J. L. Jackson, K. M. Roskin, T. D. Pham, J. Laserson, E. L. Marshall, K. Seo, J.-Y. Lee, et al. 2014. Effects of aging, cytomegalovirus infection, and EBV infection on human B cell repertoires. *J. Immunol.* 192: 603–611.
  75. Shrock, E., E. Fujimura, T. Kula, R. T. Timms, I. H. Lee, Y. Leng, M. L. Robinson, B. M. Sie, M. Z. Li, Y. Chen, et al.; MGH COVID-19 Collection & Processing Team. 2020. Viral epitope profiling of COVID-19 patients reveals cross-reactivity and correlates of severity. *Science* 370: eabd4250.
  76. Simonnet, A., I. Engelmann, A. S. Moreau, B. Garcia, S. Six, A. El Kalioubie, L. Robriquet, D. Hober, and M. Jourdain. 2021. High incidence of Epstein-Barr virus, cytomegalovirus, and human-herpes virus-6 reactivations in critically ill patients with COVID-19. *Infect. Dis. Now* 51: 296–299.
  77. Alanio, C., A. Verma, D. Mathew, S. Gouma, G. Liang, T. Dunn, D. A. Oldridge, J. E. Weaver, L. Kuri-Cervantes, M. B. Pampena, et al. 2022. Cytomegalovirus latent infection is associated with an increased risk of COVID-19-related hospitalization. *J. Infect. Dis.* 226: 463–473.
  78. Del Valle, D. M., S. Kim-Schulze, H.-H. Huang, N. D. Beckmann, S. Nirenberg, B. Wang, Y. Lavin, T. H. Swartz, D. Madduri, A. Stock, et al. 2020. An inflammatory cytokine signature predicts COVID-19 severity and survival. *Nat. Med.* 26: 1636–1643.