

Original Article

*See online Supplementary material.

Cite this article: McLaughlin AP *et al* (2022). The influence of comorbid depression and overweight status on peripheral inflammation and cortisol levels. *Psychological Medicine* **52**, 3289–3296. <https://doi.org/10.1017/S0033291721000088>

Received: 30 July 2020

Revised: 26 November 2020

Accepted: 15 January 2021

First published online: 18 March 2021

Key words:


Major depressive disorders; Overweight; Obesity; Inflammation; C-Reactive Protein; CRP; HPA axis; Cortisol

Author for correspondence:

Valeria Mondelli,

E-mail: valeria.mondelli@kcl.ac.uk

The influence of comorbid depression and overweight status on peripheral inflammation and cortisol levels

Anna P. McLaughlin^{1,2}, Naghmeh Nikkheslat¹, Caitlin Hastings¹, Maria A. Nettis^{1,2}, Melisa Kose¹, Courtney Worrell¹, Zuzanna Zajkowska¹, Nicole Mariani¹, Daniela Enache¹, Giulia Lombardo¹, Linda Pointon³, NIMA Consortium^{1,*}, Philip Cowen⁴, Jonathan Cavanagh⁵, Neil Harrison⁶, Edward Bullmore³, Carmine M. Pariante^{1,2} and Valeria Mondelli^{1,2} 

¹Department of Psychological Medicine, King's College London, Institute of Psychiatry, Psychology and Neuroscience, London, UK; ²National Institute for Health Research (NIHR) Maudsley Biomedical Research Centre, South London and Maudsley NHS Foundation Trust, King's College London, London, UK; ³Department of Psychiatry, University of Cambridge, Cambridge, UK; ⁴University Department of Psychiatry, Warneford Hospital, Oxford, UK; ⁵Mental Health and Wellbeing, Sackler Institute, Neurology block, Queen Elizabeth University hospital, Glasgow, UK and ⁶Division of Psychological Medicine and Clinical Sciences, Cardiff University Brain Research Imaging Centre (CUBRIC), Cardiff, UK

Abstract

Background. Depression and overweight are each associated with abnormal immune system activation. We sought to disentangle the extent to which depressive symptoms and overweight status contributed to increased inflammation and abnormal cortisol levels.

Methods. Participants were recruited through the Wellcome Trust NIMA Consortium. The sample of 216 participants consisted of 69 overweight patients with depression; 35 overweight controls; 55 normal-weight patients with depression and 57 normal-weight controls. Peripheral inflammation was measured as high-sensitivity C-Reactive Protein (hsCRP) in serum. Salivary cortisol was collected at multiple points throughout the day to measure cortisol awakening response and diurnal cortisol levels.

Results. Overweight patients with depression had significantly higher hsCRP compared with overweight controls ($p = 0.042$), normal-weight depressed patients ($p < 0.001$) and normal-weight controls ($p < 0.001$), after controlling for age and gender. Multivariable logistic regression showed that comorbid depression and overweight significantly increased the risk of clinically elevated hsCRP levels ≥ 3 mg/L (OR 2.44, 1.28–3.94). In a separate multivariable logistic regression model, overweight status contributed most to the risk of having hsCRP levels ≥ 3 mg/L (OR 1.52, 0.7–2.41), while depression also contributed a significant risk (OR 1.09, 0.27–2). There were no significant differences between groups in cortisol awakening response and diurnal cortisol levels.

Conclusion. Comorbid depression and overweight status are associated with increased hsCRP, and the coexistence of these conditions amplified the risk of clinically elevated hsCRP levels. Overweight status contributed most to the risk of clinically elevated hsCRP levels, but depression also contributed to a significant risk. We observed no differences in cortisol levels between groups.

Introduction

Rates of depression and obesity have risen dramatically in recent years, with each disorder separately posing a major health concern and economic cost to society (Abdelal, le Roux, & Docherty, 2017; James *et al.*, 2018; Malhi & Mann, 2018). Depression and obesity are highly comorbid disorders, and the coexistence of these conditions significantly increases the risk for developing subsequent disorders, likely due to the chronic inflammatory state that they induce (Ouaknin, Barreira, & Gois, 2018). High levels of inflammatory markers have been widely described in depression (Baumeister, Russell, Pariante, & Mondelli, 2014; Enache, Pariante, & Mondelli, 2019; Haapakoski, Mathieu, Ebmeier, Alenius, & Kivimäki, 2015; Howren, Lamkin, & Suls, 2009; Valkanova, Ebmeier, & Allan, 2013), with a recent meta-analysis finding that approximately 25% of patients with depression presented with elevated peripheral inflammation (Osimo, Baxter, Lewis, Jones, & Khandaker, 2019). However, weight gain also increases peripheral inflammation, and studies of patients with depression have found that 20% of patients to be obese and 50% to be overweight (Papakostas *et al.*, 2005). Therefore, it is unclear whether the increased inflammation observed in patients with depression is predominantly

due to the high rates of overweight and obesity in this group. As inflammation may be both a causal mechanism and potential treatment target for depressive symptoms, it is crucial to understand to what extent depression and weight gain each contribute to the increased inflammation observed in patients with depression (Ambrosio *et al.*, 2018).

Inflammation likely plays a key role in modifying the bidirectional relationship between depression and obesity, but not all patients with depression demonstrate increased inflammation. Patients with depression who have both increased inflammation and metabolic disturbances may have a distinct 'immuno-metabolic' form of depression (Milaneschi, Lamers, Berk, & Penninx, 2020). This type of depression is a significant risk factor for weight gain and subsequent obesity (Hasler *et al.*, 2004) and was specifically associated with increased peripheral levels of inflammatory markers, such as C-reactive protein (CRP) (Lamers *et al.*, 2013). Increased CRP levels have clinical relevance for depression (Miller & Raison, 2016), as CRP ≥ 3 mg/L was associated with an increased risk of developing depression later in life (Au, Smith, Garipey, & Schmitz, 2015), and a lack of response to antidepressant medication (Chamberlain *et al.*, 2019; Zhang *et al.*, 2019). Conversely, reductions in body mass index (BMI) following weight loss interventions are associated with an improvement in depressive symptoms and lower CRP levels (Capuron *et al.*, 2011; Perez-Cornago *et al.*, 2014). These data demonstrate the crucial role of inflammation, as measured by CRP, in the interface between depression and weight gain (Ambrosio *et al.*, 2018).

Depression with metabolic disturbances has also been associated with dysregulated hypothalamic-pituitary-adrenal (HPA) axis activity (Gold, 2015). The HPA axis produces the anti-inflammatory stress hormone cortisol, which is involved in regulating mood, metabolism, and circadian rhythms. Meta-analyses have suggested that patients with this subtype of depression have lower diurnal cortisol levels, meaning HPA axis hypoactivity (Jurueña, Bocharova, Agustini, & Young, 2018; Lamers *et al.*, 2013). In contrast, patients who exhibit the more classic symptoms of depression, such as reduced appetite and insomnia, tend to demonstrate HPA axis hyperactivity (Jurueña *et al.*, 2018; Lamers *et al.*, 2013). This association was mechanistically supported by longitudinal studies finding that lower cortisol levels in children are associated with higher BMI at age 18 (Ruttle *et al.*, 2014), and that blunted cortisol responses lead to further weight gain and increased inflammation (Champaneri *et al.*, 2013). HPA axis hypoactivity may therefore be indicative of a pathophysiological process, capable of influencing mood and weight gain, which is unique to depressed patients with metabolic disturbances.

Research investigating inflammatory markers and their association with depressive symptoms in individuals with obesity is typically complicated by a high incidence of other inflammatory disorders, such as atherosclerosis, diabetes, cardiovascular disease, and hypertension (Upadhyay, Farr, Perakakis, Ghaly, & Mantzoros, 2018). Once patients have developed comorbid depression and obesity, they become more vulnerable to developing subsequent inflammatory conditions. These conditions are associated with an increased risk of recurrent depressive episodes (Nigatu, Bultmann, & Reijneveld, 2015), demonstrating the vicious cycle to which these patients may be liable. Therefore, it is vital to confirm the association between depressive symptoms and inflammation in a sample free of comorbid disorders, as the presence of other disorders may confound this relationship.

It is also important to identify which subgroup of patients with depression are most vulnerable so that they can be prioritised for clinical interventions targeted at reducing their risk of developing further comorbidities.

Therefore, we investigated inflammatory mechanisms, in the form of pro-inflammatory hsCRP and anti-inflammatory cortisol, to determine how these biomarkers are associated with depression and overweight status. We also investigated whether the presence of comorbid depression and overweight increased inflammation to a clinically relevant level. Crucially, our study used a sample of patients with depression who were free of other comorbid disorders, with a comparison group of overweight controls, who were otherwise healthy.

This is the first study to explore differences in hsCRP levels and HPA axis activity in overweight individuals with and without depression, as well as normal-weight individuals with and without depression. Our cross-sectional study aimed to investigate: (1) whether overweight patients with depression had increased inflammation relative to all other groups; (2) what extent overweight status and depression status contributed to clinically elevated hsCRP levels ≥ 3 mg/L; (3) whether overweight patients with depression had lower diurnal cortisol levels relative to all other groups.

Methods

Clinical data, blood samples, and saliva samples were collected in a cross-sectional, observational design, as part of a multi-centre study investigating immune Biomarkers in Depression (BIODEP), through the Wellcome Trust Consortium for Neuroimmunology of Mood Disorders and Alzheimer's disease (NIMA). The study was approved by the Research Ethics Committee (National Research Ethics Service East of England, Cambridge Central, UK; approval number: 15/EE/0092). All procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All participants provided written informed consent and underwent eligibility screening prior to taking part in any study procedures.

Participants

Participants were aged 25–50 years inclusive and recruited at five clinical research study centres (King's College London, Oxford, Cambridge, Brighton and Glasgow) from primary and secondary NHS health services and the general population. Participants were excluded if they (1) were pregnant or breastfeeding; (2) were underweight (defined as BMI < 18); (3) were taking medication likely to compromise the interpretation of immunological data (including, but not limited to, statins, corticosteroids, antihistamines and anti-inflammatory medications); (4) met criteria for alcohol abuse, drug abuse or dependence in the last 6 months; (5) had participated in a clinical trial of an investigational drug within the last 12 months; (6) had lifetime history of any serious medical disorder likely to compromise the interpretation of immunological data; (7) had a recent infection or illness likely to compromise the interpretation of immunological data. Patients with depression were considered eligible if they met the criteria for major depressive disorder (MDD). Patients with depression were excluded if they had a lifetime history of bipolar disorder or non-affective psychosis. Healthy controls were

considered eligible if they had no personal history of MDD or treatment with a monoaminergic antidepressant for depressive symptoms or any other indication, as well as no current or lifetime history of any major psychiatric disorder as defined by Diagnostic and Statistical Manual Version 5 (DSM-5).

Sample groups

The current sample was selected from a larger sample of participants taking part in the BIODep study, where participants were recruited based on their clinical response to antidepressants as described by our group previously (Chamberlain et al., 2019; Nikkheslat et al., 2019). For the purpose of the current study, participants from the BIODep pool were grouped according to their BMI status (overweight participants with BMI ≥ 25) and then categorised these participants according to the presence of depression (MDD patients *v.* healthy controls).

Demographic and clinical measures

Age, gender, smoking status, and medical history were documented by semi-structured clinical interviews. Height and weight were measured for calculation of BMI (kg/m^2) and overweight status was defined as BMI ≥ 25 . History and diagnosis of MDD and other psychiatric disorders were assessed with the Structured Clinical Interview for DSM-5 (Kübler, 2013). The severity of depressive symptoms was assessed using 17-item Hamilton Rating Scale for Depression (HAM-D) (Hamilton, 1960).

High sensitivity C-Reactive Protein (hsCRP)

Peripheral inflammation was measured as serum levels of hsCRP, which has been demonstrated as a reliable biomarker of inflammation associated with MDD. Participants fasted for 8 h and abstained from strenuous exercise for 72 h prior to their blood draw, which was carried out between 8:00 and 10:00. Blood samples were collected in clotting tubes, allowed to coagulate at room temperature for 30–60 min, then centrifuged at 1600 Relative Centrifugal Force for 15 min. The serum samples were separated and transported to a central laboratory (Q2 solutions) where they were analysed on the day of collection. Samples were exposed to anti-CRP-antibodies on latex particles, and the increase in light absorption due to complex formation was used to quantify hsCRP levels, using Turbidimetry on Beckman Coulter AU analysers. Inter and intra-assay coefficient of variations were $<10\%$. The measure of hsCRP was calculated from one blood draw taken at the time of clinical assessment, for each participant. In line with previous studies, clinically elevated hsCRP levels were defined as hsCRP ≥ 3 mg/L (Miller & Raison, 2016; Pearson et al., 2003).

Salivary cortisol

Participants were issued with the materials and instructions for collecting the saliva samples at the time of their clinical interview. Our previously published literature describes the collection procedure for saliva samples in more detail (Nikkheslat et al., 2019). Using salivette sampling devices (Sarstedt, Leicester, UK), samples were self-collected by participants at home at six-time points throughout the day; at awakening, 15, 30, and 60 min after awakening, 12:00 and at 20:00. Individuals who described problems during sample collection in the self-recorded questionnaire, or who did not respect the time-intervals required, were

removed from the analysis. The current study included only participants who completed saliva sample collections accurately and who provided an adequate amount of saliva for cortisol measurement. Salivary cortisol levels were measured using a commercially available high-sensitivity salivary cortisol enzyme immunoassay kit from Salimetrics. SoftMax Pro 4.8 software was used to calculate the cortisol values, following a 4-parameter fit. The analytical sensitivity was set to 0.19 nmol/L. Inter and intra-assay coefficient of variations ranged from 8 to 10% and 6–10%, respectively. To investigate the activity and responsiveness of the HPA axis, we first compared the mean values at the various time points of salivary cortisol collection; and secondly, we calculated the area under the curve with respect to the increase (AUC_i) for the cortisol awakening response using the four-time points of 0, 15, 30, and 60 min after awakening; and the area under the curve with respect to the ground (AUC_g) for the diurnal cortisol using the three points: awakening, noon and 20:00. AUC_g indicated the total amount of cortisol produced and overall HPA axis activity during the day. AUC_i indicated the variation (either positive or negative) in cortisol concentration and thus signified the HPA axis reactivity and response to the stress of awakening. The formulas for the calculations of the AUC were derived from the trapezoidal formula introduced by Pruessner et al. (Pruessner, Kirschbaum, Meinlschmid, & Hellhammer, 2003).

Statistical analysis

All statistical analyses were performed using R software, version 4.0.0. Data were evaluated for normality and logarithmic transformed if required. To assess group differences in demographic variables, Chi-squared or Kruskal–Wallis tests were used, as appropriate. To address the first hypothesis, group differences in logarithmic-transformed hsCRP levels were assessed using two-way analysis of covariance (ANCOVA), with group and gender as factors and age as a covariate. Pairwise comparisons were carried out using the False Discovery Rate (FDR) correction, with the alpha set at 0.05 (5%). Effect size differences in hsCRP groups relative to the normal-weight controls were calculated using Cohen's *d*. To address the second hypothesis, two models of multivariable logistic regression were used. The first assessed how much risk comorbid depression and overweight contributed to the hsCRP levels ≥ 3 mg/L relative to other participant groups, with age and gender as covariates. The second assessed how much risk depression and overweight status (BMI ≥ 25) separately contributed to hsCRP levels ≥ 3 mg/L, with age and gender as covariates. To address the third hypothesis, group differences in cortisol awakening response (AUC_i) and logarithmic-transformed diurnal cortisol (AUC_g) were assessed using two-way ANCOVA, with adjustment for age and gender. The threshold for statistical significance of all tests was defined as two-tailed $p \leq 0.05$.

Results

Sample demographics

The demographic and clinical characteristics of the sample are presented in Table 1. Age and gender were significantly different between groups, while ethnicity and smoking status were similar across groups, in both samples. Although the proportion of smokers were not significantly different between groups, we repeated all analyses excluding smokers to determine if smoking status significantly influenced hsCRP and cortisol levels. As the direction of

Table 1. Comparison of sample groups

Variables	Groups				Group tests	
	Overweight & depressed, N = 69	Overweight control, N = 35	Normal weight & depressed, N = 55	Normal weight control, N = 57	Statistic	p
Demographics						
Median age in years, (range)	38 (25–50)	38 (25–48)	32 (25–50)	30 (24–50)	$K = 18.368$	<0.001
Gender, female (%)	43 (62.3%)	17 (48.6%)	40 (72.7%)	44 (77.2%)	$\chi^2 = 9.495$	0.023
BMI (range)	28.8 (25.3–47.8)	28.4 (25–35.6)	22.1 (18.1–24.9)	22.4 (18.4–24.9)	$K = 161.87$	<0.001
Ethnicity, white (%)	63 (91.3%)	29 (82.9%)	45 (81.8%)	45 (78.9%)	$\chi^2 = 4.102$	0.251
Smoking status, smoker (%)	11 (15.9%)	3 (8.6%)	11 (20%)	6 (10.5%)	$\chi^2 = 3.199$	0.362
Clinical						
Median HAM-D total score (range)	19 (14–31)	0 (0–5)	18 (14–26)	0 (0–7)	$K = 162.83$	<0.001
Currently on antidepressants (%)	47 (68.1%)	NA	39 (70.9%)	NA	$\chi^2 = 0.019$	0.889
Mean age of depression onset \pm s.d.	24.6 \pm 9.8	NA	25.2 \pm 9.4	NA	$F = 1.617$	0.206
hsCRP						
Median hsCRP (range)	2.2 (0.2–15.3)	1.3 (0.2–11.8)	0.7 (0.2–8.8)	0.5 (0.2–5.2)	$K = 43.725$	<0.001
Mean log hsCRP mg/L \pm s.d.	0.28 \pm 0.48	0.09 \pm 0.43	–0.12 \pm 0.42,	–0.25 \pm 0.35	$F = 16.43$	<0.001
Cohen's d effect size of hsCRP vs. normal weight controls	1.25	0.89	0.35	-		
hsCRP \geq 3 mg/L (%)	26 (37.7%)	5 (14.3%)	6 (10.9%)	3 (5.3%)	$\chi^2 = 25.96$	<0.001
Salivary cortisol						
Mean CAR AUCi (nmol min/L) \pm s.d.	100.53 \pm 248.99	75.79 \pm 240.98	41.62 \pm 240.43	86.46 \pm 299.78	$F = 0.473$	0.701
Mean log AUCg (nmol h/L) \pm s.d.	1.65 \pm 0.18	1.66 \pm 0.23	1.72 \pm 0.2	1.7 \pm 0.26	$F = 1.25$	0.293

BMI, body mass index; HAM-D, Hamilton Depression Rating Scale 17 for depressive symptoms; hsCRP, high sensitivity C-reactive protein; s.d., standard deviation; CAR, cortisol awakening response; AUCi, area under the curve with respect to increase; AUCg, area under the curve with respect to ground.

the results did not change for any of the relevant tests after excluding smokers, we have reported results for the full sample, with the data for non-smokers presented in the supplement. A similar number of overweight patients with depression and normal-weight patients with depression were currently taking antidepressant medication (47 patients *v.* 39 patients, $\chi^2 = 0.019$, $p = 0.889$) and a two-way ANCOVA showed that both groups had a similar age of onset of depressive symptoms ($F_{(1, 109)} = 1.617$, $p = 0.206$), with adjustment for age and gender.

High sensitivity C-Reactive Protein

A two-way ANCOVA showed a significant effect of group on hsCRP, with adjustment for age and gender ($F_{(3, 211)} = 16.43$, $p < 0.001$). As there was no significant interaction between group and gender on hsCRP, $F_{(3, 211)} = 1.4$, $p = 0.245$, pairwise comparisons with the FDR correction were carried out, comparing the main effects of group, but not gender. Overweight patients with depression had significantly higher hsCRP compared with overweight controls ($p = 0.042$), normal weight depressed patients ($p < 0.001$), and normal-weight controls ($p < 0.001$; see Fig. 1). Within the group of overweight patients with depression, there were six individuals with extreme BMI values ≥ 40 indicating morbid obesity. When excluding these individuals from analysis, a two-way ANCOVA showed a significant effect of group on hsCRP, with adjustment for age and gender ($F_{(3, 201)} = 15.11$, $p < 0.001$). As there was no significant interaction between group and gender on hsCRP, $F_{(3, 207)} = 1.36$, $p = 0.257$, pairwise comparisons with the FDR correction were carried out, comparing the main effects of group, but not gender. Overweight patients with depression had higher hsCRP compared with overweight controls at trend-level ($p = 0.068$), and significantly higher hsCRP compared with normal weight depressed patients ($p < 0.001$) and normal-weight controls ($p < 0.001$).

Multivariable logistic regression analysis was performed to investigate whether group status contributed to the risk of having hsCRP ≥ 3 mg/L, after controlling for age and gender. Overweight patients with depression were at a significantly increased risk of having hsCRP ≥ 3 mg/L [OR 2.44, 95% confidence interval (CI) 1.28 to 3.94, $p < 0.001$], overweight controls did not have a significantly increased risk of hsCRP ≥ 3 mg/L (OR 1.51, 95% CI -0.33 to 2.87, $p = 0.13$), normal-weight patients with depression did not have a significantly increased risk of hsCRP ≥ 3 mg/L (OR 1.08, 95% CI -0.6 to 2.4, $p = 0.28$), while normal-weight controls were at a significantly decreased risk of hsCRP ≥ 3 mg/L (OR -4.39 , 95% CI -7.04 to -1.97 , $p < 0.001$). Age and gender were not significantly associated with hsCRP ≥ 3 mg/L [OR 0.01, 95% CI -0.04 to 0.06, $p = 0.621$, OR 0.61, 95% CI -0.19 to 1.47, $p = 0.145$, respectively].

Multivariable logistic regression analysis was also performed to investigate the association between depression and overweight status on hsCRP ≥ 3 mg/L, after controlling for age and gender. Depression status was significantly associated with hsCRP ≥ 3 mg/L (OR 1.09, 95% CI 0.27 to 2.01, $p = 0.013$) and overweight status was significantly associated with hsCRP ≥ 3 mg/L (OR 1.52, 95% CI 0.7 to 2.41, $p < 0.001$). Age and gender were not significantly associated with hsCRP ≥ 3 mg/L [OR 0.62, 95% CI -0.04 to 0.06, $p = 0.647$, OR 0.01, 95% CI -0.19 to 1.47, $p = 0.137$, respectively].

Cortisol

A two-way ANCOVA showed no significant differences between groups in CARi, after adjustment for age and gender, $F_{(3, 162)} =$

0.473, $p = 0.701$. Similarly, a two-way ANCOVA showed no significant differences between groups in logarithmic-transformed AUCg, after adjustment for age and gender, $F_{(3, 167)} = 0.719$, $p = 0.542$.

Discussion

To our knowledge, this is the first study to observe significantly higher hsCRP levels in overweight patients with depression relative to overweight controls, as well as depressed and non-depressed normal-weight participants. Of clinical relevance, overweight patients with depression had the highest risk for hsCRP levels ≥ 3 mg/L, indicating clinically elevated peripheral inflammation. Our results demonstrate that coexisting depression and overweight may exacerbate peripheral inflammation and highlights the urgent need to optimise treatment strategies for these patients. We observed no differences in HPA axis activity between groups.

Our findings regarding elevated hsCRP levels in overweight patients with depression are consistent with previous studies in normal-weight patients with depression and obese patients with depression (Haapakoski et al., 2015; Rethorst, Bernstein, & Trivedi, 2014). Of note, our study design separating overweight and normal-weight participants, from those with and without depression, was able to demonstrate the separate and comorbid association between these disorders and hsCRP levels ≥ 3 mg/L. Peripheral inflammation is believed to initiate and perpetuate a range of sickness-related behaviour, such as fatigue, weakness, malaise, sleep, and disturbed appetite (Dantzer, 2006), which may further maintain the vicious cycle between weight gain and depressive symptoms. The increased risk for overweight patients with depression to have clinically elevated hsCRP ≥ 3 mg/L is concerning, as CRP is a predictor of all-cause mortality (Li et al., 2017). Increased mortality and risk of developing subsequent comorbid disorders in patients with depression place a substantial burden on healthcare infrastructure and the economy (Tremmel, Gerdtham, Nilsson, & Saha, 2017), in addition to reduced quality of life for these individuals (Nigatu, Reijneveld, de Jonge, van Rossum, & Bultmann, 2016). Answering the question of whether depression or weight gain contributed more to inflammation was identified as a key objective by a recent meta-analysis (Ambrosio et al., 2018), so that research for this field can progress into therapeutic interventions designed to break the cyclic pattern between these disorders.

Testing CRP levels in overweight patients with depression could be useful for recommending adjunct anti-inflammatory treatments or predicting their likelihood of developing additional comorbidities in the future. Indeed, several clinical trials have observed that higher BMI at baseline predicted a poorer response to antidepressant treatments (Jha et al., 2018; Uher et al., 2009), while we have previously shown that immune-metabolic status was associated with a poorer response to antipsychotic medication in individuals with psychosis (Nettis et al., 2019). Anti-inflammatory drugs improve the therapeutic action of antidepressants (Bai et al., 2020; Haroon, Raison, & Miller, 2012), which could be particularly useful in the context of chronically inflamed overweight patients with depression. Overweight patients with hsCRP ≥ 3 mg/L may prove ideal criterion to test this, as stratifying patients to specific treatments according to inflammatory markers seems to improve treatment response (Cuthbert & Insel, 2013; Kohler, Krogh, Mors, & Benros, 2016).

The lack of significant differences between groups in HPA axis activity was surprising, as HPA axis dysregulation has been




Fig. 1. Estimated marginal means of log transformed hsCRP levels for each group. A two-way ANCOVA showed a significant effect of group on log hsCRP, with adjustment for age and gender $F_{(3, 211)} = 16.43, p < 0.001$. Significance of pairwise comparisons, with the False Discovery Rate correction set at 5%, are shown within the figure.

implicated in both obesity and depression (Incollingo Rodriguez *et al.*, 2015), however, the nature of the dysregulation is not always consistent. A systematic review comparing atypical depression with the melancholic (typical) presentation, found that atypical patients demonstrated lower cortisol relative to melancholic patients, but not so low that they were significantly different from controls (Jurueña *et al.*, 2018). In contrast, our study observed no differences in cortisol levels, despite hsCRP levels varying dramatically between groups. Dysregulation of the HPA axis may change over illness course, with hypotheses suggesting that at first cortisol levels are highly responsive to peripheral inflammation, but over time this responsiveness declines (Gold, 2015; Perrin, Horowitz, Roelofs, Zunszain, & Pariante, 2019). Another explanation is that specific symptoms are responsible for changes in HPA axis activity (Iob, Kirschbaum, & Steptoe, 2020), and that antidepressant use additionally modifies HPA axis activity, as our group recently observed (Nikkheslat *et al.*, 2019). Ultimately, our sample of patients may have been too heterogeneous in terms of depressive symptoms, antidepressant use and illness course to identify group differences.

Our study design was substantially strengthened by our strict exclusion criteria, encompassing recent illnesses and comorbid disorders, such as atherosclerosis, diabetes, and cardiovascular disease, which often confounds research in patients with obesity. We also excluded patients taking medications that could influence immunological data, such as statins, corticosteroids, antihistamines, anti-inflammatory medications. Additionally, the direction of our results was maintained after we removed smokers and individuals with extreme BMI values ≥ 40 , suggesting that our findings are robust. Our study was limited by a lack of additional body composition and clinical measures, as these would have enabled us to define an immune-metabolic subgroup of patients within our sample. However, using overweight status as defined by BMI has high translational value for clinical settings. Our sample consisted of predominantly white participants, which limits the generalisability of the results. Our study could have been improved by the measurement of lifestyle factors, particularly

physical activity, sleep and diet, as these factors may confound the relationship between depression and weight gain (Schmidt *et al.*, 2015). Future studies should be designed to include more diverse participant samples and adjust for lifestyle factors as covariates as they may explain some of the heterogeneity in the association between depressive symptoms and inflammation.

Chronic inflammation is likely a key modifier in the vicious cycle between depressive symptoms and weight gain. If the cycle can be reversed, meaning that a reduction in depressive symptoms is accompanied by healthy behaviour such as exercise and healthy eating, which indirectly reduces inflammation and leads to excess weight loss, this would result in a better quality of life for patients and a vast reduction in global healthcare burdens and economic cost (Jantaratnotai, Mosikanon, Lee, & McIntyre, 2017; Nigatu *et al.*, 2016; Tremmel *et al.*, 2017). Ultimately, our data suggest that depression and overweight are separately associated with increased inflammation, but this association is amplified when these conditions coexist. Patients with depression with BMI ≥ 25 would likely benefit from further research to determine if a higher level of clinical support or anti-inflammatory treatment approaches could potentially improve depressive symptoms and reduce the risk of developing further comorbidities in this subgroup of patients.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291721000088>

Acknowledgements. This research has been supported by the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health. The authors would like to thank the research team at Brighton, Cambridge, Glasgow, King's College London and Oxford; and all the study participants without whom this work would not have been possible and completed. The members of NIMA Consortium are thanked and acknowledged (see Annex). VM is supported by MQ: Transforming Mental Health (Grant: MQBF1) and by the Medical Research Foundation (Grant: MRF-160-0005). This work was supported by

the NIHR Cambridge Biomedical Research Centre (Mental Health). This work was funded by a grant from the Wellcome Trust (Grant number: 104025/Z/14/Z) to the NIMA Consortium, which is also funded by Janssen, GlaxoSmithKline, Lundbeck and Pfizer. Recruitment of patients was supported by the National Institute of Health Research (NIHR) Clinical Research Network: Kent, Surrey and Sussex & Eastern. The work is also supported by the Medical Research Council (UK) MR/J002739/1 and the Commission of European Communities Seventh Framework Programme (Collaborative Project Grant Agreement no. 22963, Mood Inflamm); and partly funded by the NIHR/Wellcome Trust, King's Clinical Research Facility and the NIHR Biomedical Research Centre (and Dementia Unit) at South London and Maudsley NHS Foundation Trust and King's College London.

Conflict of interest. SRC consults for Cambridge Cognition and Shire; and his input in this project was funded by a Wellcome Trust Clinical Fellowship (110049/Z/15/Z). ETB was employed half-time by GlaxoSmithKline, and held stock in GSK, until May 2019; he has since worked full time for University of Cambridge; he was holding stock (not currently) in GSK. ETB is an NIHR Senior Investigator. NAH consults for GSK. PdB, DJ and WCD are employees of Janssen Research & Development, LLC., of Johnson & Johnson, and hold stock in Johnson & Johnson. The other authors report no financial disclosures or potential conflicts of interest.

References

- Abdelaal, M., le Roux, C. W., & Docherty, N. G. (2017). Morbidity and mortality associated with obesity. *The Annals of Translational Medicine*, 5(7), 161. doi: 10.21037/atm.2017.03.107
- Ambrosio, G., Kaufmann, F. N., Manosso, L., Platt, N., Ghisleni, G., Rodrigues, A. L. S., ... Kaster, M. P. (2018). Depression and peripheral inflammatory profile of patients with obesity. *Psychoneuroendocrinology*, 91, 132–141. doi: 10.1016/j.psyneuen.2018.03.005
- Au, B., Smith, K. J., Garipey, G., & Schmitz, N. (2015). The longitudinal associations between C-reactive protein and depressive symptoms: Evidence from the English Longitudinal Study of Ageing (ELSA). *International Journal of Geriatric Psychiatry*, 30(9), 976–984. doi: 10.1002/gps.4250
- Bai, S., Guo, W., Feng, Y., Deng, H., Li, G., Nie, H., ... Tang, Z. (2020). Efficacy and safety of anti-inflammatory agents for the treatment of major depressive disorder: A systematic review and meta-analysis of randomised controlled trials. *Journal of Neurology, Neurosurgery, and Psychiatry*, 91(1), 21–32. doi: 10.1136/jnnp-2019-320912
- Baumeister, D., Russell, A., Pargante, C. M., & Mondelli, V. (2014). Inflammatory biomarker profiles of mental disorders and their relation to clinical, social and lifestyle factors. *Social Psychiatry and Psychiatric Epidemiology*, 49(6), 841–849. doi: 10.1007/s00127-014-0887-z
- Capuron, L., Poitou, C., Machaux-Tholliez, D., Frochot, V., Bouillot, J. L., Basdevant, A., ... Clement, K. (2011). Relationship between adiposity, emotional status and eating behaviour in obese women: Role of inflammation. *Psychological Medicine*, 41(7), 1517–1528. doi: 10.1017/S0033291710001984
- Chamberlain, S. R., Cavanagh, J., de Boer, P., Mondelli, V., Jones, D. N. C., Drevets, W. C., ... Bullmore, E. T. (2019). Treatment-resistant depression and peripheral C-reactive protein. *British Journal of Psychiatry*, 214(1), 11–19. doi: 10.1192/bjp.2018.66
- Champaneri, S., Xu, X., Carnethon, M. R., Bertoni, A. G., Seeman, T., DeSantis, A. S., ... Golden, S. H. (2013). Diurnal salivary cortisol is associated with body mass index and waist circumference: The Multiethnic Study of Atherosclerosis. *Obesity (Silver Spring)*, 21(1), E56–E63. doi: 10.1002/oby.20047
- Cuthbert, B. N., & Insel, T. R. (2013). Toward the future of psychiatric diagnosis: The seven pillars of RDoC. *BMC Medicine*, 11, 126. doi: 10.1186/1741-7015-11-126
- Dantzer, R. (2006). Cytokine, sickness behavior, and depression. *Neurologic Clinics*, 24(3), 441–460. doi: 10.1016/j.ncl.2006.03.003
- Enache, D., Pariante, C. M., & Mondelli, V. (2019). Markers of central inflammation in major depressive disorder: A systematic review and meta-analysis of studies examining cerebrospinal fluid, positron emission tomography and post-mortem brain tissue. *Brain, Behavior, and Immunity*, 81, 24–40. doi: 10.1016/j.bbi.2019.06.015
- Gold, P. W. (2015). The organization of the stress system and its dysregulation in depressive illness. *Molecular Psychiatry*, 20(1), 32–47. doi: 10.1038/mp.2014.163
- Haapakoski, R., Mathieu, J., Ebmeier, K. P., Alenius, H., & Kivimäki, M. (2015). Cumulative meta-analysis of interleukins 6 and 1 β , tumour necrosis factor α and C-reactive protein in patients with major depressive disorder. *Brain, Behavior, and Immunity*, 49, 206–215. doi: 10.1016/j.bbi.2015.06.001
- Hamilton, M. (1960). A rating scale for depression. *Journal of Neurology, Neurosurgery, and Psychiatry*, 23, 56–62. doi: 10.1136/jnnp.23.1.56
- Haroon, E., Raison, C. L., & Miller, A. H. (2012). Psychoneuroimmunology meets neuropsychopharmacology: Translational implications of the impact of inflammation on behavior. *Neuropsychopharmacology*, 37(1), 137–162. doi: 10.1038/npp.2011.205
- Hasler, G., P. D. S., Gamma, A., Milos, G., Ajdacic, V., Eich, D., ... Angst, J. (2004). The associations between psychopathology and being overweight: A 20-year prospective study. *Psychological Medicine*, 34(6), 1047–1057.
- Howren, M. B., Lamkin, D. M., & Suls, J. (2009). Associations of depression with C-reactive protein, IL-1, and IL-6: A meta-analysis. *Psychosomatic Medicine*, 71(2), 171–186. doi: 10.1097/PSY.0b013e3181907c1b
- Incollongo Rodriguez, A. C., Epel, E. S., White, M. L., Standen, E. C., Seckl, J. R., & Tomiyama, A. J. (2015). Hypothalamic-pituitary-adrenal axis dysregulation and cortisol activity in obesity: A systematic review. *Psychoneuroendocrinology*, 62, 301–318. doi: 10.1016/j.psyneuen.2015.08.014
- Iob, E., Kirschbaum, C., & Steptoe, A. (2020). Persistent depressive symptoms, HPA-axis hyperactivity, and inflammation: The role of cognitive-affective and somatic symptoms. *Molecular Psychiatry*, 25(5), 1130–1140. doi: 10.1038/s41380-019-0501-6
- James, S. L., Abate, D., Abate, K. H., Abay, S. M., Abbafati, C., Abbasi, N., ... Murray, C. J. L. (2018). Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: A systematic analysis for the global burden of disease study 2017. *The Lancet*, 392(10159), 1789–1858. doi: 10.1016/s0140-6736(18)32279-7
- Jantaratnotai, N., Mosikanon, K., Lee, Y., & McIntyre, R. S. (2017). The inter-face of depression and obesity. *Obesity Research & Clinical Practice*, 11(1), 1–10. doi: 10.1016/j.orcp.2016.07.003
- Jha, M. K., Wakhlu, S., Dronamraju, N., Minhajuddin, A., Greer, T. L., & Trivedi, M. H. (2018). Validating pre-treatment body mass index as moderator of antidepressant treatment outcomes: Findings from CO-MED trial. *Journal of Affective Disorders*, 234, 34–37. doi: 10.1016/j.jad.2018.02.089
- Juruena, M. F., Bocharova, M., Agustini, B., & Young, A. H. (2018). Atypical depression and non-atypical depression: Is HPA axis function a biomarker? A systematic review. *Journal of Affective Disorders*, 233, 45–67. doi: 10.1016/j.jad.2017.09.052
- Kohler, O., Krogh, J., Mors, O., & Benros, M. E. (2016). Inflammation in depression and the potential for anti-inflammatory treatment. *Current Neuropharmacology*, 14(7), 732–742. doi: 10.2174/1570159x14666151208113700
- Kübler, U. (2013). Structured clinical interview for DSM-IV (SCID). In M. D. Gellman & J. R. Turner (Eds.), *Encyclopedia of behavioral medicine* (pp. 1919–1920). New York, NY: Springer New York.
- Lamers, F., Vogelzangs, N., Merikangas, K. R., de Jonge, P., Beekman, A. T., & Penninx, B. W. (2013). Evidence for a differential role of HPA-axis function, inflammation and metabolic syndrome in melancholic versus atypical depression. *Molecular Psychiatry*, 18(6), 692–699. doi: 10.1038/mp.2012.144
- Li, Y., Zhong, X., Cheng, G., Zhao, C., Zhang, L., Hong, Y., ... Wang, Z. (2017). Hs-CRP and all-cause, cardiovascular, and cancer mortality risk: A meta-analysis. *Atherosclerosis*, 259, 75–82. doi: 10.1016/j.atherosclerosis.2017.02.003
- Malhi, G. S., & Mann, J. J. (2018). Depression. *The Lancet*, 392(10161), 2299–2312. doi: 10.1016/s0140-6736(18)31948-2
- Milaneschi, Y., Lamers, F., Berk, M., & Penninx, B. (2020). Depression heterogeneity and its biological underpinnings: Toward immunometabolic depression. *Biological Psychiatry*, 88(5), 369–380. doi: 10.1016/j.biopsych.2020.01.014
- Miller, A. H., & Raison, C. L. (2016). The role of inflammation in depression: From evolutionary imperative to modern treatment target. *Nature Reviews Immunology*, 16(1), 22–34. doi: 10.1038/nri.2015.5
- Nettis, M. A., Pergola, G., Kolliakou, A., O'Connor, J., Bonaccorso, S., David, A., ... Mondelli, V. (2019). Metabolic-inflammatory status as predictor of

- clinical outcome at 1-year follow-up in patients with first episode psychosis. *Psychoneuroendocrinology*, 99, 145–153. doi: 10.1016/j.psyneuen.2018.09.005
- Nigatu, Y. T., Bultmann, U., & Reijneveld, S. A. (2015). The prospective association between obesity and major depression in the general population: Does single or recurrent episode matter? *BMC Public Health*, 15, 350. doi: 10.1186/s12889-015-1682-9
- Nigatu, Y. T., Reijneveld, S. A., de Jonge, P., van Rossum, E., & Bultmann, U. (2016). The combined effects of obesity, abdominal obesity and major depression/anxiety on health-related quality of life: The lifeLines cohort study. *PLoS One*, 11(2), e0148871. doi: 10.1371/journal.pone.0148871
- Nikkheslat, N., McLaughlin, A. P., Hastings, C., Zajkowska, Z., Nettis, M. A., Mariani, N., ... Mondelli, V. (2019). Childhood trauma, HPA axis activity and antidepressant response in patients with depression. *Brain, Behavior, and Immunity*, 87, 229–237. doi: 10.1016/j.bbi.2019.11.024
- Osimio, E. F., Baxter, L. J., Lewis, G., Jones, P. B., & Khandaker, G. M. (2019). Prevalence of low-grade inflammation in depression: A systematic review and meta-analysis of CRP levels. *Psychological Medicine*, 49(12), 1958–1970. doi: 10.1017/S0033291719001454
- Ouakinin, S. R. S., Barreira, D. P., & Gois, C. J. (2018). Depression and obesity: Integrating the role of stress, neuroendocrine dysfunction and inflammatory pathways. *Frontiers in Endocrinology*, 9, 431. doi: 10.3389/fendo.2018.00431
- Papakostas, G. I., Petersen, T., Iosifescu, D. V., Burns, A. M., Nierenberg, A. A., Alpert, J. E., ... Fava, M. (2005). Obesity among outpatients with major depressive disorder. *The International Journal of Neuropsychopharmacology*, 8(1), 59–63. doi: 10.1017/S1461145704004602
- Pearson, T. A., Mensah, G. A., Alexander, R. W., Anderson, J. L., Cannon, R. O., Criqui, M., ... Vinicor, F. (2003). Markers of inflammation and cardiovascular disease. *Circulation*, 107(3), 499–511. doi: 10.1161/01.cir.0000052939.59093.45
- Perez-Cornago, A., de la Iglesia, R., Lopez-Legarrea, P., Abete, I., Navas-Carretero, S., Lacunza, C. I., ... Zulet, M. A. (2014). A decline in inflammation is associated with less depressive symptoms after a dietary intervention in metabolic syndrome patients: A longitudinal study. *Nutrition Journal*, 13, 36. doi: 10.1186/1475-2891-13-36
- Perrin, A. J., Horowitz, M. A., Roelofs, J., Zunszain, P. A., & Pariante, C. M. (2019). Glucocorticoid resistance: Is it a requisite for increased cytokine production in depression? A systematic review and meta-analysis. *Frontiers in Psychiatry*, 10, 423. doi: 10.3389/fpsy.2019.00423
- Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., & Hellhammer, D. H. (2003). Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology*, 28(7), 916–931. doi: 10.1016/s0306-4530(02)00108-7
- Rethorst, C. D., Bernstein, I., & Trivedi, M. H. (2014). Inflammation, obesity, and metabolic syndrome in depression: Analysis of the 2009–2010 National Health and Nutrition Examination Survey (NHANES). *Journal of Clinical Psychiatry*, 75(12), e1428–e1432. doi: 10.4088/JCP.14m09009
- Ruttle, P. L., Klein, M. H., Slattery, M. J., Kalin, N. H., Armstrong, J. M., & Essex, M. J. (2014). Adolescent adrenocortical activity and adiposity: Differences by sex and exposure to early maternal depression. *Psychoneuroendocrinology*, 47, 68–77. doi: 10.1016/j.psyneuen.2014.04.025
- Schmidt, F. M., Weschenfelder, J., Sander, C., Minkwitz, J., Thormann, J., Chittka, T., ... Himmerich, H. (2015). Inflammatory cytokines in general and central obesity and modulating effects of physical activity. *PLoS One*, 10(3), e0121971. doi: 10.1371/journal.pone.0121971
- Tremmel, M., Gerdtham, U. G., Nilsson, P. M., & Saha, S. (2017). Economic burden of obesity: A systematic literature review. *International Journal of Environmental Research and Public Health*, 14(4), 435–435. doi: 10.3390/ijerph14040435
- Uher, R., Mors, O., Hauser, J., Rietschel, M., Maier, W., Kozel, D., ... Farmer, A. (2009). Body weight as a predictor of antidepressant efficacy in the GENDEP project. *Journal of Affective Disorders*, 118(1–3), 147–154. doi: 10.1016/j.jad.2009.02.013
- Upadhyay, J., Farr, O., Perakakis, N., Ghaly, W., & Mantzoros, C. (2018). Obesity as a disease. *Medical Clinics of North America*, 102(1), 13–33. doi: 10.1016/j.mcna.2017.08.004
- Valkanova, V., Ebmeier, K. P., & Allan, C. L. (2013). CRP, IL-6 and depression: A systematic review and meta-analysis of longitudinal studies. *Journal of Affective Disorders*, 150(3), 736–744. doi: 10.1016/j.jad.2013.06.004
- Zhang, J., Yue, Y., Thapa, A., Fang, J., Zhao, S., Shi, W., ... Yuan, Y. (2019). Baseline serum C-reactive protein levels may predict antidepressant treatment responses in patients with major depressive disorder. *Journal of Affective Disorders*, 250, 432–438. doi: 10.1016/j.jad.2019.03.001