




Review article

A comprehensive review on Nepalese wild vegetable food fernsGan B. Bajracharya ^{a,*}, Bashana Bajracharya ^b

^a Nepal Academy of Science and Technology (NAST), Khumaltar, Lalitpur, Nepal
^b Kirtipur Sadak 27/48, Balkhu, Tekushi, Kathmandu 14, Nepal

HIGHLIGHTS

- Ferns and fern-allies are consumed as a vegetable in many countries.
- Ferns possess several ethnomedicinal values, but are less explored scientifically.
- Flavonoids, terpenoids and steroids are major secondary metabolites found in ferns.
- This review compiles pharmacognosy and phytochemistry of 33 wild ferns eaten by Nepalese as cooked vegetable foods.

GRAPHICAL ABSTRACT**ARTICLE INFO**

Keywords:
 Bioactivity
 Biodiversity
 Food
 Pteridophytes
 Secondary metabolite
 Traditional knowledge

ABSTRACT

Ferns are used as traditional and fascinating foods in many countries. They are also considered to possess important ethnomedicinal values; however, ferns are one of the underutilized plant resources by both scientific and local communities. Pharmacognostical studies reveal that ferns and fern-allies possess several biological activities including antibacterial, antiviral, antifungal, antimarial, antidiarrheal, anthelmintic, analgesic, anti-inflammatory, antidiabetic, anticancer, neuroprotective, nephroprotective, hepatoprotective, antifertility, etc. Flavonoids and terpenoids are major secondary metabolites present in ferns. Ugonins, particularly isolated from *Helminthostachys zeylanica*, have found diverse bioactivities. Ptaquiloside, a norsesterpenoid glucoside, found in *Pteridium revolutum*, *Dryopteris cochlearia* and *Polystichum squarrosum*, is one of the hazardous metabolites of ferns which is responsible for the toxic effect. Alkaloids are reported to be present in the ferns; however, the qualitative data are uncertain. Some fern metabolites, such as cyanogenic glycosides and terpenoids, are considered to possess defensive activity against animal attacks. Some ferns are also used for manuring as biological alternative to pesticides. Nepalese have consumed at least 33 species of ferns and fern-allies belonging to 13 families, 20 genera as cooked vegetable foods. The aim of this review is compilation of information available on their distribution, ethnomedicinal values, pharmacognosy, pharmacology and phytochemistry.

1. Introduction

Nepal is a small land-locked country resides on the lap of the great Himalaya, in the north. It is sandwiched between China and India. Despite occupying only 0.1% of the global area, the country is naturally blessed and enriched with cultural, geographical and biological diversities [1]. Its cultural diversities are interlinked with more than 60

ethnic groups speaking more than 100 languages. Geographically, it has a large altitudinal variations ranging from a low land (altitude nearly 60 m in south) to the highest peak of the earth (Mt. Everest, 8,848.86 m in north) within a span of just about 200 km of aerial expansion from south to north. Its physiographic regions comprises (1) Tarai, (2) Siwaliks and Chure, (3) Mahabharat, (4) Midlands or central hills, (5) Himalayas, and (6) Inner Himalayas and Tibetan marginal mountains [2]. The country

* Corresponding author.

E-mail addresses: ganbajracharya@yahoo.com, gan.bajracharya@nast.gov.np (G.B. Bajracharya).

has 118 different types of ecosystems harboring 3.2% of the world's flora including 5.1% of gymnosperms, 3.2% of angiosperms, 5.1% of pteridophytes, 8.2% of bryophytes, 2.5% of algae, 2.6% of fungi and 2.3% of lichens [1].

For food and livelihood securities, Nepalese significantly depend on cultivated and wild vegetables [3]. In Nepal, it has been estimated that >650 species are used as food materials, and >1,000 species possess medicinal values [4]. Recently, we have listed 318 wild plant species (excluding mushrooms, fruits, spices and condiments, seeds and pulses) that are consumed by Nepalese as vegetables [5]. Wild vegetables are indispensable constituent of the human diet and are the nature's gift to mankind. They are not only cheap but also supply minerals, vitamins and certain hormone precursors. Among them, wild edible ferns play an important role in diet, and they are also a measure of income source with no investment.

Wild plants are globally used by human societies for food, medicine, ornament, defence and industrial purposes. Ferns and fern-allies have unique features in their appearance, natural habitat, food value and ethnomedicinal properties. Ferns possess a wide range of medicinal activities; unfortunately, they are poorly investigated. Comprehensive review on phytochemicals and bioactivities of ferns and fern-allies is rare. It is commonly considered that plants grown in harsh environmental conditions usually possess incredible bioactivity. Therefore, writing of this review article is envisioned to increase scientific attention on the fern species, particularly those grown in high altitudes of Nepal. Consequently, an extensive literature search was conducted. Dissertations, books, journal articles and conference proceedings on ethnobotanical surveys in Nepal were accessed in the libraries to generate a list of wild vegetable food ferns of Nepal. Taxonomic names mentioned in the original articles were verified using several efloral databases. Online literature search was conducted in Google Scholar and Pub Med by using the terms ferns, pteridophytes and individual name of the fern species. Cited references in the published research articles were also traced to obtain information on pharmacognosy, phytochemicals and bioactivities of the fern species grown across the globe.

2. Ferns and fern-allies (pteridophytes)

Fern and fern-allies (pteridophytes) are considered as a very ancient family. They are the first vascular land plants that evolved during the Devonian and Carboniferous periods of the Paleozoic era [6]. Pteridophytes dominated the earth's vegetation in the beginning of the Mesozoic era, about 280–230 million years ago [7, 8]. Ferns are consumed as vegetables in many countries across the globe and they are also used in the traditional Chinese medicines [9, 10]. However, surprisingly, compared to other vascular plants, pteridophytes are remain under-explored both in ethnobotanical and pharmacological aspects. In the last decade, some natural compounds have been isolated from different fern species and some pharmacognostical studies are begun [11]. Initially, traditional knowledge based ferns were ethnomedicinally used for some ailments, and the major studies on pteridophytes were primarily focused on taxonomic identification.

Ferns are seedless vascular plants and they reproduce by sporulation. They have many similar features of mosses and algae, but are usually differentiated from them by having xylem and phloem to transport water and nutrient materials. Like other vascular plants (gymnosperms and angiosperms), they possess stems, fronds, pinnae and roots. The life-cycle of fern is referred to as alternation of generations with two different stages—gametophyte phase (sexual) and sporophyte phase (asexual) (Figure 1). Motile male gametes (antherozoids) are produced from antheridia and non-motile female gametes (egg cells) are born singly in archegonia. Fusion of both the gametes results in the formation of a zygote, which develops into the sporophyte (diploid) after mitotic divisions. Fern sporophytes are free-living, independent on the gametophyte (prothallus), dominant and grow to a much larger size. Mature sporophyte bears sori on the underside of the blade. Sporangia release a number of non-motile spores that germinate and grow by mitotic divisions into haploid gametophytes.

It has been estimated that there are around 4,03,000 plant species in the earth including phanerogams and cryptogams [12]. About 13,271 species of pteridophytes are distributed which forms nearly 3% of the




Figure 1. Fern lifecycle.

Table 1. List of edible ferns and fern-allies of Nepal that eaten as vegetables.

SN	Botanical name [synonym]	Local name (in different community)	Habit	Vegetative parts	Altitude	Season	Availability	References
Athyriaceae								
1	<i>Athyrium atkinsonii</i> Bedd.	Niuro	Herb	Young leaf	1,200–4,000	-	Common	[25, 31, 72]
2	<i>Athyrium strigilloseum</i> (T. Moore ex E. J. Lowe) T. Moore ex Salomon	Niguro	Herb	Young frond	600–2,600	-	Common	[25, 31]
3	<i>Deparia boryana</i> (Willd.) M. Kato [<i>Dryoathyrium boryanum</i> (Willd.) Ching]	Kalo niuro, Dhengan (Tamang)	Herb	Leaf, tender shoot	400–3,300	May–Jul	Common	[26, 31, 73, 74, 75, 76]
4	<i>Diplazium esculentum</i> (Retz.) Sw.	Masino niuro, Chipley niuro, Kuturke, Pani nyuro, Piraunli	Herb	Young leaf	100–1,300	Apr–Oct	Common	[23, 26, 31, 73, 74, 77, 78]
5	<i>Diplazium kawakamii</i> Hayata	Niuro, Jire ningro (Darjiling)	Herb	Tender shoot	1,300–2,400	-	Common	[31, 59, 72]
6	<i>Diplazium maximum</i> (D. Don) C. Chr. [<i>Allantodia maxima</i> (D. Don) Ching]	Neuro	Herb	Leaf	9,00–1,800	May–Jul	Common	[26, 31, 79]
7	<i>Diplazium spectabile</i> (Wall. ex Mett.) Ching [<i>Allantodia spectabilis</i> (Wall. ex Mett.) Ching]	Saune niuro	Herb	Leaf	1,500–2,700	May–Jul	Common	[26, 31]
8	<i>Diplazium stoliczkae</i> Bedd. [<i>Allantodia stoliczkae</i> (Bedd.) Ching]	Kalo niuro, Lekh chipley niuro, Bhandhengan (Tamang)	Shrub	Tender shoot, leaf	1,300–4,000	May–Jun	Common	[26, 31, 76, 80]
Blechnaceae								
9	<i>Stenochlaena palustris</i> (Burm. f.) Bedd.	-	Climber	Young shoot	60–900	-	Rare	[25, 31, 72]
10	<i>Woodwardia unigemmata</i> (Makino) Nakai	-	Shrub	Stem	400–3,000	Jun–Jul	Common	[26, 31, 73, 74]
Cyatheaceae								
11	<i>Cyathea spinulosa</i> Wall. ex Hook. [<i>Alsophila spinulosa</i> (Wall. ex Hook.) R.M. Tryon]	Chattre, Rukh uniyu, Thulo uniyu, Jatashankari (Indian)	Shrub	Young shoot	300–1,600	Jul–Aug	Scattered to common	[23, 31, 81]
Dennstaedtiaceae								
12	<i>Pteridium revolutum</i> (Blume) Nakai [<i>Pteridium aquilinum</i> subsp. <i>wightianum</i> (J. Agardh) W.C. Shieh]	Saune niuro, Ainu	Herb	Shoot	600–3,000	Jul–Jul	Common	[8, 26, 31, 73, 74]
Dryopteridaceae								
13	<i>Dryopteris cochleata</i> (D. Don) C. Chr.	Pani niuro, Ghee niuro, Danthe niuro, Lotah niuro	Herb	Tender leaf	1,200–1,600	Mar–Oct	Common	[23, 26, 31, 72, 78, 80, 82]
14	<i>Dryopteris splendens</i> (Hook. ex Bedd.) Kuntze	Danthe neuro	Herb		<1,000	Jul–Sep	Scattered	[31, 83]
15	<i>Polystichum squarrosum</i> (D. Don) Fée [<i>Polystichum apicisterile</i> Ching & S.K. Wu]	Phusre niuro, Rato unyu	Herb	Shoot	1,900–2,400	May–Jun	Common	[26, 31, 73, 74, 80]
Lygodiaceae								
16	<i>Lygodium flexuosum</i> (L.) Sw.	Parandi sag, Lahare unyu	Climber	Young shoot, leaf	60–1,000	Whole year	Common	[23, 31, 74]
17	<i>Lygodium japonicum</i> (Thunb.) Sw.	Janai lahara, Nagbeli, Lute jhar, Ukuje jhar (Magar), Pinse (Tamang)	Climber	Leaf	60–3,900	May–Jun	Common	[23, 26, 31, 74, 84]
Marsileaceae								
18	<i>Marsilea quadrifolia</i> L.	Charpate behuli	Herb	Tender shoot	900–1,700			[85]
Nephrolepidaceae								
19	<i>Nephrolepis cordifolia</i> (L.) C. Prest	Pani amala, Bhui amala, Ras amala, Ambeli (Tamang), Pani saro (Majhi)	Herb	Tuber, shoot	60–2,400	Jul–Sep	Common	[23, 31, 74, 75, 77, 81]
Ophioglossaceae								
20	<i>Botrychium lanuginosum</i> Wall. ex Hook. & Grev.	Jaluko, Bayakhra	Herb	Shoot	1,000–3,000	May–Jun	Common	[26, 31, 74]

(continued on next page)

Table 1 (continued)

SN	Botanical name [synonym]	Local name (in different community)	Habit	Vegetative parts	Altitude	Season	Availability	References
21	<i>Helminthostachys zeylanica</i> (L.) Hook	Majurgoda (Chepang), Kamsaj (Tharu)	Herb	Tender shoot	<1,000		Scattered	[31, 75]
22	<i>Ophioglossum nudicaule</i> L.f.	Jibre sag	Herb	Leaf	1,800–4,300	Mar–Apr	Common	[26, 73, 74]
23	<i>Ophioglossum petiolatum</i> Hook.	Jibre sag	Herb	Leaf	200–3,300	Feb–Apr	Scattered	[31, 84]
24	<i>Ophioglossum reticulatum</i> L.	Jibre sag, Ek patiya (Tharu), Jibre dhap (Tamang)	Herb	Young leaf, tender shoot	1,100–4,000	Mar–Apr	Common	[26, 31, 73, 74, 77, 80]
25	<i>Ophioglossum vulgatum</i> L.	Jibre sag	Herb	Whole plant	<3,000	Mar–Apr	Common	[74]
Osmundaceae								
26	<i>Osmunda claytoniana</i> L. [Osmunda claytoniana subsp. <i>vestita</i> (Milde) Å. Löve & D. Löve]	Kuthurke	Herb	Leaf	1,600–3,400	May–Jun	Common	[26, 31, 73, 74, 77]
Pteridaceae								
27	<i>Ceratopteris thalictroides</i> (L.) Brongn.	Pani dhaniya, Dhaniya jhar	Herb	Whole plant	<1,000	Jun–Aug	Scattered	[23, 31, 74, 85, 86, 87]
28	<i>Pteris biaurita</i> L. [<i>Pteris flavigaulis</i> Hayata]	Haade unyu, Kuthurke niuro (Raji)	Herb	Young leaf, shoot	200–2,500	Aug–Oct	Common	[23, 31, 78, 88]
29	<i>Pteris vittata</i> L.	Unigar, Urakthewn	Herb	Shoot, rhizome	400–3,000	Jul–Sep	Common	[23, 31, 74]
Tectariaceae								
30	<i>Tectaria gemmifera</i> (Féé) Alston [<i>Tectaria coadunata</i> C. Chr.]	Danthe niuro, Kuthurke, Toplign degni (Tamang)	Herb	Young leaf, shoot	500–2,500	Jun–Oct	Common	[23, 26, 31, 73, 74, 78, 82, 86, 89, 90]
31	<i>Tectaria zeylanica</i> (Houtt.) Sledge [Osmunda zeylanica L.]	Mayur kutea, Dhagrjawwa (Tharu)	Herb	Leaf	100–1,000	Mar–Apr		[74, 77]
Thelypteridaceae								
32	<i>Thelypteris auriculata</i> (J. Sm.) K. Iwats [<i>Cyclosorus auriculata</i> (J. Sm.) C.M. Kuo]	Kochaya	Herb	Shoot	1,600–1,800	Mar–May	Common	[31, 74]
33	<i>Thelypteris nudata</i> (Roxb.) C.V. Morton [<i>Thelypteris multilineata</i> (Wall. ex Hook.) C.V. Morton; <i>Pronephrium nudatum</i> (Roxb.) Holttum]	Koche (Tharu)	Herb	Shoot	2,100	Jun–Jul	Common	[26, 31, 73, 74]

world flora [13]. *The Plant List* has listed 35 plant families and 568 genera of pteridophytes [14]. About 63 families, 230 genera and 2,600 species of pteridophytes have been reported in China [15]. From India, 33 families, 130 genera and 1,267 species of ferns and fern-allies are reported [16]. About 37 families, 687 species in Japan [17]; 801 species in Taiwan [18]; 39 families, 144 genera, 1,100 species in the Philippines [19]; 1,165 species in Malaysia [20], and 670 species of pteridophytes in Thailand [21] have been reported.

In Nepal, ferns are called as “Unyu/Oony” and they are distributed between 60 to 4,800 m above sea level (asl) [22, 23]. Thapa *et al.* [24] have listed 535 species of pteridophytes belonging to 35 families and 102 genera from Nepal. Recently, a book *Ferns and Fern-allies of Nepal* [25] has been published including annotated checklist and critical account of 550 species and additional 32 sub-species, altogether 582 taxa of fern and fern-allies of Nepal belonging to 32 families and 99 genera. In Nepal, *Deparia boryana*, *Diplazium esculentum*, *Dryopteris cochleata*, *Ophioglossum vulgatum*, *Pteridium revolutum*, *Diplazium maximum*, *Diplazium spectabile*, *Diplazium stoliczkae*, *Polystichum squarrosum*, *Pteris biaurita*, *Tectaria gemmifera*, etc. are popularly consumed as vegetables, and they are also sold in the markets [5, 26]. Popularity gained in consumption of ferns in Nepalese communities is not only due to their unique taste but also due to beliefs of their high nutritional contents such as vitamin C and iron [27]. Obviously, it is considered that the nutritional value and mineral content in wild edible plants are richer than that of commercial vegetables [28]; therefore, their consumption for the nutritional purpose should be

encouraged. At the same time, their poor availability and prone vulnerability are the serious issues.

3. History, expeditions and literature on Nepalese pteridophytes

Species Plantarum, originally published in 1753 by Carl Linnaeus, is the first botanical work that applied the binomial nomenclature system for the taxonomic description of 5,940 plant species, and recognized 15 fern genera [29]. The pioneering plant collection and taxonomic study of Nepalese pteridophytes were first begun in the early 19th century when Scottish Francis Buchanan (later known as Francis Hamilton) collected 433 plant specimens including 34 species of Nepalese pteridophytes in 1802–1803 and published *An Account of Kingdom of Nepal* [30, 31, 32]. In subsequent historical book publications, namely, *Tentamen Florae Nepalensis Illustratae* by Wallich [33], *Prodromus Flora Nepalensis* by Don [34], *The Flora of British India* by Hooker [35], *Index Filicum* by Moore [36], *A Priced Catalogue of Hardy Exotic and British Ferns* by Sim [37], *Catalogue of the Plants of Kumaon* by Duthie [38], *Notes from a Journey to Nepal* by Burkhill [39] and *A Plantsman in Nepal* by Lancaster [40] have documented many Nepalese ferns. Some foreign authors have also contributed in the taxonomic identification of pteridophytes of Nepal [41, 42, 43, 44].

After establishment of Botanical Survey and Herbarium Office by Nepal Government in 1961 (later renamed as National Herbarium and Plant Laboratories), Nepalese botanists have started collection of plants and

preservation of herbariums, and then they could publish several articles and books [45]. Earlier contributions are made by Gurung [46, 47, 48, 49] and Manandhar [50, 51, 52, 53] followed by others [24, 54, 55, 56, 57].

A major contribution of plant collection and taxonomic studies on Nepalese pteridophytes including Sikkim Himalayan ferns has been made by Christopher Fraser-Jenkins, a Welsh pteridologist, who resides for 40 years in Kathmandu, Nepal. Formerly, he was a research fellow at the Natural History Museum, London, and at the Royal Botanic Garden, Edinburgh. Lately, he worked in conjunction with the National Herbarium and Plant Laboratories Godawari, Nepal. He has spent more than five and half decades in studying Himalayan pteridophytes and published *Ferns and Fern-allies of Nepal*, Vols. 1–3 [58, 59, 25]; *Annotated Checklist of Indian Pteridophytes*, Parts 1–3 [60, 61, 62]; including a series of other publications [63, 64, 65].

A number of works on exploration, taxonomic identification and documentation along with rare molecular study on Nepalese pteridophytes have been made; however, not only Nepalese but also global edible pteridophytes are poorly investigated for their phytochemical constituents and pharmacological effects. Therefore, exploitation of both edible and non-edible fern flora deserves serious attention by scientific communities [66].

4. A list of wild edible ferns of Nepal

From the literature survey of the publications on Nepalese ferns by various authors, we have generated a list of 33 edible fern species (**Table 1**). The taxonomic identification of the species was followed according to Kramer and Green [67], modified by Fraser-Jenkins [65], and was cross-verified in the e-databases of *The Plant List* [68] and *Flora of China* [69]. *Cornopteris decurrenti-alata* and *Matteuccia intermedia* are rarely occurred in Nepal, while *Blechnum orientale*, *Osmunda japonica*, *Coniogramme intermedia* and *Pteris wallichiana* are common species [31]. These species are used as fern food in China [70]. We could not find proper documentation for these pteridophytes as wild edible ferns of Nepal; therefore, they are omitted in the list. The pteridophytes, which consumed for the medicinal values, but not eaten as salad or cooked vegetables, are also not included in the table [71].

5. Distribution, ethnomedicine, pharmacognosy and phytochemistry of the wild edible ferns

It has been recently reported that there are 582 taxa of pteridophytes in Nepal [72]; however, Nepalese researchers have not been focused in the ethnopteridological studies in traditional medicinal knowledge, food safety and phytoconstituents present in the pteridophytes. At the same time, surprisingly, very few scientific studies have been carried out globally in the areas of the chemical constituents and pharmacological activities of pteridophytes [71]. In this section, we accumulate available information on the distribution, ethnomedicinal uses, pharmacognosy and phytochemistry of the wild edible ferns listed in **Table 1**.

C (central), N (north), E (east), W (west) and S (south) abbreviations are used to indicate distribution of the fern species. The edible fern distribution in China and India is specified as available in different provinces.

5.1. *Athyriaceae*

5.1.1. *Athyrium atkinsonii*

5.1.1.1. Distribution. W, C and E Nepal; Chongqing, Fujian, Gansu, Guizhou, Henan, Hubei, Hunan, Jiangxi, Shaanxi, Shanxi, Sichuan, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; Myanmar; Taiwan; S Korea; Japan; etc. [31, 69].

5.1.1.2. Habitat. Damp forest at high altitude.

5.1.1.3. Phytochemistry. The plant material contains carbohydrates, proteins, alkaloids, phenolics, flavonoids, etc. [91].

5.1.2. *Athyrium strigillosum*

5.1.2.1. Distribution. W, C and E Nepal; Guangdong, Guangxi, Guizhou, Hunan, Jiangxi, Sichuan, Xizang and Yunnan; Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; Myanmar; Taiwan; Japan; etc. [31, 69].

5.1.2.2. Habitat. Wet forest and partially exposed area or along banks and beds of well-shaded ravines.

5.1.3. *Deparia boryana* [*Dryoathyrium boryanum*]

5.1.3.1. Distribution. W, C and E Nepal; Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hunan, Shaanxi, Sichuan, SE Xizang, Yunnan and Zhejiang; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Bhutan; Sri Lanka; Myanmar; Thailand; Vietnam; Malaysia; Indonesia; Philippines; Taiwan; Africa; etc. [31, 69].

5.1.3.2. Habitat. Densely forested ravine with waterfalls, moist and shade places.

5.1.3.3. Ethnomedicine. The tender shoot is used as demulcent, stomachic and laxative [92]. The powder or paste is used to treat dysentery [93]. Rhizome is used in abdominal spasm [75]. Root paste is applied externally to treat burns, injury and wounds [76].

5.1.3.4. Pharmacognosy. The ethyl acetate extract inhibits the lipid formation with 35% at 100 µg/mL in 3T3-L1 cell model [92]. Antioxidant as well as anticancer activities are explored [92, 94].

5.1.3.5. Phytochemistry. The polyphenolic content in the aerial parts is 266 µg gallic acid equivalent/mg [92]. The total flavonoids content is estimated about 145.8 mg/g [94]. By using HPLC-DAD-ESI-MS, 3-hydroxyphloretin 6'-O-hexose, querctein-7-hexoside, apigenin7-O-glucoside, luteolin 7-O-glucoside, apigenin 7-O-galactoside, acacetin 7-O-(α -D-apio-furanosyl) (1–6)- β -D-glucoside, 3-hydroxy phloretin 6-O-hexose, luteolin-6-C-glucoside, kaempferol 3-O-glucoside, etc. are tentatively identified in the plant extract [94].

5.1.4. *Diplazium esculentum*

5.1.4.1. Distribution. W, C and E Nepal; Anhui, Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hunan, Jiangxi, Sichuan, Xizang, Yunnan and Zhejiang; Arunachal; Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; Taiwan; etc. [31, 69].

5.1.4.2. Habitat. Moist, shade forest and rocks.

5.1.4.3. Ethnomedicine. Young fronds are used as tonic and eaten for laxative effect [95]. Aerial parts are used to treat fever, pain, glandular swelling, diarrhea, dermatitis, wound and measles. The rhizome is considered as anthelmintic, antidiarrheal and pest repellent [96]. It is also used in cough, asthma, fever, dyspepsia and stomachache. Decoction of rhizome and young leaves are useful in haemoptysis and constipation. Rhizomes are also used in scabies and boils [53]. A paste of leaf and stem is applied externally to treat cuts and wounds [97].

5.1.4.4. Pharmacognosy. Biological activities of the leaf extracts such as antimicrobial [98, 99, 100], antioxidant [101, 102, 103, 104], antidiabetic [103], anticancer [105], anti-inflammatory [99, 106], hepatoprotective [103], anti-Alzheimer's [104], etc. have been reported. Antibacterial [96] and anthelmintic activities of the rhizomes are reported [107].

5.1.4.5. Phytochemistry. Estimation of quercetin in the plant is performed [108]. Proximate analysis shows the presence of fat (3.4%), protein (10.7%), fiber (9.1%), total phenolics (125.6 mg gallic acid equivalent/100 g) and total flavonoids (110.8 mg quercetin equivalent/100 g) [109]. Total phenolics (167.7 mg gallic acid equivalent/100 g) and total flavonoids (70.8 mg quercetin equivalent/100 g) in the hydroalcoholic extract are separately reported by Junejo *et al.* [103]. Qualitative phytochemical screening shows the presence of carbohydrates, fats, alkaloids, anthraquinones, phenolics, saponins, cyanidins, cardiac glycosides, steroids, flavonoids, terpenoids, etc. [110]. Evaluation of vitamin C (22.2 mg/100 g), mineral contents (Ca, Cu, Fe, Mg, Mn, Na and Zn), heavy metals (Al, As, Cd, Hg, Li, Ni and Pb), etc. are performed in the leaves [27]. β -Pinene (17.2%), α -pinene (10.5%), carophyllene oxide (7.5%), sabinene (6.1%), 1,8-cineole (5.8%), α -cardinol (4.8), β -caryophyllene (4.3%) are identified as the major volatile constituents in the essential oil obtained from the leaves [111]. Pentadecanoic acid, α -linolenic acid, phytol, β -sitosterol, stigmasta-5, 22-dien-3-ol, methyl palmitate, diisobutyl phthalate, neophytadiene, 10,12-hexadecadien-1-ol, etc. are identified in the extracts of young fronds from Indian Himalaya [112].

Ascorbic acid [113], eriodictyol 5-O-methyl ether 7-O- β -D-xylosylgalactoside [114], α -tocopherol [115], quercetin [108, 116], ptaquiloside [117], pterosin [108], hopan-triterpene lactone [118], lutein [119], etc. are isolated from the edible parts of *D. esculentum*. Phenolics ((2R)-3-(4'-hydroxyphenyl)lactic acid, trans-cinnamic acid, protocatechuic acid and rutin) and ecdysteroids (amarasterone A1, makisterone C and ponasterone A) are recently isolated from the young fronds collected in Japan [120].

5.1.5. *Diplazium kawakamii*

5.1.5.1. Distribution. E Nepal; Yunnan; Arunachal, Sikkim and Darjeeling; Bhutan; Taiwan; Japan; etc. [31, 69].

5.1.5.2. Habitat. Wet, densely forested ravine and besides streamlets.

5.1.6. *Diplazium maximum* [*Allantodia maxima*]

5.1.6.1. Distribution. W, C and E Nepal; Fujian, Guangxi, Guizhou, Hainan, Jiangxi, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Afghanistan; Bhutan; Myanmar; etc. [31, 69].

5.1.6.2. Habitat. Valley, evergreen forest, besides streamlets and north-west facing slopes.

5.1.6.3. Ethnomedicine. Young shoots are used in stomach problem [79].

5.1.6.4. Pharmacognosy. Fronds exhibit antioxidant potentiality [121].

5.1.6.5. Phytochemistry. The plant material contains alkaloids, phenolics, tannins, saponins, flavonoids, terpenoids, etc. [91]. Chettri *et al.* have estimated total crude fat (2.5%), fiber (11.3%), protein (13.4%), vitamin C (25.4 mg/100 g) and several metal contents [27]. Fronds have high contents of dietary fibre (38.3%) and crude protein (25.4%) [121]. It constitutes several fatty acids and phenolics like catechin, epicatechin, procatechuric acid, myricetin, etc.

5.1.7. *Diplazium spectabile* [*Allantodia spectabilis*]

5.1.7.1. Distribution. W, C and E Nepal; Yunnan and Xizang; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Bhutan; etc. [31, 69].

5.1.7.2. Habitat. South facing forest and valley, evergreen broad-leaved forests and besides streamlets.

5.1.7.3. Phytochemistry. Crude fat (0.2%), fiber (3.4%), protein (3.6%), vitamin C (21.6 mg/100 g), mineral contents (Ca, Cu, Fe, Mg, Mn, Na and Zn), heavy metals (Al, As, Cd, Hg, Li, Ni and Pb), etc. are determined in the leaves [27].

5.1.8. *Diplazium stoliczkae* [*Allantodia stoliczkae*]

5.1.8.1. Distribution. C and E Nepal; Xizang; Arunachal; Sikkim and Darjeeling; Bhutan; etc. [31].

5.1.8.2. Habitat. Dense forest.

5.1.8.3. Ethnomedicine. Root paste is applied externally to cure burn, injury and wounds [53]. Leaf juice is given to treat diarrhea and dysentery [122].

5.1.8.4. Phytochemistry. Total phenolic content (440.6 mg gallic acid equivalent/g) and flavonoid content (625.5 quercetin equivalent/g) together with antioxidant activity in the rhizomes have been investigated [123].

5.2. *Blechnaceae*

5.2.1. *Stenochlaena palustris*

5.2.1.1. Distribution. E Nepal; Guangdong, Guangxi, Hainan and Yunnan; Arunachal, Assam, Sikkim and Darjeeling; Thailand; Laos; Vietnam; Cambodia; Malaysia; Indonesia; Australia; Pacific islands; etc. [31, 69].

5.2.1.2. Habitat. Secondary forest, open forests and interior on the hills.

5.2.1.3. Ethnomedicine. Leaves decoction cures fever [124]. Fronds are used traditionally to treat fever, diarrhea, skin diseases, cutaneous disorders and gastric ulcers [125, 126].

5.2.1.4. Pharmacognosy. Fronds extracts show antifungal [127], anti-bacterial [128], antioxidant [129, 130, 131], and butyrylcholinesterase inhibitory [126] activities. Antimalarial activity of the leaf extract is recently reported [132].

5.2.1.5. Phytochemistry. Leaves constitute stenopalustrosides A–E, kaempferol 3-O-(3"-O-E-p-coumaroyl)-(6"-O-E-feruloyl)- β -D-glucopyranoside, kaempferol 3-O-(3",6"-di-O-E-p-coumaroyl)- β -D-glucopyranoside, kaempferol 3-O-(3"-O-E-p-coumaroyl)- β -D-glucopyranoside, kaempferol 3-O-(6"-O-E-p-coumaroyl)- β -D-glucopyranoside, kaempferol 3-O- β -D-glucopyranoside, (4S*,5R*)-4-[{(Z)-2,13-di-(O- β -D-glucopyranosyl)-5,9,10-trimethyl-8-oxo-9-tetradecene-5-yl]-3,3,5-trimethylcyclohexanone, 1-O- β -D-glucopyranosyl-(2S*,3R*,4E,8Z)-2-N-[(2R)-hydroxytetracosanoyl]octadecasphinga-4,8-dienine, 3-oxo-4,5-dihydro- α -ionyl β -D-glucopyranoside, 3-formylindole, lutein and β -sitosterol-3-O- β -D-glucopyranoside [128, 133].

Fronds contain kaempferol 3-O- β -glucopyranoside, kaempferol 3-O-(3"-O-E-p-coumaroyl)-(6"-O-E-feruloyl)- β -glucopyranoside, kaempferol 3-O-(3",6"-di-O-E-p-coumaroyl)- β -glucopyranoside, kaempferol 3-O-(6"-O-E-p-coumaroyl)- β -glucopyranoside, kaempferol 3-O-(3"-O-E-p-coumaroyl)- β -glucopyranoside, kaempferol 3-O- α -rhamnopyranoside and kaempferol 3-O-(6"-O- α -rhamnopyranoside)- β -glucopyranoside [126].

5.2.2. *Woodwardia unigemmata* [*Woodwardia biserrata*]

5.2.2.1. Distribution. W, C and E Nepal; Fujian, Gansu, Guangdong, Guangxi, Guizhou, Hubei, Hunan, Jiangxi, Shanxi, Sichuan, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; Myanmar; Vietnam; Philippines; Taiwan; Japan; etc. [31, 69].

5.2.2.2. Habitat. Densely forested gully and slopes below trees, exposed and shaded places or wet localities along roadsides.

5.2.2.3. Ethnomedicine. Fronds are used in dysentery, skin diseases and infirility [134, 135, 136]. Leaf is used for abdominal pain, constipation and sore throat [135]. Rhizome is used as purgative.

5.2.2.4. Pharmacognosy. Antioxidant [137, 138] and antibacterial [138] activities of the aerial parts are reported. Anti-HIV activity has been shown by the rhizome extracts [139, 140].

5.2.2.5. Phytochemistry. Total phenolic and flavonoid contents in the aerial parts are reported [137, 138]. GC-MS analysis of the aerial parts shows the presence of catechol (22%), glycerol (20.2%), n-pentadecanoic acid (6.9%), glyceryl monoacetate (6.3%), ethyl acetimidate (5.4%) and 3-hydroxy-2,3-dihydromalton (5.4%) in the aqueous extract; β -sitosterol (17.4%), pentadecanoic acid (9.8%), vitamin E (7.8%) and glycerol (7.0%) in the methanolic extract; and γ -sitosterol (33.4%), vitamin E (10.0%) and campesterol (7.3%) in the hexane extract [138]. From the roots, crassirhizomosides B-C, sutchuenoside A, kaempferol 3-O- α -L-(3-O-acetyl) rhamopyranoside-7-O- α -L-rhanopyranoside, kaempferol 3-O- α -L-(2-O-acetyl) rhamopyranoside-7-O- α -L-rhanopyranoside, kaempferol 3-O- α -L-rhamopyranoside-7-O- α -L-rhanopyranoside and kaempferol 3-O- α -L-(3,4-O-di-acetyl) rhamopyranoside-7-O- α -L-(4-O-acetyl)rhanopyranoside are isolated [11].

5.3. Cyatheaceae

5.3.1. Cyathea spinulosa [*Alsophila spinulosa*]

5.3.1.1. Distribution. W, C and E Nepal; Chongqing, Fujian, Guangdong, Guangxi, Guizhou, Hainan, Jiangxi, Sichuan, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling and Uttarakhand; Bhutan; Bangladesh; Sri Lanka; Myanmar; N Thailand; Taiwan; S Japan; etc. [31, 69].

5.3.1.2. Habitat. Densely forested ravine with waterfalls and shade places.

5.3.1.3. Ethnomedicine. It is useful as hair tonic, sudorific and aphrodisiac [141]. Young leaf is taken orally in excessive bleeding during menstrual period [97]. Pith decoction is given in pains and bodyaches [142]. Rhizomes are used againts snake bites [143].

5.3.1.4. Pharmacognosy. Leaves possess antioxidant and anti-Alzheimer's potentiality [144].

5.3.1.5. Phytochemistry. From the fronds, hegovlavones A-B are isolated [145]. From the leaves, (2S,3S,4R)-2-[(2'R)-2'-hydroxytetracosanoylamino]-1,3,4-octadecanetriol, 1-O- β -D-glucopyranosyl-(2S,3R,4E,8Z)-2-[(2-hydroxyoctadecanoyl) amido]-4,8-octadecadiene-1,3-diol, 3 α -hydroxyfilic-4(23)-ene, 2-oxofilic-3-ene, hop-22(29)-ene fern-7-ene, cycloaudenyl palmitate, daucosterol, dryocrasol, ergosterol, fern-9(11)-ene, filic-3-ene, hopan-29,17 α -olide, hopan-17 α ,29-epoxide, hydroxyhopane, protocatechuic aldehyde, sitostanol, sitosterol, sitosteryl palmitate, stigmast-3,6-dione, stigmast-4-ene-3,6-dione and tetrahymanol are isolated [146, 147]. From the stalks, (E)-4-O- β -D-glucopyranosyl caffeic acid, 1-O-hexadecanolenin, 3,4,6-trihydroxy-1-ring vinyl carboxylic acid, 30-O- β -D-xylopyranosyl-dryocrassol, 4-O- β -D-glucopyranosyl-p-coumaric acid, 6 β -hydroxy-24-ethyl-cholest-4-en-3-one, 9 α -hydroxy-1 β -methoxyaryolanol, clovandiol, cyathenosin A, daucosterol, decumbic acid, iso-rientin, n-tetracosane, pimaric acid, protocatechualdehyde, protocatechuic acid, stigmastane-3,6-dione, vitexin and β -sitosterol are isolated [148, 149].

5.4. Dennstaedtiaceae

5.4.1. *Pteridium revolutum* [*Pteridium aquilinum* subsp. *Wightianum*]

5.4.1.1. Distribution. W, C and E Nepal; Gansu, Guangdong, Guangxi, Guizhou, Henan, Hubei, Hunan, Jiangxi, Shaanxi, Sichuan, Xizang, Yunnan and Zhejiang; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; Taiwan; N Australia; etc. [31, 69]. *P. aquilinum*, known as bracken fern, is one of the most commonly distributed weed species that grows in many types of soil with extreme high rate of reproduction [150].

5.4.1.2. Habitat. Exposed and shaded places, and open humus-rich forests.

5.4.1.3. Ethnomedicine. It is an ornamental fern that planted for indoor decoration and borders [151]. It is traditionally used for animal bedding [52]. The rhizomes are astringent and useful in diarrhea, intestinal inflammation and wounds. The rhizomes are also used for making breads and brews. The plant juice is considered antibacterial.

5.4.1.4. Pharmacognosy. It contains poisonous cyanogen glycosides which are antithiamine and carcinogenic. It has been demonstrated that feeding of the plant induces urinary bladder tumors, illeum sarcomas, pulmonary adenomas and leukemia [152, 153]. However, Yoshihira *et al.* performed toxicity tests and found that none of the plant extracts and the fractions could induce tumors under the conditions studied [154].

5.4.1.5. Phytochemistry. The rhizome constitutes the bitter starch [151]. It contains carbohydrate (51%), protein (9.5%), fat (1.2%), fibre (20%) and ash (8.3%). The fronds contain protein (1%), fat (0.1%), fibre (1.4%), minerals (0.6%), β -carotene (1 mg/100 g) and iodine (900 μ g/kg). Phytochemical screening of the fronds grown in the Philippines reveals the presence of alkaloids, flavonoids and tannins [99]. Prunasin, a main cyanogen component in the fern that can release HCN to induce cyanogenesis, has been estimated about 10.4–61.3 mg/g in fresh plant tissue [155]. Ptaquiloside and quercetin are estimated in the plant material [108].

Several sesquiterpenes have been isolated from the fronds that include pterosins A-G, I-J, K-L, N, O, Z, etc. [156, 157, 158, 159]. Thiaminases 1 and 2 are isolated, which cause vitamin B1 deficiency in animals [160]. Steroids α - and β -ecdysones are identified in the bracken fern [161].

5.5. Dryopteridaceae

5.5.1. *Dryopteris cochleata*

5.5.1.1. Distribution. W, C and E Nepal; Guizhou, Sichuan and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; Bangladesh; Myanmar; Thailand; Indonesia; Philippines; etc. [31, 69].

5.5.1.2. Habitat. Moist and shady slopes, and roadsides.

5.5.1.3. Ethnomedicine. Decoction of rhizome, stem and stripe is given for blood purification [162]. Dried rhizome powder is internally taken to cure diarrhea [134]. Root decoction is used in cuts and throat problems. It is antibacterial, antiepileptic, and effective in rheumatism, amoebic dysentery and leprosy [163]. The juice of fronds is used to treat muscular and rheumatic pain [164].

5.5.1.4. Pharmacognosy. Antibacterial [165] and antioxidant [166] activities of the leaf extracts, and antimicrobial [167] activity of the rhizome extracts have been reported.

5.5.1.5. Phytochemistry. Leaves contain fat (3.6%), fiber (11.2%), protein (20.1%), vitamin C (23.1 mg/100 g) and minerals (Ca, Cu, Fe, Mg, Mn, Mo, Na, Zn, Al, As, Cd, Hg, Li, Ni and Pb) [27]. Preliminary phytochemical screenings and GC-MS analyses of the rhizomes and leaves extracts are reported [166, 167, 168]. The amounts of ptaquiloside and quercetin are estimated in the plant material [108].

5.5.2. *Dryopteris splendens*

5.5.2.1. Distribution. E. Nepal; Yunnan; Arunachal, Sikkim and Darjeeling; Bhutan; etc. [31, 69].

5.5.2.2. Habitat. Densely forested gully and slopes below trees, and evergreen and deciduous broad-leaved forests.

5.5.3. *Polystichum squarrosum* [*Polystichum apicisterile*]

5.5.3.1. Distribution. W, C and E Nepal; S Xizang; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; etc. [31, 69].

5.5.3.2. Habitat. Near streams and tree shades.

5.5.3.3. Pharmacognosy. Leaf extracts show antifungal [169] and antioxidant [165] activities. *P. squarrosum* fed rats show moderate mortality, splenomegaly and develop neoplastic lesions awareing its toxic effects [170].

5.5.3.4. Phytochemistry. The amounts of ptaquiloside and quercetin are estimated in the leaves [108].

5.6. *Lygodiaceae*

5.6.1. *Lygodium flexuosum*

5.6.1.1. Distribution. W, C and E Nepal; Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hunan and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand and Himachal; Bhutan; Sri Lanka; Thailand; Vietnam; Malaysia; Philippines; S Japan; E Australia; etc. [31, 69].

5.6.1.2. Habitat. Interior exposed area.

5.6.1.3. Ethnomedicine. The plant is used as an expectorant [171]. Root is smoked as cigarette [172]. Fresh root is boiled with mustard oil and then used for massage to treat rheumatism and sprains. It is also used for the treatment of jaundice, dysmenorrhea, wounds and eczema. Plant ash is used for treating gonorrhea, herpes, and other foot and mouth diseases. The whole plant is used in sprain, fracture, cuts and wounds [173].

5.6.1.4. Pharmacognosy. Antifertility [174, 175], antiproliferative, apoptotic, hepatoprotective [176, 177], antibacterial properties [178, 179] are reported.

5.6.1.5. Phytochemistry. It contains lygodinolide, coumaryl dryocassol, tectoquinone, kaempferol, kaempferol 3-β-D-glucoside, β-sitosterol and stigmasterol [180, 181, 182]. Several anhidiogens are identified from the spores [183].

5.6.2. *Lygodium japonicum*

5.6.2.1. Distribution. W, C and E Nepal; Anhui, Chongqing, Fujian, Gansu, Guangdong, Guangxi, Guizhou, Hainan, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Shaanxi, Shanghai, Sichuan, Xizang, Yunnan and Zhejiang; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu

and Kashmir; Pakistan; Bhutan; Sri Lanka; Indonesia (Java); Philippines; Taiwan; Korea; Japan; Australia, N America; etc. [31, 69].

5.6.2.2. Habitat. Moist and exposed area, and roadsides.

5.6.2.3. Ethnomedicine. It has been used in China for the treatment of calculi, edema, dysentery, hepatitis, pneumonia, gonorrhea, etc. [184]. Vegetative parts and spores decoction is used for diuretic or cathartic effects [171]. The plant is also used as an expectorant. The plant juice is applied in scabies, boils and wounds, and the paste is used in joint pains [51, 185]. The whole plant is applied to treat whitlow [52].

5.6.2.4. Pharmacognosy. The methanolic extract of the spores shows neuroprotective effect [186].

5.6.2.5. Phytochemistry. The amount of quercetin present in *L. japonicum* is estimated [108]. GC-MS analysis of the essential oil from the whole herb leads to identify 52 compounds including 3-methyl-1-pentanol (4%), 2-(methylacetyl)-3-carene (4.2%), cyclooctenone (7.6%), (E)-2-hexenoic acid (12.8%) and 1-undecyne (28.6%) [187].

Roots contain 1-hentriacontanol, hexadecanoic acid, 2,3-dihydroxypropyl ester, kaempferol, kaempferol-7-O- α -L-rhamnopyranoside, lygodiumsteroside B, p-coumaric acid, tiliatin, daucosterol and β -sitosterol [188, 189]. From the aerial parts, 2-O-caffeyl-D-glucopyranose, 3-O-caffeyl-D-glucopyranose, 4-O-caffeyl-D-glucopyranose, 4-O-pcoumaroyl-D-glucopyranose, 6-O-caffeyl-D-glucopyranose, 6-O-p-coumaroyl-D-glucopyranose, caffamic acid and p-coumaric acid are isolated [190]. Antheridiogens are identified in the spores [191, 192].

Interestingly, *L. japonicum* grown in the metal-enriched conditions can accumulate copper in the cell wall pectin [193].

5.7. *Marsileaceae*

5.7.1. *Marsilea quadrifolia*

5.7.1.1. Distribution. C and E Nepal; Gansu, Hebei, Heilongjiang, Henan, Jilin, Liaoning, Nei Mongol, Ningxia, Qinghai, Shaanxi, Shandong and Shanxi; Arunachal, Sikkim and Darjeeling; Bhutan; Korea; Japan; America; Europe; etc. [69, 85].

5.7.1.2. Habitat. Ponds and paddy fields.

5.7.1.3. Ethnomedicine. It produces aphrodisiac effect [124]. Leaf juice is anti-inflammatory, diuretic, depurative, febrifuge and refrigerant [194]. The plant is useful to relieve hypertension, sleep disorder, headache, cough, convulsion, respiratory troubles, nervous problems, constipation, dyspepsia, etc. [195, 196]. It is also used in the treatment of leprosy, haemorrhoids, fever and insomnia [197]. It is a famine food and only used during food scarcity.

5.7.1.4. Pharmacognosy. The plant extracts exhibit antibacterial [196, 198, 199, 200], antidiarrheal [194], antifungal [201], analgesic [194], anti-inflammatory [202], antioxidant [203, 204], antidiabetic [205], anti-Alzheimer's [206, 207], antiepileptic [208], anticonvulsive [209], nephroprotective [210] and antitumor [211] activities.

5.7.1.5. Phytochemistry. The plant material contains carbohydrate (19.5% g), protein (4.9%), fat (0.2%), amino acid (3%), flavonoids (0.3%) and saponins (0.3%) [212]. Seasonal variations in the compositions of sodium, potassium, calcium, phosphorus and β -carotene contents in the whole plant have been reported [213]. Total phenolics, flavonoids, alkaloids, tannins and saponins in both leaves and stems have been separately evaluated [214].

Leaves possess alkaloids, steroids, tannins, terpenoids, flavonoids, etc. [200, 202, 208, 214]. 2,3,7,8-Tetrachlorodibenzofuran,

3-hydroxy-triacontan-11-one, chlorogenic acid, hentriaccontan-6-ol, marsileagenin A, marsilin (1-triicontanol-cerotate), methylamine, pentachlorophenylacetate, quercetin, triacontyl hexacosanoate, tridecyl iodide, β -sitosterol, etc. are identified in *M. quadrifolia* [198, 202, 204].

Zhang *et al.* have isolated (\pm)-(E)-4b-methoxy-3b,5b-dihydroxy-scirpusin A, (\pm)-(E)-cyperusphenol A, (\pm)-(E)-scirpusin A, (E)-caffeic acid 4-O- β -D-glucopyranoside, (E)-ethyl caffeoate, 4-methy-3'-hydroxypsilotinin, hegovflavone A, kaempferol 3-O-(2''-O-E-caffeooyl)- β -D-glucopyranoside, kaempferol 3-O-(3''-O-E-caffeooyl)- α -L-arabinopyranoside, kaempferol 3-O-(4''-O-E-caffeooyl)- α -L-arabinopyranoside, kaempferol 3-O- α -L-arabinopyranoside, kaempferol 3-O- β -D-glucopyranoside, kaempferol, mesocyperusphenol A, quercetin, quercetin 3-O-(6''-O-E-caffeooyl)- β -D-glucopyranoside, quercetin 3-O- α -L-arabinopyranoside and quercetin 3-O- β -D-glucopyranoside from the ethanolic extract of *M. quadrifolia*, and their antioxidant capacity is evaluated [215].

5.8. Nephrolepidaceae

5.8.1. *Nephrolepis cordifolia* [*Nephrolepis auriculata*]

5.8.1.1. Distribution. W, C and E Nepal; Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hunan, Taiwan, Xizang, Yunnan and Zhejiang; Arunachal, Sikkim, Darjeeling and Uttarakhand; Pakistan; Bhutan; Bangladesh; Sri Lanka; Myanmar; Thailand; Laos; Vietnam; Cambodia; Malaysia; Singapore; Indonesia; Philippines; Korea; Japan; Australia; N and S America; Africa; etc. [31, 69].

5.8.1.2. Habitat. Moist place and roadsides.

5.8.1.3. Ethnomedicine. Tuberous root is used in indigestion, cold, cough, fever, jaundice and burning urination [52, [216]]. It is eaten to cure hypertension and inflammation [185]. Root paste is used in boils and skin problems [217]. Tuber juice is useful in dehydration [218]. Whole plant is used to cure renal, liver and skin disorders [219]. It is useful in chest congestion, rheumatism, jaundice and wounds [220]. Cough is cured by a decoction of the fresh fronds [221].

5.8.1.4. Pharmacognosy. Frond possesses antibacterial and antifungal potentials [99, 222]. Fruit and leaf are evaluated for the 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical scavenging, α -amylase inhibitory and glucose diffusion inhibition activities [223]. Anthelmintic activity of the aqueous extract of leaves is claimed as it paralyzes earthworms (*Pheretima posthuma*) at the concentration of 10 mg/mL [224]. Diuretic potential of the rhizome juice is shown [225].

5.8.1.5. Phytochemistry. Proximate analysis of different parts of *N. cordifolia* grown in Nepal and Nigeria has been reported [226, 227]. Tannins (11.5 mg/100 g), alkaloids (9.1 mg/100 g), flavonoids (16.5 mg/100 g), phenols (8.3 mg/100 g), saponin (1.2 mg/100 g) and glycosides (5.4 mg/100 g) are estimated in the leaflets [227]. Essential oil obtained from the aerial parts constitutes nonanal (10.6%), β -ionone (8%), eugenol (7.2%) and anethol (4.6%) [228]. Sequoyitol, β -sitosterol, fern-9(11)-ene, oleanolic acid, myristic acid octadecylester, hentriaccontanoic acid and triacontanol are isolated from the plant material [229].

5.9. Ophioglossaceae

5.9.1. *Botrychium lanuginosum*

5.9.1.1. Distribution. W, C and E Nepal; Guangxi, Guizhou, Hunan, Sichuan, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand and Himachal; Bhutan; Sri Lanka; Myanmar; Thailand; Vietnam; Malaysia; Indonesia; Philippines; Taiwan; New Guinea; etc. [31, 69].

5.9.1.2. Habitat. Moist and shady forest floor.

5.9.1.3. Ethnomedicine. It is used in dysentery and wounds [134, 163, 220, 230]. Rhizomes are used in pneumonia, catarrh and cough [124]. The paste is used as a facial mask, and it is applied in boils on the tongue and on the forehead to relieve headache [50]. Shoots are used in body ache [84].

5.9.1.4. Phytochemistry. Apigenin, luteolin, luteolin-7-O-glucoside, daucosterol, β -sitosterol, (6'-O-palmitoyl)-sitosterol-3-O- β -D-glucoside, thunberginol A, 30-nor-21 β -hopan-22-one, 1-O- β -D-glucopyranosyl-(2S,3R,4E,8Z)-2-[{(2R-hydroxy hexadecanoyl) amino]-4,8-octadecadiene-1,3-diol and sucrose are isolated from *B. lanuginosum* [231].

5.9.2. *Helminthostachys zeylanica*

5.9.2.1. Distribution. W, C and E Nepal; Guangdong, Hainan and Yunnan; Arunachal, Assam, Darjeeling, Bihar and Uttarakhand; Sri Lanka; Thailand; Laos; Vietnam; Cambodia; Philippines; Taiwan; Japan; Australia; Pacific islands; etc. [31, 69].

5.9.2.2. Habitat. Forest, edges of marshes and swampy places.

5.9.2.3. Ethnomedicine. Fronds are used as aphrodisiac [232]. The plant is considered as intoxicant and anodyne. It is useful in impotency, sciatica, dysentery, malaria and to relieve blisters on the tongue [124]. The rhizome is regarded as a tonic [233]. It is also used in dysentery, catarrh and phthisis. The rhizomes are popularly used as an antipyretic and anti-inflammatory Chinese herbal medicine, *Daodi-Ugon* [234, 235].

5.9.2.4. Pharmacognosy. Rhizomes show hepatoprotective [236, 237], anti-inflammatory [238, 239, 240], antihyperuricemia [241], cytotoxic [242], anticancer [243], antidiabetic [244] and antibacterial [245] activities.

5.9.2.5. Phytochemistry. The plant material contains fat (0.6%), crude fibre (1%), calcium (47.9 mg/100 g), phosphorus (91.5 mg/100 g), iron (1.8 mg/100 g), carotene (2.1 mg/100 g), riboflavin (0.1 mg/100 g), niacin (0.9 mg/100 g) and ascorbic acid (45.9 mg/100 g) [233].

From the rhizomes, flavonoids *viz* ugonins E-Y, quercetin, quercetin-4'-O- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside, quercetin-3-O- β -D-glucopyranosyl-4'-O- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside, 2-(3,4-dihydroxyphenyl)-6-((2,2-dimethyl-6-methylenecyclohexyl)methyl)-5,7-dihydroxy-chroman-4-one, 2-(3,4-dihydroxy-2-[(2,6,6-trimethylcyclohex-2-enyl)methyl]phenyl)-3,5,7-trihydroxy-4-H-chromen-4-one and 5,4'-dihydroxy-4'',4''-dimethyl-5''-methyl-5''H-dihydrofuran-2'',3''-6,7]flavanone are isolated by Huang *et al.* [235, 246, 247, 248]. Quality of the root and rhizome of *H. zeylanica* can be determined by using ugonins as markers [249]. Geranyl stilbenes ugonstilbenes A-C and 3-hydroxyacetophenone are isolated from the rhizomes by Chen *et al.* [234]. Ugonstilbenes A-C exhibit moderate antioxidant activity. 4'-O- β -D-Glucopyranosyl-quercetin-3-O- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranoside and 4'-O- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl-quercetin-3-O- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranoside are isolated by Yamauchi *et al.* [250]. These glycosides display melanogenic acceleratory effect.

From the whole plant, neougonins A-B, ugonins D-E, ugonin J, ugonin L, ugonin N, ugonin S, 2-(3,4-dihydroxyphenyl)-6-((2,2-dimethyl-6-methylenecyclohexyl)methyl)-5,7-dihydroxy-chroman-4-one, etc. are isolated [251].

5.9.3. *Ophioglossum nudicaule*

5.9.3.1. Distribution. Nepal; SW Sichuan, Xizang, C and NW Yunnan; Darjiling, Sikkim, Himanchal, Jharkand, Karnataka, Kerala, Tamilnadu, Jammu and Kashmir; Indonesia; Malaysia; Thailand; etc. [69, 252].

5.9.3.2. Habitat. Rainforest, impermeable soil in open places and by culverts on roadsides in the hills.

5.9.3.3. Ethnomedicine. Fronds are used as a tonic and styptic [253]. It is used in wounds and old skin diseases [52, 254].

5.9.4. *Ophioglossum petiolatum*

5.9.4.1. Distribution. C and E Nepal; Fujian, Guizhou, Hainan, Hubei, Sichuan, Yunnan and Xizang; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Uzbekistan; Bhutan; Sri Lanka; Thailand; Indonesia; Philippines; Taiwan; Japan; Australia; New Zealand; N America; England; Fiji; etc. [31, 69, 252].

5.9.4.2. Habitat. Moist exposed grassy areas.

5.9.4.3. Ethnomedicine. It is used for checking of nose bleeding [84]. It has been used as a detoxifying agent and found effective against hepatitis [255, 256].

5.9.4.4. Phytochemistry. Ophioglolin, ophioglolin 7-O- β -D-glucopyranoside, ophioglolin, ophioglolin prenyl ether, ophioglolin 4'-O- β -D-glucopyranoside, isophioglolin 7-O- β -D-glucopyranoside, quercetin, luteolin, kaempferol, 3,5,7,3',4'-pentahydroxy-8-prenylflavone, and quercetin 3-O-methyl ether are isolated from the whole plant [256].

5.9.5. *Ophioglossum reticulatum*

5.9.5.1. Distribution. W, C and E Nepal; Fujian, Gansu, Guizhou, Henan, Hubei, Jiangxi, Shaanxi, Sichuan, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; Sri Lanka; Philippines; Korea; S America; Madagascar; Africa; etc. [31, 69, 252].

5.9.5.2. Habitat. Roadsides, under beneath of tree in the forests and grassy plateaus.

5.9.5.3. Ethnomedicine. It is used as cooling agent in burns, tonic and wounds remedy [230]. Fronds are used as styptic in contusions and haemorrhages [163]. Leaf extract is vulnerary [124]. Leaf juice is useful against heart spasms [257]. Rhizome decoction is used in boils and the extract is used as an antidote in snake bite.

5.9.6. *Ophioglossum vulgatum*

5.9.6.1. Distribution. Nepal; Anhui, Fujian, Guangdong, Guizhou, Henan, Hubei, Hunan, Jiangxi, S Shaanxi, Sichuan, Xizang, Yunnan and Zhejiang; India; Sri Lanka; Korea; Japan; Australia, N America; Europe; etc. [69].

5.9.6.2. Habitat. Naturally grow in damp grasslands and semi-evergreen forests, particularly after fire [258].

5.9.6.3. Ethnomedicine. It possesses detergent, antiseptic, styptic and vulnerary properties [259]. Rhizome is used to treat boils. Pulverized rhizome is externally applied to sores, burns and wounds [135]. Decoction is useful in heart diseases [134]. It is useful in clearing heat, detoxifying toxins, activating blood circulation, treating hepatitis and pneumonia, etc. [260].

5.9.6.4. Pharmacognosy. *O. vulgatum* exhibits wound healing [261, 262], antidiarrheal [263] and antiviral [264] properties.

5.9.6.5. Phytochemistry. The whole plant material constitutes total flavonoid and total polyphenols in 2.7 and 1.5%, respectively [263].

Flavonoids, glycerides and amino acids are the constituents in *O. vulgatum* [265]. From the aerial parts, 3-O-methylquercetin 7-O-diglycoside 4'-O-glycoside, quercetin-3-O-[(6-caffeoyl)- β -glucopyranosyl(1 \rightarrow 3) α -rhamnopyranoside]-7-O- α -rhamnopyranoside, kaempferol-3-O-[(6-caffeoyl)- β -glucopyranosyl (1 \rightarrow 3) α -rhamnopyranoside]-7-O- α -rhamnopyranoside and quercetin-3-O-methyl ether have been isolated [261, 266].

5.10. *Osmundaceae*

5.10.1. *Osmunda claytoniana* [*Osmunda claytoniana* subsp. *Vestita*]

5.10.1.1. Distribution. W, C and E Nepal; Chongqing, Guizhou, Hubei, Hunan, Liaoning, Sichuan, Taiwan, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Pakistan; Bhutan; Korea; Japan; E Russia; N America; etc. [31, 69].

5.10.1.2. Habitat. At high altitude above in meadows, moist and shady forest area.

5.10.1.3. Ethnomedicine. Plant paste is applied on wounds externally [134]. The rhizome and stipe bases are used for adulteration as a substitute of *Filix-mas* for vermifuge effect and flavor [267].

5.10.1.4. Pharmacognosy. α -Glucosidase inhibitory activity [268] and antitrypanosomal activity [269] of the plant extracts are evaluated.

5.10.1.5. Phytochemistry. The plant material consists of phenolics, tannins, flavonoids, saponins, terpenoids, etc. [91].

5.11. *Pteridaceae*

5.11.1. *Ceratopteris thalictroides*

5.11.1.1. Distribution. W and C Nepal; Anhui, Fujian, Guangdong, Guangxi, Guizhou (Liping), Hainan, Hubei, Jiangsu, Jiangxi, Shandong, Sichuan, Yunnan and Zhejiang; Assam, W Bengal, Bihar, Uttarakhand, Himachal and C India; Sri Lanka; Myanmar; Thailand; Vietnam; Malaysia; Indonesia; Philippines; Taiwan; Japan; New Guinea; Australia; C, N and S America; Madagascar; Pacific islands; West Indies; Africa; etc. [31, 69].

5.11.1.2. Habitat. Marshy areas.

5.11.1.3. Ethnomedicine. The fronds are used as a poultice for skin problems [163, 270, 271]. Its paste with turmeric is applied in cuts and wounds. It is also used as a tonic and styptic [272], and it is also reported to be toxic [163].

5.11.1.4. Pharmacognosy. The plant extracts possess antibacterial [273], anti-inflammatory [274], antioxidative [275] and antidiabetic activities [276].

5.11.1.5. Phytochemistry. The leaves contain protein (13.4%), carbohydrate (54.3%), fat (3.2%), vitamin C (6.1 mg/100 g) and minerals (eg. Na, K, P, Mg, Ca, Mn, Fe, Zn and Cu) [277]. The plant materials showed the presence of alkaloids, steroids, coumarins, tannins, saponins, flavonoids, quinones, anthroquinones, phenols, proteins, xanthoproteins, carbohydrates, glycosides, catachins, sugars and terpenoids [270, 277]. 2-Nitroethyl benzene, γ -sitosterol, n-hexadecanoic acid and linalool are the major phytochemicals present in the leaves.

5.11.2. *Pteris biaurita* [*Pteris flavigaulis*]

5.11.2.1. Distribution. W, C and E Nepal; Guangdong, Guangxi (Baise), Guizhou, Hainan, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling and

Table 2. Summary of the medicinal ailments of the vegetable food ferns.

SN	Medicinal ailments	Ferns	Parts used	References
1	Antibacterial	<i>D. esculentum</i> <i>S. palustris</i> <i>W. unigemmata</i> <i>D. cochleata</i> <i>L. flexuosum</i> <i>M. quadrifolia</i> <i>N. cordifolia</i> <i>H. zeylanica</i> <i>O. vulgatum</i> <i>C. thalictroides</i> <i>P. vittata</i> <i>T. gemmifera</i>	Rhizome, leaf Frond Aerial part Leaf, rhizome Frond, whole plant Stem, leaf, aerial part Frond Rhizome Frond, leaf and stem Leaf Frond Rhizome, leaf, frond	[96, 98, 99, 100] [128] [138] [165, 167] [178, 179] [196, 198, 199, 200] [99, 222] [245] [261, 262] [273] [285, 286] [293, 294, 295, 296]
2	Antiviral	<i>W. unigemmata</i> <i>O. vulgatum</i>	Rhizome Not specified	[139, 140] [264]
3	Antifungal	<i>S. palustris</i> <i>P. squarrosum</i> <i>M. quadrifolia</i> <i>N. cordifolia</i> <i>P. biaurita</i>	Frond Leaf Whole part Frond Leaf	[127] [169] [201] [222] [281]
4	Antimalarial	<i>S. palustris</i>	Leaf	[132]
5	Antidiarrheal	<i>M. quadrifolia</i> <i>O. vulgatum</i>	Aerial part Whole part	[194] [263]
6	Anthelmintic	<i>D. esculentum</i> <i>N. cordifolia</i>	Rhizome Leaf	[107] [224]
7	Analgesic and anti-inflammatory	<i>D. esculentum</i> <i>M. quadrifolia</i> <i>H. zeylanica</i> <i>C. thalictroides</i>	Leaf Aerial part, leaf Rhizome Spore	[99, 106] [194, 202] [238, 239, 240] [274]
8	Antidiabetic	<i>D. esculentum</i> <i>M. quadrifolia</i> <i>H. zeylanica</i> <i>O. claytoniana</i> <i>C. thalictroides</i> <i>T. gemmifera</i>	Leaf Aerial part Rhizome Frond, rhizome Whole part Frond, rhizome	[103] [205] [244] [268] [276] [299, 301]
9	Antioxidant and anticancer	<i>D. boryana</i> <i>D. esculentum</i> <i>D. maximum</i> <i>S. palustris</i> <i>W. unigemmata</i> <i>C. spinulosa</i> <i>D. cochleata</i> <i>P. squarrosum</i> <i>M. quadrifolia</i> <i>N. cordifolia</i> <i>H. zeylanica</i> <i>C. thalictroides</i> <i>P. biaurita</i> <i>T. gemmifera</i> <i>T. zeylanica</i>	Aerial part Leaf Frond Frond Aerial part Leaf Leaf Leaf Aerial part, leaf Fruit, leaf Rhizome Frond Frond Rhizome, leaf, twig Leaf	[92, 94] [101, 102, 103, 104, 105] [121] [129, 130, 131] [137, 138] [144] [166] [165] [203, 204, 211] [223] [242, 243] [275] [278] [295, 297, 298] [303]
10	Neuroprotective	<i>D. esculentum</i> <i>C. spinulosa</i> <i>L. japonicum</i> <i>M. quadrifolia</i>	Leaf Leaf Spore Whole plant, leaf	[104] [144] [186] [206, 207, 208, 209]
11	Nephroprotective	<i>M. quadrifolia</i> <i>N. cordifolia</i>	Leaf Rhizome	[210] [225]
12	Hepatoprotective	<i>D. esculentum</i> <i>L. flexuosum</i> <i>H. zeylanica</i>	Leaf Whole plant Rhizome	[103] [176, 177] [236, 237]
13	Antifertility	<i>L. flexuosum</i>	Not specified	[174, 175]

Uttarakhand; Bhutan; Bangladesh; Sri Lanka; Thailand; Laos; Malaysia; Indonesia; Philippines; C and S Taiwan; etc. [31, 69].

5.11.2.2. Habitat. Moist, shady and exposed areas, and roadsides.

5.11.2.3. Ethnomedicine. The rhizome and frond decoction is used to treat chronic disorders [278]. To relieve body pain, the rhizome paste is applied [279]. Frond juice and paste are applied on cuts and bruises [53, 280].

5.11.2.4. Pharmacognosy. The leaves exhibit antimicrobial activity [281]. The fronds are evaluated for the antioxidant activity [278].

5.11.2.5. Phytochemistry. Fronds constitute alkaloids (16.4 mg/g), phenolics (13.2 mg/g), saponins (11.1 mg/g), tannins (6.2 mg/g), flavonoids (17.5 mg/g) as well as steroids, triterpenoids, anthraquinones, amino acids and sugars [282]. Leaves contain flavonoids (14.5 mg/g), total phenol (10.4 mg/g), ascorbate (1.6 mg/g), α -tocopherol (0.9 mg/g) and total sugars (125 mg/g) [278]. Total protein (110.5 mg/g), total chlorophyll (0.9 mg/g) and carotenoids (62.3 μ g/g) in the fresh material are also estimated. Eicosenes and heptadecanes present in the leaves are attributed for the antimicrobial activity [281].

5.11.3. *Pteris vittata*

5.11.3.1. Distribution. W, C and E Nepal; Anhui, Fujian, SE Gansu (Kangxian), Guangdong, Guangxi, Guizhou, SW Henan, Hubei, Hunan, Jiangxi, Shaanxi, Sichuan, Xizang, Yunnan and Zhejiang; Arunachal, Sikkim, Darjeeling and Uttarakhand; Bhutan; Taiwan; etc. [31, 69]. It is commonly known as brake fern.

5.11.3.2. Habitat. Moist and shady areas, roadsides and tropical semi-evergreen forests.

5.11.3.3. Ethnomedicine. It is a poisonous plant and therefore it is planted on village borders to keep enemies away [283]. It is used in tongue sores, burns and glandular swellings and for population control [136, 220, 279]. Herb juice is used for treating dysentery and diarrhea [284]. Young fronds are used in manuring [134]. Plants are used as fodder and thatching [52].

5.11.3.4. Pharmacognosy. Its antibacterial property is explored [285, 286]. It has been reported that *P. vittata* hyperaccumulates arsenic [287, 288].

5.11.3.5. Phytochemistry. Leaves contain leucocyanidin, leucodelphinidin, apigenin 7-O-p-hydroxybenzoate, leutolin, kaempferol, quercetin, rutin, etc. [285, 289, 290].




Figure 2. Some flavonoids isolated from *S. palustris*.

5.12. *Tectariaceae*

5.12.1. *Tectaria gemmifera* [*Tectaria coadunata*; *Tectaria macrodonta*]

5.12.1.1. Distribution. W, C and E Nepal; Guangdong, Guangxi, Guizhou, Sichuan, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Bhutan; Sri Lanka; Myanmar; Thailand; Laos; Vietnam; Malaysia; Taiwan; Madagascar; Africa; etc. [31, 69].

5.12.1.2. Habitat. North-west facing moist and rocky slopes, exposed areas and roadsides or in rock crevices near water channels.

5.12.1.3. Ethnomedicine. Rhizome decoction is given to treat stomachache and gastrointestinal disorders [291]. Root juice is used to treat diarrhea and dysentery [142]. Leaf decoction is given to treat asthma and bronchitis [253]. Leaf paste is applied in stings of honeybee, centipedes, etc. Whole plant is used in eczema, scabies and jaundice [254, 292]. Leaves are used as fodder [134].

5.12.1.4. Pharmacognosy. Antibacterial [293, 294, 295, 296], antioxidant [295, 297, 298] and antidiabetic [299] activities are reported. Cholinesterases and tyrosinase inhibitory activities together with cytotoxic effect on the 2008 and BxPC3 cell lines of the plant extract from twigs and rhizomes are reported [297].

5.12.1.5. Phytochemistry. Crude fat (0.2%), fiber (2.3%), protein (12.1%), vitamin C (22.2 mg/100 g), mineral contents (Ca, Cu, Fe, Mg, Mn, Na and Zn), heavy metals (Al, As, Cd, Hg, Li, Ni and Pb), etc. are determined in *T. gemmifera* [27]. The fronds constituted of anthraquinones, coumarins, flavonoids, steroids, alkaloids, tannins, phenolics, etc. [296, 300], and the rhizomes constitutes alkaloids, coumarins, phenolics, flavonoids, quinones, etc. [294, 296, 301]. GC-MS analysis of the methanolic extract of leaves show the presence of palmitic acid, methyl palmitate, phytol, etc. as the major constituents [295], while the rhizome constitutes fatty acids [295, 302]. The qualitative and quantitative analyses reveal that the twig and rhizome contain a large amount of procyanidins as well as eriodictyol-7-O-glucuronide and luteolin-7-O-glucuronide [297].

5.12.2. *Tectaria zeylanica* [*Osmunda zeylanica*]

5.12.2.1. Distribution. W, C and E Nepal; Fujian, Guangdong, Guangxi, Guizhou, Hainan and Yunnan; S India; Sri Lanka; Thailand; Laos; Vietnam; Malaysia; Indonesia; Philippines; Taiwan; Indian Ocean islands (Mauritius); Pacific islands (Polynesia); etc. [69].

5.12.2.2. Habitat. Muddy rocks in forests, near streams and on steep banks.




Figure 3. Ugonins isolated from *H. zeylanica* rhizomes.

5.12.2.3. Pharmacognosy. Ethanolic extract of the leaves shows antioxidant activity by scavenging DPPH free radical with the IC₅₀ value of 0.78 mg/mL [303].

5.12.2.4. Phytochemistry. Leaf constitutes alkaloids, terpenoids, flavonoids, tannins, etc. [303]. Fronds constitute phenolics, saponins, steroids, tannins, coumarins, etc. [304].

5.13. *Thelypteridaceae*

5.13.1. *Thelypteris auriculata* [*Cyclosorus auriculata*]

5.13.1.1. Distribution. W, C and E Nepal; Guangxi, Guizhou, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand and Himachal; Bhutan; Sri Lanka; Thailand; Vietnam; etc. [31, 69].




Figure 4. Neougonin A isolated from *H. zeylanica*.

5.13.1.2. Habitat. Along river banks and forest margins.

5.13.2. *Thelypteris nudata* [*Thelypteris multilineata*; *Pronephrium nudatum*]

5.13.2.1. Distribution. W, C and E Nepal; Guizhou, Xizang and Yunnan; Arunachal, Sikkim, Darjeeling, Uttarakhand, Himachal, Jammu and Kashmir; Bhutan; Indonesia; Vietnam; etc. [31, 69].

5.13.2.2. Habitat. Shaded sparse forests on slopes.

5.13.2.3. Pharmacognosy. The frond extract exhibits synergistic antibacterial effect against methicillin resistant *Staphylococcus aureus* (MRSA) [305].

6. Altitudinal zones-based categorization of Nepalese wild edible ferns

The distribution of ferns is varied with different altitudinal zones. Nepal is particularly considered to be divisible into three altitudinal zones, namely, (a) Tarai (plain area), (b) Pahad (hilly area) and (c) Himal (mountainous area). Accordingly, based on the species found at different altitudes, the Nepalese wild edible vegetable ferns can be categorized into three:

6.1. Lower altitude species (60–1,200 m asl)

The species grown in low-lands, i.e. Tarai, Siwalik and Chure areas of Nepal, include: *A. strigillossum*, *D. boryana*, *D. esculentum*, *S. palustris*, *W. unigemmata*, *C. spinulosa*, *P. revolutum*, *D. maximum*, *D. splendens*, *D. cochleata*, *L. flexuosum*, *L. japonicum*, *N. cordifolia*, *B. lanuginosum*, *H. zeylanica*, *O. petiolatum*, *O. reticulatum*, *C. thalictroides*, *P. biaurita*, *P. vittata*, *T. gemmifera* and *T. zeylanica*.

6.2. Mid-hills species (1,200–2,500 m asl)

The species grown in mid-lands, i.e. Chure and Mahabharat areas of Nepal, include: *A. atkinsonii*, *A. strigillossum*, *D. boryana*, *D. esculentum*, *D. kawakamii*, *D. spectabile*, *D. stoliczkae*, *W. unigemmata*, *C. spirulosa*, *P. revolutum*, *D. maximum*, *D. cochleata*, *P. squarrosum*, *M. quadrifolia*, *L. japonicum*, *N. cordifolia*, *B. lanuginosum*, *B. lanuginosum*, *H. zeylanica*, *O. nudicaule*, *O. petiolatum*, *O. reticulatum*, *O. vulgatum*, *O. claytoniana*, *P. biaurita*, *P. vittata*, *T. gemmifera*, *T. auriculata* and *T. multilineata*.





Figure 5. Some bioactive terpenoids obtained from food ferns.

Table 3. Cytotoxicity (HeLa Cell) of pterosins and pterosides isolated from *Pteridium aquilinum* var. *latiusculum*.



Compound	R ¹	R ²	R ³	R ⁴	Cytotoxicity IC ₅₀ (µg/mL)
Pterosin A	—CH ₂ OH	—CH ₃	H	—CH ₂ OH	320
Pterosin B	—CH ₃	H	H	—CH ₂ OH	100
Pterosin C	H	—CH ₃	—OH	—CH ₂ OH	320
Pterosin D	—CH ₃	—CH ₃	—OH	—CH ₂ OH	>320
Pterosin E	—CH ₃	H	H	—CO ₂ H	30
Pterosin F	—CH ₃	H	H	—CH ₂ Cl	65
Pterosin G	—CH ₂ OH	H	H	—CH ₂ OH	180
Pterosin J	H	—CH ₃	—OH	—CH ₂ Cl	>100
Pterosin K	—CH ₂ OH	—CH ₃	H	—CH ₂ Cl	>100
Pterosin L	—CH ₂ OH	—CH ₃	—OH	—CH ₂ OH	>100
Pterosin N	—CH ₃	—OH	H	—CH ₂ OH	180
Pterosin O	—CH ₃	H	H	—CH ₂ OCH ₃	30
Pterosin Z	—CH ₃	—CH ₃	H	—CH ₂ OH	10
Pteroside A	—CH ₂ OH	—CH ₃	H	—CH ₂ O glucose	>320
Pteroside B	—CH ₃	H	H	—CH ₂ O glucose	>100
Pteroside C	H	—CH ₃	—OH	—CH ₂ O glucose	>320

6.3. High altitude species (2,500–4,000 m asl)


The species grown in upper-lands, i.e. Himalayas or mountainous areas of Nepal, include: *D. boryana*, *D. spectabile*, *D. stoliczkae*, *P. squarrosum*, *L. japonicum*, *W. unigemmata*, *P. revolutum*, *B. lanuginosum*, *O. nudicaule*, *O. petiolatum*, *O. reticulatum*, *O. vulgatum* and *O. claytoniana*.

7. Inhabitutions-based distribution of Nepalese wild edible ferns

Virtually, the ferns are inhabited where flowering plants are found [32]. Humid, moist and shady forests are more suitable for their growth. Based on the species found at different ecological habitats, the edible vegetable ferns of Nepal are categorized into epiphytes, lithophytes, terrestrials, climbers and hydrophytes as given below. Noteworthy, some species could be found in more than one habitat.

7.1. Epiphytes

These species grow on bases, trunks and branches of trees, and are covered with mosses and liverworts. Rainy season, hill-side evergreen forest and moist forest are more suitable for their growth. In middle elevation, altitude up to 2,000 m, the distribution of epiphytes is very few. Examples of epiphytes are: *N. cordifolia*, *B. lanuginosum* and *P. vittata*.






Figure 6. Steroids found in food ferns.

7.2. Lithophytes

These species grow in rock crevices; humus rich rocks in shady areas; moist, shaded and exposed forests; crevices of brick walls; and muddy rocks in streamlets. Examples of lithophytes are: *W. unigemmata*, *D. cochleata*, *N. cordifolia*, *B. lanuginosum*, *P. biaurita*, *P. vittata* and *T. gemmifera*.

7.3. Terrestrials

They grow in evergreen and semi-evergreen forests that enriched with humus and organic nutrients. They also grow in shady areas, stream banks, moist hill slopes and shaded roadsides. Examples of terrestrial species are: *A. strigillosum*, *D. boryana*, *D. esculentum*, *D. maximum*, *D. stoliczkae*, *W. unigemmata*, *C. spinulosa*, *P. revolutum*, *D. cochleata*, *D. splendens*, *P. squarrosum*, *N. cordifolia*, *B. lanuginosum*, *H. zeylanica*, *O. nudicaule*, *O. petiolatum*, *O. reticulatum*, *O. claytoniana*, *P. biaurita*, *P. vittata*, *T. gemmifera*, *T. zeylanica* and *T. nudata*.

7.4. Climbers

They grow on rich humus soil. They climb other trees growing inside shaded forests, forest edges and tropical areas. Examples of climber species include: *S. palustris*, *L. flexuosum* and *L. japonicum*.

7.5. Hydrophytes

They are water loving species. They occur on wet or marshy border of ponds, lakes and waterfalls. They also occur in paddy fields. Examples of hydrophytes include: *M. quadrifolia* and *C. thalictroides*.

8. Medicinal ailments categorization of Nepalese wild edible ferns

Based on pharmacognosy of the plant extracts of ferns and fern-allies reported by several authors, the medicinal ailments categorization of Nepalese wild edible vegetable food ferns is summarized in Table 2.

9. Important phytochemicals present in the wild edible ferns

The major bioactive compounds present in the ferns and fern-allies are flavonoids, terpenoids, steroids and alkaloids.

9.1. Flavonoids

The total flavonoid content has been reported in the edible parts of some ferns, for examples, 145.8 mg/g in *D. boryana* (arial parts) [94], 110.8 mg quercetin equivalent/100 g in *D. esculentum* (arial parts) [109], 151 mg quercetin equivalent/g in *W. unigemmata* (arial parts) [138], 3.5 mg/g in *M. quadrifolia* (arial parts) [212], 7.5 mg/g in *M. quadrifolia* (leaves) [214], 6.4 mg/g *M. quadrifolia* (stems) [214], 27.4 mg/g in *O. vulgatum* (whole plant) [263], 17.5 mg/g in *P. biaurita* (fronds) [282], 14.5 mg/g in *P. biaurita* (leaves) [278], etc. A higher total flavonoid content in the plant material is usually correlated with a more antioxidant capacity and thus potentially anticarcinogenic [306, 307].

Several flavonoid glycosides have been reported from the leaves [128] and fronds of *S. palustris* [126] (Figure 2). The isolated flavonoids viz. kaempferol 3-O-(3''-O-E-p-coumaroyl)-(6''-O-E-feruloyl)-β-glucopyranoside and kaempferol 3-O-(3'',6''-di-O-E-p-coumaroyl)-β-glucopyranoside moderately inhibit butyrylcholinesterase (BChE) with IC₅₀ values of 113.66 and 85.37 μM, respectively [126]; and kaempferol 3-O-(6''-O-E-p-coumaroyl)-β-glucopyranoside strongly inhibits acetylcholinesterase (AChE) with IC₅₀ value of 23.5 μM [308] exhibiting potentiality against Alzheimer's disease.

Huang *et al.* reported isolation of several flavonoids (ugonins E-Y) from the rhizomes of *H. zeylanica* [235, 246, 247, 248] (Figure 3). Ugonins J-L are found more active than trolox, with IC₂₀ values of 5.29 ± 0.32, 7.23 ± 0.22 and 7.93 ± 0.31 μM, respectively, in antioxidative activity using the DPPH assay; ugonins L,M,O,Q,S and T are found anti-inflammatory; and ugonin K is found antiosteoporosis. Ugonins J,K, L,M,S and U possess antidiabetic activity by inhibiting α-glucosidase [309, 310]. Ugonin J is found effective in inhibition of neointima formation [311]. Being a novel viral protease inhibitor, ugonin J inhibits severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) 3CLpro

activity and hence is therapeutically potential for SARS-CoV-2 infection [312]. Ugonin K induces apoptosis [313]. Neuroprotective [314] and osteogenesis [315] effects of ugonin K are reported. Ugonin M is considerably potential to develop hepatoprotective and anti-inflammatory natural agents [316]. Ugonin U shows immunomodulatory effect in human neutrophils via stimulation of phospholipase C [317], and induces Ca^{2+} mobilization and eventually activates the NLRP3 inflammasome [318].

From the whole plant of *H. zeylanica*, Su *et al.* have isolated several prenylated flavonoids [251]. Among them, neougonin A (Figure 4) inhibited NO production in lipopolysaccharide-treated RAW264.7 cells with an IC_{50} value of 3.32 μM , and it displayed a moderate cytotoxicity with IC_{50} value of 16.13 μM in the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay.

9.2. Terpenoids

Ptaquiloside (Figure 5) is one of the main hazardous metabolites found in the fern that is considered to be responsible for the toxicological problems to ruminant and non-ruminant animals, and then to human alike through the milk and meat [150]. The amounts of ptaquiloside have been estimated in *P. revolutum*, *D. cochleata* and *P. squarrosum* (leaves) that collected from different parts of India [108]. *P. revolutum* is considered as carcinogenic [152]. It has been demonstrated that feeding of bracken fern induces urinary bladder tumors, ileum sarcomas, pulmonary adenomas and leukemia [153]. On the other hand, Yoshihira *et al.* have isolated several sesquiterpenes, namely, pterosins A–G, J–L, N, O, Z, pterosides A–C, etc., from the fronds of Japanese bracken fern and performed cytotoxicity test (Table 3) [154]. They have mentioned that although these indanone derivatives showed some cytotoxicity to HeLa cells; however, none of them could induce tumors under the conditions studied.

Lygodinolide (Figure 5), present as major constituent in *L. flexuosum*, is attributed for the wound healing property [181,182].

From *N. cordifolia*, oleanolic acid and fern-9(11)-ene have been isolated (Figure 5) [229]. Fern-9(11)-ene has also been isolated from *C. spinulosa* leaves [146]. Oleanolic acid shows antimicrobial activity against vancomycin resistant *enterococci*, *Streptococcus pneumoniae* and MRSA [319]. It is also found to exhibit anticancer activity [320]. Fern-9(11)-ene is found to be susceptible against *Salmonella typhi* and *Pseudomonas aeruginosa* [321].

9.3. Steroids

Stigmast-3,6-dione, β -sitosterol and daucosterol in the leaves and stalks of *C. spinulosa* [147, 148, 149]; α - and β -ecdysones in the aerial parts of *P. revolutum* [161]; and daucosterol, β -sitosterol, (6'-O-palmitoyl)-stigmasterol-3-O- β -D-glucoside in *B. lanuginosum* [231] are some examples of alkaloids present in different edible parts of the ferns and fern-allies (Figure 6).

9.4. Alkaloids

Total alkaloids of 6.1 mg/g in *M. quadrifolia* (leaves) [214], 5.9 mg/g *M. quadrifolia* (stems) [214], 0.1 mg/g in *N. cordifolia* (leaves) [227] and 16.4 mg/g in *P. biaurita* (fronds) [282] are reported.

10. Conclusion

Taxonomic and ethnobotanical studies on the ferns of Nepal are exploited; however, chemical and biological investigations on Nepalese ferns and fern-allies are limited. This review primarily briefs historical background on expeditious journey on Nepalese pteridophytes, updates number of ferns consumed as vegetables by Nepalese, outlines their ethnomedicinal uses, and compiles information on their pharmacognosy,

pharmacology and phytochemistry that reported from elsewhere. In Nepal, some species of ferns that primarily collected from nearby forests are sold in the markets. Some species are consumed by rural Nepalese as the last option of food and there is no proper information transformation to the next generation about ferns as food materials and their potential medicinal values. *C. decurrenti-alata*, *M. intermedia*, *B. orientale*, *O. japonica*, *C. intermedia* and *P. wallichiana* are available in Nepal, but there is no precedented report for their consumption by Nepalese; however, they are consumed in China. The traditional ethnopharmacological knowledge on ferns and fern-allies is neglected both by locals and scientific community.

11. Future recommendation

Most of the ferns and fern-allies, collected for vegetable food by Nepalese, grow in their natural habitat. Lack of sharing of traditional ethnomedicinal knowledge to new generation and unsustainable collection practice making them vulnerable. Only in the recent past, a very few species bearing economical value are agricultured in local farms. From the literature evidences, we believe that investigations on unfocused ferns can provide new research opportunities and potent bioactive phytochemicals.

Parts of some species such as *P. revolutum*, *P. squarrosum*, *C. thalictroides*, *P. vittata*, etc. are considered to be toxic, thus they warrant immediate further investigations. *P. revolutum* constitutes cyanogen glycosides. *P. squarrosum* constitutes prunasin that can release HCN. *P. vittata* hyperaccumulates arsenic. *P. revolutum* and *P. squarrosum* can induce carcinogenesis. On the other hand, the antioxidant capacity of the extracts of *D. boryana*, *D. esculentum*, *D. maximum*, *S. palustris*, *W. unigemmata*, *D. cochleata*, *P. squarrosum*, *M. quadrifolia*, *N. cordifolia*, *P. biaurita*, *T. gemmifera*, *T. zeylanica*, etc. has been reported indicating their role in cancer chemoprevention. Further investigations of these plant materials to isolate and identify effective anticancer agents are highly desirable.

Rhizomes of *C. spinulosa* and *O. reticulatum* have found their traditional uses against snake bites. Ugonins, particularly isolated from the rhizomes of *H. zeylanica*, have found diverse bioactivities. Recently, ugonin J has reported to be effective against coronavirus disease 2019 (COVID-19). Hence, not only the vegetative parts of the ferns but also other parts should be investigated in order to search for potential drug candidates.

Declarations

Author contribution statement

All authors listed have significantly contributed to the development and the writing of this article.

Funding statement

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Data availability statement

Data included in article/supp. material/referenced in article.

Declaration of interest's statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

Acknowledgements

We thank Nepal Academy of Science and Technology (NAST), Khamaltar, Lalitpur, Nepal for providing a platform to write this article. Tribhuvan University Central Library, Kirtipur, Kathmandu, Nepal is acknowledged for providing literature access.

References

- [1] GoN/MoFSC, Nepal National Biodiversity Strategy and Action Plan: 2014–2020, Ministry of Forests and Soil Conservation, Government of Nepal, Kathmandu, 2014. http://mofc.gov.np/downloadfile/29_Strategy%20and%20action%20plan_1526382225.pdf. (Accessed 11 March 2021).
- [2] P.K. Paudel, B.P. Bhattacharai, P. Kindlmann, An overview of the biodiversity in Nepal, in: P. Kindlmann (Ed.), *Himalayan Biodiversity in the Changing World*, Springer, Dordrecht, 2012.
- [3] S.L. Shyaula, G.B. Bajracharya, S.M. Shakya, D. Subba, Comprehensive Insights in Vegetables of Nepal, Nepal Academy of Science and Technology, Khamaltar, Lalitpur, Nepal, 2021. <https://nast.gov.np/documentfile/Comprehensive-Insights-in-Vegetables-Book.pdf>.
- [4] Background Information about Nepal. <http://www.florafornepal.org/countryinformation>. (Accessed 18 March 2021).
- [5] G.B. Bajracharya, B. Bajracharya, Diversity, availability, using status, nutritional and pharmacological aspects of wild vegetables of Nepal, in: S.L. Shyaula, G.B. Bajracharya, S.M. Shakya, D. Subba (Eds.), Comprehensive Insights in Vegetables of Nepal, Nepal Academy of Science and Technology, Khamaltar, Lalitpur, Nepal, 2021, pp. 157–200. <https://nast.gov.np/documentfile/Comprehensive-Insights-in-Vegetables-Book.pdf>.
- [6] S.N. Pandey, P.S. Trivedi, S.P. Mishra, *A Textbook of Botany II*, Vikas Publishing House Pvt. Ltd., New Delhi, India, 1977.
- [7] P.B. Khare, Ferns and fern allies - their significance and fantasies, *Appl. Bot. Abstr.* 16 (1) (1996) 50–61.
- [8] C. Thakur, S. Rajbhandary, Fern and fern allies of panchase protected forest, central Nepal, *J. Plant Res.* 16 (1) (2018) 39–45.
- [9] Z.P. Gao, R.F. Li, B.H. Wang, Y.R. Lu, Progress in chemical constituents of genus *Dryopteris*, *Chin. J. Exp. Tradit. Med. Formulae* 9 (3) (2003) 50–55.
- [10] L. Wang, Z.R. He, Advances in medical components of pteridophytes, *Chin. Wild Plant Resour.* 25 (2006) 1–4.
- [11] R. Ma, H. Pan, T. Shen, P. Li, Y. Chen, Z. Li, X. Di, S. Wang, Interaction of flavonoids from *Woodwardia unigemmata* with bovine serum albumin (BSA): application of spectroscopic techniques and molecular modeling methods, *Molecules* 22 (2017) 1317.
- [12] R.P. Chaudhary, *Biodiversity in Nepal: Status and Conservation*, Tec Press Books, Bangkok, 1998.
- [13] M. Hassler, World ferns: checklist of ferns and lycophtyes of the world (version Apr. 2018), in: Y. Roskov, L. Abucay, T. Orrell, D. Nicolson, N. Bailly, P.M. Kirk, T. Bourgoin, R.E. De Walt, W. Decock, A. De Wever, E. van Nieukerken, J. Zaruccchi, L. Penev (Eds.), *Species 2000 and ITIS Catalogue of Life*, 2018 Annual Checklist, Species 2000, Naturalis, Leiden, the Netherlands, 2018. <http://www.catalogueoflife.org/annual-checklist/2018>.
- [14] The Pteridophytes (Ferns and Fern Allies). <http://www.theplantlist.org/browse/P/>. (Accessed 27 March 2022).
- [15] S.G. Lu, *Pteridology*, Higher Education Press, Beijing, 2007.
- [16] Pteridophytes. <https://bsi.gov.in/page/en/pteridophytes>. (Accessed 27 March 2022).
- [17] A. Ebihara, J.H. Nitta, An update and reassessment of fern and lycophtye diversity data in the Japanese Archipelago, *J. Plant Res.* 132 (2019) 723–738.
- [18] R. Knapp, T.C. Hsu, *Ferns and Fern Allies of Taiwan*, Second supplement, KBCC Press, Taiwan, 2017.
- [19] M.D.A. delos Angeles, I.E. Buot Jr., Orders and families of Philippine pteridophytes, *J. Nat. Stud.* 11 (1–2) (2012) 19–33. <https://www.journalofnaturestudies.org/files/19-33-delos-angeles-full.pdf>.
- [20] U.K. Yusuf, *Ferns of Malaysian Rain Forest. A Journey through the Fern World*, Universiti Putra Malaysia Press, Serdang, 2010.
- [21] S. Lindsay, D.J. Middleton, T. Boonkerd, S. Suddee, Towards a stable nomenclature for Thai ferns, *Thai For. Bull. (Bot.)* 37 (2009) 64–106. <http://li01.tci-thaijo.org/index.php/ThaiForestBulletin/article/view/24338>.
- [22] K.R. Bhattacharai, Ethnobotanical survey on plants used in Ilam Municipality of Ilam district, eastern Nepal, *Banko Janakari* 30 (2) (2020) 11–35. http://bankojanakari.gov.np/articles/2-BJ_Ethnobotanical_Bhattacharai_final_1606748171.pdf.
- [23] I.M. Bhagat, S. Shrestha, Fern and fern-allies of eastern terai, Nepal, *Our Nat.* 8 (2010) 359–362.
- [24] N. Thapa, Pteridophytes of Nepal, in: M.S. Bista, M.K. Adhikari, K.R. Rajbhandary (Eds.), *Bulletin of Department of Plant Resources No. 19*, His Majesty's Government of Nepal, Ministry of Forests and Soil Conservation, Department of Plant Resources, National Herbarium and Plant Laboratories, Lalitpur, 2002, pp. 1–175.
- [25] D.R. Kandel, C.R. Fraser-Jenkins, *Ferns and Fern-Allies of Nepal 3*, Ministry of Forests and Environment, Department of Plant Resources, National Herbarium and Plant Laboratories, Godawari, Lalitpur, Nepal, 2020.
- [26] N. Joshi, K. Kehlenbeck, B.L. Maass, Traditional, neglected vegetables of Nepal: their sustainable utilization for meeting human needs, in: Conference on International Agricultural Research for Development, Tropentag 2007, University of Kassel-Witzenhausen and University of Göttingen, October 9–11, 2007. <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.624.5143&rep=rep1&type=pdf>.
- [27] S. Chettri, S. Manivannan, V.R. Muddarsu, Nutrient and elemental composition of wild edible ferns of the Himalaya, *Am. Fern J.* 108 (3) (2018) 95–106.
- [28] T. Seal, Evaluation of nutritional potential of wild edible plants, traditionally used by the tribal people of Meghalaya State in India, *Am. J. Plant Nutr. Fertilization Technol.* 2 (2012) 19–26.
- [29] M.K. Aulakh, N. Kaur, M.I.S. Saggo, Bioactive phytoconstituents of Pteridophytes - a review, *Indian Fern J.* 36 (2019) 37–79.
- [30] F. Buchanan-Hamilton, *An Account of the Kingdom of Nepal*, Asian Educational Service, New Delhi, 1986.
- [31] C.R. Fraser-Jenkins, Nepal's Little Known Pteridophytes, the Hidden Works of David Don, and the Geography and Distribution of Indo-Himalayan Ferns, with State Lists, Website Version, updated 31 Dec. 2010, 2011, www.groups.yahoo.com/group/Indian-Ferns, 2010. (Accessed 20 July 2020).
- [32] S. Rajbhandary, Fern and fern allies of Nepal, in: P.K. Jha, M. Siwakoti, S. Rajbhandary (Eds.), *Frontiers of Botany*, Central Department of Botany, Tribhuvan University, Kirtipur, Kathmandu, 2016, pp. 124–150.
- [33] N. Wallich, *Tentamen Flora Neapensis Illustratae Consisting of Botanical Descriptions and Lithographic Figures of Select Nipal Plants*, Asiatic Lithographical Press, Calcutta, Serampore, 1826. <https://www.biodiversitylibrary.org/page/58005015#page/7/mode/1up>.
- [34] D. Don, *Prodromus Florae Nepalensis*, J. Gale, London, England, 1825.
- [35] J.D. Hooker, *The Flora of British India 1–7*, L. Reeve and Co., London, 1872.
- [36] T. Moore, *Index Filicum*, William Pamplin, London, 1857.
- [37] R. Sim, A Priced Catalogue with Brief Descriptive and Cultural Remarks on the Extensive Collection of Stove, Greenhouse and Hardy Exotic and British Ferns, Grown for Sale by Robert Sim, Nurseyman, Foot's Cray, Kent, London, 1859. <http://books.google.com.ng/books?id=VNRRbAAAQAAJ&pg=PA1&hl=gv=PA1&dq=v=onepage&pg=q&f=false>.
- [38] R. Strachey Sir, J.F. Duthie, Catalogue of the Plants of Kumaon and of the Adjacent Portions of Garhwal and Tibet, Based on the Collections Made by Strachey and Winterbottom during the Years 1846 to 1849 and on the Catalogue Originally Prepared in 1852 by Sir Richard Strachey, Lovell Reeve and Co. Ltd., London, 1906.
- [39] I.H. Burkhill, Notes from a Journey to Nepal, Superintendent Government Printing, Calcutta, India, 1910.
- [40] C.R. Lancaster, *A Plantsman in Nepal*, 1981. London.
- [41] M.B. Raizada, K.M. Vaid, *Fern of Nepal*, India Forester 78 (1952) 576–581.
- [42] B.D. Panday, Some aspects of the vegetation of Nepal, *Bull. Bot. Survey India* 4 (4) (1962) 135–140.
- [43] M. Tagawa, Contribution to the ferns of Annapurna-Dhaulagiri range, central Nepal, *J. Bombay Nat. Hist. Soc.* 72 (3) (1975) 728–731.
- [44] K. Iwatsuki, An enumeration of the pteridophytes of Nepal, in: H. Ohba, S.B. Malla (Eds.), *The Himalayan Plants*, University of Tokyo Press, Tokyo, 1988, pp. 231–339. <http://umdb.um.u-tokyo.ac.jp/DKankoub/Bulletin/no31/no31018.html>.
- [45] K.R. Rajbhandari, History of botanical explorations in Nepal, *J. Bombay Nat. Hist. Soc.* 73 (1976) 468–481.
- [46] V.L. Gurung, Pteridophyta, in: S.B. Malla, A.B. Shrestha, S.B. Rajbhandari, T.B. Shrestha, P.M. Adhikari, S.R. Adhikari (Eds.), *Catalogue of Nepalese Vascular Plants*, Bulletin of the Department of Medicinal Plant Nepal No. 7, His Majesty's Government of Nepal, Ministry of Forests, Department of Medicinal Plants, Thapathali, Kathmandu, Nepal, 1976, pp. 1–27.
- [47] V.L. Gurung, Ferns, in: T.C. Majapuria (Ed.), *Nepal - Nature's Paradise*, White Lotus Co. Ltd., Bangkok, Thailand, 1984, pp. 194–211.
- [48] R. Shrestha, V.L. Gurung, A study on ecology of euphorbiaceous ferns of Nepal Himalaya, in: Proceedings of National Conference on Science and Technology, April 24–29, 1988, Royal Nepal Academy of Science and Technology, Kathmandu, Nepal, 1989, pp. 359–364.
- [49] V.L. Gurung, Ferns: the Beauty of Nepalese Flora, Sahayogi Press Pvt. Ltd., Kathmandu, 1991.
- [50] N.P. Manandhar, A survey of medicinal plants of Jajarkot district, Nepal, *J. Ethnopharmacol.* 48 (1995) 1–6.
- [51] N.P. Manandhar, An inventory of some herbal drugs of Myagdi district, Nepal, *Econ. Bot.* 49 (4) (1995) 371–379.
- [52] N.P. Manadhar, Ethnobotanical observation on ferns and fern allies of Nepal, in: J.K. Maheshwari (Ed.), *Ethnobotany in South Asia*, Scientific Publishers, Jodhpur, India, 1996, pp. 414–422.
- [53] N.P. Manandhar, *Plants and People of Nepal*, Timber Press, Portland, Oregon, 2002.
- [54] D.K. Subba, B.K. Rai, M.R. Dhakal, Food value of some edible ferns from Dharan, south eastern Nepal, *J. Bombay Nat. Hist. Soc.* 98 (3) (2001) 499–502.
- [55] S.R. Baral, P.P. Kurmi, *A Compendium of Medicinal Plants in Nepal*, Rachana Sharma, Maijubhala, Kathmandu, Nepal, 2006.
- [56] N. Phuyal, G.D. Bhatta, S. Maharjan, K.K. Pokharel, A contribution to the pteridophytic flora of Makwanpur district, central Nepal, *Bull. Dept. Plant Resour.* 33 (2011) 45–59.
- [57] M. Pathak, N. Phuyal, L.R. Tharu, Inventory of the pteridophytic flora of Sankhuwasabha district, eastern Nepal with notes on medicinal values, *Bull. Dept. Plant Resour.* 34 (2012) 47–55.
- [58] C.R. Fraser-Jenkins, D.R. Kandel, S. Pariyar, *Ferns and Fern-Allies of Nepal 1*, Ministry of Forests and Soil Conservation, Department of Plant Resources, National Herbarium and Plant Laboratories, Nepal, 2015.

- [59] C.R. Fraser-Jenkins, D.R. Kandel, Ferns and Fern-Allies of Nepal 2, Ministry of Forests and Soil Conservation, Department of Plant Resources, National Herbarium and Plant Laboratories, Nepal, 2019.
- [60] C.R. Fraser-Jenkins, K.N. Gandhi, B.S. Kholia, An Annotated Checklist of Indian Pteridophytes Part-1 (Lycopodiaceae to Thelypteridaceae), Bishen Singh Mahendra Pal Singh, Dehra Dun, India, 2016.
- [61] C.R. Fraser-Jenkins, K.N. Gandhi, B.S. Kholia, An Annotated Checklist of Indian Pteridophytes Part-2 (Woodsiaceae to Dryopteridaceae), Bishen Singh Mahendra Pal Singh, Dehra Dun, India, 2018.
- [62] C.R. Fraser-Jenkins, K.N. Gandhi, B.S. Kholia, D.R. Kandel, An Annotated Checklist of Indian Pteridophytes Part-3 (Lomariopsidaceae to Salviniaceae), Bishen Singh Mahendra Pal Singh, Dehra Dun, India, 2020.
- [63] C.R. Fraser-Jenkins, New Species Syndrome in Indian Pteridology and the Ferns of Nepal, International Book Distributors, Dehra Dun, India, 1997.
- [64] C.R. Fraser-Jenkins, Taxonomic Revision of Three Hundred Indian Subcontinental Pteridophytes with a Revised Census-List, Bishen Singh Mahendra Pal Singh, Dehra Dun, India, 2008.
- [65] C.R. Fraser-Jenkins, A brief comparison of modern pteridophyte classifications (families and genera in India), Indian Fern J. 26 (2010) 107–131.
- [66] T.B. Shrestha, B. Bhandari, N. Belbase, Nepal Country Report on Biological Diversity, IUCN Nepal, Kathmandu, 1999.
- [67] K.U. Kramer, P.S. Green, Pteridophytes, in: K. Kubitzki (Ed.), The Families and Genera of Vascular Plants 1, Pteridophytes and Gymnosperms, Springer-Verlag, 1990, pp. 12–279.
- [68] The Plant List. <http://www.theplantlist.org/>. (Accessed 18 March 2021).
- [69] eFloras. <http://www.efloras.org/>. (Accessed 18 March 2021).
- [70] Y. Liu, W. Wujsiguleng, C. Long, Food uses of ferns in China: a review, Acta Soc. Bot. Pol. 81 (4) (2012) 263–270.
- [71] R. Ojha, H.P. Devkota, Edible and medicinal pteridophytes of Nepal: a review, Ethnobot. Res. Appl. 22 (2021) 16.
- [72] D.R. Kandel, Pteridophytes of Nepal, in: M. Siwakoti, P.K. Jha, S. Rajbhandary, S.K. Rai (Eds.), Plant Diversity of Nepal, Botanical Society of Nepal, Kathmandu, 2020, pp. 10–22.
- [73] D. Shrestha, Indigenous vegetables of Nepal for biodiversity and food security, Int. J. Biodivers. Conserv. 5 (3) (2013) 98–108.
- [74] D.R. Dangol, K.L. Maharjan, S.K. Maharjan, A.K. Acharya, Wild edible plants in Nepal, in: B.K. Joshi, H.B. KC, A.K. Acharya (Eds.), Conservation and Utilization of Agricultural Plant Genetic Resources in Nepal, Proceedings of 2nd National Workshop, 22–23 May 2017, Dhulikhel, National Agriculture Genetic Resources Center, Food Development Directorate, Department of Agriculture, Ministry of Agricultural Development, Kathmandu, Nepal, 2017, pp. 390–407. https://nepalindata.com/media/resources/bulkuploaded/Plant_Genetic_Resources CUAPG_R_Nepal_OCT17.pdf.
- [75] A. Rijal, Surviving on knowledge: ethnobotany of Chepang community from mid-hills of Nepal, Ethnobot. Res. Appl. 9 (2011) 181–215. <https://ethnobotanyjournal.org/index.php/era/article/view/249>.
- [76] M.B. Rokaya, Z. Münzbergová, B. Timsina, Ethnobotanical study of medicinal plants from the Humla district of western Nepal, J. Ethnopharmacol. 130 (2010) 485–504.
- [77] Y. Uprety, R.C. Poudel, K.K. Shrestha, S. Rajbhandary, N.N. Tiwari, U.B. Shrestha, H. Asselin, Diversity of use and local knowledge of wild edible plant resources in Nepal, J. Ethnobiol. Ethnomed. 8 (2012) 16.
- [78] N. Joshi, M. Siwakoti, Wild vegetables used by local community of Makawanpur district and their contribution to food security and income generation, Nepal J. Sci. Technol. 13 (1) (2012) 59–66.
- [79] S.K. Rana, P.S. Oli, H.K. Rana, Traditional botanical knowledge (TBK) on the use of medicinal plants in Sikles area, Nepal, Asian J. Plant Sci. Res. 5 (11) (2015) 8–15.
- [80] P.M. Shrestha, S.S. Dhillion, Diversity and traditional knowledge concerning wild food species in a locally managed forest in Nepal, Agrofor. Syst. 66 (2006) 55–63.
- [81] S. Bhattrai, S. Rajbhandary, Pteridophyte flora of manaslu conservation area, central Nepal, Am. J. Plant Sci. 8 (4) (2017) 680–687.
- [82] D.R. Kandel, M. Pathak, Documentation of ferns from subtropical forests of Pyuthan district, western Nepal, J. Dept. Plant Resour. 35 (2013) 46–49.
- [83] S. Shrestha, S.K. Rai, Survey of marketable vegetables and edible fruits in Dharan, eastern Nepal, Nepalese J. Biosci. 2 (2012) 134–147. <https://www.nepjol.info/index.php/NJBS/article/view/7501>.
- [84] A.G. Singh, Survey of some medicinally important leafy vegetables in Rupandehi district of western Nepal, Int. J. Appl. Sci. Biotechnol. 3 (2015) 111–118.
- [85] A.R. Joshi, K. Joshi, Indigenous uses of wetland plant diversity of two valleys (Kathmandu and Pokhara) in Nepal, Ethnobotany 21 (2009) 11–17.
- [86] D.R. Dangol, Traditional uses of plants of commonland habitats in western Chitwan, Nepal, J. Inst. Agric. Anim. Sci. 29 (2008) 71–78.
- [87] M.D. Bhatt, A. Tewari, S.P. Singh, Floristic composition of weeds in paddy fields in Mahendranagar, Nepal, Ecoprint 16 (2009) 15–19.
- [88] L.B. Thapa, T.M. Dhakal, R. Chaudhary, Wild edible plants used by endangered and indigenous Raji tribes in western Nepal, Int. J. Appl. Sci. Biotechnol. 2 (3) (2014) 243–252. <https://www.nepjol.info/index.php/IJASBT/article/view/10969/9012>.
- [89] K.P. Aryal, Å. Berg, B. Ogle, Uncultivated plants and livelihood support - a case study from the Chepang people of Nepal, Ethnobot. Res. Appl. 7 (2009) 409–422. <https://ethnobotanyjournal.org/index.php/era/article/view/255>.
- [90] P. Luni, K. Maharjan, N. Joshi, Forest and food security of indigenous people: a case of Chepangs in Nepal, J. Int. Dev. Cooperation 17 (1) (2011) 113–135.
- [91] S.A. Mir, A.K. Mishra, Z.A. Reshi, M.P. Sharma, Preliminary phytochemical screening of some pteridophytes from district Shopian (J & K), Int. J. Pharm. Pharmaceut. Sci. 5 (S4) (2013) 632–637.
- [92] A. Poudel, S.G. Kim, D.K. Kim, Y.K. Kim, Y.S. Lee, G.W. Lee, B.S. Min, H.J. Jung, Antioxidative and antioesity activity of Nepalese wild herbs, Nat. Prod. Sci. 17 (2011) 123–129. <https://www.koreascience.or.kr/article/JAKO201121055567508.page>.
- [93] K.P. Acharya, M.B. Rokaya, Ethnobotanical survey of medicinal plants traded in the streets of Kathmandu valley, Sci. World 3 (3) (2005) 44–48.
- [94] J. Cao, X. Xia, X. Dai, J. Xiao, Q. Wang, K. Andrae-Marobela, H. Okatch, Flavonoids profiles, antioxidant, acetylcholinesterase inhibition activities of extract from *Dryoathyrium boryanum* (Willd.) Ching, Food Chem. Toxicol. 55 (2013) 121–128.
- [95] G.K. Dash, S.K.J. Khadidi, A.F. Shamsuddin, Pharmacognostic studies on *Diplazium esculentum* (retz.) sw, Der Pharm. Lett. 9 (2017) 113–120. <https://www.scholarsresearchlibrary.com/articles/pharmacognostic-studies-on-diplazium-esculentum-retz-sw.pdf>.
- [96] A. Semwal, S. Kaushik, S.P. Bhatt, A. Negi, Antibacterial activity of *Diplazium esculentum* (retz.) sw, Pharmacogn. J. 3 (2011) 77–79.
- [97] K.R. Bhattacharai, M.K. Khadka, Ethnobotanical survey of medicinal plants from Ilam district, east Nepal, Our Nat. 14 (1) (2016) 78–91.
- [98] S. Akter, Md M. Hossain, I. Ara, P. Akhtar, Investigation of in vitro antioxidant, antimicrobial and cytotoxic activity of *Diplazium esculentum* (RETZ) SW, Int. J. Adv. Pharm. Biol. Chem. 3 (3) (2014) 723–733. <http://www.ijapbc.com/files/32-3369.pdf>.
- [99] V.B. Amoroso, D.A. Antesa, D.P. Buenavista, F.P. Coritico, Antimicrobial, antipyretic, and anti-inflammatory activities of selected Philippine medicinal pteridophytes, Asian J. Biodivers. 5 (2014) 18–40. <http://www.asianscientificjournals.com/new/publication/index.php/ajob/article/viewFile/479>.
- [100] M.O. Ullah, M. Haque, K.F. Urmia, A.H. Md Zulfikar, E.S. Anita, M. Begum, K. Hamid, Antibacterial activity and brine shrimp lethality bioassay of methanolic extracts of fourteen different edible vegetables from Bangladesh, Asian Pac. J. Trop. Biomed. 3 (2013) 1–7.
- [101] S. Roy, B. Hazra, N. Mandal, T.K. Chaudhuri, Assessment of the antioxidant and free radical scavenging activities of methanolic extract of *Diplazium esculentum*, Int. J. Food Prop. 16 (2013) 1351–1370.
- [102] A. Kaushik, C. Jijta, J.J. Kaushik, R. Zeray, A. Ambesajir, L. Beyene, FRAP (Ferric reducing ability of plasma) assay and effect of *Diplazium esculentum* (Retz) Sw. (a green vegetable of north India) on central nervous system, Indian J. Nat. Prod. Resour. 3 (2012) 228–231. <http://nopr.niscair.res.in/bitstream/123456789/4404/1/IJNPR%203%282%29%2020228-231.pdf>.
- [103] J.A. Junejo, G. Gogoi, J. Islam, M. Rudrapal, P. Mondal, H. Hazarika, K. Zaman, Exploration of antioxidant, antidiabetic and hepatoprotective activity of *Diplazium esculentum* - a wild edible plant from north eastern India, Future J. Pharm. Sci. 4 (2018) 93–101.
- [104] T. Kunkeaw, U. Suttiisanee, D. Trachootham, J. Karinchai, B. Chantong, S. Potikanond, W. Inthachat, P. Pitchakarn, P. Temviriyankul, *Diplazium esculentum* (Retz.) Sw. reduces BACE-1 activities and amyloid peptides accumulation in *Drosophila* models of Alzheimer's disease, Sci. Rep. 11 (2021), 23796.
- [105] A. Rahmat, Determination of total antioxidant activity in three types of local vegetables shoots and the cytotoxic effect of their ethanolic extracts against different cancer cell lines, Asia Pac. J. Clin. Nutr. 12 (3) (2003) 308–311.
- [106] A. Kaushik, J.J. Kaushik, A. Das, S. Gemal, D. Gaim, Preliminary studies on anti-inflammatory activities of *Diplazium esculentum* in experimental animal models, Int. J. Pharm. Sci. Res. 2 (2011) 1251–1253.
- [107] A. Semwal, M.S. Farswan, In-vitro anthelmintic activity of *Diplazium esculentum* (Retz.) Swiss rhizome extract, J. Pharmacogn. Phytochem. 1 (2012) 84–87. <https://www.phytojournal.com/archives/2012/vol1issue4/PartA/11.1.pdf>.
- [108] S. Pathania, P. Kumar, S. Singh, S. Khatoon, A.K.S. Rawat, N. Punetha, D.J. Jensen, D.R. Lauren, R. Somvanshi, Detection of ptaquiloside and quercetin in certain Indian ferns, Curr. Sci. 102 (2012) 1683–1691. <https://www.currentscience.ac.in/Volumes/102/12/1683.pdf>.
- [109] J.V.V. Tongco, R.A.P. Villaber, R.M. Aguda, R.A. Razal, Nutritional and phytochemical screening, and total phenolic and flavonoid content of *Diplazium esculentum* (Retz.) Sw. from Philippines, J. Chem. Pharmaceut. Res. 6 (8) (2014) 238–242. <https://www.jocpr.com/articles/nutritional-and-phytochemical-screening-and-total-phenolic-and-flavonoid-content-of-diplazium-esculentum-retz-sw-from-ph.pdf>.
- [110] P. Semwal, S. Painuli, K.M. Painuli, G. Antika, T.B. Turner, A. Thapliyal, W.N. Setzer, M. Martorell, Md M. Alshehri, Y. Taheri, S.D. Dastan, S.A. Ayatollahi, A.T. Petkoska, J. Sharifi-Rad, W.C. Cho, *Diplazium esculentum* (Retz.) Sw.: ethnomedicinal, phytochemical, and pharmacological overview of the Himalayan ferns, Oxid. Med. Cell. Longev. (2021), 1917890.
- [111] E.E. Essien, R. Ascrizzi, G. Flaminii, Characterization of volatile compounds of *Diplazium esculentum*, Chem. Natl. Compd. 55 (2019) 958–959.
- [112] B. Naik, V.K. Maurya, V. Kumar, V. Kumar, S. Upadhyay, S. Gupta, Phytochemical analysis of *Diplazium esculentum* reveals the presence of medically important components, Curr. Nutr. Food Sci. 17 (2021) 210–215.
- [113] M.J. Caldwell, Ascorbic acid content of Malaysian leaf vegetables, Ecol. Food Nutr. 1 (1972) 313–317.
- [114] S.K. Srivastava, S.D. Srivastava, V.K. Saksena, S.S. Nigam, A flavanone glycoside from *Diplazium esculentum*, Phytochemistry (Oxf.) 20 (1981) 862.
- [115] L.S. Ching, S. Mohamed, Alpha-tocopherol content in 62 edible tropical plants, J. Agric. Food Chem. 49 (2001) 3101–3105.

- [116] K.H. Miean, S. Mohamed, Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants, *J. Agric. Food Chem.* 49 (2001) 3106–3112.
- [117] R. Somvanshi, D.R. Lauren, B.L. Smith, R.K. Dawra, O.P. Sharma, V.K. Sharma, A.K. Singh, N.K. Gangwar, Estimation of the fern toxin, ptaquiloside, in certain Indian ferns other than bracken, *Curr. Sci.* 91 (2006) 1547–1552. <https://www.currentscience.ac.in/Volumes/91/11/1547.pdf>.
- [118] M.D. Astuti, E.M. Kuntorini, F.E.P. Wisuda, Isolasi dan identifikasi terpenoid Dari fraksi n-butanol herba lampasau (*Diplazium esculentum* Swartz), *Jurnal Kimia Valensi* 4 (2014) 20–24.
- [119] A. Wali, S. Sharma, M. Walia, P. Kumar, S. Thakur, A. Kumari, B. Lal, V.K. Agnihotri, Two edible ferns of western Himalaya: a comparative in vitro nutritional assessment, antioxidant capacity and quantification of lutein by UPLC-DAD, *Int. J. Food Nutr. Sci.* 5 (9) (2016) 9–18.
- [120] M. Watamabe, T. Miyashita, H.P. Devkota, Phenolic compounds and ecdysteroids of *Diplazium esculentum* (Retz.) Sw. (Athyriaceae) from Japan and their chemotaxonomic significance, *Biochem. Systemat. Ecol.* 94 (2021), 104211.
- [121] B. Sareen, A. Bhattacharya, V. Srivatsan, Nutritional characterization and chemical composition of *Diplazium maximum* (D. Don) C. Chr, *J. Food Sci. Technol.* 58 (2021) 844–854.
- [122] B. Malla, R.B. Chhetri, Indigenous knowledge on ethnobotanical plants of Kavrepalanchowk district, Kathmandu Univ. *J. Sci. Eng. Technol.* 5 (2) (2009) 96–109.
- [123] S. Parajuli, N.T. Pun, S. Parajuli, N. Jamarkattel-Pandit, Antioxidant activity, total phenol and flavonoid contents in some selected medicinal plants of Nepal, *J. Health Allied Sci.* 2 (2012) 27–31.
- [124] H.K. Goswami, K. Sen, R. Mukhopadhyay, Pteridophytes: evolutionary boon as medicinal plants, *Plant Genetic Resour.* 14 (4) (2016) 328–355.
- [125] Adenan, E. Suhartono, *Stenochlaena palustris* aqueous extract reduces hepatic peroxidative stress in *Marmota caligata* with induced fever, *Univ. Medicina* 29 (3) (2010) 123–128.
- [126] N.J.Y. Chear, K.Y. Khaw, V. Murugaiyah, C.S. Lai, Cholinesterase inhibitory activity and chemical constituents of *Stenochlaena palustris* fronds at two different stages of maturity, *J. Food Drug Anal.* 24 (2016) 358–366.
- [127] V. Sumathy, S.J. Lachumy, Z. Zuraini, S. Sasidharan, Effects of *Stenochlaena palustris* leaf extract on growth and morphogenesis of food borne pathogen *Aspergillus niger*, *Malays. J. Nutr.* 16 (2010) 439–446.
- [128] H. Liu, J. Orjala, O. Sticher, T. Rali, Acylated flavonol glycosides from leaves of *Stenochlaena palustris*, *J. Nat. Prod.* 62 (1999) 70–75.
- [129] T.T. Chai, E. Panirchellvum, H.C. Ong, F.C. Wong, Phenolic contents and antioxidant properties of *Stenochlaena palustris*, an edible medicinal fern, *Bot. Stud.* 53 (2012) 439–446.
- [130] Y. Ponnusamy, N.J.Y. Chear, S. Ramanathan, V. Murugaiyah, C.S. Lai, Antioxidant and antibacterial properties of Malaysian ferns used traditionally against infection, *J. Nat. Prod. Plant Resour.* 3 (6) (2013) 14–18. <https://www.scholarsresearchlibrary.com/articles/antioxidant-and-antibacterial-properties-of-malaysian-ferns-use-d-traditionallyagainst-infection.pdf>.
- [131] T.T. Chai, M.T. Kwek, H.C. Ong, F.C. Wong, Water fraction of edible medicinal fern *Stenochlaena palustris* is a potent α-glucosidase inhibitor with concurrent antioxidant activity, *Food Chem.* 186 (2015) 26–31.
- [132] L.N. Azizah, P. Wardhani, H. Arwati, Antimalarial activity of ethanol extract of kelakai leaves (*Stenochlaena palustris*) to parasitemia and splenomegaly in BALB/c mice infected with *Plasmodium berghei* ANKA, *J. Ilmiah Mahasiswa Kedokteran Univ. Airlangga* 13 (1) (2022).
- [133] H. Liu, J. Orjala, T. Rali, O. Sticher, Glycosides from *Stenochlaena palustris*, *Phytochemistry (Oxf.)* 49 (1998) 2403–2408.
- [134] R.D. Gaur, B.P. Bhatt, Folk utilization of some pteridophytes of deoprayag area in garhwal Himalaya: India, econ, *Bot* 48 (1994) 146–151.
- [135] M.O. Nwosu, Ethnobotanical studies on some pteridophytes of southern Nigeria, *Econ. Bot.* 56 (2002) 255–259.
- [136] S.K. Verma, S. Kanwar, Medicinal pteridophytes used in the treatment of various diseases by the inhabitants of Sarkaghat Tehsil, Mandi district, Himachal Pradesh, *J. Pharmaceut. Sci. Res.* 12 (3) (2020) 360–364. <https://www.jpsr.pharmainfo.in/Documents/Volumes/vol12issue03/jpsr12032004.pdf>.
- [137] Z.T. Ding, Y.S. Fang, Z.G. Tai, M.H. Yang, Y.Q. Xu, F. Li, Q.E. Cao, Phenolic content and radical scavenging capacity of 31 species of ferns, *Fitoterapia* 79 (2008) 581–583.
- [138] P. Takuli, K. Khulbe, P. Kumar, A. Parki, A. Syed, A.M. Elgorban, Phytochemical profiling, antioxidant and antibacterial efficacy of a native Himalayan fern: *Woodwardia unigemmata* (Makino) Nakai, *Saudi J. Biol. Sci.* 27 (2020) 1961–1967.
- [139] R.S. Chang, H.W. Yeung, Inhibition of growth of human immunodeficiency virus in vitro by crude extracts of Chinese medicinal herbs, *Antivir. Res.* 8 (1988) 163–176.
- [140] R.A. Collins, T.B. Ng, W.P. Fong, C.C. Wan, H.W. Yeung, A comparison of human immunodeficiency virus type 1 inhibition by partially purified aqueous extracts of Chinese medicinal herbs, *Life Sci.* 60 (1997) 345–351.
- [141] A.V. Singh, Edible and medicinal ferns of Lakhimpur, Assam, *Int. J. Eng. Sci. Math.* 6 (3) (2017) 1–16. https://www.ijesm.co.in/abstract.php?article_id=3101&title=EDIBLE%20AND%20MEDICINAL%20FERNS%20OF%20LAKHIMPUR,%20ASSAM.
- [142] M. Adhikari, R. Thapa, R.M. Kunwar, H.P. Devkota, P. Poudel, Ethnomedicinal uses of plant resources in the machhapuchhre rural municipality of kaski district, Nepal, *Medicines* 6 (2) (2019) 69.
- [143] J. Barukial, J.N. Sarmah, Ethnomedicinal plants used by the people of Golaghat district, Assam, India, *Int. J. Med. Aromatic Plants* 1 (3) (2011) 203–211.
- [144] N. Yan, H. Zhang, Z. Zhang, H. Zhang, L. Zhou, T. Chen, S. Feng, C. Ding, M. Yuan, The extraction, antioxidant and against β-amyloid induced toxicity of polyphenols from *Alsophila spinulosa* leaves, *Arab. J. Chem.* 15 (2022), 103707.
- [145] H. Wada, T. Satake, T. Murakami, T. Kojima, Y. Saiki, C.M. Chen, Chemische und chemotaxonomische untersuchungen der pterophyten. LIX. Chemische untersuchungen der inhaltsstoffe von *Alsophila spinulosa* Tryon, *Chem. Pharm. Bull.* 33 (1985) 4182–4187.
- [146] Y. Arai, N. Koide, F. Ohki, H. Ageta, L.L. Yang, K.Y. Yen, Fern constituents: triterpenoids isolated from leaflets of *Cyathea spinulosa*, *Chem. Pharm. Bull.* 42 (1994) 228–231.
- [147] J.S. Jiang, Z.L. Zhan, Z.M. Feng, Y.N. Yang, P.C. Zhang, Study on the chemical constituents from *Cyathea spinulosa*, *J. Chin. Med. Mater.* 35 (4) (2012) 568–570.
- [148] Y. Cheng, F.Z. Chen, X.J. He, Isolation of three chemical constituents from *Alsophila spinulosa* stalks for the first time, *Med. Plants* 2 (11) (2011) 5–7.
- [149] R. M Lu, M. Cao, P.Y. Liao, J.H. Wei, Chemical constituents in stems of Zhuang medicine *Alsophila spinulosa*, *Chin. Tradit. Herb. Drugs* 44 (16) (2013) 2195–2199.
- [150] J. Vetter, A biological hazard of our age: bracken fern [*Pteridium aquilinum* (L.) Kuhn] – a review, *Acta Vet. Hung.* 57 (1) (2009) 183–196.
- [151] CSIR, The Wealth of India, A dictionary of Indian raw materials and industrial products. Raw Materials VIII, Ph-Re, Publications and Information Directorate, Council of Scientific and Industrial Research, New Delhi, 1969, pp. 299–300.
- [152] I.A. Evans, J. Mason, Carcinogenic activity of bracken, *Nature* 208 (1965) 913–914.
- [153] A.M. Pamukcu, E. Ertük, J.M. Price, G.T. Bryan, Lymphatic leukemia and pulmonary tumors in female Swiss mice fed bracken fern (*Pteris aquilina*), *Cancer Res.* 32 (1972) 1442–1445. <https://aacrjournals.org/cancerres/article/32/7/1442/478915>.
- [154] K. Yoshihira, M. Fukuoka, M. Kuroyanagi, S. Natori, M. Umeda, T. Morohoshi, M. Enomoto, M. Saito, Chemical and toxicological studies on bracken fern, *Pteridium aquilinum* var. *latiusculum*. I. Introduction, extraction and fractionation of constituents, and toxicological studies including carcinogenicity tests, *Chem. Pharm. Bull.* 26 (1978) 2346–2364.
- [155] M.E. Alonso-Amelot, A. Oliveros, A method for the practical quantification and kinetic evaluation of cyanogenesis in plant material. Application to *Pteridium aquilinum* and *Passiflora capsularis*, *Phytochem. Anal.* 11 (2000) 309–316.
- [156] H. Hikino, T. Takahashi, S. Arihara, T. Takemoto, Structure of pteroside B, glycoside of *Pteridium aquilinum* var. *latiusculum*, *Chem. Pharm. Bull.* 18 (1970) 1488–1489.
- [157] H. Hikino, T. Takahashi, T. Takemoto, Structure of pterosides Z and D, glycosides of *Pteridium aquilinum* var. *latiusculum*, *Chem. Pharm. Bull.* 19 (1971) 2424–2425.
- [158] H. Hikino, T. Takahashi, T. Takemoto, Structure of pteroside A and C, glycosides of *Pteridium aquilinum* var. *latiusculum*, *Chem. Pharm. Bull.* 20 (1972) 210–212.
- [159] M. Fukuoka, M. Kuroyanagi, K. Yoshihira, S. Natori, Chemical and toxicological studies on bracken fern, *Pteridium aquilinum* var. *latiusculum*. II. Structures of pterosins, sesquiterpenes having 1-indanone skeleton, *Chem. Pharm. Bull.* 26 (8) (1978) 2365–2385.
- [160] P. Meyer, Thiaminase activities and thiamine content of *Pteridium aquilinum*, *Equisetum arvense*, *Malva parviflora*, *Pennisetum clandestinum* and *Medicago sativa*, *Ondersteopoort J. Vet. Res.* 56 (1989) 145–146.
- [161] P. Selvaraj, A.J. de Britto, K. Sahayaraj, Phytoecdysone of *Pteridium aquilinum* (L.) Kuhn (Dennstaedtiaceae) and its pesticidal property on two major pests, *Arch. Phytopathol. Plant Protect.* 38 (2005) 99–105.
- [162] S. Shweta, R.D. Dixit, T.R. Sahu, Ethnomedical uses of pteridophytes of Amarkatak, Madhya Pradesh, *Int. J. Tradit. Knowl.* 4 (4) (2005) 392–395. <http://nopr.niscair.res.in/handle/123456789/8542>.
- [163] A. Banjamin, V.S. Manickam, Medicinal pteridophytes from the western ghats, *Indian J. Tradit. Knowl.* 6 (4) (2007) 611–618. <http://nopr.niscair.res.in/handle/123456789/1013>.
- [164] J. Kunjani, A.R. Joshi, Ethnobotanical studies on some lower plants of the central development region, Nepal, *Ethanobot. Leafl.* 12 (1) (2008) 832–840. <https://open.lib.siu.edu/ebl/vol2008/iss1/113>.
- [165] P. Parihar, L. Parihar, A. Bohra, In vitro antibacterial activity of fronds (leaves) of some important pteridophytes, *J. Microbiol. Antimicrob.* 2 (2) (2010) 19–22. <https://academicjournals.org/journal/JMA/article-full-text-pdf/A1C09D59652>.
- [166] A. Kathirvel, V. Sujatha, Phytochemical studies, antioxidant activities and identification of active compounds using GC-MS of *Dryopteris cochleata* leaves, *Arab. J. Chem.* 9 (S2) (2016) S1435–S1442.
- [167] A. Kathirvel, A.K. Rai, G.S. Maurya, V. Sujatha, *Dryopteris cochleata* rhizome: a nutritional source of essential elements, phytochemicals, antioxidants and antimicrobials, *Int. J. Pharm. Pharmaceut. Sci.* 6 (S2) (2014) 179–188.
- [168] K. Dubal, S. Patil, M. Dongare, M. Kale, Investigation of chemical composition from *Dryopteris cochleata* (D. Don) C. Chr. (Dryopteridaceae), *Asian J. Pharmaceut. Clin. Res.* 8 (2015) 311–314. <https://innovareacademics.in/journals/index.php/ajpcr/article/view/6821>.
- [169] A. Tapwal, Nisha, S. Garg, N. Gautam, R. Kumar, In vitro antifungal potency of plant extracts against five phytopathogens, *Braz. Arch. Biol. Technol.* 54 (6) (2011) 1093–1098.
- [170] S. Sivasankar, R. Somavanshi, Pathological evaluation of *Polystichum squarrosum* (D. Don) fern in laboratory rats, *Indian J. Exp. Biol.* 39 (2001) 772–776. <http://hdl.handle.net/123456789/23855>.
- [171] CSIR, The Wealth of India, A dictionary of Indian raw materials and industrial products. Raw Materials VI, L-M, Council of Scientific and Industrial Research, New Delhi, 1962, pp. 199–200.

- [172] E. Yadav, M. Mani, P. Chandra, N. Sachan, A.K. Ghosh, A review on therapeutic potential of *Lygodium flexuosum* Linn, Phcog. Rev. 6 (2012) 107–114.
- [173] K.R. Bhattacharai, O.L. Vetaas, J.A. Grytness, Fern species richness along a central Himalayan elevation gradient, Nepal, J. Biogeogr. 31 (2004) 389–400.
- [174] B.B. Gaitonde, R.T. Mahajan, Antifertility activity of *Lygodium flexuosum*, Indian J. Med. Res. 12 (1980) 597–604.
- [175] V.G. Nangare, R.L. Jadhav, Evaluation of contraceptive efficacy of ethanolic extract of *Lygodium flexuosum* in male Albino rat: a histological approach, J. Glob. Biosci. 10 (2021) 8968–8979. https://www.mutagens.co.in/jgb/vol.10/09/1_00901.pdf.
- [176] P.J. Wills, V.V. Asha, Protective effect of *Lygodium flexuosum* (L.) Sw. extract against carbon tetrachloride-induced acute liver injury in rats, J. Ethnopharmacol. 108 (2006) 320–326.
- [177] P.J. Wills, V.V. Asha, Chemopreventive action of *Lygodium flexuosum* extract in human hepatoma PLC/PRF/5 and Hep 3B cells, J. Ethnopharmacol. 122 (2009) 294–303.
- [178] N. Nayak, S. Rath, M.P. Mishra, G. Ghosh, R.N. Padhy, Antibacterial activity of the terrestrial fern *Lygodium flexuosum* (L.) Sw. against multidrug resistant enteric- and uro-pathogenic bacteria, J. Acute Dis. 2 (4) (2013) 270–276.
- [179] M.V. Jeeshesha, Antimicrobial activities and phytochemical investigation of fern, *Lygodium flexuosum* (Linn) Sw., Kongunadu Res. J. 4 (2017) 87–91.
- [180] B. Achari, K. Basu, C.R. Saha, S.C. Prakashi, A new triterpene ester, an anthraquinone and other constituents of the fern *Lygodium flexuosum*, Planta Med. 52 (1986) 329–330.
- [181] A. Basudev, C. Chaudhuri, C.R. Saha, S.C. Prakashi, D.R. McPhail, A.T. McPhail, X-ray crystal structure of lygodinolide: a novel spiro furopyran-perhydrophenanthrene derivatives from *Lygodium flexuosum*, J. Org. Chem. 55 (1990) 4977–4978.
- [182] S. Rao, Pharmacognostical studies on *Lygodium flexuosum*, J. Pharm. Res. 3 (2010) 1976–1978.
- [183] T. Yamauchi, N. Oyama, H. Yamane, N. Murofushi, H. Schraudolf, M. Pour, M. Furber, L.N. Mander, Identification of antheridiogens in *Lygodium circinnatum* and *Lygodium flexuosum*, Plant Physiol 111 (1996) 741–745.
- [184] K.R. Wang, C. Nan, R. Ding, J. Lin, D.J. Hu, S.H. Zhong, R. Gu, A cross-national comparaison of medicinal plants used by the Miao, Yi and Lisu ethnic groups in Yanbian, China, Res. Square (2020).
- [185] B. Malla, D.P. Gauchan, R.B. Chhetri, An ethnobotanical study of medicinal plants used by ethnic people in Parbat district of western Nepal, J. Ethnopharmacol. 165 (2015) 103–107.
- [186] J.B. Won, C.J. Ma, Neuroprotective activities of some medicinal plants against glutamate-induced neurotoxicity in primary cultures of rat cortical cells, Nat. Prod. Sci. 15 (2009) 125–129. https://www.koreascience.or.kr/article/JAKO2009293_49389228.pdf.
- [187] S.F. Ni, Y.J. Pan, P. Wu, Y.C. Chen, C.X. Fu, H. Xu, Analysis of volatile oil from *Lygodium japonicum* by GC-MS, Chin. Pharmaceut. J. 39 (2) (2004) 99–100. <http://hdl.handle.net/10397/62568>.
- [188] L.H. Zhang, Z.Q. Yin, W.C. Ye, S.X. Zhao, L. Wang, F. Hu, Studies on the chemical constituents in herb of *Lygodium japonicum*, China, J. Chin. Mater. Med. 30 (19) (2005) 1522–1524.
- [189] L. Zhu, G. Zhang, S. Wang, L. Li, P. Li, S. Wu, X. Wang, L. Chen, C. Pan, W. Mi, A new compound from *Lygodium japonicum* (Thunb.) Sw, Nat. Prod. Res. 23 (2009) 1284–1288.
- [190] Y.H. Duan, Y. Dai, R.R. He, H. Kurihara, Y.L. Li, X.S. Yao, A new phenylpropanoid glucoside from the aerial parts of *Lygodium japonicum*, J. Asian Nat. Prod. Res. 14 (2012) 286–292.
- [191] H. Yamane, N. Takahashi, K. Takeno, M. Furuya, Identification of gibberellin A₉ methyl ester as a natural substance regulating formation of reproductive organs in *Lygodium japonicum*, Planta 147 (1979) 251–256.
- [192] H. Yamane, Y. Satoh, K. Nohara, M. Nakayama, N. Murofushi, N. Takahashi, K. Takeno, M. Furuya, M. Furber, L.N. Mander, The methyl ester of a new gibberellin, GA73: the principal antheridiogen in *Lygodium japonicum*, Tetrahedron Lett. 29 (1988) 3959–3962.
- [193] H. Konno, T. Nakato, S. Nakashima, K. Katoh, *Lygodium japonicum* fern accumulates copper in the cell wall pectin, J. Exp. Bot. 56 (2005) 1923–1931.
- [194] L. Nahar, F.A. Ripa, R. Zahar, I.J. Bulbul, I. Parvej, Evaluation of analgesic and antidiarrhoeal effects of methanolic extract of *Marsilea quadrifolia* in rat, J. Pharm. Res. 4 (12) (2011) 4374–4376.
- [195] P. Soni, L. Singh, *Marsilea quadrifolia* Linn. - a valuable culinary and remedial fern in Jaduguda, Jharkhand, India, Int. J. Life Sci. Pharm. Res. 2 (3) (2012) L99–L104. https://www.ijlpr.com/admin/php/uploads/106_pdf.pdf.
- [196] K. Gopalakrishnan, R. Udayakumar, Antimicrobial activity of *Marsilea quadrifolia* (L.) against some selected pathogenic microorganisms, Microbiol. Res. J. Int. 4 (9) (2014) 1046–1056. <https://www.journalmriji.com/index.php/MRJI/article/view/6788>.
- [197] K.S. Reddy, C.S. Reddy, S. Ganapaty, Psychopharmacological studies of hydro alcoholic extract of whole plant of *Marsilea quadrifolia*, J. Sci. Res. 4 (2012) 279–285.
- [198] F.A. Ripa, L. Nahar, M. Haque, Md M. Islam, Antibacterial, cytotoxic and antioxidant activity of crude extract of *Marsilea quadrifolia*, Eur. J. Sci. Res. 33 (1) (2009) 123–129.
- [199] P. Sethi, Antimicrobial activities of *Turbinaria conoides* (J. Agardh) kutzng and *Marsilea quadrifolia* linn, Asian J. Plant Sci. Res. 4 (6) (2014) 36–40. <https://www.imedpub.com/articles/antimicrobial-activities-of-turbinaria-conoides-j-agardh-kutzng-andmarsilea-quadrifolia-linn.pdf>.
- [200] R.K. Nivas, M. Boominathan, Antimicrobial evaluation of selected south Indian medicinal plants against *Streptococcus pneumoniae*, Int. J. Curr. Microbiol. Appl. Sci. 4 (2015) 835–840. <https://www.ijcmas.com/vol-4-3/R.%20Kadhar%20Nivas%20and%20M.%20Boominathan.pdf>.
- [201] T.G. Gini, G.J. Jothi, In vitro screening of antibacterial and antifungal activity of *Marsilea quadrifolia* (Marsileaceae) Linn. extract, Am. J. Phytomed. Clin. Ther. 3 (4) (2015) 313–329.
- [202] M. Vijayalakshmi, R. Kiruthika, K. Bharathi, K. Ruckmani, Phytochemical screening by LC-MS analysis and invitro anti-inflammatory activity of *Marsilea quadrifolia* plant extract, Int. J. Pharm. Tech. Res. 8 (2015) 148–157. [https://sphinxsa1.com/2015/ph_vol8_no9/1/\(148-157\)V8N9PT.pdf](https://sphinxsa1.com/2015/ph_vol8_no9/1/(148-157)V8N9PT.pdf).
- [203] T. Mathangi, P. Prabhakaran, DPPH free radical scavenging activity of the extracts of the aquatic fern *Marsilea quadrifolia* Linn, Int. J. Pharm. Pharmaceut. Sci. 2 (2013) 534–536. <https://www.ijcmas.com/vol-2-10/T.%20Mathangi%20and%20Prabhakaran.pdf>.
- [204] A. Chowdhury, T. Panneerselvam, K. Suthendran, C. Bhattachjee, S. Balasubramanian, S. Murugesan, B. Suraj, K. Selvaraj, Optimization of microwave-assisted extraction of bioactive polyphenolic compounds from *Marsilea quadrifolia* L. using RSM and ANFIS modelling, Indian J. Nat. Prod. Resour. 9 (3) (2018) 204–221. <http://nopr.niscair.res.in/bitstream/123456789/45556/1/JINPR%20%283%29%20204-221.pdf>.
- [205] S.S. Dongare, A.P. Maske, S.M. Patil, R.P. Umbare, G.S. Mate, Antidiabetic activity of *Marsilea quadrifolia* Linn in alloxan-diabetic rats, Res. J. Pharmacogn. Phytochem. 1 (2009) 15–17. <https://rjpponline.org/AbstractView.aspx?PID=2010-2-6-14>.
- [206] G. Ashwin, P. Pranay, G. Thrinath, T. Karnaker Reddy, V.S. Giri Prasad, Pharmacological evalution of *Marsilea quadrifolia* plant extracts against Alzheimer's disease, Int. J. Drug Dev. Res. 4 (2) (2012) 153–158. <https://www.ijddr.in/drug-d-development/pharmacological-evaluation-of-marsilea-quadrifolia-plant-extractsagainst-alzheimers-disease.pdf>.
- [207] S. Bhadra, P.K. Mukherjee, A. Bandyopadhyay, Cholinesterase inhibition activity of *Marsilea quadrifolia* Linn. an edible leafy vegetable form West Bengal, India, Nat. Prod. Res. 26 (16) (2012) 1519–1522.
- [208] A. Snehunu, N. Mukunda, M.C.S. Kumar, N. Sadhana, S.N. Narayanan, K.V. Kappal, H. Avinash, B.R. Chandrashekhar, K.R. Rao, B.S. Nayak, Evaluation of anti-epileptic property of *Marsilea quadrifolia* Linn. in maximal electroshock and pentylenetetrazole-induced rat models of epilepsy, Brain Inj. 27 (2013) 1707–1714.
- [209] S. Sahu, G. Dutta, N. Mandal, A.R. Goswami, T. Ghosh, Anticonvulsant effect of *Marsilea quadrifolia* Linn. on pentylenetetrazole induced seizure: a behavioral and EEG study in rats, J. Enthnopharmacol. 141 (2012) 537–541.
- [210] D.E. Vallabhi, V. Elango, Antioxidant activity of *Cyatthula prostrata*, *Ipomea obscura* and *Marsilea quadrifolia* in nephrotoxicity induced male Albino rats, World J. Pharmaceut. Res. 5 (8) (2016) 1000–1006.
- [211] R. Uma, B. Pravin, Invitro cytotoxic activity of *Marsilea quadrifolia* Linn of MCF-7 cells of human breast cancer, Int. Res. J. Med. Sci. 1 (2013) 10–13. http://www.isca.in/MEDI_SCI/Archive/v1/i1/2.ISCA-IRJMedS-2013-006.pdf.
- [212] S.K. Agarwal, S. Roy, P. Pramanick, P. Mitra, R. Gobato, A. Mitra, *Marsilea quadrifolia*: a floral species with unique medicinal properties, Parana J. Sci. Edu. 4 (5) (2018) 15–20.
- [213] A. Dewanjhi, S. Matai, L. Si, S. Barik, A. Nag, Chemical composition of two semi-aquatic plants for food use, Plant Foods Hum. Nutr. 44 (1993) 11–16.
- [214] K. Gopalakrishnan, R. Udayakumar, Phytochemical content of leaf and stem of *Marsilea quadrifolia* (L.), J. Plant Sci. Phytopathol. 1 (2017) 26–37.
- [215] Y. Zhang, H.Y. Tian, Y.F. Tan, Y.L. Wong, H.Y. Wu, J.F. Jia, G.E. Wang, J.J. Gao, Y.F. Li, H. Kurihara, P.C. Shaw, R.W. Jiang, Isolation and identification of polyphenols from *Marsilea quadrifolia* with antioxidant properties in vitro and in vivo, Nat. Prod. Res. 30 (2016) 1404–1410.
- [216] A.G. Singh, A. Kumar, D.D. Tewari, K.A. Bharati, New ethnomedicinal claims from Magar community of Palpa district, Nepal, Indian J. Tradit. Knowl. 17 (3) (2018) 499–511. <http://nopr.niscair.res.in/handle/123456789/44583>.
- [217] S.P. Pradhan, R.P. Chaudhary, S. Sigdel, B.P. Pandey, Ethnobotanical knowledge of Khandadevi and Gokulganga rural municipality of Ramechhap district of Nepal, Ethnobot. Res. Appl. 20 (2020) 1–32. <https://ethnobotanyjournal.org/index.php/era/article/view/2027>.
- [218] G.R. Silwal, Medicinal plants and their traditional uses in Ramkot village, Kathmandu Nepal, Patan Pragya 6 (1) (2020) 180–190. <https://www.nepjol.info/index.php/pragya/article/view/34431>.
- [219] A.K. Dhiman, Ethnomedicinal uses of some pteridophytic species in India, Indian Fern J. 15 (1–2) (1998) 61–64.
- [220] H.B. Singh, Potential medicinal pteridophytes of India and their chemical constituents, J. Econ. Taxon. Bot. 23 (1) (1999) 63–78.
- [221] CSIR, The Wealth of India, A dictionary of Indian raw materials and industrial products, Raw Materials VII, N-Pe, Publications and Information Directorate, Council of Scientific and Industrial Research, New Delhi, 1966, pp. 14–15.
- [222] D. Rani, P.B. Khare, P.K. Dantu, In vitro antibacterial and antifungal properties of aqueous and non-aqueous frond extracts of *Psilotum nudum*, *Nephrolepis biserrata* and *Nephrolepis cordifolia*, Indian J. Pharmaceut. Sci. 72 (6) (2010) 818–822.
- [223] R. Baral, L. Subedi, M. Gurung, S. Ojha, B. Shrestha, D. Chaudhary, N. Jammarkattel, Phytochemical screening, free radical scavenging activity, in-vitro alpha-amylase enzyme and glucose diffusion inhibition activity of ethyl acetate and water extracts of selected medicinal plants of Nepal, Int. J. Herb. Med. 9 (3) (2021) 18–27. <https://www.florajournal.com/archives/2021/vol9Issue3/Parta/8-5-64-832.pdf>.
- [224] R. Pal, In-vitro evaluation of antihelmintic activity of *Nephrolepis cordifolia* leaves, J. Pharm. Exp. Med. 1 (2021) 176.
- [225] A. Rajasekaran, V. Sivakumar, Evaluation of diuretic potential of *Nephrolepis cordifolia* rhizome juice in Wistar rats, Sains Malaysiana 38 (1) (2009) 57–59.

- [226] D.P. Gauchan, D. Manandhar, N. Shrestha, S.K. Suwal, Nutrient analysis of *Nephrolepis cordifolia* (L.) C. Presl, Kathmandu univ. . J. Sci. Eng. Technol. 4 (1) (2008) 68–72.
- [227] A.O. Adebiyi, Phytochemical constituents and proximate composition of *Nephrolepis cordifolia* (L.) C. Presl grown in Nigeria, New York Sci. J. 9 (2) (2016) 79–82. http://www.sciencepub.net/newyork/ny090216/013_30082nys090216_79_82.pdf.
- [228] M.E. El-Tantawy, M.M. Shams, S. Afifi, Chemical composition and biological evaluation of the volatile constituents from the aerial parts of *Nephrolepis exaltata* (L.) and *Nephrolepis cordifolia* (L.) C. Presl grown in Egypt, Nat. Prod. Res. 30 (2016) 1197–1201.
- [229] Z. Liang, X. Yang, H. Zhu, X. Hao, Chemical constituents of *Nephrolepis cordifolia*, *Guang Xi Zhi Wu* 28 (2008) 420–421.
- [230] B.P. Singh, R. Upadhyay, Medicinal pteridophytes of Madhya Pradesh, J. Pharmacogn. Phytochem. 3 (2014) 173–176. https://www.phytojournal.com/vol3Issue3/Issue_sep_2014/19.1.pdf.
- [231] D. Wang, X.Q. Liu, C.S. Yao, W.S. Fang, Studies on chemical constituents from herbs of *Botrychium lanuginosum*, China, J. Chin. Mater. Med. 33 (22) (2008) 2627–2629.
- [232] K. Upreti, J.S. Jalal, L.M. Tewari, G.C. Joshi, Y.P.S. Pangtey, G. Tewari, Ethnomedical uses of pteridophytes of kumaun Himalaya, Uttarakhand, India, J. Am. Sci. 5 (2009) 167–170. http://www.jofamericanmedicine.org/journals/amsci/0504/24_0797_pteridophyte_am0504.pdf.
- [233] CSIR, The Wealth of India, A dictionary of Indian raw materials and industrial products. Raw Materials V, H-K, Council of Scientific and Industrial Research, New Delhi, 1959, p. 31.
- [234] C.C. Chen, Y.L. Huang, P.Y. Yeh, J.C. Ou, Cyclized geranyl stilbenes from the rhizomes of *Helminthostachys zeylanica*, Planta Med. 69 (2003) 964–967.
- [235] Y.L. Huang, P.Y. Yeh, C.C. Shen, C.C. Chen, Antioxidant flavonoids from the rhizomes of *Helminthostachys zeylanica*, Phytochemistry (Oxf.) 64 (2003) 1277–1283.
- [236] T.C. Chang, H. Chiang, Y.H. Lai, Y.L. Huang, H.C. Huang, Y.C. Liang, H.K. Liu, C. Huang, *Helminthostachys zeylanica* alleviates hepatic steatosis and insulin resistance in diet-induced obese mice, BMC Compl. Alternative Med. 19 (2019) 368.
- [237] S.R. Suja, P.G. Latha, P. Pushpangadan, S. Rajasekharan, Evaluation of hepatoprotective effects of *Helminthostachys zeylanica* (L.) Hook against carbon tetrachloride-induced liver damage in Wistar rats, J. Ethnopharmacol. 92 (2004) 61–66.
- [238] H.L. Hsieh, S.H. Yang, T.H. Lee, J.Y. Fang, C.F. Lin, Evaluation of anti-inflammatory effects of *Helminthostachys zeylanica* extracts via inhibiting bradykinin-induced MMP-9 expression in brain astrocytes, Mol. Neurobiol. 53 (2016) 5995–6005.
- [239] C.J. Liou, Y.L. Huang, W.C. Huang, K.W. Yeh, T.Y. Huang, C.F. Lin, Water extract of *Helminthostachys zeylanica* attenuates LPS-induced acute lung injury in mice by modulating NF-κB and MAPK pathways, J. Ethnopharmacol. 199 (2017) 30–38.
- [240] W.C. Huang, N.C. Ting, Y.L. Huang, L.C. Chen, C.F. Lin, C.J. Liou, *Helminthostachys zeylanica* water extract ameliorates airway hyperresponsiveness and eosinophil infiltration by reducing oxidative stress and Th2 cytokine production in a mouse asthma model, Mediat. Inflamm. (2020), 1702935.
- [241] Fitrya, Muhamni, An antihyperuricemia effect of ethanol extract of Tunjuk Langit rhizome (*Helminthostachys Zeylanica* Linn Hook) on Swiss male mice, Tradit. Med. J. 19 (1) (2014) 14–18.
- [242] F.D.L. Anwar, Uji aktivitas antikanker secara in vitro dengan sel murine P-388 senyawa flavonoid Dari fraksi etilasetat akar tumbuhan tunjuk langit (*Helminthostachys zeylanica* (Linn) Hook), J. Penelitian Sains. 12 (1) (2009), 12106.
- [243] M.M. Tsai, H.C. Lin, M.C. Yu, W.J. Lin, M.Y. Chu, C.C. Tsai, C.Y. Cheng, Anticancer effects of *Helminthostachys zeylanica* ethyl acetate extracts on human gastric cancer cells through downregulation of the TNF-α-activated COX-2-cPLA2-PGE2 Pathway, J. Cancer 12 (2021) 7052–7068. <https://www.jcancer.org/v12p7052.htm>.
- [244] F.E. Ridhasya, H.Y. Teruna, R. Hendra, Md Almurdani, Natural antidiabetic of Tunjuk Langit (*Helminthostachys zeylanica*) rhizome extracts, Pharmacol. Clin. Pharm. Res. 4 (3) (2019) 67–70.
- [245] T.W. Yenn, L.C. Ring, K.A. Zahan, Md S.A. Rahman, W.N. Tan, B.J.S. Alaudin, Chemical composition and antimicrobial efficacy of *Helminthostachys zeylanica* against foodborne *Bacillus cereus*, Nat. Prod. Sci. 24 (1) (2018) 66–70.
- [246] Y.C. Huang, T.L. Hwang, C.S. Chang, Y.L. Yang, C.N. Shen, W.Y. Liao, S.C. Chen, C.C. Liaw, Anti-inflammatory flavonoids from the rhizomes of *Helminthostachys zeylanica*, J. Nat. Prod. 72 (2009) 1273–1278.
- [247] Y.C. Huang, T.L. Hwang, Y.L. Yang, S.H. Wu, M.H. Hsu, J.P. Wang, S.C. Chen, L.J. Huang, C.C. Liaw, Acetogenin and prenylated flavonoids from *Helminthostachys zeylanica* with inhibitory activity on superoxide generation and elastase release by neutrophils, Planta Med. 76 (2010) 447–453.
- [248] Y.L. Huang, C.C. Shen, Y.C. Shen, W.F. Chiou, C.C. Chen, Anti-inflammatory and antiosteoporosis flavonoids from the rhizomes of *Helminthostachys zeylanica*, J. Nat. Prod. 80 (2017) 246–253.
- [249] K.C. Wu, C.P. Kao, Y.L. Ho, Y.S. Chang, Quality control of the root and rhizome of *Helminthostachys zeylanica* (Daodi-Ugon) by HPLC using quercetin and ugonins as markers, Molecules 22 (2017) 1115.
- [250] K. Yamauchi, T. Mitsunaga, I. Batubara, Novel quercetin glucosides from *Helminthostachys zeylanica* root and acceleratory activity of melanin biosynthesis, J. Nat. Med. 67 (2013) 369–374.
- [251] L.H. Su, Y.P. Li, H.M. Li, W.F. Dai, D. Liu, L. Cao, R.T. Li, Anti-inflammatory prenylated flavonoids from *Helminthostachys zeylanica*, Chem. Pharm. Bull. 64 (2016) 497–501.
- [252] S. Patil, M. Dongare, The genus *Ophioglossum* from western ghats of India, Indian Fern J. 31 (2014) 17–24.
- [253] P. Parihar, L. Parihar, Some pteridophytes of medicinal importance from Rajasthan, Nat. Product. Radiance 5 (4) (2006) 297–301.
- [254] B.P. Singh, R. Upadhyay, Pteridophytes diversity of Satpura hills, Int. J. Curr. Microbiol. Appl. Sci. 3 (2014) 162–168. https://www.ijcmas.com/vol-3-9/Bale_ndra%20Pratap%20Singh%20and%20Ravi%20Upadhyay.pdf.
- [255] Editorial Committee of the Flora of Taiwan, Flora of Taiwan second ed., 1, 1994, p. 71. Taipei, Taiwan.
- [256] Y.L. Lin, C.C. Shen, Y.J. Huang, Y.Y. Chang, Homoflavonoids from *Ophioglossum petiolatum*, J. Nat. Prod. 68 (2005) 381–384.
- [257] W.J. van der Burg, *Ophioglossum reticulatum* L, in: G.J.H. Gruppen, O.A. Denton (Eds.), Plant Resources of Tropical Africa 2: Vegetables, 2004, pp. 173–175. Prota Foundation, Wageningen, Netherlands.
- [258] R.C. Benedict, What is the habitat of *Ophioglossum vulgatum*, Am. Fern J. 4 (1914) 121–122.
- [259] CSIR, The Wealth of India, A dictionary of Indian raw materials and industrial products VII, N-Pe, Publications and Information Directorate, Council of Scientific and Industrial Research, New Delhi, 1966, pp. 97–98.
- [260] J.Z. Zhao, Treatment of snake bite with *Ophioglossum vulgatum* L. of traditional Chinese medicine, Gansu Chin. Med. 14 (2001) 51–52.
- [261] M. Clericuzio, S. Tinello, B. Burlando, E. Ranzato, S. Martinotti, L. Cornara, A. La Rocca, Flavonoid oligoglycosides from *Ophioglossum vulgatum* L. having wound healing properties, Planta Med. 78 (2012) 1639–1644.
- [262] M. Clericuzio, B. Burlando, G. Gandini, S. Tinello, E. Ranzato, S. Martinotti, L. Cornara, Keratinocyte wound healing activity of galactoglycerolipids from the fern *Ophioglossum vulgatum* L, J. Nat. Med. 68 (2014) 31–37.
- [263] J. Zhang, L. Yang, Y. Fang, J. Li, C. Gu, Q. Zhou, Q. Zheng, W. Chen, Y. Zhou, M. Zhuang, Q. Liu, C. Li, Y. Zhang, Antidiarrheal activity of ethanol extract of *Ophioglossum vulgatum* in mice and spasmyolytic effect on smooth muscle contraction of isolated jejunum in rabbits, Res. Square (2020).
- [264] F. Herrmann, M.R. Romero, A.G. Blazquez, D. Kaufmann, Md L. Ashour, S. Kahl, J.J.G. Marin, T. Efferth, M. Wink, Diversity of pharmacological properties in Chinese and European medicinal plants: cytotoxicity, antiviral and antitrypanosomal screening of 82 herbal drugs, Diversity 3 (2011) 547–580.
- [265] W.C. Hu, Q.Y. Meng, C.X. Wan, Fingerprint analysis of *Ophioglossum* HPLC, Chin. J. Exp. Tradit. Med. Form. 22 (6) (2016) 61–65.
- [266] K.R. Markham, T.J. Mabry, B. Voisin, 3-O-methylquercetin 7-O-diglucoside 4'-O-glucoside from the fern, *Ophioglossum vulgatum*, Phytochem. 8 (1969) 469–472.
- [267] CSIR, The Wealth of India, A dictionary of Indian raw materials and industrial products. Raw Materials VII, N-Pe, Publications and Information Directorate, Council of Scientific and Industrial Research, New Delhi, 1966, pp. 193–194.
- [268] N.R. Kim, L.W. Chi, C.H. Lee, Alpha-glucosidase inhibition activity of methanol extracts obtained from nine pteridophyte species native to Korea, Korean J. Polar Res. 26 (4) (2013) 411–416.
- [269] S. Jain, M. Jacob, L. Walker, B. Tekwani, Screening North American plant extracts in vitro against *Trypanosoma brucei* for discovery of new antitrypanosomal drug leads, BMC Compl. Alternative Med. 16 (2016) 131.
- [270] V. Smitha, V. Vadivel, Phytochemical screening for active compounds in *Ceratopteris thalictroides* (L.) Brogn, J. Pharmacogn. Phytochem. 8 (2019) 3556–3559. <https://www.phytojournal.com/archives/2019/vol8Issue3/PartAZ/8-3-382-884.pdf>.
- [271] M.D. Bhatt, Y.P. Adhikari, R.M. Kunwar, Ethnomedicinal values of weeds in Kanchanpur district, far-western Nepal, Ethnobot. Res. Appl. 21 (2021) 1–19. <https://ethnobotanyjournal.org/index.php/era/article/view/2613>.
- [272] CSIR, The wealth of India: a dictionary of Indian raw materials and industrial products, Raw Materials II (1950) 122. Publications and Information Directorate, Council of Scientific and Industrial Research, New Delhi.
- [273] P.V. Anto, K.V. Greeshma, A.S. Neenu, Antibacterial activity of five selected species of pteridophytes, Int. J. Innov. Res. Dev. 4 (2015). http://www.internationaljournalcorner.com/index.php/ijird_ojs/article/view/135914.
- [274] J.M.A. Antonysamy, P. Ramakrishnan, S. Perumal, T. Shibila, Anti inflammatory activity of selected pteridophytes from western ghats, Int. J. Complement. Alt. Med. 9 (2017) 307.
- [275] V.B. Amoroso, R.A. Mendez, A.P. Villalobos, Bringing back the lost values of Phillipine edible ferns: their antioxidant, proteins and utilization, Int. J. Adv. Res. 5 (2017) 757–770.
- [276] C.Y. Chen, F.U. Chiu, Y. Lin, W.J. Haung, P.S. Hsieh, F.L. Hsu, Chemical constituents analysis and antidiabetic activity validation of four fern species from Taiwan, Int. J. Mol. Sci. 16 (2015) 2497–2516.
- [277] A.O. Adebiyi, S.D. Oyeyemi, P.O. Tedela, V.I. Ojo, Chemical composition of the leaves of *Ceratopteris thalictroides* (L.) Brongn, Scholars Acad. J. Biosci. 7 (2019) 377–386.
- [278] N. Jaishee, U. Chakraborty, Evaluation of in-vitro antioxidant activities of *Pteris biaurita* L, Int. J. Pharm. Pharmaceut. Sci. 6 (S2) (2014) 413–421.
- [279] S.D. Rout, T. Panda, N. Mishra, Ethnomedicinal studies on some pteridophytes of simlipal biosphere reserve, Orissa, India, Int. J. Med. Med. Sci. 1 (2009) 192–197.
- [280] R. Ale, B.B. Raskoti, K. Shrestha, Ethnobotanical knowledge of Magar community in Siluwa VDC, Palpa district, Nepal, J. Nat. Hist. Mus. 24 (2009) 58–71.
- [281] A.K. Dalli, G. Saha, U. Chakraborty, Characterization of antimicrobial compounds from a common fern, *Pteris biaurita*, Indian J. Exp. Biol. 45 (2007) 285–290. <http://hdl.handle.net/123456789/5249>.
- [282] D.H.S. Gracelin, A.J. de Britto, P.B.J.R. Kumar, Qualitative and quantitative analysis of phytochemicals in five *Pteris* species, Int. J. Pharm. Pharmaceut. Sci. 5 (S1) (2013) 105–107.
- [283] C.A. Lans, Ethnomedicines used in Trinidad and Tobago for urinary problems and diabetes mellitus, J. Ethnobiol. Ethnomed. 2 (2006) 45.

- [284] K. Thulsi Rao, R.N. Reddy, C. Pattanaik, C.S. Reddy, Ethnomedicinal importance of pteridophytes used by Chenchus of Nallamalais, Andhra Pradesh, India, *Ethnobot. Leafl.* 11 (2007) 6–10. <https://opensiuc.lib.siu.edu/ebli/vol2007/iss1/2/>.
- [285] M. Singh, R. Govindarajan, A.K.S. Rawat, P.B. Khare, Antimicrobial flavonoid rutin from *Pteris vittata* L. against pathogenic gastrointestinal microflora, *Am. Fern J.* 98 (2008) 98–103.
- [286] K. Nath, M.K. Bhattacharya, S. Ka, Antimicrobial potential of ethnomedicinal ferns of southern Assam, India, *Indian J. Pharmaceut. Sci.* 80 (3) (2018) 556–560.
- [287] L.Q. Ma, K.M. Komar, C. Tu, W. Zhang, Y. Cai, E.D. Kennelly, A fern that hyperaccumulated arsenic, *Nature* 409 (2001) 579–580.
- [288] L.T. Danh, P. Truong, R. Mammucari, N. Foster, A critical review of the arsenic uptake mechanisms and phytoremediation potential of *Pteris vittata*, *Int. J. Phytoremediation* 16 (2014) 429–453.
- [289] M.L.F. Salatino, J. Prado, Flavonoid glycosides of pteridaceae from Brazil, *Biochem. Systemat. Ecol.* 26 (1998) 761–769.
- [290] F. Imperato, The new flavone ester apigenin-7-O-p-hydroxybenzoate and three 3-di-C-glycosylflavones from *Pteris vittata*, *Am. Fern J.* 96 (2006) 62–65.
- [291] K. Joshi, R. Joshi, A.R. Joshi, Indigenous knowledge and uses of medicinal plants in Macchegaun, Nepal, *Indian J. Tradit. Knowl.* 10 (2) (2011) 281–286. <http://noprin.niscain.res.in/handle/123456789/11504>.
- [292] B. Subba, C. Srivastav, R.C. Kandel, Scientific validation of medicinal plants used by Yakkha community of Chanuwa VDC, Dhankuta, Nepal, *SpringerPlus* 5 (2016) 155.
- [293] V.P. Masal, M.M. Dongare, Studies on ethnomedicinal uses of pteridophytes of Ratnagiri district (Maharashtra, India), *Indian Fern J.* 27 (12) (2010) 88–93.
- [294] R.S. Neel, A.S. Jadhao, A.S. Bhuktar, Antibacterial activity of rhizomes and leaf of *Tectaria gemmifera* (Fee.) Alston, *Int. J. Bot. Stud.* 2 (5) (2017) 89–92.
- [295] A.B. Marahatta, B. Poudel, R.C. Basnyat, The phytochemical and nutritional analysis and biological activity of *Tectaria coadunata* Linn, *Int. J. Herb. Med.* 7 (2019) 42–50. <https://www.florajournal.com/archives/?year=2019&vol=7&issue=2&part=A&ArticleId=567>.
- [296] A.P. Thomas, N. Sindhu, S. Dileesh, Comparative phytochemical and antibacterial efficacy of *Tectaria coadunata* and *Tectaria wightii*, *Int. J. Res. Anal. Rev.* 8 (2021) 152–158. <https://ijrar.org/papers/IJRARIcep024.pdf>.
- [297] S.S. Shrestha, S. Sut, S.B. Di Marco, G. Zengin, V. Gandin, M. De Franco, D.R. Pant, Md F. Mahomoodally, S. Dall'Acqua, S. Rajbhandary, Phytochemical fingerprinting and in vitro bioassays of the ethnomedicinal fern *Tectaria coadunata* (J. Smith) C. Christensen from central Nepal, *Molecules* 24 (2019) 4457.
- [298] M.N. Reddy, Md Adnan, M.M. Alreshidi, Md Saeed, M. Patel, Evaluation of anticancer, antibacterial and antioxidant properties of a medicinally treasured fern *Tectaria coadunata* with its phytoconstituents analysis by HR-LCMS, *Anti Cancer Agents Med. Chem.* 20 (2020) 1845–1856.
- [299] D.R. Pant, B. Aryal, D. Pun, S. Sharma, G.P. Joshi, Inhibition of α -amylase and α -glucosidase activities in vitro by extracts of selected medicinal plants, *Biodiversitas* 22 (2021) 1187–1193.
- [300] S.G. Pawar, S.Y. Kamble, S.R. Patil, P.S. Sawant, E.A. Singh, Preliminary phytochemical investigations of three species of traditional medicinal plants of tribal regions of Maharashtra (India), *Int. J. Pharmacogn. Phytochem. Res.* 8 (5) (2016) 742–745. <http://impactfactor.org/PDF/IJPPR/8/IJPPR,Vol8,Issue5,Article6.pdf>.
- [301] J. Rai, K.R. Sharma, Y.R. Pokharel, Antioxidant and alpha amylase inhibitory activity of Nepalese medicinal plants from Gorkha district, *J. Pharmacogn. Phytotherapy* 12 (2) (2020) 28–35.
- [302] K.N. Dubal, P.N. Ghorpade, M.V. Kale, Studies on bioactive compounds of *Tectaria coadunata* (wall. Ex hook. & grev.) C. Chr, *Asian J. Pharmaceut. Clin. Res.* 6 (S2) (2013) 186–187. <https://innovareacademics.in/journal/ajpcr/Vol6Suppl2/1723.pdf>.
- [303] M. Maridass, B. Ravichandran, The screening of phytochemical and antioxidant activity of two endemic ferns, *Mycodium exertum* (Wall. Ex Hook) Copel and *Tectaria Zeilanica* (Holtt.) Sledge, *Pharmacologyonline* 1 (2009) 56–63. <https://pharmacologyonline.silae.it/files/newsletter/2009/vol1/9.Maridass.pdf>.
- [304] S. Sukumaran, M. Mahesh, S. Jeeva, Bioactive constituents of oak leaf fern - *Tectaria zeylanica* (Houtt.) Sledge from southern western ghats, *Asian Pac. J. Trop. Biomed.* 2 (S1) (2012) S64–S66.
- [305] K. Nath, A.D. Talukdar, M.K. Bhattacharya, D. Bhowmik, S. Chettri, D. Choudhury, A. Mitra, A. Bhattacharjee, Antibacterial activity of certain ferns against multi drug resistant organisms, *J. Nat. Remedies* 17 (2017).
- [306] J. Cao, X. Xia, X. Chen, J. Xiao, Q. Wang, Flavonoids contents, antioxidant and anticancer activities of 72 species of ferns from China, *Biomed. Pap.* 156 (S1) (2012) S33. https://biomed.papers.upol.cz/inpdfs/pri-99000-0200_10_026.pdf.
- [307] J. Cao, X. Xia, X. Chen, J. Xiao, Q. Wang, Characterization of flavonoids from *Dryopteris erythrosora* and evaluation of their antioxidant, anticancer and acetylcholinesterase inhibition activities, *Food Chem. Toxicol.* 51 (2013) 242–250.
- [308] M. Jung, M. Park, Acetylcholinesterase inhibition by flavonoids from *Agrimony pilosa*, *Molecules* 12 (2007) 2130–2139.
- [309] F.E. Ridhasya, N. Rahim, Md Almurdani, R. Hendra, H.Y. Teruna, Antidiabetic constituents from *Helminthostachys zeylanica* (L) hook (ophioglossaceae), *pharmacogn.* J 12 (2) (2020) 223–226.
- [310] A.B. Shah, S. Yoon, J.H. Kim, K. Zhumanova, Y.J. Ban, K.W. Lee, K.H. Park, Effectiveness of cyclohexyl functionality in ugonins from *Helminthostachys zeylanica* to PTP1B and α -glucosidase inhibitions, *Int. J. Biol. Macromol.* 165 (2020) 1822–1831.
- [311] C.H. Wu, W.T. Yeh, P.C. Li, Ugonin J, a bioactive compound isolated from *Helminthostachys zeylanica* (L) Hook. (Ophioglossaceae), demonstrates inhibitory effects on neointima formation in the rat carotid artery, *Exp. Biol.* 27 (S1) (2013) 922.
- [312] W.C. Chiou, H.F. Lu, N.Y. Hsu, T.Y. Chang, Y.F. Chin, P.C. Liu, J.M. Lo, Y.B. Wu, J.M. Yang, C. Huang, Ugonin J acts as a SARS-CoV-2 3C-like protease inhibitor and exhibits anti-inflammatory properties, *Front. Pharmacol.* 12 (2021), 720018.
- [313] L.P. Chan, T.H. Chou, G.H. Wang, Y.P. Tseng, P.J. Chen, Da L. Cheng, C.H. Liang, Ugonin K induces cell cycle arrest and apoptosis through reactive oxygen species/apoptosis signal pathway in human skin cancer cells, in: X. Liu, K. Zhang, M. Li (Eds.), *Adv. Mater. Res.* 690–693 (2013) 1422–1425. Trans Tech Publications Ltd.
- [314] Y.C. Lin, Y.C. Huang, S.C. Chen, C.C. Liaw, S.C. Kuo, L.J. Huang, P.W. Gean, Neuroprotective effects of ugonin K on hydrogen peroxide-induced cell death in human neuroblastoma SH-SY5Y cells, *Neurochem. Res.* 34 (2009) 9223–9930.
- [315] C.H. Lee, Y.L. Huang, J.F. Liao, W.F. Chiou, Ugonin K-stimulated osteogenesis involves estrogen receptor-dependent activation of non-classical Src signaling pathway and classical pathway, *Eur. J. Pharmacol.* 676 (2012) 26–33.
- [316] K.C. Wu, Y.L. Ho, Y.H. Kuo, S.S. Huang, G.J. Huang, Y.S. Chang, Hepatoprotective effect of ugonin M, a *Helminthostachys zeylanica* constituent, on acetaminophen-induced acute liver injury in mice, *Molecules* 23 (2018) 2420.
- [317] C.Y. Chen, C.C. Liaw, Y.H. Chen, W.Y. Chang, P.J. Chung, T.L. Hwang, A novel immunomodulatory effect of ugonin U in human neutrophils via stimulation of phospholipase C, *Free Radic. Biol. Med.* 72 (2014) 222–231.
- [318] C.Y. Chen, C.H. Yang, Y.F. Tsai, C.C. Liaw, W.Y. Chang, T.L. Hwang, Ugonin U stimulates NLRP3 inflammasome activation and enhances inflammasome-mediated pathogen clearance, *Redox Biol.* 11 (2017) 263–274.
- [319] H. Kumiko, S. Shiota, T. Hatano, T. Yoshida, T. Kuroda, T. Tsuchiya, Antimicrobial activity of oleanolic acid from *Salvia officinalis* and related compounds on vancomycin-resistant Enterococci (VRE), *Biol. Pharm. Bull.* 30 (6) (2007) 1147–1149.
- [320] X. Wang, H. Bai, X. Zhang, J. Liu, P. Cao, N. Liao, W. Zhang, Z. Wang, C. Hai, Inhibitory effect of oleanolic acid on hepatocellular carcinoma via ERK-p53-mediated cell cycle arrest and mitochondrial-dependent apoptosis, *Carcinogenesis* 34 (2013) 1323–1330.
- [321] V.L. N Reddy, V. Ravikanth, T.P. Rao, P.V. Diwan, Y. Venkateswarlu, A new triterpenoid from the fern *Adiantum lunulatum* and evaluation of antibacterial activity, *Phytochemistry (Oxf.)* 56 (2001) 173–175.