



Laminin-511 Activates the Human Induced Pluripotent Stem Cell Survival via $\alpha 6\beta 1$ Integrin-Fyn-RhoA-ROCK Signaling

Yoshiki Nakashima and Masayoshi Tsukahara

In human induced pluripotent stem cells (hiPSCs), laminin-511/ $\alpha 6\beta 1$ integrin interacts with E-cadherin, an intercellular adhesion molecule, to induce the activation of the phosphatidylinositol 3-kinase (PI3K)-dependent signaling pathway. The interaction between laminin-511/ $\alpha 6\beta 1$ integrin and E-cadherin, an intercellular adhesion molecule, results in protection against apoptosis through the proto-oncogene tyrosine-protein kinase Fyn(Fyn)-RhoA-ROCK signaling pathway and the Ras homolog gene family member A (RhoA)/Rho kinase (ROCK) signaling pathway (the major pathway for cell death). In this article, the impact of laminin-511 on hiPSC on $\alpha 6\beta 1$ integrin-Fyn-RhoA-ROCK signaling is discussed and explored along with validation experiments. *PIK3CA* mRNA (mean [standard deviation {SD}]: iMatrix-511, 1.00 [0.61]; collagen+MFGE8, 0.023 [0.02]; $**P < 0.01$; $n = 6$) and *PIK3R1* mRNA (mean [SD]: iMatrix-511, 1.00 [0.79]; collagen+MFGE8, 0.040 [0.06]; $*P < 0.05$; $n = 6$) were upregulated by iMatrix-511 resulting from an increased expression of *Integrin $\alpha 6$* mRNA (mean [SD]: iMatrix-511, 1.00 [0.42]; collagen+MFGE8, 0.23 [0.05]; $**P < 0.01$; $n = 6$). The iMatrix-511 increased the expression of p120-*Catenin* mRNA (mean [SD]: iMatrix-511, 1.00 [0.71]; collagen+MFGE8, 0.025 [0.03]; $**P < 0.01$; $n = 6$) and *RAC1* mRNA (mean [SD]: iMatrix-511, 1.00 [0.28]; collagen+MFGE8, 0.39 [0.15]; $**P < 0.01$; $n = 6$) by increasing the expression of *E-cadherin* mRNA (mean [SD]: iMatrix-511, 1.00 [0.38]; collagen+MFGE8, 0.16 [0.11]; $**P < 0.01$; $n = 6$). As a result, iMatrix-511 increased the expression of P190 *RhoGAP* (*GTPase-activating proteins*) mRNA, such as *ARHGAP1* mRNA (mean [SD]: iMatrix-511, 1.00 [0.57]; collagen+MFGE8, 0.032 [0.03]; $**P < 0.01$; $n = 6$), *ARHGAP4* mRNA (mean [SD]: iMatrix-511, 1.00 [0.56]; collagen+MFGE8, 0.039 [0.049]; $**P < 0.01$; $n = 6$), and *ARHGAP5* mRNA (mean [SD]: iMatrix-511, 1.00 [0.39]; collagen+MFGE8, 0.063 [0.043]; $**P < 0.01$; $n = 6$). Western blotting showed that phospho-Rac1 remained in the cytoplasm and phospho-Fyn showed nuclear transition in iPSCs cultured on iMatrix-511. Proteome analysis showed that PI3K signaling was enhanced and cytoskeletal actin was activated in iPSCs cultured on iMatrix-511. In conclusion, laminin-511 strongly activated the cell survival by promoting $\alpha 6\beta 1$ integrin-Fyn-RhoA-ROCK signaling in hiPSCs.

Keywords: Laminin-511, human induced pluripotent stem cells, signaling pathway, ROCK inhibitor (Y-27632)

Introduction

IN 2016, A NEW type of induced pluripotent stem cell (iPSC) culture medium (named StemFit) [1] was developed in a joint venture between Professor Yamanaka of Kyoto University (the creator of iPSCs) and Ajinomoto Co., Ltd. The laminin-511 scaffold is capable of binding the $\alpha 3\beta 1$, $\alpha 6\beta 1$,

and $\alpha 6\beta 4$ integrins and maintaining hPSCs in an undifferentiated state in serum-free and xeno-free medium under feeder cell-free culture conditions [2–6]. StemFit medium uses laminin-511 as a scaffold material for cells and is currently the standard medium for culturing human iPSCs (hiPSCs) for medical use in Japan. At the time, we described in detail the intracellular signaling pathways at work in the

Kyoto University Center for iPS Cell Research and Application Foundation (CiRA Foundation), Facility for iPS Cell Therapy (FiT), Kyoto, Japan.

© Yoshiki Nakashima *et al.*, 2022; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons License [CC-BY] (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This article has been updated on September 12, 2022 after first online publication of July 26, 2022 to reflect Open Access, with copyright transferring to the author(s), and a Creative Commons License (CC-BY) added (<http://creativecommons.org/licenses/by/4.0>).

culture conditions of these revolutionary clinical iPSCs [7]. At present, >6 years after its development, no clinical culture material has been developed that surpasses the combination of StemFit and laminin-511 as a scaffold material. In this study, we attempted to identify the effects of laminin-511 scaffolds by examining their effects on signaling pathways.

Laminin-511-based culture systems were found to inhibit cell death through $\alpha6\beta1$ integrin-Fyn-RhoA-ROCK signaling. However, because trypsin (used during cell passaging) transiently cleaves the binding of the laminin-511 scaffold to $\alpha6\beta1$ integrin in the StemFit–laminin-511 culture system and blocks the cell signaling pathway, iPSCs require a localized ROCK inhibitor (Y-27632) during the passaging process. Furthermore, the repair of trypsin-cleaved laminin-511 scaffold binding to $\alpha6\beta1$ integrin requires 24 h after passaging [8]. In this study, we meticulously detail the effects of the laminin-511 scaffold and the $\alpha6\beta1$ integrin-Fyn-RhoA-ROCK signaling pathway on hiPSC culture operations.

Materials and Methods

Reagents

StemFit AK03N was obtained from AJINOMOTO HEALTHY SUPPLY CO., INC. (Tokyo, Japan). The iMatrix-511 was obtained from Matrixome, Inc. (Osaka, Japan). A 10-mM Y-27632 Solution, D-PBS (–), and 0.5 M-EDTA Solution (pH 8.0) were obtained from Nacalai Tesque (Kyoto, Japan). TrypL™ Select Enzyme (1×) was obtained from Thermo Fisher Scientific K.K. (Kanagawa, Japan). Recombinant human MFGE8, Anti-Human E-Cadherin Affinity Purified Polyclonal Ab, and Goat IgG HRP-conjugated Antibody were purchased from R&D Systems (Minneapolis, MN). Anti-beta-actin (C4), Anti-Fyn (D-1), phospho Thr12, Anti-Rock-1 (B-1), and Anti-Rock-2 (D-11) were purchased from Santa Cruz Biotechnology, Inc., (Dallas, TX). GLS250 Gelatin Solution (1.0 mg/g) was purchased from Nitta Gelatin, Inc. (Osaka, Japan). Anti-Integrin $\alpha6$ -CD49f Antibody clone 5I3 ZooMAb(R), Anti-Integrin $\beta1$ Antibody clone 1A27 ZooMAb(R), and Anti-phospho-RAC1 (pSer71) antibody were purchased from Sigma-Aldrich Co. LLC (St. Louis, MO).

Fyn Antibody, p190-B RhoGAP Antibody and p190-A RhoGAP Antibody, Anti-rabbit IgG, HRP-linked Antibody, Anti-mouse IgG, and HRP-linked Antibody were purchased from Cell Signaling Technology, Inc., (Danvers, MA). Anti Rac1, Anti p120 Catenin, and Anti RHOA were purchased from Proteintech Group, Inc., (Rosemont, IL). Anti-PI3-kinase (p85 α) mAb was purchased from Medical & Biological Laboratories Co., Ltd. (Tokyo, Japan). Anti-PI3 Kinase Antibody, p110 α was purchased from Merck & Co., Inc. (Land Hessen, Germany). Goat Anti-Mouse IgM HRP Conjugate was purchased from Tokyo Chemical Industry Co., Ltd. (Tokyo, Japan).

Maintenance culture of human induced pluripotent stem cells

The hiPSC line 201B7 was established by Shinya Yamanaka (CiRA Foundation) and obtained from CiRA Foundation (Kyoto, Japan). To culture iPSCs, a publicly available method (CiRA_Ff-iPSC_protocol_Eng_v140310) was used (<https://www.cira.kyoto-u.ac.jp/j/research/img/>

protocol/Ff-iPSC-culture_protocol_E_v140311.pdf). Collagen coating was performed by adding 200 μ L of GLS250 Gelatin Solution (1.0 mg/g) and 100 μ L (5 μ g) of recombinant human MFGE8 (50 μ g/mL) to one well of a six-well plate.

Real-time PCR

RNA was prepared using a SuperPREP II Cell Lysis & RT Kit for quantitative PCR (TOYOBO CO., LTD., Osaka, Japan) according to the manufacturer's instructions. Real-time PCR was performed using a StepOnePlus system (Life Technologies, Carlsbad, CA). Luna Universal qPCR Master Mix (New England Biolabs, Inc.) was used according to the manufacturer's instructions. For the design of human β -actin, human integrin $\alpha6$, human integrin $\beta1$, and human cadherin 1 (CDH1) primers, the gene names were retrieved from the U.S. National Library of Medicine NIH website. The human β -actin, human integrin $\alpha6$, human integrin $\beta1$, and human cadherin 1 (CDH1) primers were designed using the Primer 3 Plus application. Other primers were purchased from TaKaRa Bio, Inc. (Shiga, Japan). The primers used for PCR were as follows:

human integrin $\alpha6$ (NM_000210.3) 195 bp
(forward) GTTTTGTTTCCTCCCCTATCTGTAT
(reverse) GCTCCCCATATAACTTAACATTGTG

human integrin $\beta1$ (NM_002211.3) 179 bp
(forward) CTGAAGACTATCCCATTGACCTCTA
(reverse) GCTAATGTAAGGCATCACAGTCTTT

human cadherin 1 (CDH1) (NM_004360.4) 191 bp
(forward) GCCACATCTTGACTAGGTATTGTCT
(reverse) GCAGCACTTTAGGCACTATTCTAAG

human β -actin (NM_001101.3) 224 bp
(forward) GTGACATTAAGGAGAAGCTGTGCTA
(reverse) CTTCATGATGGAGTTGAAGGTAGTT

HA348692 (FYN)
(forward) CAATCCGTAGCCAGCTGCTC
(reverse) CCATGGAAGTTCGTCAGCTTCA

HA370128 (PIK3CA)
(forward) TCAGTCCTCAGACCTAATTGGGTTG
(reverse) TCTTGATAMGCTTCCAGGTGAACA

HA377172 (PIK3R1)
(forward) AGCATTGGGACCTCACATTACACA
(reverse) TACTGGAAACACAGTCCATGCACA

HA320879 (CTNND1)
(forward) TGCTCCTGATGATGGTCCTGTC
(reverse) CCTGGTCTCACTAGCCCATGAA

HA356749 (RAC1)
(forward) CCTGTAGTCGCTTTGCCTATTGA
(reverse) AGGGTCCCACGCTGTATTCTC

HA356425 (ARHGAP1)
(forward) CCGAGAGCTACAGCGTGTGA
(reverse) GGGCCAGTGTGACTCCGTAA

HA144198 (ARHGAP4)
(forward) GCAACACGTGGAGGTGGATAA
(reverse) GTGGTAGATGCTGGCCCAAG

HA134083 (ARHGAP5)
(forward) TTTGAGCTGTGGCTAGACATTCTT
(reverse) GACCTCACAGATCCAGACTGAC

HA390884 (RHOA)
(forward) CAGCTGCMGGTACTCTGGTGA
(reverse) CTCTGCCACAGCTGCATGAA

HA271276 (ROCK1P1)
(forward) ACAAATATCACAGGCTTCAGGGTTA
(reverse) TGTAGGCAAACCCGCGATA

HA355736 (ROCK1)
(forward) TGCAACTGGAACCAACCAAGAA
(reverse) GCTGGCCAACTGCATCTGAA

HA359025 (ROCK2)
(forward) GCAAGTCACTGCCGAGCTTC
(reverse) GCTGTCACACAGTGCTTATGTTCA

Western blotting

Western blotting analyses using ATTO Products, EzRIPA Lysis kit, EzSubcell Extract, cPAGE Twin, myPower II 300, HorizeBLOT 2M-R, EzApply, EzStandard PrestainBlue, c-PAGEL 10%, EzRun, P plus membranes, Filter paper, EzFastBlot, EzBlock Chemi, EzTBS, and EzWestBlue, were performed according to the manufacturer's instructions (ATTO, Tokyo, Japan). Blots were probed using specific antibodies for E-Cadherin, integrin $\alpha 6$, integrin $\beta 1$, Fyn, phospho-Fyn, Rac1, phospho-Rac1, p120 catenin, PI3-kinase (p85 α), PI3-kinase (p110 α), p190-A RhoGAP, p190-B RhoGAP, RhoA, ROCK-1, ROCK-2, and β -actin.

Standard data-independent acquisition proteome analyses

The following sample preparation procedures were performed as pretreatment for the proteome analysis: (1) Add chloroform to the sample containing TRIzol, mix, and centrifuge (15,000g, 4°C, 15 min). (2) Remove the aqueous layer, add acetonitrile to the remaining TRIzol solution, and precipitate the protein. (3) Add 100 mM Tris-HCL pH 8.5, 2% SDS to the precipitate, and dissolve the protein using a sealed ultrasonic disruption machine. (4) Measure the protein concentration by a bicinchoninic acid (BCA) assay, adjusted with 100 mM Tris-HCL pH8.5, 2% SDS to make protein concentration 0.5 μ g/ μ L. (5) Cleave the S-S bond of the protein by adding Tris(2-carboxyethyl)phosphine to the protein lysate (20 μ g protein) to a final concentration of 20 mM, and then incubate at 80°C for 10 min. (6) Alkylate cysteine residues by adding iodoacetamide to a final concentration of 30 mM, and then incubate at room temperature (light shielded) for 30 min. (7) Mix Sera-Mag SpeedBead Carboxylate-Modified Magnetic Particles (Hydrophilic) from Cytiva and Sera-Mag Carboxylate-Modified Magnetic Particles (Hydrophobic) at a 1:1 (v/v) ratio and wash three times with distilled water to 15 μ g solids/ μ L in distilled

water (SP3 beads). (8) Put 20 μ L of SP3 beads into the alkylated sample, add another 2.5 times the sample liquid volume of ethanol, and mix at room temperature for 20 min. (9) After washing the beads twice with 80% ethanol, add 100 μ L of 50 mM Tris-HCL pH 8.0 and mix. (10) Add 500 ng of Trypsin/Lys-C Mix (Promega) for peptide fragmentation and incubate at 37°C overnight. (11) Add 20 μ L of 5% trifluoroacetic acid (TFA) and process in a sample-sealing ultrasonic disruption machine. (12) Desalt using a reversed-phase spin column (GL-Tip SDD; GL Sciences) and dry using a centrifugal evaporator. (13) Add 2% acetonitrile (ACN)-0.1% TFA and dissolve the peptides in a sample sealed sonication system. (14) Measure the peptide concentration by a BCA assay and adjust with 2% ACN-0.1% TFA to reach 200 ng/ μ L peptide concentration. (15) Conduct an LC-MS analysis.

The prepared samples were subjected to nanoLC-MS/MS under appropriate analytical conditions, as follows: injected peptide volume, 300 ng; nanoLC used, UltiMate 3000 RSLCnano LC System (Thermo Fisher Scientific); Column size, 75 μ m i.d. \times 120 mm length (Nikkoy Technos); column temperature, 40°C; solvents, Solvent A, distilled water with 0.1% formic acid; Solvent B, 80% ACN with 0.1% formic acid.

The MS data obtained were analyzed using Scaffold data-independent acquisition (DIA) under the following conditions: software program used, Scaffold DIA (Proteome Software); Protein Sequence Database, Human UniProtKB/Swiss-Prot database (Proteome ID UP000005640, 20588 entries, downloaded on November 26, 2021); Spectral Library, created by Prosit (<https://www.proteomicsdb.org/prosit/>) from the above sequence database.

Data analyses were performed in Perseus (<https://maxquant.net/perseus/>). The quantitative values of the proteome analysis results were Log2 transformed and missing values (quantitative values of 0) were randomly assigned at low values that were below the detection limit.

Statistical analyses

Statistical analyses were performed using Student's *t*-test to compare two samples. *P* values of <0.05 were considered statistically significant for all tests.

Institutional review board

This research does not involve human subjects, and does not need to obtain Institutional Review Board (IRB).

Results

Elucidation of the signaling pathways involved in the process of promoting iPSC viability in laminin-511 culture

Laminin-511, a scaffolding component composed of $\alpha 5$, $\beta 1$, and $\gamma 1$ chains [9], binds to integrins on the cell membrane ($\alpha 3\beta 1$ [10], $\alpha 6\beta 1$ [11], and $\alpha 6\beta 4$ [12]), which induces the activation of the phosphatidylinositol 3-kinase (PI3K)/AKT-dependent [13] and Ras/MAPK-dependent signaling pathways [7]. In hiPSCs, the interaction between laminin-511/ $\alpha 6\beta 1$ integrin and E-cadherin (a cell-cell adhesion molecule) provides protection against apoptosis through the proto-oncogene tyrosine-protein kinase Fyn (Fyn)-RhoA-ROCK signaling pathway [7] and the Ras homolog gene family member A




FIG. 1. Fyn-RhoA-ROCK signaling (on laminin-511). In cultures using laminin-511 scaffolds, the E-cadherin- $\alpha 6 \beta 1$ integrin signaling pathway regulates hiPSC cell death. PI3K signaling promotes the expression of Rac1, which binds to cadherin, inhibiting cadherin endocytosis and increasing the expression of cadherin on the plasma membrane. When $\alpha 6 \beta 1$ integrin binds to laminin-511, Fyn-RhoA-ROCK signaling is induced. As a result, ROCK-induced hiPSC cell death is inhibited. This figure is a modified version of an illustration in our previous article [7]. hiPSC, human induced pluripotent stem cells; PI3K, phosphatidylinositol 3-kinase.

(RhoA)/Rho kinase (ROCK) signaling pathway, which is the major pathway for cell death [14] (Fig. 1). An older scaffold material that does not use iMatrix-511 is gelatin. Gelatin was used in the feeder culture method with mouse embryonic fibroblasts (MEFs) [7]. In addition, we have previously reported that MFGE8 (Milk Fat Globule EGF and Factor V/VIII Domain Containing), secreted by MEFs themselves, promotes the cellular adhesion of hiPSCs [15].

Therefore, we chose to use gelatin+MFGE8 as a scaffold material, which is the least likely to affect the intracellular signaling pathways, as a control for iMatrix-511 in this study. As a result, in comparison to iMatrix-511, colonies were observed in gelatin+MFGE8, similarly to the ES cells of iPSCs, but there were fewer colonies. This result indicates that iMatrix-511 has the effect of promoting cell survival activity (Fig. 2).

Laminin receptors (integrin $\alpha 6 \beta 1$ and $\alpha 6 \beta 4$)

PSCs interact with laminin-511 through $\beta 1$ -integrins (predominantly $\alpha 6 \beta 1$). $\alpha 6 \beta 1$ has broad specificity and is capable of binding to a number of laminin isoforms. However, laminin-511 and laminin-521 are the only isoforms that maintain the pluripotency of hPSCs after $\alpha 6 \beta 1$ integrin signaling and thereby induce the PI3K/AKT signaling pathway [16,17]. In hPSCs, laminin-511 prevents the induction of apoptosis by ROCK in vitro. In cultures that utilize a laminin-511 scaffold, the interaction between laminin-511 and $\alpha 6 \beta 1$ integrin therefore partly compensates for Y-27632

(a ROCK inhibitor). A number of studies have indicated that $\alpha 6 \beta 1$ -integrin is highly expressed and/or have described the role of this integrin in PSC adhesion [18–21] and self-renewal [22]. We examined the mRNA levels of integrin $\alpha 6$ and $\beta 1$ expressed by hiPSCs when iMatrix-511 and collagen+MFGE8 were used as scaffolds.

The results showed that the expression of integrin $\alpha 6$ was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [standard deviation {SD}]: iMatrix-511 1.00 [0.42]; collagen+MFGE8, 0.23 [0.047]; $**P < 0.01$; $n = 6$) (Fig. 3A). On the other hand, the mRNA expression of integrin $\beta 1$ was similar when iMatrix-511 and collagen+MFGE8 were used as scaffolds (mean [SD]: iMatrix-511, 1.00 [0.45]; collagen+MFGE8, 1.25 [0.57]; $n = 6$) (Fig. 3B). The $\alpha 6 \beta 1$ integrin on the cell membrane surface may decrease if trypsin is used to harvest cells.

Fyn-p85-PI3K signaling

Fyn, a proto-oncogene tyrosine-protein kinase encoded by the FYN gene [23], is associated with the p85 subunit of PI3K. There are few reports on the role of FYN in hiPSCs. We first examined whether or not hiPSCs express FYN mRNA and confirmed that hiPSC expresses FYN mRNA. The Fyn-RhoA-ROCK signaling cascade [24] has received attention as Fyn is directly associated with $\alpha 6 \beta 1$ -integrin [25,26]. We examined the mRNA levels of FYN expressed by hiPSCs when iMatrix-511 and collagen+MFGE8 were used as scaffolds. The results




FIG. 2. Cell observation. Photographs of hiPSCs taken by optical microscope at 6 days after seeding on each well coated with iMatrix511 (A) or collagen+MFGE8 (B). Scale bar = 500 μ m.




FIG. 3. mRNA expression analysis. (A) The integrin $\alpha 6$ and (B) *integrin $\beta 1$* mRNA expression from hiPSCs on iMatrix511 or collagen+MFGE8 was measured by real-time PCR. Values indicate the relative value obtained by converting the calculated value of iMatrix511 to 1 ($n=6$). The data are presented as the mean \pm SD; standard error. * $P < 0.05$, ** $P < 0.01$. SD, standard deviation.

showed that the expression of FYN was decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.73]; collagen+MFGE8, 0.32 [0.50]; $n=6$) (Fig. 4A).

Previous reports have shown that the PI3K/AKT signaling pathway has an essential role in the survival of iPSCs [27]. PI3K is a heterodimer consisting of catalytic subunits (p110 α , p110 β , and p110 δ) and regulatory subunits (p85 α , p55 α , p50 α , p85 β , and p55 γ). The catalytic subunit p110 α is encoded by PIK3CA, and the regulatory subunit p85 α is encoded by PIK3R1. This means that a decrease in the expression of PIK3CA or PIK3R1 leads to a weakening of the PI3K/AKT signaling pathway. Furthermore, we previously reported that the reduced expression of PIK3CA makes it extremely difficult for hiPSCs to survive [15]. The results showed that the expression of PIK3CA (Phosphatidylinositol-4,5-Bisphosphate 3-Kinase Catalytic Subunit Alpha; also called the p110 α protein), which is a class of PI3K catalytic subunit [28,29], was significantly decreased when colla-

gen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.61]; collagen+MFGE8, 0.024 [0.02]; ** $P < 0.01$; $n=6$) (Fig. 4B).

The results showed that the expression of PIK3R1 (Phosphoinositide-3-Kinase Regulatory Subunit 1) was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.79]; collagen+MFGE8, 0.04 [0.06]; * $P < 0.05$, $n=6$) (Fig. 4C). These results showed that the scaffold material iMatrix-511 enhanced the PI3K/AKT pathway of hiPSCs.

E-cadherin-catenin-Rac1-p120-P190 RhoGAP signaling

Cadherin-1, which is also known as CAM 120/80 or epithelial cadherin (E-cadherin) or uvomorulin, is a protein that is encoded by the CDH1 gene in humans. E-cadherin, a Ca^{2+} -dependent cell-cell adhesion molecule [30,31], is stabilized at the cell surface through its link—through β -catenin and α -catenin—to the cytoskeleton of actin. It is essential for the intercellular adhesion and colony formation of PSCs [32–35]. The loss of this E-cadherin-dependent intercellular adhesion can cause cell death [36]. The results showed that the expression of CDH1 was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.38]; collagen+MFGE8, 0.16 [0.11]; ** $P < 0.01$; $n=6$) (Fig. 5A). The results showed that the scaffold material, iMatrix-511, significantly enhanced the E-cadherin expression of hiPSCs. Endocytosis and recycling regulate the level of E-cadherin at adherens junctions; activated Rac1 reduces E-cadherin endocytosis, thereby increasing the E-cadherin level on the cell surface and consequently promoting cell-cell adhesion.

The ectopic overexpression of E-cadherin also increases the survival of dissociated hPSCs [37]. However, dissociated hPSCs grown on E-cadherin-coated plates form membrane protrusions and show a lower survival rate. Taken together, E-cadherin interactions are not the sole factor affecting cell survival [36]. It appears that transcription factors for the expression of E-cadherin act downstream in various signaling pathways [eg, TGF- β , FGF2, nuclear factor κ B (NF κ B), and integrin cascades] [35].




FIG. 4. mRNA expression analysis. The (A) FYN, (B) PIK3CA, and (C) *PIK3R1* mRNA expression from hiPSCs on iMatrix511 or collagen+MFGE8 was measured by real-time PCR. Indicates the relative value obtained by converting the calculated value of iMatrix511 to 1 ($n=6$). The data are presented as the mean \pm SD; standard error. * $P < 0.05$, ** $P < 0.01$.




FIG. 5. mRNA expression analysis. (A) CDH1, (B) CTNND1, (C) RAC1, (D) ARHGAP1, (E) ARHGAP4, and (F) ARHGAP5 mRNA expression from hiPSC on iMatrix511 or collagen+MFGE8 was measured using real-time PCR. Values indicate the relative value obtained by converting the calculated value of iMatrix511 to 1 ($n=6$). The data are presented as the mean \pm SD: standard error. * $P < 0.05$, ** $P < 0.01$.

The CTNND1 gene provides instructions for making a protein called p120-catenin, also known as delta 1 catenin. E-cadherin-mediated adhesion has been reported to stimulate PI3K/AKT signaling and to have an association with β -catenin signaling [38,39]. The promotion of cell survival by the PI3K/AKT pathway indicates that the Rho family GTPases have a wider role through p120-catenin [40,41]. p120-catenin may also influence cytoskeletal organization in the cytoplasm through the regulation of the opposing activities of Rho and Rac GTPase organization [42]. The results showed that the expression of CTNND1 was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.71]; collagen+MFGE8, 0.025 [0.03]; ** $P < 0.01$; $n=6$) (Fig. 5B). Many of the activities of p120-catenin are associated with increased cellular proliferation and alteration of the cell cycle [43]. As a result, the use of iMatrix-511 as a scaffold material accelerates the increase in cell proliferation through the activation of p120-catenin.

Rho-GTPases, small G-proteins that mediate cellular proliferation and motility [44], are involved in regulating cell survival or death decisions. Functional interaction between Rac1, a small Rho family GTPase, and E-cadherin have been demonstrated to be responsible for the regulation of hPSC self-renewal [45,46]. The results showed that the expression of RAC1 was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.28];

collagen+MFGE8, 0.39 [0.15]; ** $P < 0.01$; $n=6$) (Fig. 5C). According to previous reports, cadherin engagement can inhibit RhoA activity and activate Rac1 [47].

P190 RhoGAP (GTPase-activating protein) was demonstrated to be an important binding partner of p120 [48]. The RhoGAP domain is located at the C-terminal end of p190RhoGAP proteins. Both p190A and B mainly have catalytic activity toward RhoA [49,50]. The inactivation of RhoA by p190 RhoGAP regulates the spreading and migration of cells through the promotion of membrane polarity and membrane protrusion [51]. ARHGAP1, also known as RhoGAP, RhoGAP1, CDC42GAP, and p50rhoGAP, is officially named Ras homology (Rho) GTPase-activating protein 1, which is one of the key members of the RhoGAPs [52]. The results showed that the expression of ARHGAP1 was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.57]; collagen+MFGE8, 0.03 [0.03]; ** $P < 0.01$; $n=6$) (Fig. 5D).

Rho GTPase-activating protein p115 or ARHGAP4 is encoded by the gene ARHGAP4 and is a member of the RHO GTPase-activating proteins (rhoGAP) family of proteins. It has been reported that ARHGAP4 regulates the β -catenin pathway. The results showed that the expression of ARHGAP4 was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.56]; collagen+MFGE8, 0.039 [0.049]; ** $P < 0.01$; $n=6$) (Fig. 5E).

The ARHGAP5 gene provides instructions for making a protein called Rho GTPase-activating protein 5 (Rho-type GTPase-activating protein 5) (p190-B). The results showed that the expression of ARHGAP5 was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.39]; collagen+MFGE8, 0.06 [0.04]; $**P < 0.01$; $n = 6$) (Fig. 5F). These results showed that the expression of p190 RhoGAP, which deactivates RhoA, was significantly higher in hiPSCs cultured with iMatrix-511 as scaffold material.

RhoA-ROCK signaling

RhoA is a 22-kDa small GTPase, a member of the Ras superfamily and Rho subfamily of GTPases. The results showed that the expression of RhoA was significantly decreased when




FIG. 6. mRNA expression analysis. The (A) RHOA, (B) ROCK1P1, (C) ROCK1, and (D) ROCK2 mRNA expression from hiPSCs on iMatrix511 or collagen+MFGE8 was measured by real-time PCR. Values indicate the relative value obtained by converting the calculated value of iMatrix511 to 1 ($n = 6$). The data are presented as the mean \pm SD: standard error. $*P < 0.05$, $**P < 0.01$.

collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.16]; collagen+MFGE8, 0.52 [0.11]; $**P < 0.01$; $n = 6$) (Fig. 6A).

The effect of ROCK1P1 (Rho-associated coiled-coil containing protein kinase 1 pseudogene 1) on hiPSCs is unknown, but its mRNA expression was observed. The results showed that the expression of ROCK1P1 was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.40]; collagen+MFGE8, 0.17 [0.13]; $**P < 0.01$; $n = 6$) (Fig. 6B).

The small molecule Rho-associated kinase (ROCK) proteins consist of two subunits, ROCK1 and ROCK2. Rho kinases (ROCK1 and ROCK2) function downstream of the small GTPase RhoA to drive actomyosin cytoskeletal remodeling [53]. Cell adhesion, cell morphology, and cytoskeletal tension regulate the activation of ROCK by RhoA [54]. The ROCK inhibitor Y-27632, or a combination of ROCK inhibitors, has been shown to enhance hPSC survival after passage as single cells [55,56]. The cell-cell interaction mediated by ROCK inhibitor seems to be associated with the stabilization of the cell surface by E-cadherin. The results showed that the expression of ROCK1 was decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.35]; collagen+MFGE8, 0.59 [0.38]; $n = 6$) (Fig. 6C). The results showed that the ROCK1 mRNA expression of ROCK1 was similar when iMatrix-511 and collagen+MFGE8 were used as scaffolds.

In a prior experiment with $n = 3$ specimens, hiPSCs cultured on the scaffold material collagen+MFGE8 showed significantly higher ROCK1 mRNA expression than hiPSCs cultured on iMatrix-511 (data not shown). In this experiment conducted with $n = 6$ specimens, hiPSCs cultured on iMatrix-511 had their samples collected on day 5, whereas hiPSCs cultured on collagen+MFGE8 had their samples collected on day 8 due to slow cell growth. The expression of ROCK1 mRNA seems to be affected by the timing of sample collection after cell seeding.

The results showed that the expression of ROCK2 was significantly decreased when collagen+MFGE8 was used as a scaffold material in comparison to iMatrix-511 (mean [SD]: iMatrix-511, 1.00 [0.31]; collagen+MFGE8, 0.30 [0.16]; $**P < 0.01$; $n = 6$) (Fig. 6D). Since the ROCK inhibitor Y-27632 inhibits targets of both ROCK1 and ROCK2, it was assumed that the effect of iMatrix-511 would reduce the levels of both ROCK1 and ROCK2, but in fact, when the expression of ROCK2 mRNA was checked, iMatrix-511 was found to promote the expression of ROCK2 mRNA. This result suggests that only ROCK1—and not ROCK2—affects the cell death of hiPSCs.

Western blotting

Western blotting was performed for various factors involved in the Fyn-RhoA-ROCK signaling pathway. Unfortunately, no experimental results are available for integrin $\alpha 6$, ROCK-1, ROCK-2, PI3-kinase (p85 α), P190-A RhoGAP, and P190-B RhoGAP because nonspecific bands were identified.

First, we investigated the mechanism by which iMatrix-511 activates PI3K through integrin $\alpha 6\beta 1$ and Fyn for the




FIG. 7. Western blotting. **(A)** Western blot assay image of E-Cadherin, integrin α 6, integrin β 1, Fyn, phospho-Fyn, Rac1, phospho-Rac1, p120 catenin, PI3-kinase (p85 α), PI3-kinase (p110 α), p190-A RhoGAP, p190-B RhoGAP, RhoA, ROCK-1, ROCK-2, and β -actin bands photosensitively detected on the membrane. **(B)** Proteins of iPSCs cultured on iMatrix-511 were fractionated into four groups (cytoplasm, plasma membrane, nucleus, and cytoskeleton). Western blot assay image of E-Cadherin, integrin α 6, integrin β 1, Fyn, phospho-Fyn, Rac1, phospho-Rac1, p120 catenin, PI3-kinase (p85 α), PI3-kinase (p110 α), p190-A RhoGAP, p190-B RhoGAP, RhoA, ROCK-1, ROCK-2, and β -actin bands photo-sensitively detected on the membrane of each lanes (fractionated into cytoplasm, plasma membrane, nucleus, and cytoskeleton). The samples flowed into each lane (10 μ l) of the membrane.

Fyn-RhoA-ROCK signaling pathway. The integrin β 1 band was detected in whole cell proteins of iPSCs cultured on iMatrix-511 and collagen+MFGE8 (Fig. 7A). When the proteins of iPSCs cultured on iMatrix-511 were fractionated into cytoplasm, plasma membrane, nucleus, and cytoskeleton, the integrin β 1 band was detected in the plasma membrane. A narrow band of phosphorylated Fyn was detected in iPSCs cultured on iMatrix-511 in whole cells. In contrast, no band of phosphorylated Fyn was detected in iPSCs cultured on collagen+MFGE8 in whole cells (Fig. 7A). When proteins were fractionated into cytoplasm, plasma membrane, nucleus, and cytoskeleton, iPSCs cultured on iMatrix-511 showed thin bands of Fyn in the plasma membrane and nucleus, whereas thin bands of phosphorylated Fyn were detected in the plasma membrane, and thick bands were detected in the nucleus (Fig. 7B).

The band of PI3-kinase (p110 α) was detected as a thin band in both iPSCs cultured on iMatrix-511 and all protein samples obtained from iPSCs cultured on collagen+MFGE8 (Fig. 7A). The band of PI3-kinase (p110 α) from iPSCs cultured on iMatrix-511 in whole cells showed a narrow band in the nucleus (Fig. 7B). In conclusion, iMatrix-511 phosphorylates Fyn on the cell surface through integrin α 6 β 1. Phosphorylated Fyn is translocated to the nucleus, and nuclear translocation of PI3K seems to occur together with the phosphorylation of Fyn.

Both E-cadherin and P120 catenin bands were detected in total proteins of iPSCs cultured on iMatrix-511 and collagen+MFGE8 (Fig. 7A). When the proteins of iPSCs cultured on iMatrix-511 were fractionated into four groups (cytoplasm, plasma membrane, nucleus, and cytoskeleton), E-cadherin and P120 catenin were both detected in the plasma membrane and nucleus (Fig. 7B).

Rac1 bands were detected in total proteins of iPSCs cultured on iMatrix-511 and collagen+MFGE8 (Fig. 7A). No bands of phospho-RAC1 were detected in total protein of iPSCs cultured in collagen+MFGE8, and only in total protein of iPSCs cultured with iMatrix-511 was a thin band of phospho-RAC1 detected (Fig. 7A). When the proteins of iPSCs cultured on iMatrix-511 were fractionated into four groups (cytoplasm, plasma membrane, nucleus, and cytoskeleton), Rac1 and phospho-RAC1 were both detected in the cytoplasm. Given the above, it appears that iMatrix-511 promotes phosphorylation of RAC1 in the cytoplasm of iPSCs. In addition, E-cadherin and P120 catenin on the plasma membrane can be activated by phospho-RAC1.

RhoA bands were detected in total proteins of iPSCs cultured on iMatrix-511 and collagen+MFGE8 (Fig. 7A). When the proteins of iPSCs cultured on iMatrix-511 were fractionated into four groups (cytoplasm, plasma membrane, nucleus, and cytoskeleton), RhoA was detected in the membrane (Fig. 7B).





FIG. 8. Standard DIA proteome analysis findings. **(A)** The correlation coefficient (Pearson r) between samples is calculated from the overall quantitative values, and a heatmap is created to cluster and present the differences between samples. **(B)** Proteins with $P < 0.05$ (two groups; T -test) were selected to a Z-score, and a heatmap was varied more than twofold, and the difference between groups were $P < 0.05$ (two groups; T -test). Each quantitative value was converted to a Z-score, and a heatmap was created. **(C)** The results of the GO analysis of increased protein groups in cluster 1 (*upper panel*) or increased protein groups in cluster 2 (*lower panel*), showing the bar graph of the top 10 in the result of Biological Process is shown. **(D)** The results of the Pathway analysis in the increased protein groups in cluster 1 (*upper panel*) or increased protein groups in cluster 2 (*lower panel*), showing the bar graph of the top 10 in the result of Pathway analysis is shown. DIA, data-independent acquisition.

B



D




FIG. 8. (Continued).

Proteome analyses

Proteins were extracted from samples of hiPSCs cultured on iMatrix-511 and collagen+MFGE8 scaffolds, fragmented into peptide fragments by digestive enzymes, and subjected to a DIA analysis by LC-MS. The obtained data were analyzed using the DIA proteome analysis software program (Scaffold DIA) to identify proteins and peptides with both Peptide false discovery rate (FDR) and Protein FDR <1% and to calculate quantitative values. The identification and quantification results from the analysis software program (Scaffold DIA) are shown. Protein quantification values were calculated from the DIA analysis data, with “0” meaning that the protein could not be detected (missing value). Normalization was performed based on the median of the quantitative values (Supplementary Table S1). The quantitative values of this DIA analysis datum were Log2 transformed, and missing values (quantitative values of 0) were randomly assigned at low values that were below the detection limit. A heatmap was created by determining the correlation coefficient (Pearson r) between samples from the overall quantitative values, and differences between samples were clustered (Fig. 8A). Next, proteins were selected that satisfied the following conditions: (1) the mean value of each group varied by a factor of ≥ 2 , and (2) the difference between groups was $P < 0.05$ (two groups: T -test, ≥ 3 groups: analysis of variance).

Each quantitative value was converted to a Z-score, and a heatmap was created (Fig. 8B). The 444 proteins in cluster 1 comprised a group in which the protein expression by iPSCs cultured on iMatrix-511 was higher than that by iPSCs cultured on collagen+MFGE8. A list of the 444 proteins in cluster 1 is shown in Supplementary Table S2. The 649 proteins in cluster 2 comprised group in which the protein expression by iPSCs cultured on collagen+MFGE8 was higher than that by iPSCs cultured on iMatrix-511. A list of the 649 proteins in cluster 2 is shown in Supplementary Table S3.

The protein groups with increased protein quantification in cluster 1 (average protein quantification in cluster 1) increased more than twofold compared with cluster 2, and the difference between the groups was $P < 0.05$ (two groups: T -test). Analyses by GO, Pathway, and an Upstream analysis of GeneXplain were performed. iPSCs cultured on iMatrix-511 showed a greater expression of a group of proteins related to the regulation of the cytoskeleton, mainly actin, than iPSCs cultured on collagen+MFGE8 (Fig. 8C, upper panel) (Supplementary Table S4). iPSCs cultured on iMatrix-511 showed a greater expression of a group of proteins related to the regulation of the PI3K signaling than iPSCs cultured on collagen+MFGE8 (Fig. 8D, upper panel) (Supplementary Table S6). These results indicate that iMatrix-511 activates PI3K signaling, which is reported to be essential for the survival [15] of iPSCs. In addition, the mechanism of actin, a cytoskeletal protein, was shown to be active.

The protein groups with increased protein quantification in cluster 2 (the average protein quantification in cluster 2) increased more than twofold compared with cluster 1 and the difference between the two groups was $P < 0.05$ (two groups: T -test), analyzed by GO and Pathway, and Upstream analysis of GeneXplain were performed. The iPSCs cultured in collagen+MFGE8 showed higher expression of a group of

proteins related to neural differentiation, glycosaminoglycan metabolism, aminoglycan metabolism, RNA splicing, and RNA processing compared with iPSCs cultured in iMatrix-511 (Fig. 8C, lower panel) (Supplementary Table S5). The iPSCs cultured in collagen+MFGE8 showed a higher expression of a group of proteins related to degradation of glycosphingolipids, sialogangliosyl ceramide, the Notch pathway, YAP, cyclin D3, and cyclin D1 than iPSCs cultured in iMatrix-511 (Fig. 8D, lower panel) (Supplementary Table S7).

These results indicate that Notch signaling of iPSCs and YAP are activated in the absence of iMatrix-511. They also indicate that neural developmental mechanisms are active. The iMatrix-511 has been previously reported to inhibit YAP [57]. Notch signaling is known to be involved in many differentiation processes, including neural, hematopoietic, vascular, and somatic differentiation.

Discussion

While xeno-free media, which are alternatives to animal serum, contain no animal-derived components, they may contain human-derived components. A combination of collagen IV, fibronectin, laminin, and vitronectin can be used instead of Matrigel to derive and expand hPSCs under specific culture conditions [58]. However, for culture materials that can be used to produce clinical iPSCs, to give top priority to safety, it is necessary to select materials that conform to the Japanese Standards for Biological Ingredients (JSBI). The JSBI stipulate measures to be taken to ensure the quality, efficacy, and safety of pharmaceuticals, quasi-drugs, cosmetics, medical devices, and regenerative medicine when raw materials used in these products are derived from other living organisms other than humans and plants (<https://www.pmda.go.jp/files/000204341.pdf>).

In 2008, it was reported that the self-renewal ability of mouse ES cells was enabled by the use of laminin-511, but not laminin-332, -111, or -411, as a scaffold material [16]. In 2010, it was reported that long-term self-renewal of human pluripotent cells was enabled by using recombinant protein laminin-511 as a scaffold material [17].

Laminin 511, a recombinant protein from CHO cells, was developed in 2014 as a product (iMatrix-511: Nippi, Tokyo, Japan) that meets the safety standards for culture materials used for clinical cell production under Good Manufacturing Practice manufacturing control and the JSBI by incorporating sterility and virus negation tests into the manufacturing process. The product was developed in 2014. The iMatrix-511 is used as a scaffold material to produce master cell banks, which are the most fundamental cell material for the current production of iPSCs for clinical use [1]. The laminin-511 scaffold can bind the $\alpha 3\beta 1$, $\alpha 6\beta 1$, and $\alpha 6\beta 4$ integrins. In a feeder cell-free culture system, it can be applied to maintain hPSCs in an undifferentiated state under serum and xeno-free conditions [2–6].

Laminin isoforms, laminin-511 and -521, are expressed by human embryonic stem cells (hESCs) [59]. Therefore, before the development of laminin-511 as a scaffold material, ES cell and iPSC researchers tried to prevent cell death by passaging hiPSCs in clumps by physically dissociating the colonies or weakly activating trypsin. By passaging

hiPSCs in clumps, ES cell and iPSC researchers were able to manipulate hiPSCs without cleaving $\alpha 6\beta 1$ integrin-Fyn-RhoA-ROCK signaling.

In addition, it was possible to passage hiPSCs without using ROCK inhibitors when hiPSCs colonies became dense. However, to systematically cultivate high-quality hiPSCs using a culture method that relies on the state of viable cells, ES cell and iPSC researchers needed to be skilled in hiPSC colony mass size and colony density. However, with the development of laminin-511, the difficulty of culturing hiPSCs has been greatly simplified, the stable supply of high-quality hiPSC has become possible, and the hiPSC manufacturing process has reached a technological level that is suitable for industrialization.

Conclusion

Binding of $\alpha 6\beta 1$ with laminin-511 and interaction with E-cadherin stimulates Fyn-RhoA-ROCK signaling to deactivate ROCK signaling, which indicates that the signaling pathways induced by laminin-511 scaffolds are dependent on PI3K signaling through E-cadherin-mediated cell-cell adhesion, and further, demonstrate an association between the Fyn-RhoA-ROCK signaling cascade and $\alpha 6\beta 1$ integrin-laminin-511 adhesion. Therefore, the ROCK inhibitor Y-27632 is unnecessary when the expression of ROCK1 is suppressed by the effect of the scaffold material iMatrix511. However, if the Fyn-RhoA-ROCK signaling in the cells is broken down and ROCK1 is activated by exposure to trypsin, Y-27632 must be added to the culture medium during the hiPSC culture period, until Fyn-RhoA-ROCK signaling is restored.

Author Disclosure Statement

The authors have received no payment for the preparation of this article and state no financial and nonfinancial conflicts of interest.

Funding Information

This research was supported by AMED under Grant Number JP22bm0104001.

Supplementary Material

Supplementary Table S1
Supplementary Table S2
Supplementary Table S3
Supplementary Table S4
Supplementary Table S5
Supplementary Table S6
Supplementary Table S7

References

- Nakagawa M, Y Taniguchi, S Senda, N Takizawa, T Ichisaka, K Asano, A Morizane, D Doi, J Takahashi, et al. (2014). A novel efficient feeder-free culture system for the derivation of human induced pluripotent stem cells. *Sci Rep* 4:3594.
- Gao J, MC DeRouen, CH Chen, M Nguyen, NT Nguyen, H Ido, K Harada, K Sekiguchi, BA Morgan, et al. (2008). Laminin-511 is an epithelial message promoting dermal papilla development and function during early hair morphogenesis. *Genes Dev* 22:2111–2124.
- Hakala H, K Rajala, M Ojala, S Panula, S Areva, M Kellomaki, R Suuronen and H Skottman. (2009). Comparison of biomaterials and extracellular matrices as a culture platform for multiple, independently derived human embryonic stem cell lines. *Tissue Eng Part A* 15:1775–1785.
- International Stem Cell Initiative C, V Akopian, PW Andrews, S Beil, N Benvenisty, J Brehm, M Christie, A Ford, V Fox, et al. (2010). Comparison of defined culture systems for feeder cell free propagation of human embryonic stem cells. *In Vitro Cell Dev Biol Anim* 46:247–258.
- Chin AC, J Padmanabhan, SK Oh and AB Choo. (2010). Defined and serum-free media support undifferentiated human embryonic stem cell growth. *Stem Cells Dev* 19:753–761.
- Rajala K, H Hakala, S Panula, S Aivio, H Pihlajamaki, R Suuronen, O Hovatta and H Skottman. (2007). Testing of nine different xeno-free culture media for human embryonic stem cell cultures. *Hum Reprod* 22:1231–1238.
- Nakashima Y and T Omasa. (2016). What kind of signaling maintains pluripotency and viability in human-induced pluripotent stem cells cultured on Laminin-511 with serum-free medium? *Biores Open Access* 5:84–93.
- Claassen DA, MM Desler and A Rizzino. (2009). ROCK inhibition enhances the recovery and growth of cryopreserved human embryonic stem cells and human induced pluripotent stem cells. *Mol Reprod Dev* 76:722–732.
- Arimori T, N Miyazaki, E Mihara, M Takizawa, Y Taniguchi, C Cabanas, K Sekiguchi and J Takagi. (2021). Structural mechanism of laminin recognition by integrin. *Nat Commun* 12:4012.
- Polisetti N, L Sorokin, N Okumura, N Koizumi, S Kinoshita, FE Kruse and U Schlotzer-Schrehardt. (2017). Laminin-511 and -521-based matrices for efficient ex vivo-expansion of human limbal epithelial progenitor cells. *Sci Rep* 7:5152.
- Takizawa M, T Arimori, Y Taniguchi, Y Kitago, E Yamashita, J Takagi and K Sekiguchi. (2017). Mechanistic basis for the recognition of laminin-511 by $\alpha 6\beta 1$ integrin. *Sci Adv* 3:e1701497.
- Sugawara Y, K Hamada, Y Yamada, J Kumai, M Kanagawa, K Kobayashi, T Toda, Y Negishi, F Katagiri, et al. (2019). Characterization of dystroglycan binding in adhesion of human induced pluripotent stem cells to laminin-511 E8 fragment. *Sci Rep* 9:13037.
- Rodin S, L Antonsson, C Niaudet, OE Simonson, E Salmela, EM Hansson, A Domogatskaya, Z Xiao, P Damdimopoulou, et al. (2014). Clonal culturing of human embryonic stem cells on laminin-521/E-cadherin matrix in defined and xeno-free environment. *Nat Commun* 5:3195.
- Smith AL, MR Dohn, MV Brown and AB Reynolds. (2012). Association of Rho-associated protein kinase 1 with E-cadherin complexes is mediated by p120-catenin. *Mol Biol Cell* 23:99–110.
- Nakashima Y, C Miyagi-Shiohira, H Noguchi and T Omasa. (2018). Atorvastatin inhibits the HIF1 α -PPAR axis, which is essential for maintaining the function of human induced pluripotent stem cells. *Mol Ther* 26:1715–1734.

16. Domogatskaya A, S Rodin, A Boutaud and K Tryggvason. (2008). Laminin-511 but not -332, -111, or -411 enables mouse embryonic stem cell self-renewal in vitro. *Stem Cells* 26:2800–2809.
17. Rodin S, A Domogatskaya, S Strom, EM Hansson, KR Chien, J Inzunza, O Hovatta and K Tryggvason. (2010). Long-term self-renewal of human pluripotent stem cells on human recombinant laminin-511. *Nat Biotechnol* 28:611–615.
18. Hayashi Y, MK Furue, T Okamoto, K Ohnuma, Y Myoishi, Y Fukuhara, T Abe, JD Sato, R Hata and M Asashima. (2007). Integrins regulate mouse embryonic stem cell self-renewal. *Stem Cells* 25:3005–3015.
19. Meng Y, S Eshghi, YJ Li, R Schmidt, DV Schaffer and KE Healy. (2010). Characterization of integrin engagement during defined human embryonic stem cell culture. *FASEB J* 24:1056–1065.
20. Cooper HM, RN Tamura and V Quaranta. (1991). The major laminin receptor of mouse embryonic stem cells is a novel isoform of the alpha 6 beta 1 integrin. *J Cell Biol* 115:843–850.
21. Hierck BP, S Thorsteinsdottir, CM Niessen, E Freund, LV Iperen, A Feyen, F Hogervorst, RE Poelmann, CL Mummery and A Sonnenberg. (1993). Variants of the alpha 6 beta 1 laminin receptor in early murine development: distribution, molecular cloning and chromosomal localization of the mouse integrin alpha 6 subunit. *Cell Adhes Commun* 1:33–53.
22. Cattavarayane S, R Palovuori, J Tanjore Ramanathan and A Manninen. (2015). alpha6beta1- and alphaV-integrins are required for long-term self-renewal of murine embryonic stem cells in the absence of LIF. *BMC Cell Biol* 16:3.
23. Semba K, M Nishizawa, N Miyajima, MC Yoshida, J Sukegawa, Y Yamanashi, M Sasaki, T Yamamoto and K Toyoshima. (1986). yes-related protooncogene, syn, belongs to the protein-tyrosine kinase family. *Proc Natl Acad Sci U S A* 83:5459–5463.
24. Baer AS, YA Syed, SU Kang, D Mitteregger, R Vig, C Ffrench-Constant, RJ Franklin, F Altmann, G Lubec and MR Kotter. (2009). Myelin-mediated inhibition of oligodendrocyte precursor differentiation can be overcome by pharmacological modulation of Fyn-RhoA and protein kinase C signalling. *Brain* 132:465–481.
25. Liang X, NA Draghi and MD Resh. (2004). Signaling from integrins to Fyn to Rho family GTPases regulates morphologic differentiation of oligodendrocytes. *J Neurosci* 24:7140–7149.
26. Colognato H, S Ramachandrapa, IM Olsen and C Ffrench-Constant. (2004). Integrins direct Src family kinases to regulate distinct phases of oligodendrocyte development. *J Cell Biol* 167:365–375.
27. Hossini AM, AS Quast, M Plotz, K Grauel, T Exner, J Kuchler, H Stachelscheid, J Eberle, A Rabien, E Makrantonaki and CC Zouboulis. (2016). PI3K/AKT signaling pathway is essential for survival of induced pluripotent stem cells. *PLoS One* 11:e0154770.
28. Samuels Y, Z Wang, A Bardelli, N Silliman, J Ptak, S Szabo, H Yan, A Gazdar, SM Powell, et al. (2004). High frequency of mutations of the PIK3CA gene in human cancers. *Science* 304:554.
29. Karakas B, KE Bachman and BH Park. (2006). Mutation of the PIK3CA oncogene in human cancers. *Br J Cancer* 94:455–459.
30. Takeichi M. (1995). Morphogenetic roles of classic cadherins. *Curr Opin Cell Biol* 7:619–627.
31. Gumbiner BM. (2005). Regulation of cadherin-mediated adhesion in morphogenesis. *Nat Rev Mol Cell Biol* 6:622–634.
32. Larue L, C Antos, S Butz, O Huber, V Delmas, M Dominis and R Kemler. (1996). A role for cadherins in tissue formation. *Development* 122:3185–3194.
33. Dang SM, S Gerecht-Nir, J Chen, J Itskovitz-Eldor and PW Zandstra. (2004). Controlled, scalable embryonic stem cell differentiation culture. *Stem Cells* 22:275–282.
34. Chen T, D Yuan, B Wei, J Jiang, J Kang, K Ling, Y Gu, J Li, L Xiao and G Pei. (2010). E-cadherin-mediated cell-cell contact is critical for induced pluripotent stem cell generation. *Stem Cells* 28:1315–1325.
35. Soncin F and CM Ward. (2011). The function of e-cadherin in stem cell pluripotency and self-renewal. *Genes (Basel)* 2:229–259.
36. Ohgushi M, M Matsumura, M Eiraku, K Murakami, T Aramaki, A Nishiyama, K Muguruma, T Nakano, H Suga, et al. (2010). Molecular pathway and cell state responsible for dissociation-induced apoptosis in human pluripotent stem cells. *Cell Stem Cell* 7:225–239.
37. Rizzino A. (2010). Stimulating progress in regenerative medicine: improving the cloning and recovery of cryopreserved human pluripotent stem cells with ROCK inhibitors. *Regen Med* 5:799–807.
38. De Santis G, S Miotti, M Mazzi, S Canevari and A Tomassetti. (2009). E-cadherin directly contributes to PI3K/AKT activation by engaging the PI3K-p85 regulatory subunit to adherens junctions of ovarian carcinoma cells. *Oncogene* 28:1206–1217.
39. Pece S, M Chiariello, C Murga and JS Gutkind. (1999). Activation of the protein kinase Akt/PKB by the formation of E-cadherin-mediated cell-cell junctions. Evidence for the association of phosphatidylinositol 3-kinase with the E-cadherin adhesion complex. *J Biol Chem* 274:19347–19351.
40. Goodwin M, EM Kovacs, MA Thoreson, AB Reynolds and AS Yap. (2003). Minimal mutation of the cytoplasmic tail inhibits the ability of E-cadherin to activate Rac but not phosphatidylinositol 3-kinase: direct evidence of a role for cadherin-activated Rac signaling in adhesion and contact formation. *J Biol Chem* 278:20533–20539.
41. Thiery JP. (2003). Cell adhesion in development: a complex signaling network. *Curr Opin Genet Dev* 13:365–371.
42. Wildenberg GA, MR Dohn, RH Carnahan, MA Davis, NA Lobdell, J Settleman and AB Reynolds. (2006). p120-catenin and p190RhoGAP regulate cell-cell adhesion by coordinating antagonism between Rac and Rho. *Cell* 127:1027–1039.
43. McCreia PD and JI Park. (2007). Developmental functions of the P120-catenin sub-family. *Biochim Biophys Acta* 1773:17–33.
44. Cavallaro U and G Christofori. (2004). Cell adhesion and signalling by cadherins and Ig-CAMs in cancer. *Nat Rev Cancer* 4:118–132.
45. Li L, SA Wang, A Jezierski, L Moalim-Nour, K Mohib, RJ Parks, SF Retta and LS Wang. (2010). A Unique interplay between Rap1 and E-cadherin in the endocytic pathway regulates self-renewal of human embryonic stem cells. *Stem Cells* 28:247–257.

46. Ohgushi M and Y Sasai. (2011). Lonely death dance of human pluripotent stem cells: ROCKing between metastable cell states. *Trends Cell Biol* 21:274–282.
47. Noren NK, CM Niessen, BM Gumbiner and K Burridge. (2001). Cadherin engagement regulates Rho family GTPases. *J Biol Chem* 276:33305–33308.
48. Brouns MR, SF Matheson, KQ Hu, I Delalle, VS Caviness, J Silver, RT Bronson and J Settleman. (2000). The adhesion signaling molecule p190 RhoGAP is required for morphogenetic processes in neural development. *Development* 127:4891–4903.
49. Burbelo PD, S Miyamoto, A Utani, S Brill, KM Yamada, A Hall and Y Yamada. (1995). p190-B, a new member of the Rho GAP family, and Rho are induced to cluster after integrin cross-linking. *J Biol Chem* 270:30919–30926.
50. Ridley AJ, AJ Self, F Kasmi, HF Paterson, A Hall, CJ Marshall and C Ellis. (1993). rho family GTPase activating proteins p190, bcr and rhoGAP show distinct specificities in vitro and in vivo. *EMBO J* 12:5151–5160.
51. Arthur WT and K Burridge. (2001). RhoA inactivation by p190RhoGAP regulates cell spreading and migration by promoting membrane protrusion and polarity. *Mol Biol Cell* 12:2711–2720.
52. Li JP, Y Liu and YH Yin. (2017). ARHGAP1 overexpression inhibits proliferation, migration and invasion of C-33A and SiHa cell lines. *Onco Targets Ther* 10:691–701.
53. Priya R, X Liang, JL Teo, K Duszyc, AS Yap and GA Gomez. (2017). ROCK1 but not ROCK2 contributes to RhoA signaling and NMIIA-mediated contractility at the epithelial zonula adherens. *Mol Biol Cell* 28:12–20.
54. Bhadriraju K, M Yang, S Alom Ruiz, D Pirone, J Tan and CS Chen. (2007). Activation of ROCK by RhoA is regulated by cell adhesion, shape, and cytoskeletal tension. *Exp Cell Res* 313:3616–3623.
55. Damoiseaux R, SP Sherman, JA Alva, C Peterson and AD Pyle. (2009). Integrated chemical genomics reveals modifiers of survival in human embryonic stem cells. *Stem Cells* 27:533–542.
56. Watanabe K, M Ueno, D Kamiya, A Nishiyama, M Matsumura, T Wataya, JB Takahashi, S Nishikawa, S Nishikawa, K Muguruma and Y Sasai. (2007). A ROCK inhibitor permits survival of dissociated human embryonic stem cells. *Nat Biotechnol* 25:681–686.
57. Shibata S, R Hayashi, T Okubo, Y Kudo, T Katayama, Y Ishikawa, J Toga, E Yagi, Y Honma, et al. (2018). Selective laminin-directed differentiation of human induced pluripotent stem cells into distinct ocular lineages. *Cell Rep* 25:1668–1679.e5.
58. Ludwig TE, ME Levenstein, JM Jones, WT Berggren, ER Mitchen, JL Frane, LJ Crandall, CA Daigh, KR Conard, et al. (2006). Derivation of human embryonic stem cells in defined conditions. *Nat Biotechnol* 24:185–187.
59. Pook M, I Teino, A Kallas, T Maimets, S Ingerpuu and V Jaks. (2015). Changes in laminin expression pattern during early differentiation of human embryonic stem cells. *PLoS One* 10:e0138346.

Address correspondence to:

*Prof. Masayoshi Tsukahara
Kyoto University Center for iPS Cell Research
and Application Foundation (CiRA Foundation)
Facility for iPS Cell Therapy (FiT)
Kyoto 606-8397
Japan*

E-mail: masayoshi.tsukahara@cira-foundation.or.jp

*Dr. Yoshiki Nakashima
Kyoto University Center for iPS Cell Research
and Application Foundation (CiRA Foundation)
Facility for iPS Cell Therapy (FiT)
Kyoto 606-8397
Japan*

E-mail: yoshiki.nakashima@cira-foundation.or.jp

Received for publication January 13, 2022

Accepted after revision June 17, 2022

Prepublished on Liebert Instant Online June 20, 2022