

AUTHORS' CORRECTIONS

cagA-Positive *Helicobacter pylori* Populations in China and The Netherlands Are Distinct

ARIE VAN DER ENDE, ZHI-JUN PAN, ALDERT BART, RENÉ W. M. VAN DER HULST,
MONIQUE FELLER, SHU-DONG XIAO, GUIDO N. J. TYTGAT, AND JACOB DANKERT

*Departments of Medical Microbiology and Gastroenterology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands, and
Shanghai Institute of Digestive Disease, Shanghai Second Medical University, Shanghai, People's Republic of China*

Volume 66, no. 5, p. 1822–1826, 1998. Page 1823, column 2: The following paragraph should be inserted at the end of Materials and Methods:

Nucleotide sequence accession numbers. The nucleotide sequences of *cagA* and *glmM* have been deposited in the GenBank database under accession no. AJ252963 to AJ252986 and AJ252987 to AJ253010, respectively.

Genomic Analysis Reveals Variation between *Mycobacterium tuberculosis* H37Rv and the Attenuated *M. tuberculosis* H37Ra Strain

ROLAND BROSCH, WOLFGANG J. PHILLIPP, EVANGELOS STAVROPOULOS, M. JOSEPH COLSTON, STEWART T. COLE,
AND STEPHEN V. GORDON

*Unité de Génétique Moléculaire Bactérienne, Institut Pasteur, 75724 Paris Cedex 15, France, and Division of Mycobacterial Research, National
Institute for Medical Research, London NW7 1AA, United Kingdom*

Volume 67, no. 11, p. 5768–5774, 1999. Page 5773, column 2, “Acknowledgments”: Because of an administrative error, the second paragraph was incomplete and should read as follows:

“Financial support for this work was provided by the Wellcome Trust, the Biomed Program of the European Community (grant BMH4/CT97/2277), the Institut Pasteur, and l'Association Française Raoul Follereau. S. V. Gordon was the recipient of a Wellcome Trust International Travelling fellowship.”