Hindawi BioMed Research International Volume 2022, Article ID 9794832, 3 pages https://doi.org/10.1155/2022/9794832

Corrigendum

Corrigendum to "Investigation of Hepatoprotective Activity of Induced Pluripotent Stem Cells in the Mouse Model of Liver Injury"

Chih-Hung Chiang, Ching-Chih Chang, Hui-Chun Huang, Sing Yi-Jen Chen, Sing-Hsing Tsai, Shaw-Yeu Jeng, Shuen-Iu Hung, Jung-Hung Hsieh, Hsu-Shan Huang, Shih-Hwa Chiou, Fa-Yauh Lee, Sing and Shou-Dong Lee, Sing Shih-Hwa Chiou, Sing Fa-Yauh Lee, Sing Shih-Hwa Chiou, Sing Shih-

Correspondence should be addressed to Shih-Hwa Chiou; shchiou@vghtpe.gov.tw and Fa-Yauh Lee; fylee@vghtpe.gov.tw

Received 31 October 2022; Accepted 31 October 2022; Published 26 November 2022

Copyright © 2022 Chih-Hung Chiang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled "Investigation of Hepatoprotective Activity of Induced Pluripotent Stem Cells in the Mouse Model of Liver Injury" [1], a duplication within Figure 1(b) was mistakenly introduced. The authors have stated that this was due to an error when preparing the files for publication and have now repeated the experiment to validate the accuracy and reproducibility of their study.

To take into account the continuity of the hepatocyte differentiation from iPSCs to mature iPSC-derived hepatocytes, the authors have provided new images for Figures 1(a) and 1(b). Additionally, due to the poor resolution of Figure 2 in [1], the authors have provided a revised figure. New Figures 1 and 2 are shown below.

¹Institute of Pharmacology, National Yang-Ming University, No. 155, Section 2, Linong Street, Taipei 11221, Taiwan ²Division of Urology, Department of Surgery, Taipei Veterans General Hospital and Su-Ao & Yuan-Shan Branch, No. 201, Section 2, Shih-Pai Road, Taipei 11217, Taiwan

³School of Medicine, National Yang-Ming University, No. 155, Section 2, Linong Street, Taipei 11221, Taiwan

⁴Division of General Medicine, Taipei Veterans General Hospital, No. 201, Section 2, Shih-Pai Road, Taipei 11217, Taiwan

⁵Division of Gastroenterology, Taipei Veterans General Hospital, No. 201, Section 2, Shih-Pai Road, Taipei 11217, Taiwan

⁶Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, No. 201, Section 2, Shih-Pai Road, Taipei 11217, Taiwan

⁷Graduate Institute of Pharmacy, National Defense Medical Center, No. 161, Section 6, Minquan E. Road, Taipei 11490, Taiwan ⁸Department of Medical Research & Education, Taipei Veterans General Hospital, No. 201, Section 2, Shih-Pai Road, Taipei 11217, Taiwan

2 BioMed Research International




FIGURE 1: In vitro differentiation of iPSCs into iPSC-Heps. iPSCs were seeded at 2×10^4 cells/cm², maintained in Dulbecco's modified Eagle's medium. (a) Left: morphology of iPSC colonies. Right: iPSC colonies were positive for alkaline phosphate stain (purple). (b) The hepatogenic differentiation was induced by a 2-step procedure as described in Section 2. Morphology of undifferentiated and differentiated iPSCs was evaluated at different days after hepatogenic differentiation.

BioMed Research International




FIGURE 2: Immunofluorescence staining for several hepatocyte-specific markers in iPSC-Heps. Immunostaining imaging (800x) results showed that several hepatocyte-specific markers were detected by using (a) anti-AFP antibody and (b) anti-albumin antibody in iPSC-Heps. Blue signal indicated DAPI-positive cells.

References

[1] C.-H. Chiang, C.-C. Chang, H.-C. Huang et al., "Investigation of Hepatoprotective Activity of Induced Pluripotent Stem Cells in the Mouse Model of Liver Injury," *Journal of Biomedicine and Biotechnology*, vol. 2011, Article ID 219060, 11 pages, 2011.