

## Research Article

# Exploration of the Mechanism of Linoleic Acid Metabolism Dysregulation in Metabolic Syndrome

Yan Wen, Yawen Shang , and Qing Wang 

Department of Endocrinology, China-Japan Union Hospital of Jilin University, 126 Xian-tai Street, Changchun, Jilin 130033, China

Correspondence should be addressed to Yawen Shang; 1298602050@qq.com and Qing Wang; wang\_qing@jlu.edu.cn

Received 23 June 2022; Revised 11 October 2022; Accepted 27 October 2022; Published 28 November 2022

Academic Editor: Vasudevan Ramachandran

Copyright © 2022 Yan Wen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We aimed to explore the mechanism of the linoleic acid metabolism in metabolic syndrome (MetS). RNA-seq data for 16 samples with or without MetS from the GSE145412 dataset were collected. Gene set variation analysis (GSVA), gene set enrichment analysis (GSEA), and gene differential expression analysis were performed. Expression data of differentially expressed genes (DEGs) involved in the linoleic acid metabolism pathway were mapped to the pathway by using Pathview for visualization. There were 19 and 10 differentially expressed biological processes in the disease group and healthy group, respectively. 9 KEGG pathways were differentially expressed in the disease group. Linoleic acid metabolism was the only differentially expressed pathway in the healthy group. The GSVA enrichment score of the linoleic acid metabolism pathway in the disease group was markedly lower than that in the healthy group. The GSEA result showed that the linoleic acid metabolism pathway was significantly downregulated in the disease group. JMJD7-PLA2G4B, PLA2G1B, PLA2G2D, CYP2C8, and CYP2J2 involved in the pathway were significantly downregulated in the disease group. This study may provide novel insight into MetS from the point of linoleic acid metabolism dysregulation.

## 1. Introduction

Metabolic syndrome (MetS) is a common complex entity characterized by a set of metabolic abnormalities that serve as risk factors for the development of cardiovascular disease (CVD) and type 2 diabetes mellitus (T2DM) [1]. MetS doubles the risk of CVD and increases the risk of diabetes five-fold [2, 3]. The main characteristic components of MetS include central obesity, hypertension, hyperglycemia, hypertriglyceridemia, and a low level of high-density lipoprotein cholesterol (HDL-C) [1]. The existence of any three or more of these traits constitutes a clinical diagnosis of MetS. The prevalence of MetS ranges from 10% to 40% and keeps rising with the increasing rate of obesity worldwide [4]. The complicated pathogenesis of MetS results from the interaction of multiple biological processes (BPs) and pathways, as well as genetic variations and environmental and nutritional factors; however, the exact underlying mechanisms have not been fully understood [5].

As an essential nutrient, linoleic acid is the most abundant n-6 polyunsaturated fatty acid (PUFA) in the

human diet [6]. PUFAs are the basic components of cellular membranes and serve as cellular signaling molecules. Linoleic acid usually plays an active role in human health. Linoleic acid is correlated with a reduced incidence of CVD and T2DM, which may be attributed to its action on risk factors for diseases, for instance, linoleic acid's reduction of blood cholesterol levels and its impact on insulin and glucose metabolisms [7]. Therefore, linoleic acid may be associated with MetS.

Gene set enrichment methods focus on gene sets in the analysis of gene expression data. Gene set variation analysis (GSVA) calculates gene set enrichment scores for each sample and assesses the variation of gene set enrichment over the samples. It identifies the differential pathway activity in an unsupervised manner [8]. Gene set enrichment analysis (GSEA) utilizes the predefined gene sets and ranks of genes to determine the significant pathways that are related to the phenotypic distinction [9]. Compared with single-gene analysis, gene set enrichment methods have several benefits, including dimensionality reduction and interpretability [10]. In the

present study, we performed GSEA, GSEA, and gene differential expression analysis on RNA-seq data of 16 samples from the GSE145412 expression profile [11]. 16 samples were divided into four groups according to their metabolic health status and body mass index (BMI), including lean (BMI < 25) with the MetS group (MetS lean) and obese (BMI > 30) with the MetS group (MetS obese), the healthy obese group, and the healthy lean group [11]. We found that the enrichment score of the linoleic acid metabolism pathway in the MetS samples was markedly lower than that in the healthy controls, suggesting the dysregulation of the linoleic acid metabolism pathway in MetS, and JMJD7-PLA2G4B, PLA2G1B, PLA2G2D, CYP2C8, and CYP2J2 involved in the pathway were significantly downregulated. This study may provide novel insight into MetS from the point of linoleic acid metabolism dysregulation.

## 2. Materials and Methods

**2.1. Data Collection and Preprocessing.** Raw data from the GSE145412 dataset [11] were downloaded from the Sequence Read Archive (SRA). FASTX-Toolkit was used to trim off the low-quality bases of the raw reads. After quality control by FastQC, clean reads were mapped to the reference genome via TopHat2 (*Homo sapiens*, GRCh38, version 23). Single-mapped reads were utilized to calculate the read count and TPM for each gene. TPM normalization was performed by the `normalizeBetweenArrays` function of the `limma` package [12].

**2.2. GSEA and GSEA.** Gene sets in *Homo sapiens* grouped by KEGG pathways and gene ontology BPs were downloaded from the Molecular Signatures Database (MSigDB) [13] by using the `msigdb` R package. The enrichment scores of each sample on gene sets were computed by the GSEA R package. Differentially expressed gene sets were identified by the `limma` package. The thresholds of differentially expressed BPs were set as  $|\log_2FC| > 0.5$  and  $p < 0.05$ , while those of differentially expressed pathways were set as  $|\log_2FC| > 0.1$  and  $p < 0.05$ . Significantly enriched pathways were identified by the GSEA method with the cutoff values of NOM  $p$  value  $< 0.05$  and FDR  $q$  value  $< 0.25$ .

**2.3. Identification of Differentially Expressed Genes (DEGs).** The gene set related to the linoleic acid metabolism pathway was extracted from MSigDB to construct a protein-protein interaction (PPI) network by using STRING, in which genes with the top 10 highest degrees were regarded as hub genes [14]. DEGs between groups were identified by using the Wilcoxon rank sum test with the threshold as  $p < 0.05$ . Expression data of DEGs were mapped to the linoleic acid metabolism pathway for visualization by using Pathview [15], and proteins encoded by DEGs were colored in the graph to display their roles in the linoleic acid metabolism pathway.

**2.4. Statistical Analysis.** All the statistical analyses were performed by R Statistical Software (version 4.1.2). The Wilcoxon rank sum test was applied to compare the differences between groups.  $p < 0.05$  was statistically significant.

## 3. Results

**3.1. Significantly Downregulated Linoleic Acid Metabolism Pathway in MetS.** 16 samples were divided into four groups, including MetS lean, MetS obese, healthy obese, and healthy lean. Differentially expressed gene sets are shown in Figure S1. There were 19 differentially expressed BPs in the disease group, including regulation of nodal signaling pathways involved in asymmetry, copper ion transmembrane transport, Purkinje myocyte to ventricular cardiac muscle cell signaling, Toll-like receptor 2 signaling pathway, regulation of metanephros development, negative regulation of translational elongation, regulation of endosome size, protein repair, sialic acid transport, the neutrophil-mediated killing of gram-negative bacteria, acylglycerol transport, fructose 2,6-bisphosphate metabolic process, negative regulation of endoplasmic reticulum stress, cellular response to bacterial lipoprotein, sphingomyelin biosynthetic process, response to diacyl bacterial lipopeptide, nucleotide-binding oligomerization domain signaling pathway, regulation of barbed end actin filament capping, and regulation of branching morphogenesis of a nerve (Figure 1(a)). In the healthy group, there were 10 differentially expressed BPs, including lipid hydroxylation, anterior head development, trigeminal nerve development, the somatostatin receptor signaling pathway, histone H3 H4 dimethylation, double-strand break repair involved in meiotic recombination, viral translational termination reinitiation, PML body organization, DNA methylation on cytosine, and epithelial cell proliferation involved in wound healing (Figure 1(a)). On the other hand, differentially expressed pathways in the disease group involved pantothenate and coA biosynthesis, biosynthesis of unsaturated fatty acids, amyotrophic lateral sclerosis, the Toll-like receptor signaling pathway, the P53 signaling pathway, apoptosis, snare interactions in vesicular transport, the TGF beta signaling pathway, and the NOD-like receptor signaling pathway, while in the healthy group, linoleic acid metabolism was the only differentially expressed pathway (Figure 1(b)). The GSEA enrichment score of the linoleic acid metabolism pathway was markedly lower in the disease group than that in the healthy group (Figure 1(c)). The GSEA result revealed that the linoleic acid metabolism pathway was significantly downregulated in the disease group (Figure 1(d)).

We next examined the differences in the GSEA enrichment score of the linoleic acid metabolism pathway between obese and lean samples in the disease group and the healthy group. As depicted in Figure 2, there was no significant difference in the GSEA enrichment score of the linoleic acid metabolism pathway between obese and lean samples in both groups, suggesting that linoleic acid metabolism dysregulation was not significantly associated with obesity.




FIGURE 1: Gene set enrichment analysis. (a) Differentially expressed BPs in the disease group and the healthy group. (b) Differentially expressed pathways in the disease group and the healthy group. (c) Comparison of the GSVAscore of the linoleic acid metabolism pathway between the disease group and the healthy group. (d) GSEA result of the linoleic acid metabolism pathway in the disease group and the healthy group.

29 genes involved in the linoleic acid metabolism pathway were obtained, and the PPI network is shown in Figure S2A. The hub genes with the top 10 highest degrees were ALOX15, CYP2J2, CYP2C8, CYP3A4, CYP2C9, CYP1A2, CYP2C19, PLA2G3, PLA2G12A, and CYP2E1 (Figure S2B). Several genes were found to be significantly downregulated in the linoleic acid metabolism pathway, including JMJD7-PLA2G4B, PLA2G1B, PLA2G2D, CYP2C8, and CYP2J2 (Figure 3(a)), and DEG-encoded proteins participated in the linoleic acid metabolism pathway as enzymes (Figure 3(b)).

#### 4. Discussion

MetS has become a global health concern currently. Except for heart disease and diabetes, MetS also has a link with cancer [16]. The complex interplay of multiple causative factors makes it hard to treat MetS [17]. A healthy lifestyle (including diet and physical activity), weight loss, and control of comorbidities are the therapeutic targets of MetS, and long-term goals include reducing the risks of CVD, T2DM, and other MetS-related conditions [18]. The enhancement of knowledge on mechanisms of MetS




FIGURE 2: Comparison of the GSVA enrichment score of the linoleic acid metabolism pathway between obese and lean samples in the disease group (a) and the healthy group (b).

contributes to improving interventions and treatments for MetS. Fatty acids are a major source of energy. As mediators for cell signaling, they play a role in the etiology of MetS [19].


In the present study, we applied GSVA and GSEA to the RNA-seq data from patients with or without MetS and found that the enrichment score of the linoleic acid metabolism pathway was significantly lower in MetS patients, suggesting the dysregulation of the linoleic acid metabolism pathway in MetS. Through integrated proteomics and metabolomics analysis, Chen et al. also revealed that the linoleic acid metabolism pathway was significantly perturbed in the MetS rat model [20]. Szczuko et al. observed a significant decrease in the content of linoleic acid in MetS compared with the healthy control [21]. Linoleic acid has a correlation with metabolic diseases [22]. Previous studies have revealed that lower linoleic acid content is associated with an increased risk for CVD and T2DM [23, 24]. Yary et al. found that higher linoleic acid concentrations were associated with a lower risk of developing MetS [25].

We also found that several genes involved in the linoleic acid metabolism pathway were significantly downregulated in MetS patients, including JMJD7-PLA2G4B, PLA2G1B, PLA2G2D, CYP2C8, and CYP2J2. PLA2G4B, PLA2G1B, and PLA2G2D encode members of the phospholipase A2 family that generally hydrolyze the sn-2 acyl bond in phospholipids to release fatty acids and lysophospholipids. In the linoleic acid metabolism pathway, lecithin in the cell membrane is converted by phospholipase A2 enzymes to linoleate. PLA2G4B was associated with age-related changes in phospholipids and decreased energy metabolism in monocytes [26]. PLA2G2D increased energy expenditure and thermogenesis by promoting adipocyte browning, thereby improving diet-induced metabolic disorders [27]. PLA2G1B deficiency or inactivation gave protection from diet-induced obesity, insulin resistance, hyperglycemia,


hyperlipidemia, and atherosclerosis [28–30], and PLA2G1B inhibitors suppressed diet-induced obesity and diabetes effectively in mice [31], implying that PLA2G1B inhibition may be a viable therapeutic option for metabolic diseases.

CYP2C8 and CYP2J2 encode members of the cytochrome P450 (CYP) superfamily of enzymes. CYP monooxygenases participate in the metabolism of various endogenous substrates, including PUFAs. In a CYP-dependent manner, linoleic acid is metabolized to produce 9,10-epoxyoctadecenoic acid (9,10-EpOME) and 12,13-epoxyoctadecenoic acid (12,13-EpOME) [32]. CYP2C8 and CYP2J2 are two of the main CYP isoforms that catalyze this conversion. Subsequently, these epoxides are hydrolyzed by soluble epoxide hydrolase (sEH) to yield 9,10-dihydroxyoctadecenoic acid (9,10-DiHOME) and 12,13-dihydroxyoctadecenoic acid (12,13-DiHOME) [32]. These metabolites of CYP produced from linoleic acid are associated with health and disease in many ways [33]. 9,10-EpOME and 9,10-DiHOME induce oxidative stress under 90  $\mu$ M concentration by activating NF- $\kappa$ B and AP-1 transcription factors that mediate inflammation [34]. EpOMEs and DiHOMEs exhibited cardiotoxicity [35, 36], while sEH inhibitors showed a cardioprotective effect [37–39]. In addition, the application of sEH inhibitors in MetS treatment has been explored. In a diet-induced MetS rat model, oral treatment with an sEH inhibitor alleviated the symptoms of MetS, indicating that sEH inhibitors have therapeutic potential for MetS [40].

The study has several limitations. First, the sample size was small. Second, the bioinformatic analysis was restricted to the RNA-seq data of MetS, while proteomics and metabolomics may provide novel insight into the underlying mechanisms of MetS. Third, there is a lack of experiments to detect the linoleic acid metabolism pathway's activity. Therefore, future studies should be conducted with



(a)



Data on KEGG graph  
Rendered by Pathview

(b)

FIGURE 3: Linoleic acid metabolism pathway visualization. (a) Significantly downregulated genes involved in the linoleic acid metabolism pathway. (b) Linoleic acid metabolism pathway visualization with mapped data of DEGs.

a larger sample size and in a more comprehensive way, combining multidimensional omics data analysis and experimental verification to enhance the understanding of MetS.

## 5. Conclusion

In conclusion, we found the dysregulation of the linoleic acid metabolism pathway in MetS, and JMJD7-PLA2G4B, PLA2G1B, PLA2G2D, CYP2C8, and CYP2J2 involved in the linoleic acid metabolism pathway were significantly downregulated.

## Data Availability

The data used and/or analyzed during the current study are available from the corresponding author upon request.

## Conflicts of Interest

The authors declare that there are no conflicts of interest.

## Supplementary Materials

Supplementary figures depict the differentially expressed gene sets (Figure S1) and the PPI network (Figure S2). (*Supplementary Materials*)

## References

- [1] H. H. Wang, D. K. Lee, M. Liu, P. Portincasa, and D. Q.-H. Wang, "Novel insights into the pathogenesis and management of the metabolic syndrome," *Pediatric Gastroenterology, Hepatology and Nutrition*, vol. 23, no. 3, pp. 189–230, 2020.
- [2] E. S. Ford, "Risks for all-cause mortality, cardiovascular disease, and diabetes associated with the metabolic syndrome: a summary of the evidence," *Diabetes Care*, vol. 28, no. 7, pp. 1769–1778, 2005.
- [3] S. Mottillo, K. B. Filion, J. Genest et al., "The metabolic syndrome and cardiovascular risk: a systematic review and meta-analysis," *Journal of the American College of Cardiology*, vol. 56, no. 14, pp. 1113–1132, 2010.
- [4] S. M. Grundy, "Metabolic syndrome update," *Trends in Cardiovascular Medicine*, vol. 26, no. 4, pp. 364–373, 2016.
- [5] U. Zafar, S. Khaliq, H. U. Ahmad, S. Manzoor, and K. P. Lone, "Metabolic syndrome: an update on diagnostic criteria, pathogenesis, and genetic links," *Hormones*, vol. 17, no. 3, pp. 299–313, 2018.
- [6] J. Whelan and K. Fritsche, "Linoleic acid," *Advances in Nutrition*, vol. 4, no. 3, pp. 311–312, 2013.
- [7] F. Marangoni, C. Agostoni, C. Borghi et al., "Dietary linoleic acid and human health: focus on cardiovascular and cardiometabolic effects," *Atherosclerosis*, vol. 292, pp. 90–98, 2020.
- [8] S. Hänzelmann, R. Castelo, and J. Guinney, "GSVA: gene set variation analysis for microarray and RNA-seq data," *BMC Bioinformatics*, vol. 14, no. 1, pp. 7–15, 2013.
- [9] A. Subramanian, P. Tamayo, V. K. Mootha et al., "Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles," *Proceedings of the National Academy of Sciences*, vol. 102, no. 43, pp. 15545–15550, 2005.
- [10] M. Ackermann and K. Strimmer, "A general modular framework for gene set enrichment analysis," *BMC Bioinformatics*, vol. 10, no. 1, pp. 47–20, 2009.
- [11] M. Paczkowska-Abdulsalam, M. Niemira, A. Bielska et al., "Evaluation of transcriptomic regulations behind metabolic syndrome in obese and lean subjects," *International Journal of Molecular Sciences*, vol. 21, no. 4, p. 1455, 2020.
- [12] M. E. Ritchie, B. Phipson, D. Wu et al., "Limma powers differential expression analyses for RNA-sequencing and microarray studies," *Nucleic Acids Research*, vol. 43, no. 7, 2015.
- [13] A. Liberzon, C. Birger, H. Thorvaldsdóttir, M. Ghandi, J. Mesirov, and P. Tamayo, "The molecular signatures database hallmark gene set collection," *Cell Systems*, vol. 1, no. 6, pp. 417–425, 2015.
- [14] D. Szklarczyk, A. Franceschini, M. Kuhn et al., "The STRING database in 2011: functional interaction networks of proteins, globally integrated and scored," *Nucleic Acids Research*, vol. 39, pp. D561–D568, 2010.
- [15] W. Luo, G. Pant, Y. K. Bhavnasi, S. G. Blanchard Jr., and C. Brouwer, "Pathview Web: user friendly pathway visualization and data integration," *Nucleic Acids Research*, vol. 45, no. W1, pp. W501–W508, 2017.
- [16] M. Uzunlulu, O. Telci Kalkili, and A. Oguz, "Association between metabolic syndrome and cancer," *Annals of Nutrition and Metabolism*, vol. 68, no. 3, pp. 173–179, 2016.
- [17] H. Xu, X. Li, H. Adams, K. Kubena, and S. Guo, "Etiology of metabolic syndrome and dietary intervention," *International Journal of Molecular Sciences*, vol. 20, no. 1, p. 128, 2018.
- [18] C. A. Aguilar-Salinas and T. Viveros-Ruiz, "Recent advances in managing/understanding the metabolic syndrome," *F1000Research*, vol. 8, 2019.
- [19] S. J. Wakil and L. A. Abu-Elheiga, "Fatty acid metabolism: target for metabolic syndrome," *Journal of Lipid Research*, vol. 50, pp. S138–S143, 2009.
- [20] M. Chen, Z. Yang, H. Gan, Y. Wang, C. Li, and Y. Gao, "Investigation into potential mechanisms of metabolic syndrome by integrative analysis of metabolomics and proteomics," *PLoS One*, vol. 17, no. 7, Article ID e0270593, 2022.
- [21] M. Szczuko, M. Kaczkan, A. Drozd et al., "Comparison of fatty acid profiles in a group of female patients with chronic kidney diseases (CKD) and metabolic syndrome (MetS)-similar trends of changes, different pathophysiology," *International Journal of Molecular Sciences*, vol. 20, no. 7, p. 1719, 2019.
- [22] B. Choque, D. Catheline, V. Rioux, and P. Legrand, "Linoleic acid: between doubts and certainties," *Biochimie*, vol. 96, pp. 14–21, 2014.
- [23] W. S. Harris, W. C. Poston, and C. K. Haddock, "Tissue n-3 and n-6 fatty acids and risk for coronary heart disease events," *Atherosclerosis*, vol. 193, no. 1, pp. 1–10, 2007.
- [24] S. M. Mousavi, Y. Jalilpiran, E. Karimi et al., "Dietary intake of linoleic acid, its concentrations, and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of prospective cohort studies," *Diabetes Care*, vol. 44, no. 9, pp. 2173–2181, 2021.
- [25] T. Yary, S. Voutilainen, T. P. Tuomainen, A. Ruusunen, T. Nurmi, and J. K. Virtanen, "Omega-6 polyunsaturated fatty acids, serum zinc, delta-5- and delta-6-desaturase activities and incident metabolic syndrome," *Journal of Human Nutrition and Dietetics*, vol. 30, no. 4, pp. 506–514, 2017.
- [26] M. Saare, L. Tserel, L. Haljasmägi et al., "Monocytes present age-related changes in phospholipid concentration and decreased energy metabolism," *Aging Cell*, vol. 19, no. 4, Article ID e13127, 2020.

- [27] H. Sato, Y. Taketomi, Y. Miki, R. Murase, K. Yamamoto, and M. Murakami, "Secreted phospholipase PLA2G2D contributes to metabolic health by mobilizing  $\omega$ 3 polyunsaturated fatty acids in WAT," *Cell Reports*, vol. 31, no. 5, Article ID 107579, 2020.
- [28] K. W. Huggins, A. C. Boileau, and D. Y. Hui, "Protection against diet-induced obesity and obesity-related insulin resistance in Group 1B PLA2-deficient mice," *American Journal of Physiology, Endocrinology and Metabolism*, vol. 283, no. 5, pp. E994–E1001, 2002.
- [29] E. D. Labonté, R. J. Kirby, N. M. Schildmeyer, A. M. Cannon, K. W. Huggins, and D. Y. Hui, "Group 1B Phospholipase A2-mediated lysophospholipid absorption directly contributes to postprandial hyperglycemia," *Diabetes*, vol. 55, no. 4, pp. 935–941, 2006.
- [30] N. I. Hollie, E. S. Konanah, C. Goodin, and D. Y. Hui, "Group 1B phospholipase A2 inactivation suppresses atherosclerosis and metabolic diseases in LDL receptor-deficient mice," *Atherosclerosis*, vol. 234, no. 2, pp. 377–380, 2014.
- [31] D. Hui, M. Cope, E. Labonte et al., "The phospholipase A2 inhibitor methyl indoxam suppresses diet-induced obesity and glucose intolerance in mice," *British Journal of Pharmacology*, vol. 157, no. 7, pp. 1263–1269, 2009.
- [32] J. W. Newman, C. Morisseau, and B. D. Hammock, "Epoxide hydrolases: their roles and interactions with lipid metabolism," *Progress in Lipid Research*, vol. 44, no. 1, pp. 1–51, 2005.
- [33] K. Hildreth, S. D. Kodani, B. D. Hammock, and L. Zhao, "Cytochrome P450-derived linoleic acid metabolites EpOMEs and DiHOMEs: a review of recent studies," *The Journal of Nutritional Biochemistry*, vol. 86, Article ID 108484, 2020.
- [34] S. Viswanathan, B. D. Hammock, J. W. Newman, P. Meerarani, M. Toborek, and B. Hennig, "Involvement of CYP 2C9 in mediating the proinflammatory effects of linoleic acid in vascular endothelial cells," *Journal of the American College of Nutrition*, vol. 22, no. 6, pp. 502–510, 2003.
- [35] A. Fukushima, M. Hayakawa, S. Sugiyama et al., "Cardiovascular effects of leukotoxin (9,10-epoxy-12-octadecenoate) and free fatty acids in dogs," *Cardiovascular Research*, vol. 22, no. 3, pp. 213–218, 1988.
- [36] M. R. Siegfried, N. Aoki, A. M. Lefer, E. M. Elisseou, and R. E. Zipkin, "Direct cardiovascular actions of two metabolites of linoleic acid," *Life Sciences*, vol. 46, no. 6, pp. 427–433, 1990.
- [37] O. Islam, P. Patil, S. K. Goswami et al., "Inhibitors of soluble epoxide hydrolase minimize ischemia-reperfusion-induced cardiac damage in normal, hypertensive, and diabetic rats," *Cardiovascular Therapeutics*, vol. 35, no. 3, Article ID e12259, 2017.
- [38] P. Sirish, N. Li, J.-Y. Liu et al., "Unique mechanistic insights into the beneficial effects of soluble epoxide hydrolase inhibitors in the prevention of cardiac fibrosis," *Proceedings of the National Academy of Sciences*, vol. 110, no. 14, pp. 5618–5623, 2013.
- [39] D. Xu, N. Li, Y. He et al., "Prevention and reversal of cardiac hypertrophy by soluble epoxide hydrolase inhibitors," *Proceedings of the National Academy of Sciences*, vol. 103, no. 49, pp. 18733–18738, 2006.
- [40] A. Iyer, K. Kauter, M. A. Alam et al., "Pharmacological inhibition of soluble epoxide hydrolase ameliorates diet-induced metabolic syndrome in rats," *Experimental Diabetes Research*, vol. 2012, Article ID 758614, 11 pages, 2012.