




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Brief Communication



Long-term persistence of IgG antibodies in recovered COVID-19 individuals at 18 months post-infection and the impact of two-dose BNT162b2 (Pfizer-BioNTech) mRNA vaccination on the antibody response: Analysis using fixed-effects linear regression model

Puya Dehgani-Mobaraki^{a,*}, Chao Wang^b, Alessandro Floridi^c, Emanuela Floridi^c, Sunny Dawoodi^d, Asiya K. Zaidi^e

^a Associazione Naso Sano, Umbria Regional Registry of Volunteer Activities, Via Luca Benincasa 2, San Mariano, 06073, Perugia, Italy

^b Health & Social Care Statistic, Faculty of Health, Social Care and Education, Kingston University and St George's, University of London, London, SW17 0RE, UK

^c Centro Ricerche Analisi Biochimico Specialistiche, Perugia, Italy

^d RIPAS Hospital, Brunei

^e Associazione Naso Sano, Umbria Regional Registry of Volunteer Activities and Research, San Mariano, Italy

A B S T R A C T

This era of emerging variants needs a thorough evaluation of data on the long-term efficacy of immune responses in vaccinated as well as recovered individuals, to understand the overall evolution of the pandemic. In this study, we aimed to assess the dynamics of IgG response over 18 months in $n = 36$ patients from the Umbria region in Italy, who had a documented history of COVID-19 infection in March 2020, and then compared the impact of two-dose BNT162b2 (Pfizer-BioNTech) vaccination on the antibody responses of these patients with the ones who did not receive any dose of vaccine. This is the longest observation (March 2020–September 2021) for the presence of antibodies against SARS-CoV-2 in recovered individuals along with the impact of 2 dose-BNT162b2 vaccination on these responses. Fixed-effect regression models were used for statistical analysis which could be also used to predict future titer trends. At 18 months, 97% participants tested positive for anti-NCP hinting towards the persistence of infection-induced immunity even for the vaccinated individuals. Our study findings demonstrate that while double dose vaccination boosted the IgG levels in recovered individuals 161 times, this “boost” was relatively short-lived. The unvaccinated recovered individuals, in contrast, continued to show a steady decline but detectable antibody levels. Further studies are required to re-evaluate the timing and dose regimen of vaccines for an adequate immune response in recovered individuals.

1. Introduction

This era of emerging variants needs a thorough evaluation of data on the long-term efficacy of immune responses in vaccinated as well as recovered individuals, to understand the overall pandemic evolution. Vaccine equity is an evolving global challenge that needs prompt alternate strategies for equal allocation across all countries in line with WHO's campaign for Vaccine Equity (Who.int, 2021). Recent data has suggested post-infection immunity to be durable, long-acting, and protective. (Gaebler et al., 2021; Abu-Raddad et al., 2021; Anand et al., 2021; Egbert et al., 2021; Turner et al., 2021; Gallais et al., 2021; Dehgani-Mobaraki et al., 2021). However, recovered individuals have also been advised to receive complete vaccination similar to naïve individuals with no previous history of COVID-19 infection. Whether two

doses are required in recovered individuals or a single dose could offer adequate protection, is still a matter of debate. There are countries that have given three/four doses of vaccines to all individuals (recovered/naïve).

Real-time data indicates that recovered individuals have protective immunity that could be sufficiently enhanced by a single dose of the vaccine rather than a double dose. Unfortunately, neither recall responses nor ideal vaccine dosing regimens have been studied in them (Saadat et al., 2021; Ebinger et al., 2021). When compared to natural infection, vaccination elicits a more specific response which is greater in magnitude and is largely focused on the spike-receptor binding domain (S-RBD) rather than Nucleocapsid (NCP). Recent data suggests impaired protection, predominantly viral neutralization, and vaccine effectiveness against the variants of concern (Gregory Hames et al., 2021).

* Corresponding author. Associazione Naso Sano, Education & Research, Italy.

E-mail addresses: dehganipuya@gmail.com (P. Dehgani-Mobaraki), C.Wang@sgul.kingston.ac.uk (C. Wang), direzione@crabion.it (A. Floridi), emanuela@crabion.it (E. Floridi), sunnydawoodi@gmail.com (S. Dawoodi), asiyazaidia@gmail.com (A.K. Zaidi).

<https://doi.org/10.1016/j.virol.2022.12.003>

Received 2 September 2022; Received in revised form 21 November 2022; Accepted 2 December 2022

Available online 6 December 2022

0042-6822/© 2022 Elsevier Inc. All rights reserved.

In this study, we aimed to assess the dynamics of IgG response over 18 months in patients who had a documented history of COVID-19 infection in March 2020, and then compared the impact of two-dose BNT162b2 (Pfizer-BioNTech) vaccination on the antibody responses of these patients with the ones who did not receive any dose of vaccine (NCT05038475). (*Clinical and Immunological Responses After*).

2. Materials and methods

2.1. Patient cohort

A monocentric longitudinal observational study was conducted in patients based in the Umbria region of Italy who had tested positive for SARS-CoV-2 in March 2020 by RT-qPCR, aimed to analyze the presence of antibodies against SARS-CoV-2. These RT-PCR tests were performed by the Local health regulatory authorities according to the national guidelines and standard operating protocols. The patients were managed as per the set protocols by the treating doctor, based on the severity of the disease. On recovery, all subjects were informed about this study for seroprevalence and were invited for voluntary participation. After detailed written informed consent, serological samples were collected at timed intervals and antibody levels were analyzed using the MAGLUMI® 2019-nCoV IgM/IgG chemiluminescent analytical system (CLIA) assay and the MAGLUMI® SARS-CoV-2 S-RBD IgG CLIA. (New Industries Biomedical Engineering Co., Ltd [Snibe], Shenzhen, China). Both these immunoassays were granted Emergency Use Authorization by the US Food and Drug Administration, anti-NCP during the initial months of the pandemic followed by anti-S-RBD later in 2020 (MAGLUMI. nCoV IgM, 2019).

Using a standardized questionnaire, the participants were asked to provide information about their COVID-19 clinical history along with symptoms and treatment undertaken. Although reminders were sent for follow-up periodically, participation in the study was completely voluntary. The study participants did not receive any compensation or any other benefit but were informed individually about their antibody status.

The blood samples were collected after informed consent by the patients and with the approval of the ethics committee of the Associazione Naso Sano (Document number ANS-2020/001) at an accredited lab (Laboratory of Nuclear Lipid BioPathology, CRABION, Perugia, Italy). Data collection and analysis were masked from the main principal investigator, who was also a part of the study sample to avoid observer bias (World Health Organization, 2020; Xydakis et al., 2020). The study was conducted in accordance with the Declaration of Helsinki and national and institutional standards.

3. Patient selection

Study timeline and assays used: The antibody levels were analyzed in two phases, based on the availability of immunoassays at the time, i.e. from May 2020 to January 2021, antibodies against Nucleocapsid (NCP) were analyzed using the MAGLUMI® 2019-nCoV IgM/IgG CLIA assay through sequential serum samples. Time was treated as a factor and six different time points were defined; (T0-T5). The first blood sample was collected in May 2020, 2 months after infection (march), and was defined as T0. Consecutive serological samples were analyzed at three months (T1), five months (T2), seven months (T3), eight months (T4) ten months (T5) post-infection in June, August, October, November of 2020, and January 2021 respectively.

The second phase comprised of the introduction of a different immunoassay; MAGLUMI® SARS-CoV-2 S-RBD IgG CLIA to detect antibodies against Spike-Receptor Binding Domain from Feb 2021 to September 2021.

Analysis using both the immunoassays (anti-NCP and anti-S-RBD) for each patient of the study group could not be done throughout 18 months due to practical challenges associated with it. Italy was one of the first

countries to be affected by COVID-19 in early 2020 and the anti-S-RBD immunoassay received emergency approval only later in 2020.

From the original cohort, in $n = 24$ individuals, anti-NCP was analyzed from May 2020 to January 2021 and then anti-S-RBD was analyzed from February 2021 to September 2021. For $n = 12$ (8 female and 4 male) participants, who met the eligibility criteria for participation (documented history of COVID-19 in March 2020) that were enrolled in this study in Feb 2021, only anti-S-RBD could be analyzed from February 2021 to September 2021 making the final sample size of $n = 36$ participants.

4. Introduction of the vaccinated subgroup

Since the legal provisions adopted by the Italian Ministry of Health advised vaccination, irrespective of previous disease status, $n = 21$ participants were gradually vaccinated from mid-March 2021. Out of these, $n = 19$ participants continued to follow up voluntarily for antibody titer assessment at monthly intervals.

Therefore at 18 months, the study participants were divided into two groups:

Group A: Recovered and received two doses of BNT162b2 vaccine ($n = 21$). From this recovered vaccinated group, $n = 2$ (P23 and P30 in the figure) failed to follow up post-vaccination making the final sample size of $n = 19$.

Group B: Recovered individuals who were unvaccinated ($n = 15$).

For the month of September 2021 (18 months post-infection), the immunoassay to detect anti-NCP was reintroduced so that even vaccinated individuals could be analyzed for the persistence of infection-induced immunity not generated by mRNA spike-based vaccination. Only qualitative assessment could be done for the presence of anti-NCP antibodies at this point.

Although the study sample is small, the presented data can be useful in providing information on the longer-term dynamics of relevant antibodies since the beginning of the pandemic and the impact of vaccination on antibody responses.

5. Analytical system used

For analysis at 18 months, the study participants were divided into two groups based on their vaccination status.

Group A: recovered and received two doses of BNT162b2 vaccine ($n = 19$) and Group B: recovered individuals who were unvaccinated ($n = 15$).

As per the Specifications, the anti-SARS-CoV-2 S-RBD IgG assay had a sensitivity of 100% with CI [99.9%–100.0%] at ≥ 15 days post symptom onset, and specificity of 99.6%; CI [98.7%–100.0%]. High-concentration samples were diluted automatically by analyzers and the recommended dilution was 1:9 with the diluent in the kit. The sample, buffer, and magnetic microbeads coated with S-RBD recombinant antigen were mixed thoroughly and incubated, forming immune complexes. After precipitation, decanting of supernatant, and performing a wash cycle, ABEI labeled with anti-human IgG antibody was added, and incubated to form complexes. Again after precipitation in a magnetic field, decanting of supernatant, and performing another wash cycle, the Starter 1 + 2 were added to initiate a chemiluminescent reaction. The light signal was measured by a photomultiplier as relative light units (RLUs), which is proportional to the concentration of SARS-CoV-2 S-RBD IgG presented in the sample. The measurements and interpretation of results were made according to the manufacturer's instructions. The analyzer automatically calculates the concentration in each sample using a calibration curve which is generated by a 2-point calibration master curve procedure. The results were expressed in AU/mL. A result less than 1.00 AU/mL (< 1.00 AU/mL) was considered to be non-reactive while a result greater than or equal to 1.00 AU/mL (≥ 1.00 AU/mL) was considered to be reactive (<https://www.fda>.

gov/medical-devices/coronavir-us-disease-2019-covid-19-emergency-use-authorizations-medical-devices/vitro-diagnostics-euas., 2019; https://www.snibe.com/zh_en/en_newsView.aspx?id = 647).

6. Statistical analysis

To model how antibody response evolves over time, we used a fixed-effects linear regression while adjusting for the testing method/immunoassay used (NCP or S-RBD), vaccination status, and temporal effects. To model the non-linear effects of time, a restricted cubic spline function for time (month) was applied in the regression model using the method by Orsini and Greenland (2011) to visualize and interpret the results (Orsini and Greenland, 2011). To account for serial correlation, the error term is modeled as a first-order autoregression (AR) process.

7. Results

Fig. 1 shows how the IgM and IgG levels have evolved over time for the recovered subjects and the impact of two-dose BNT162b2 (Pfizer-BioNTech) vaccination on them.

Fig. 1(a) shows the IgM response (red) over 18 months in each subject as per the patient number that was assigned randomly. The blue vertical line represents the time of vaccination. Fig. 1(b) shows the IgG response (log scale) (red) over 18 months in each subject as per the





Fig. 1. Fig. 1 shows how the IgM and IgG levels have evolved over 18 months for n = 21 of Group A: recovered subjects and the impact of two-dose BNT162b2 (Pfizer-BioNTech) vaccination on the antibody responses, where Fig. 1(a) shows the IgM titer trend (red) and Fig. 1(b) shows the IgG titer trend (red line) over 18 months in each subject as per the patient number that was assigned randomly. The blue vertical line represents the time of vaccination. From this recovered vaccinated group, n = 2 (P23 and P30 in the figure) failed to follow up post-vaccination.

patient number that was assigned randomly. The blue vertical line represents the time of vaccination. The graphs demonstrate that vaccination has an enormous impact on antibody responses. It was interesting to note that, although there was a rapid and steep rise in the antibody responses immediately post-vaccination, this rise was short-lived. This explains the role of boosters that might, perhaps “boost” the titer levels for a brief duration. Whether the titer trend after vaccination drops back to pre-vaccination levels in a few weeks or in fact, lowers below the baseline needs further analysis on a larger population.

Fig. 2(a) visualizes the modeling results for the four scenarios, based on immunoassay employed (anti-NCP or anti-S-RBD) and completion of two doses of BNT162b2 (Pfizer–BioNTech) mRNA vaccine: 1) unvaccinated and tested against NCP; 2) vaccinated and tested against NCP; 3) unvaccinated and tested against S-RBD; and 4) vaccinated and tested against S-RBD. IgG values were log-transformed to avoid negative values.

Fig. 2(b) describes the differences in antibody responses between a given month and the baseline month (month 2), holding all the other variables (vaccination status and testing method) constant. This allows visualization of antibody response dynamics regardless of other factors since this model assumes an independent temporal effect. Statistical models (especially *Fixed-effect* regression models) are flexible to *diminish* the *impact* of confounding by (unmeasured) fixed/time-invariant factors.

Fig. 2(c) describes the modeling results to determine the effect of vaccination on antibody responses (IgM and IgG) in recovered subjects. IgM responses against NCP are lower by an amount of 0.324, relative to the response against spike-RBD. Similarly, IgG response against NCP is lower than that against spike-RBD by a factor of 0.119.

A double-dose vaccination with BNT162b2 in recovered individuals induced an enormous but short-lived increase in antibody levels. While the IgM response increased by an amount of 0.773, the IgG response increased 161 times.

However, we do agree that a caveat should be noted that the model assumed that anti-S-RBD responses only affect the level but not the trend/shape of the curve relative to anti-NCP. Further analysis of the interaction between time and type of antibody is not statistically significant which supports our original model.

It was interesting to note that all participants from Group B (n = 15) tested positive for anti-S-RBD for 18 months post-infection, with no case reported of reinfection despite multiple waves of mutant strains encountered in the region since March 2020. Also, 33 out of the 34 participants (97%) tested positive for anti-NCP at 18 months. We hypothesize that anti-NCP immunoassays could be a valuable tool for the assessment of persisting infection-induced immunity and should be re-adopted for analysis even for vaccinated individuals. In terms of strain, whether the infection with the wild-type conferred a more robust protective immunity against re-infection as compared to infection with other mutant strains needs further large-scale analysis.

8. Discussion

In this study, we longitudinally analyzed the antibody responses in patients that recovered from COVID -19, 18 months after SARS-CoV-2 infection (wild type), and divided them into two groups for comparison based on vaccination with two doses of BNT162b2. Group A: Recovered and received two doses of the BNT162b2 vaccine (n = 21). From this recovered vaccinated group, n = 2 (P23 and P30 in the figure) failed to follow up post-vaccination making the final sample size of n = 19, and Group B: Recovered individuals who were unvaccinated (n = 15).

To our knowledge, at the time of writing this manuscript, this is one of the longest observational studies that report the persistency of antibodies against SARS-CoV-2 at 18 months and the impact of 2 doses of BNT162b2 vaccine in seropositive convalescent patients.

Our observations have been in line with previously published studies suggesting the presence and persistence of long-term protective




Fig. 2. Fig. 2(a) visualizes the modeling results for the four scenarios, based on immunoassay employed (anti-NCP or anti-S-RBD) and completion of two doses of BNT162b2 (Pfizer-BioNTech) mRNA vaccine: 1) unvaccinated and tested against NCP; 2) vaccinated and tested against NCP; 3) unvaccinated and tested against S-RBD; and 4) vaccinated and tested against S-RBD. IgG levels were log-transformed to avoid negative values. Fig. 2(b) describes the differences in antibody responses between a given month and the baseline month (month 2), holding all the other variables (vaccination status and testing method) constant and Fig. 2(c) describes the modeling results to determine the effect of vaccination on antibody levels (IgM and IgG) in recovered subjects. The IgM levels against NCP are lower, by an amount of 0.324, relative to spike-RBD IgM levels. Similarly, IgG levels against NCP are lower than spike-RBD IgG levels by a factor of 0.119. The vaccines lead to a large increase in antibody responses: vaccination increases IgM by amount of 0.773, and IgG by 161 times.

immunity after SARS-CoV-2 infection. A study by Gaebler et al. concluded that “memory B cell response to SARS-CoV-2 evolves between 1.3 and 6.2 months after infection in a manner that is consistent with antigen persistence” (Gaebler et al., 2021). The famous Qatar study by Abu-Raddad LJ et al., observed the estimated efficacy of immunity against reinfection to be 95% over 7 months (Abu-Raddad et al., 2021). This finding was similar to the findings by Anand et al. that anti-S-RBD IgM in plasma decayed rapidly, whereas the reduction of IgG was less prominent up to 8 months (Anand et al., 2021). Thereafter, a study was conducted on 3015 healthcare workers by Egbert ER. et al. demonstrating spike antibodies to SARS-CoV-2 to be durable for up to 10 months after natural infection (Egbert et al., 2021). Later, Turner et al. (n = 77, mild infection) observed that anti-S-IgG was detectable at 11 months. This could be attributed to the fact that S-binding bone marrow plasma cells (BMPCs) are quiescent, indicating that they are part of a long-lived compartment (Turner et al., 2021). When it comes to reduction in risk of infection, these antibodies might offer protection for up to 13 months (Gallais et al., 2021). Our previous study for infection-acquired immunity against SARS-CoV-2 at 14 months has been published (Dehghani-Mobaraki et al., 2021).

Vaccination-induced immunity in individuals with no history of previous infection has been relatively short-lived (half-life of 69–173 days) with requirements of boosters to augment antibody responses against subsequent variants of concern, especially the Omicron variant (Levin et al., 2021). This variant harbors more mutations compared to prior variants, and therefore “efficiently” escapes humoral immunity

induced by primary vaccination. It was observed by Garcia-Beltran et al. that additional mRNA vaccine doses (boosters) tend to demonstrate cross-neutralizing responses against Omicron as well. This could be due to “affinity maturation of the existing antibodies” or new shared epitopes of variants serving as targets (Garcia-Beltran et al., 2022a). The authors also observed a 9-fold increase in the geometric mean neutralization titer (GMNT) for Delta pseudoviruses in individuals that received three-doses of BNT162b as compared to two doses, and an increase in cross-neutralization of the Omicron variant by 27-fold for individuals who received three doses of BNT162b vaccine. This neutralization signal could be correlated with antibody levels against SARS-CoV-2 spike-RBD measured by anti-spike-RBD immunoassays that are widely available (Garcia-Beltran et al., 2021). Therefore, our focus on antibody levels against spike-RBD is valuable.

Our study findings demonstrate that while double-dose vaccination boosted the IgG levels in recovered individuals 161 times, this “boost” was relatively short-lived. The unvaccinated recovered individuals, in contrast, continued to show a steady decline but detectable antibody levels. Whether or not the present vaccines induce sterilizing immunity remains unclear. They trigger serological IgA response but do not generate mucosal IgA at the virus site of entrance. Vaccine breakthrough infections may be because of deficient local preventive immunity due to absence of mucosal IgA (Piano Mortari et al., 2021).

Upon mRNA BNT162b2 vaccination, neutralizing antibodies are boosted significantly in the serum, but not at the mucosal sites and saliva. This indicates poor activation of oral mucosal immunity that fails

to limit the viral attachment to target cells at the entry site (Azzi et al., 2021).

In contrast, convalescent individuals that recovered from COVID-19 infection (also known as hybrid immunity) demonstrated the presence of specific IgA indicating that natural infection conferred mucosal immunity. (Piano Mortari et al., 2021; Azzi et al., 2021; Sterlin et al., 2021).

Vaccination-induced immunity in individuals with a history of prior infection was associated with high levels of wild-type neutralization titers even when vaccinated distantly. (>6 months) (Garcia-Beltran et al., 2022b).

In line with a study by Hall et al., our study results demonstrate that a previous COVID-19 infection (infection-acquired immunity) appeared to elicit robust, long-term, and sustained levels of SARS-CoV-2 antibodies in individuals who received two doses of BNT162b2. It was also observed that individuals with hybrid immunity developed the “highest and most durable protection”. (Hall et al., 2021).

Our study had some limitations. First, a small sample size. Second, ideally, simultaneous antibody titer detection of each patient at each time point using both NCP and S-RBD assay would have given the best results for comparative analysis but the S-RBD assay received emergency approval only later in 2020.

The nucleocapsid protein (NCP) of the SARS-CoV-2 is considered a biomarker associated with natural exposure. Since this NCP is not present in spike-based vaccines, there is no vaccine-induced response against it in vaccinated individuals. The type of antibodies produced and the type of cell-mediated immune response triggered depends on the mechanism of virus/vaccine product exposure (infection or vaccination). For example, mRNA and adenovirus vector vaccines induce antibodies against the spike protein of SARS-CoV-2; therefore a positive test for spike protein specific IgM or IgG in vaccinated individuals could indicate prior infection or vaccination. To differentiate between these two situations, a positive Nucleocapsid-protein assay could be used to indicate prior infection. Therefore, testing for anti-NCP could be an important tool to analyze the type of existing immunity in an individual (vaccine-induced/natural/hybrid). Moreover, Vaccinated individuals who were previously exposed usually test positive for antibodies against NCP (hybrid immunity) while the vaccinated individuals who have never been exposed lack anti-NCP antibodies (vaccine-induced immunity) (Assis et al., 2021; Krutikov et al., 2022). The most recent study in 2022, by Di Chiara et al. has also suggested that anti-S-RBD antibodies persist up to 18 months after infection, even in children bringing hope for the future of antibody dynamics and protection against reinfection.

The major strengths of our study include the longest observation (March 2020–September 2021) for the presence of antibodies against SARS-CoV-2 in recovered individuals along with the impact of 2 dose-BNT162b2 vaccination on these levels. Secondly, mathematical models: fixed-effect regression models were used for statistical analysis that diminishes the impact of confounding by (unmeasured) fixed/time-invariant factors. These modeling results could also be used to predict future titer trends. Thirdly, since the immunoassay to detect anti-NCP was reintroduced at 18 months, even the vaccinated individuals could be analyzed for the persistence of infection-induced immunity not generated by mRNA spike-based vaccination. Finally, combining the data for both was necessary because this makes it possible to explore the differences between anti-NCP and anti-S-RBD (as also shown by the regression results in Fig. 2(c)). This is similar to an interrupted time series (ITS) analysis which typically looks at how an ‘intervention’ affects the temporal pattern between before and after periods using whole datasets. Our data is rich in that it is both cross-sectional (multiple patients) and time-series which allows us to adopt complex longitudinal models.

9. Conclusion

This work provides further evidence of long-term persistence of

immune response in individuals recovered from SARS-CoV-2 wild-type infection and the impact of two-dose BNT162b2 (Pfizer-BioNTech) mRNA vaccination on these antibody responses which could help in vaccination strategies and prevention policies.

CRedit authorship contribution statement

Puya Dehgani-Mobaraki: Conceptualization, Writing – review & editing. **Chao Wang:** Software, Methodology, Formal analysis, Visualization. **Alessandro Floridi:** Methodology, Investigation, Resources. **Emanuela Floridi:** Methodology, Investigation, Resources. **Sunny Dawoodi:** Writing – review & editing. **Asiya K. Zaidi:** Data curation, Writing – original draft, preparation, Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgment

The authors would like to acknowledge the assistance provided by the Rotary Club Fortebraccio Montone for the successful conduction of this research study.

References

- Abu-Raddad, L.J., Chemaitelly, H., Coyle, P., Malek, J.A., Ahmed, A.A., Mohamoud, Y.A., Younuskuju, S., Ayoub, H.H., Al Kanaani, Z., Al Kuwari, E., Butt, A.A., Jeremijenko, A., Kaleeckal, A.H., Latif, A.N., Shaik, R.M., Abdul Rahim, H.F., Nasrallah, G.K., Yassine, H.M., Al Kuwari, M.G., Al Romaihi, H.E., Al-Thani, M.H., Al Khal, A., Bertollini, R., 2021. SARS-CoV-2 antibody-positivity protects against reinfection for at least seven months with 95% efficacy. *EClinicalMedicine* 35, 100861. <https://doi.org/10.1016/j.eclim.2021.100861>. May Epub 2021 Apr 28. PMID: 33937733; PMCID: PMC8079668.
- Anand, S.P., Prévost, J., Nayrac, M., et al., 2021. Longitudinal analysis of humoral immunity against SARS-CoV-2 Spike in convalescent individuals up to 8 months post-symptom onset. *Cell Rep Med* 2 (6), 100290. <https://doi.org/10.1016/j.xcrm.2021.100290>.
- Assis, R., Jain, A., Nakajima, R., et al., 2021. Distinct SARS-CoV-2 antibody reactivity patterns elicited by natural infection and mRNA vaccination. *npj Vaccines* 6, 132. <https://doi.org/10.1038/s41541-021-00396-3>.
- Azzi, L., Dalla Gasperina, D., Veronesi, G., Shalal, M., Ietto, G., Iovino, D., Baj, A., Gianfagna, F., Maurino, V., Focosi, D., Maggi, F., Ferrario, M.M., Dentali, F., Carcano, G., Tagliabue, A., Maffioli, L.S., Accolla, R.S., Forlani, G., 2021. Mucosal immune response in BNT162b2 COVID-19 vaccine recipients. *EBioMedicine* 75, 103788. <https://doi.org/10.1016/j.ebiom.2021.103788>. Dec 23 Epub ahead of print. PMID: 34954658; PMCID: PMC8718969.
- Clinical and Immunological Responses after SARS-CoV-2 Infection Causing COVID-19 - Full Text View, 2020. [ClinicalTrials.gov](https://www.clinicaltrials.gov).
- Dehgani-Mobaraki, P., Zaidi, A.K., Yadav, N., Floridi, A., Floridi, E., 2021. Longitudinal observation of antibody responses for 14 months after SARS-CoV-2 infection. *Clin. Immunol.* 230, 108814. <https://doi.org/10.1016/j.clim.2021.108814>. Sep Epub 2021 Jul 31. PMID: 34343708; PMCID: PMC8325385.
- Di Chiara, C., Cantarutti, A., Costenaro, P., et al., 2022. Long-term immune response to SARS-CoV-2 infection among children and adults after mild infection. *JAMA Netw. Open* 5 (7), e2221616. <https://doi.org/10.1001/jamanetworkopen.2022.21616>.
- Ebinger, J.E., Fert-Bober, J., Printsev, I., et al., 2021. Antibody responses to the BNT162b2 mRNA vaccine in individuals previously infected with SARS-CoV-2. *Nat. Med.* 27, 981–984. <https://doi.org/10.1038/s41591-021-01325-6>.
- Egbert, E.R., Xiao, S., Colantuoni, E., et al., 2021. Durability of spike immunoglobulin G antibodies to SARS-CoV-2 among health care workers with prior infection. *JAMA Netw. Open* 4 (8), e2123256. <https://doi.org/10.1001/jamanetworkopen.2021.23256>.
- Gaebler, C., Wang, Z., Lorenzi, J.C.C., Muecksch, F., Finkin, S., et al., 2021. Evolution of antibody immunity to SARS-CoV-2. *Nature* 591 (7851), 639–644. <https://doi.org/10.1038/s41586-021-03207-w>. Mar Epub 2021 Jan 18. PMID: 33461210; PMCID: PMC8221082.
- Gallais, F., Gantner, P., Bruel, T., Velay, A., Planas, D., Wendling, M.J., Bayer, S., Solis, M., Laugel, E., Reix, N., Schneider, A., Gladly, L., Panaget, B., Collongues, N., Partisani, M., Lessinger, J.M., Fontanet, A., Rey, D., Hansmann, Y., Kling-

- Pillitteri, L., Schwartz, O., De Sèze, J., Meyer, N., Gonzalez, M., Schmidt-Mutter, C., Fafi-Kremer, S., 2021. Evolution of antibody responses up to 13 months after SARS-CoV-2 infection and risk of reinfection. *EBioMedicine* 71, 103561. <https://doi.org/10.1016/j.ebiom.2021.103561>. Sep Epub 2021 Aug 27. PMID: 34455390; PMCID: PMC8390300.
- Garcia-Beltran, W.F., Lam, E.C., Astudillo, M.G., Yang, D., Miller, T.E., Feldman, J., Hauser, B.M., Caradonna, T.M., Clayton, K.L., Nitido, A.D., Murali, M.R., Alter, G., Charles, R.C., Dighe, A., Branda, J.A., Lennerz, J.K., Lingwood, D., Schmidt, A.G., Iafate, A.J., Balazs, A.B., 2021. COVID-19-neutralizing antibodies predict disease severity and survival. *Cell* 184 (2), 476–488. Jan 21.
- Garcia-Beltran, W.F., St Denis, K.J., Hoelzemer, A., Lam, E.C., Nitido, A.D., Sheehan, M. L., Berrios, C., Ofoman, O., Chang, C.C., Hauser, B.M., Feldman, J., Roederer, A.L., Gregory, D.J., Poznansky, M.C., Schmidt, A.G., Iafate, A.J., Naranbhai, V., Balazs, A. B., 2022a. mRNA-based COVID-19 vaccine boosters induce neutralizing immunity against SARS-CoV-2 Omicron variant. *Cell*. <https://doi.org/10.1016/j.cell.2021.12.033>. Jan 6:S0092-8674(21)01496-3 Epub ahead of print. PMID: 34995482; PMCID: PMC8733787.
- Garcia-Beltran, W.F., St Denis, K.J., Hoelzemer, A., Lam, E.C., Nitido, A.D., Sheehan, M. L., Berrios, C., Ofoman, O., Chang, C.C., Hauser, B.M., Feldman, J., Roederer, A.L., Gregory, D.J., Poznansky, M.C., Schmidt, A.G., Iafate, A.J., Naranbhai, V., Balazs, A. B., 2022b. mRNA-based COVID-19 vaccine boosters induce neutralizing immunity against SARS-CoV-2 Omicron variant. *Cell*. <https://doi.org/10.1016/j.cell.2021.12.033>. Jan 6:S0092-8674(21)01496-3 Epub ahead of print. PMID: 34995482; PMCID: PMC8733787.
- Gregory, M., Hames, Thomas, et al., 2021. Does infection with or vaccination against SARS-CoV-2 lead to lasting immunity? *Lancet Respir. Med.* [https://doi.org/10.1016/S2213-2600\(21\)00407-0](https://doi.org/10.1016/S2213-2600(21)00407-0). Published Online October 21, 2021.
- Hall, V., Foulkes, S., Insalata, F., et al., 2021. Effectiveness and durability of protection against future SARS-CoV-2 infection conferred by COVID-19 vaccination and previous infection; findings from the UK SIREN prospective cohort study of healthcare workers March 2020 to September 2021. *medRxiv* 11 (29), 21267006. <https://doi.org/10.1101/2021.11.29.21267006>.
- <https://www.fda.gov/medical-devices/coronavirus-disease-2019-covid-19-emergency-use-authorizations-medical-devices/vitro-diagnostics-euas>.
- https://www.snibe.com/zh_en/en_newsView.aspx?id=647.
- Krutikov, M., Palmer, T., Tut, G., Fuller, C., Azmi, B., Giddings, R., Shrotri, M., Kaur, N., Sylla, P., Lancaster, T., Irwin-Singer, A., Hayward, A., Moss, P., Copas, A., Shallcross, L., 2022. Prevalence and duration of detectable SARS-CoV-2 nucleocapsid antibodies in staff and residents of long-term care facilities over the first year of the pandemic (VIVALDI study): prospective cohort study in England. *Lancet Healthy Longev* 3 (1), e13–e21. [https://doi.org/10.1016/S2666-7568\(21\)00282-8](https://doi.org/10.1016/S2666-7568(21)00282-8). Jan Epub 2021 Dec 16. PMID: 34935001; PMCID: PMC8676418.
- Levin, E.G., Lustig, Y., Cohen, C., Fluss, R., Indenbaum, V., Amit, S., Doolman, R., Asraf, K., Mendelson, E., Ziv, A., Rubin, C., Freedman, L., Kreiss, Y., Regev-Yochay, G., 2021. Waning immune humoral response to BNT162b2 covid-19 vaccine over 6 months. *e84 N. Engl. J. Med.* 385 (24). <https://doi.org/10.1056/NEJMoa2114583>. Epub 2021 Oct 6 PMID: 34614326; PMCID: PMC8522797.
- MAGLUMI. nCoV IgM/IgG - Letter of Authorization, 2019. [fda.gov](https://www.fda.gov) (Google Scholar).
- Orsini, N., Greenland, S., 2011. A procedure to tabulate and plot results after flexible modeling of a quantitative covariate. *STATA J.* 11 (1), 1–29.
- Piano Mortari, E., Russo, C., Vinci, M.R., Terreri, S., et al., 2021. Highly specific memory B cells generation after the 2nd dose of BNT162b2 vaccine compensate for the decline of serum antibodies and absence of mucosal IgA. *Cells* 10 (10), 2541. <https://doi.org/10.3390/cells10102541>. Sep 26 PMID: 34685521; PMCID: PMC8533837.
- Saadat, S., Rikhtegaran Tehrani, Z., Logue, J., et al., 2021. Binding and neutralization antibody titers after a single vaccine dose in health care workers previously infected with SARS-CoV-2. *JAMA* 325 (14), 1467–1469. <https://doi.org/10.1001/jama.2021.3341>.
- Sterlin, D., Mathian, A., Miyara, M., et al., 2021. IgA dominates the early neutralizing antibody response to SARS-CoV-2. *Sci. Transl. Med.* 13 (577), eabd2223 <https://doi.org/10.1126/scitranslmed.abd2223>.
- Turner, J.S., Kim, W., Kalaidina, E., Goss, C.W., Rauseo, A.M., Schmitz, A.J., Hansen, L., Haile, A., Klebert, M.K., Pusic, I., O'Halloran, J.A., Presti, R.M., Ellebedy, A.H., 2021. SARS-CoV-2 infection induces long-lived bone marrow plasma cells in humans. *Nature* 595 (7867), 421–425. <https://doi.org/10.1038/s41586-021-03647-4>. Jul Epub 2021 May 24. PMID: 34030176.
- Who.int, 2021. Vaccine equity declaration [online] Available at: <https://www.who.int/campaigns/vaccine-equity/vaccine-equity-declaration>>. (Accessed 7 November 2021).
- World Health Organization, 2020. International guidelines for certification and classification (coding) of COVID-19 as cause of death. Available from: https://www.who.int/classifications/icd/Guidelines_Cause_of_Death_COVID-19-20200420-EN.pdf?ua=1 Document Number: WHO/HQ/DDI/DNA/CAT. June 1.
- Xydakis, M.S., Dehghani-Mobaraki, P., Holbrook, E.H., et al., 2020. Smell and taste dysfunction in patients with COVID-19. *Lancet Infect. Dis.* 20 (9), 1015–1016. [https://doi.org/10.1016/S1473-3099\(20\)30293-0](https://doi.org/10.1016/S1473-3099(20)30293-0). Sep Epub 2020 Apr 15. PMID: 32304629; PMCID: PMC7159875.