

Research

**Cite this article:** Divín D *et al.* 2022Cannabinoid receptor 2 evolutionary gene loss makes parrots more susceptible to neuroinflammation. *Proc. R. Soc. B* **289**: 20221941.<https://doi.org/10.1098/rspb.2022.1941>

Received: 28 September 2022

Accepted: 9 November 2022

Subject Category:

Neuroscience and cognition

Subject Areas:

evolution, genetics, immunology

Keywords:

neuroimmunology, avian immunology, cannabinoid receptors, neural inflammation, gene loss, cannabinoid receptor pseudogenization

Author for correspondence:

Michal Vinkler

e-mail: michal.vinkler@natur.cuni.czElectronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6307543>.

Cannabinoid receptor 2 evolutionary gene loss makes parrots more susceptible to neuroinflammation

Daniel Divín¹, Mercedes Gómez Samblas¹, Nithya Kuttiyarthu Veetil¹, Eleni Voukali¹, Zuzana Świdarská¹, Tereza Krajcingrová¹, Martin Těšický¹, Vladimír Beneš², Daniel Elleder³, Oldřich Bartoš⁴ and Michal Vinkler¹¹Faculty of Science, Department of Zoology, Charles University, Viničná 7, Prague 128 44, Czech Republic²Genomics Core Facility, European Molecular Biology Laboratory Heidelberg, Meyerhofstrasse 1, 69117 Heidelberg, Germany³Institute of Molecular Genetics, Czech Academy of Sciences, Vídeňská 1083, 142 20 Prague 4, Czech Republic⁴Military Health Institute, Military Medical Agency, Tychonova 1, 160 01 Prague 6, Czech Republic

ID DD, 0000-0001-8513-7214; NKV, 0000-0002-1202-4902; MV, 0000-0003-3572-9494

In vertebrates, cannabinoids modulate neuroimmune interactions through two cannabinoid receptors (CNRs) conservatively expressed in the brain (CNR1, syn. CB1) and in the periphery (CNR2, syn. CB2). Our comparative genomic analysis indicates several evolutionary losses in the *CNR2* gene that is involved in immune regulation. Notably, we show that the *CNR2* gene pseudogenized in all parrots (Psittaciformes). This *CNR2* gene loss occurred because of chromosomal rearrangements. Our positive selection analysis suggests the absence of any specific molecular adaptations in parrot *CNR1* that would compensate for the *CNR2* loss in the modulation of the neuroimmune interactions. Using transcriptomic data from the brains of birds with experimentally induced sterile inflammation we highlight possible functional effects of such a *CNR2* gene loss. We compare the expression patterns of CNR and neuroinflammatory markers in *CNR2*-deficient parrots (represented by the budgerigar, *Melopsittacus undulatus* and five other parrot species) with *CNR2*-intact passerines (represented by the zebra finch, *Taeniopygia guttata*). Unlike in passerines, stimulation with lipopolysaccharide resulted in neuroinflammation in the parrots linked with a significant upregulation of expression in proinflammatory cytokines (including interleukin 1 beta (*IL1B*) and 6 (*IL6*)) in the brain. Our results indicate the functional importance of the *CNR2* gene loss for increased sensitivity to brain inflammation.

1. Introduction

Psychiatric and neurodegenerative disorders have increased in incidence globally in the human population [1,2]. Surprisingly, similar psychological (in animals referred to as behavioural) disorders have been frequently recognized in some cognitively advanced animals, namely the parrots [3–5]. Like in human depression, in parrots symptoms like anxiety, apathy, over-eating, indifference and self-damage (feather plucking) are observed and diagnosed by veterinarians [6–8]. While in parrots little is presently known about the possible causes of the behavioural disorders, in humans they have been recently linked with neural inflammation [9]. Brain neuroinflammation can be induced by signals from the periphery, where pathogens and tissue damage trigger immune responses through stimulation of pattern recognition receptors [10]. Resultant cytokine signalling may modulate central nervous system functioning through the activation of brain microglia and astrocytes [11] interfering with healthy brain neuronal regulation [12,13]. Proinflammatory cytokines, such as interleukin 1 beta (*IL1B*) or 6 (*IL6*) [14,15] become overexpressed in the brain,

being key neuroinflammation markers [16,17]. Mental health depends on finely balanced regulation of the neuroimmune interplay. Among neuronal modulators interlinking the nervous and immune systems, cannabinoids recognized through cannabinoid receptors (CNRs) have been shown to provide important anti-neuroinflammatory effects in humans [18–21]. Compared to humans, in animals the immunological factors affecting behavioural disorders are far less clear and interspecific variation in the neuroimmune regulatory networks remains unknown. Parrots with their advanced cognitive abilities [22], dense neuronal networks [23] and common psychopathologies [6,7] could help us to understand the general principles of neuroinflammation effects on behaviour.

Gene loss is a widespread phenomenon responsible for evolutionary changes in organisms, including immunity and other physiological functions [24]. It may be involved in adaptive responses to environmental or pathogen-driven changes in selective pressures [25] or represent a random shift in gene content with deleterious effects insufficient to be prevented by negative selection [26]. Genomic chromosomal rearrangement is probably an important source of the gene loss events. Massive chromosomal alterations have profoundly affected vertebrate evolution in general [27], as well as in certain lineages [28] including parrots in particular [29–31]. Recent advances in genomic research have allowed thorough mapping of evolutionary gene loss events affecting immune signalling [24,32–35]. In the present study, we performed a comparative genomic database search of negative regulators of neural inflammation which indicated an interesting pattern of loss in the *CNR2* gene in parrots.

The endocannabinoid system regulating both the neural and immune functions consists of CNRs, their ligands (endocannabinoids) and enzymes synthesizing and degrading cannabinoids [36]. Two CNR paralogues are known in all vertebrates [37]: *CNR1*, which is mainly expressed in cells of the nervous system, and *CNR2*, which is mainly expressed in immune cells, including microglia in the brain [38,39]. *CNR1* is involved in the regulation of emotions, memory, motor activity, feelings, attention, neuropeptide synthesis, gastrointestinal tract functions, metabolism and—in birds—singing [40–44]. *CNR2* affects immunosuppression and decreases inflammation, pain and the expression of proinflammatory cytokines, playing an important role in negative feedback regulation [45–47]. *CNR2* expression has been shown to increase with the activation of immune cells related to higher expression of proinflammatory cytokines [39]. In mice, *CNR2* expression in brain-based microglia was upregulated during neurological inflammation, contributing to the suppression of the inflammatory response [45].

Since our comparative genomic search suggested that among negative regulators of inflammation, parrots consistently miss only the *CNR2* gene, here we use genomic and transcriptomic data to map the putative *CNR2* loss events across vertebrate phylogeny. Subsequently, exemplified in the budgerigar (*Melopsittacus undulatus*) and kakapo (*Strigops habroptila*) genomes we reconstruct the *CNR2* loss events in parrots. Using analysis of positive selection, we search for compensatory adaptations in *CNR1* in species lacking *CNR2*. Finally, by comparing parrot and passerine neuroinflammation marker expression patterns in the brain during an immune response, we assess the consequences of *CNR2* loss on neuroimmune regulation in parrots.

2. Methods

(a) Identification of cannabinoid receptor-loss events

To identify the candidate genomic differences between passerines and parrots that could result in parrot susceptibility to neuroinflammatory disorders, we first checked for the patterns of gene representation in these two taxa. The gene set of negative regulators of inflammatory responses (GO:0050728) was retrieved from the AmiGO database, accessed August 2022 [48], based on human (*Homo sapiens*) annotation ($n = 154$) and then chicken (*Gallus gallus*) orthologues, identified using the ENSEMBL BioMart tool [49]. This allowed us to shortlist 122 avian genes involved in the regulation. Using the Avian Immunome database (AVIMM; accessed August 2022 [50]) we identified 53 immune genes to which information on presence/absence could be retrieved across six parrot and 13 passerine species covering the Psittacopasserae phylogeny: *Melopsittacus undulatus*, *Eolophus roseicapillus*, *Probosciger aterrimus*, *Amazona guildingii*, *Agapornis roseicollis*, *Nestor notabilis*, *Corvus moneduloides*, *Ficedula albicollis*, *Hirundo rustica*, *Lepidothrix coronata*, *Lonchura striata*, *Molothrus ater*, *Parus major*, *Passer domesticus*, *Serinus canaria*, *Sturnus vulgaris*, *Taeniopygia guttata*, *Zonotrichia albicollis*, *Zosterops hypoxanthus*. In this list, we checked for cases of consistent absence of a regulating gene in parrots and its consistent presence in passerines. Only a single gene, *CNR2*, fulfilled this criterion (electronic supplementary material, S2, table S21).

For the phylogenetic analysis of the two related CNR genes, *CNR1* and *CNR2*, we first downloaded all available tetrapod CNR coding DNA sequences from the Ensembl genome browser database (release 103, www.ensembl.org; last accessed on 22 January 2021). Based on a comparison of lists of species with annotated *CNR1* and *CNR2*, we identified all cases of putative *CNR1* or *CNR2* absence. For these species, we performed a targeted search through the NCBI databases (<https://www.ncbi.nlm.nih.gov>, release 236) using blastx and tblastn (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) to find the missing orthologues. Using this complete sequence dataset, supplemented with *CNR1* sequences from five other parrots species represented in the parrots' experiment (E2) obtained by NextSeq Illumina transcriptomic sequencing (see below), we reconstructed the CNR phylogenetic tree (based on 318 sequences) in the online tool iTOL to verify the sequence gene-specific orthology [51,52]. For a list of all species, including their *CNR1* and *CNR2* sequence accession numbers, see the electronic supplementary material, S1, table S1. The final dataset consisted of 160 orthologues of zebra finch (*T. guttata*)/budgerigar *CNR1* and 158 orthologues of zebra finch *CNR2* (electronic supplementary material, table S1). The position of *CNR2* in the zebra finch, chicken and human karyotypes was checked in Ensembl and the neighbouring coding genes were identified in parrots with karyotype information available in Ensembl (the budgerigar and kakapo). Based on this data, we reconstructed the genomic changes leading to *CNR2* pseudogenization.

(b) Selection analysis

We examined the evidence for positive selection acting on vertebrate CNRs in order to infer whether loss of *CNR2* could be linked to any alteration to *CNR2* functioning in the clade of parrot-related taxa, and whether it might have resulted in any compensatory evolution in parrot *CNR1*. First, we used the tool CONSURF (<http://consurf.tau.ac.il>; [53]) to identify non-conservative regions on the CNR surface. Next, we adopted a combination of tools for detecting positive selection available on the Datamonkey server (<https://www.datamonkey.org/>): FUBAR [54], MEME [55], aBSREL [56] and RELAX [57]; for details see the electronic supplementary material. We then used the online tools PROVEAN (Protein Variation Effect Analyzer, <http://provean.jcvi.org/>; [58]) and SIFT (Scale Invariant Feature Transform, <https://sift.bii.a-star.edu.sg/>; [59]) to predict functional effects of the amino acid substitutions observed at sites under positive selection.

(c) Experimental procedures

In budgerigars (experiment 1 (E1), $n=30$), in the six selected parrot species (experiment 2 (E2), $n=36$, i.e. the red-rumped parrot *Psephotus haematonotus*, the rosy-faced lovebird *Ag. roseicollis*, the elegant parrot *Neophema elegans*, the budgerigar, the cockatiel *Nymphicus hollandicus* and the pacific parrotlet *Forpus coelestis*, six individuals per species) and in the zebra finches (experiment 3 (E3), $n=24$) we used standardized methodology to map the *CNR* and *IL1B* expression trajectories during acute immune response (see the electronic supplementary material, S1, table S2). All birds from all experiments (E1–3) were obtained from local hobby breeders and housed in pairs in cages $100 \times 50 \times 50$ cm. The birds had access to food and water ad libitum and were kept under a 12 L : 12 D controlled light/dark cycle with a regulated temperature of $22 \pm 2^\circ\text{C}$. Treatment individuals were intra-abdominally injected with lipopolysaccharide (LPS; *Escherichia coli* O55:B5; Sigma-Aldrich, cat. no. L2880) in a dose equivalent to $6 \mu\text{g}$ per gram body weight and compared to controls injected with a sterile Dulbecco's phosphate-buffered saline (Sigma-Aldrich, cat. no. D5652). The LPS dose was chosen based on previous studies in other small-sized birds inducing a measurable non-specific immune response [60]. In the first experiment with budgerigars (E1) the experimental birds were euthanized at different time points, i.e. at 3, 6, 12, 24 and 48 h post-treatment ($n=3$ per time point and treatment) to check for the immune response dynamics, while in the second experiment with the different parrot species (E2) and in the third experiment with the zebra finches (E3) (consistent with the results from E1) the time interval for the immune response was set to 24 h. In all birds, tissue samples of the small intestine (ileum) and the brain hyperpallial area were collected as necropsies after euthanasia, placed immediately into RNA later (cat. no. 76106, Qiagen, Hilden, Germany) and stored at -80°C until RNA extraction. The research was approved by the Ethical Committee of Charles University, Faculty of Science (permits 13882/2011-30 and MSMT-30397/2019-5) and was carried out in accordance with the current laws of the Czech Republic and the European Union.

(d) Transcriptomic search for *CNR1* and *CNR2* genes in parrots

Small intestine transcriptomes for the six parrot species were obtained from sequencing libraries prepared using the NEB-Next Ultra II Directional RNA Library Prep Kit for Illumina (cat. no. E7760, San Diego, CA, USA) in the European Molecular Biology Laboratory (EMBL), Heidelberg (NCBI accession numbers: SAMN23963146, SAMN23963147, SAMN23963148, SAMN23963149, SAMN23963150, SAMN23963151). Paired-end sequencing (80 bp from each end) was performed on the NextSeq 500 system (Illumina) at a sequencing depth of 13–19 million reads per library. Forward and reverse reads were merged, and low-quality reads and adaptor sequences discarded, using BBsuite ('BBMap' n.d.). *De novo* transcriptome assembly was performed by TRINITY [61] under default settings. To obtain sets of non-redundant transcripts, we applied two filtering steps. First, we used TRANSDCODER [62] to identify the longest open reading frame of each transcript for each species individually, and second, redundancy was further reduced in the remaining transcript sets by clustering highly similar sequences with CD-HIT [63], using a sequence identity threshold of 0.9. Completeness of the six assembled transcript sets against a set of highly conserved single-copy orthologues was assessed using BUSCO (Benchmarking Universal Single-Copy Orthologs v.4.1.4; [64]). To identify *CRN1* and *CRN2* coding sequences for each species, reference budgerigar (for *CRN1*, Ensembl transcript ID: ENSMUNT00000010298.1) and zebra finch (for *CRN2*, Ensembl transcript ID: ENSTGUG00000001188) sequences were searched using BLASTN [65] and compared against

raw reads and the sequences obtained for positive selection analysis, and further against transcriptome assemblies.

(e) Brain transcriptomic gene expression analysis

As an initial check for the differential gene expression in selected cytokines in zebra finch and budgerigar brains, we used the QuantSeq 3' end sequencing approach [66]. Samples were first barcoded with Illumina TruSeq adapters and sequencing was undertaken on the Illumina HiSeq 2500 platform at EMBL, Heidelberg. The sequenced samples (NCBI accession number: PRJNA751848 and PRJNA879979) were then analysed using the BAQCOM pipeline (<https://github.com/hanielcedraz/BAQCOM>), the adapters being removed using the TRIMMOMATIC tool (<http://www.usadellab.org/cms/>). The samples were then aligned to the zebra finch reference genome (downloaded from Ensemble) using STAR aligner (<https://github.com/alexdobin/STAR>), the featureCounts, Subread R package being used to assign read counts to the genes. Given their low representation in the transcriptomic data, specific inflammatory markers, *IL1B*, *IL6*, *IL8*, *IL12B*, *IL15*, *IL17B*, *IL18* and *IL22* were selected based on the literature review [67] and their 3' annotation available in the Ensemble. In *IL6*, *IL17B* and *IL22* we did not obtain sufficient read coverage to proceed further with a quantitative analysis. In order to normalize the expression data in the rest of the target genes, we first divided the total number of reference (cytokine)-aligned reads by the total number of reads in the sample (C_n). To scale the data, we then multiplied each of the normalized read counts by 10 million (approx. 10 million was the average number of reads per sample in our dataset). The cytokine expression was quantified as the scaled-normalized number of reads per treatment individual divided by the mean scaled-normalized number of reads in all the control birds: relative differential gene expression = $(C_n \times 10^6)_{\text{Treatment}} / (\Sigma(C_n \times 10^6) / N)_{\text{Control}}$.

(f) Real-time quantitative polymerase chain reaction gene expression analysis

Designing conserved primers based on avian interspecific alignments we amplified the partial coding regions of *IL1B*, *IL6*, *CNR1* and *CNR2* and Sanger sequenced these in genomic DNA (gDNA) extracted from 12 blood samples representing different parrot species, 10 budgerigar samples and 12 zebra finch samples to assess intraspecific genetic variability and to design conserved real-time quantitative polymerase chain reaction (RT-qPCR) primers (electronic supplementary material, S1, table S3). The sequences were analysed using GENEIOUS (<http://www.geneious.com>, [68]).

Total RNA was extracted from parrot and zebra finch brain samples using the High Pure RNA Tissue Kit (cat. no. 12033674001; Roche, Basel, Switzerland), the concentration and quality of the RNA being measured on the NanoDrop 1000 Spectrophotometer (Thermo Fisher Scientific). The RNA was diluted in molecular water enriched with carrier transfer RNA (Qiagen, cat. no. 1068337) in the ratio 1 : 5 for target genes or 1 : 500 for *28S rRNA*. To calculate the efficiency of each primer pair, a calibration curve was constructed with synthetic DNA standard (gBlocks; IDT, Coralville, IA, USA; electronic supplementary material, S1, table S4) using a dilution series of 10^8 – 10^2 copies μl^{-1} , estimated according to Vinkler *et al.* [69]. The RNA samples and standards were amplified using the Luna Universal Probe One-Step RT-PCR Kit (E3006, BioLabs Inc, Ipswich, MA, USA), with 0.6 mM primer and 0.2 mM probe concentrations (electronic supplementary material, S1, table S5). RT-qPCR quantification was conducted using a LIGHTCYCLER 480 PCR platform (Roche) set with the cycling conditions shown in the electronic supplementary material, S1, table S6. All assays were performed with template-free negative controls and block positive controls in a freshly prepared dilution series, using *28S rRNA* as a reference gene. Relative quantification (R) was calculated from the crossing point (Cp) values determined




Figure 1. Phylogenetic tree showing gene-specific clustering of *CNR1* (blue) and *CNR2* (green). Lamprey (orange) shows the root of the tree as a common ancestor of the genes. Terminal triangles represent collapsed taxon-specific branches. The red colour highlights the presence of species with missing receptors (i.e. cases where the receptors were not revealed in the database search). A fully expanded tree is provided in the electronic supplementary material, figure S1).

by the second derivative maximum [70], using E and Cp data calculated using LightCycler480 software v.1.5.1. To test for gene expression changes between treatment and control birds, we quantified relative gene expression as standardized relative quantities (Qst; [69]). For the RT-qPCR efficiencies (E) see the electronic supplementary material, S1, table S7; for the final RT-qPCR data see the electronic supplementary material, S2, table S22).

(g) Statistical analysis

The statistical analysis was performed in RSTUDIO v.2021.09.0 [71]. First, the initial transcriptomic cytokine expression data were tested for the differences in inflammatory genes expression between the budgerigars and zebra finches using a Wilcoxon paired test and the results were plotted in a heatmap generated using the pheatmap package. Next, the RT-qPCR verification of these results was performed. Given their non-Gaussian distribution, the Qst values were normalized using decadic logarithms ($\log Qst$). The effects of experimental treatment on gene expression changes were assessed using the linear models (LMs) in the 'lme4' package, where gene expression (continuous) served as a response variable. For the budgerigar (E1) dataset, the full model contained treatment, sex and time as explanatory variables. For the comparative parrot (E2) dataset, the full model contained treatment, sex and species as explanatory variables. Based on the E1 results, for the zebra finch (E3) dataset, only treatment was used as an explanatory variable in the full model.

Minimum adequate models (here defined as models with all terms significant at $p \leq 0.05$) were selected by backward elimination of non-significant terms from the full models. All backward elimination steps in the models were verified by changes in deviance with an accompanying change in degrees of freedom (ANOVA) and Akaike information criterion, using *F*-statistics. The Pearson correlation test was used to assess the relationship between the expression of the *CNR* genes and *IL1B* and *IL6*.

3. Results

(a) Identification of the cannabinoid receptor genes in parrot genomes

Searching through genomic databases, we identified a single negative regulator of neuroinflammation consistently missing in parrots, but consistently present in passerines, the *CNR2*. To confirm this pattern, we used the tetrapod *CNR* sequence data retrieved from Ensembl supplemented with the NCBI BLAST-search results (electronic supplementary material, S1, table S1) to construct a *CNR* phylogenetic tree showing *CNR1* and *CNR2* presence and absence (figure 1). We failed to identify the *CNR2* gene in any parrot species, though it was present in all parrot relatives: falcons (Falconiformes), seriemas (Cariamiformes) and passerines (Passeriformes). According to




Figure 2. Schematic of the *CNR2* locus position and its neighbourhood in the human, chicken, budgerigar, zebra finch and kakapo genomes. Chromosome sequence is schematically indicated with the grey arrowed line over which positions of individual genes are indicated by colourful arrows (each colour represents one group of genes sitting in proximity in the ancestral state); *CNR2* and its closest human, chicken and zebra finch neighbouring genes, *FUCA1* and *PNRC2*, are marked in red and linked by red shades between the species. Each gene is labelled with its symbol above the arrow and its chromosomal location is marked below. White areas with the dotted grey lines indicate longer interspacing regions containing additional genes that are not important for the reconstruction of the chromosomal rearrangements in parrots. The recombination breakpoint in *CNR2* is indicated by a broken line, a curved arrow indicates the inversion event that occurred in the budgerigar evolutionary lineage, while a straight arrow indicates the translocation event that putatively occurred independently in the kakapo evolutionary lineage.

Ensembl, the *CNR2* gene is located on the 23rd chromosome in the zebra finch and chicken genomes, being directly adjacent to the *FUCA1* gene (upstream) and the *PNRC2* gene (downstream; figure 2). In the budgerigar genome, we found both these genes on chromosome 14; however, there was an approximately 5.5 Mbp insertion with inverted gene order directly between *FUCA1* and *PNRC2* (figure 2). Using BLAST, we identified short gene fragments showing 28% similarity to the barn owl (*Tyto alba*) *CNR2* and 10% similarity to the blue-crowned manakin (*Lepidothrix coronata*) *CNR2*, 6072 bp downstream of *PNRC2*. Interestingly, in the kakapo genome, different genes were situated downstream of the *PNRC2* gene (on the 15th chromosome; figure 2) and there was no sign of any remaining *CNR2* gene or pseudogene. To confirm the absence of the *CNR2* gene in parrot genomes, we designed sequence-conserved *CNR1* and *CNR2*-specific primers and sequenced the parrot gDNA-derived PCR amplicons. By contrast to the zebra finch, we found no evidence for the *CNR2* presence in budgerigar or any other parrot gDNA. Finally, our whole transcriptome complementary DNA sequencing in inflamed small intestine tissue failed to reveal *CNR2* in budgerigars, or in any of the other five parrot species analysed. We take this as conclusive evidence for the absence of functional *CNR2* in parrots.

(b) Positive selection in cannabinoid receptors

We next questioned the hypotheses that the loss of *CNR2* could be linked to its altered function in the parrot-related taxa and that *CNR1* took over the functional role of *CNR2* when lost in the parrots. Across tetrapods, the test for selection relaxation was not significant in *CNR1* ($K=0.66$, $p=0.822$, $LR=0.05$) or *CNR2* ($K=1.03$, $p=0.964$, $LR<0.001$). Using CONSURF, we identified 67 non-conservative sites in *CNR1* and 61 non-conservative sites in *CNR2* (electronic supplementary material, figure S2; S1, tables S8 and S9). In *CNR1*, the FUBAR test failed to identify any positively selected sites, while the MEME test identified seven sites under episodic positive selection (electronic supplementary material, S1, table S10). In *CNR2*, one positively selected site was identified by FUBAR and 15 sites were revealed as under branch-specific positive selection by MEME (electronic supplementary material, table S10). However, no specific non-synonymous substitutions with a putatively compensatory role were identified in *CNR1* in parrots and there was no indication of any changes in *CNR2* function in parrot relatives. Also, aBSREL found no evidence of any episodic diversifying selection in parrot phylogeny in the *CNR1* gene or in parrot-related species (i.e. zebra finch, common kestrel) in the




Figure 3. Heatmap showing differences in relative expression changes of proinflammatory cytokines in brains of LPS-stimulated budgerigars and zebra finches. MeUnX = budgerigar, X = number of individual; TaGuY = zebra finch, Y = number of individual. The heatmap colour ranges from blue (lowest upregulation in expression) to red (highest upregulation in expression).

CNR2 gene. PROVEAN used to identify significant changes in function caused by any amino acid variation, failed to indicate any important changes. Finally, SIFT predicted functional changes in *CNR1* at the sites D466R (with a score of 0.04) and T468I (score 0.04), and in *CNR2* at site V342I (score 0.05), but none of these changes proved important in birds. As such, we consider both *CNR1* and *CNR2* to be functionally conserved in the taxa where these genes are present.

(c) Transcriptomic evidence for *CNR2*-associated variation in inflammatory marker expression changes in brain during an immune response

We used transcriptomic data from zebra finch and budgerigar brains to check for the *CNR2*-linked functional variation in neuroinflammatory responsiveness. Checking for expression changes in the expression of proinflammatory cytokines after *in vivo* stimulation with LPS, we detected statistically significant differences between these two species in *IL1B* (Wilcoxon paired test; $p = 0.021$), *IL8* ($p = 0.037$), *IL12B* ($p = 0.001$) and *IL18* ($p = 0.020$) responses. For *IL6*, *IL17B* and *IL22* there was insufficient read representation to perform the statistical test and for *IL15* we found no significant difference in gene expression between the two species ($p = 0.717$). The results are shown in figure 3.

To verify these transcriptomic patterns indicating functional effects of *CNR2* loss on neuroimmune functioning in parrots, we followed the *CNR1/2*, *IL1B* and *IL6* expression on

messenger RNA (mRNA) level across the two taxa using RT-qPCR. In the budgerigar (E1), relative *IL1B* and *IL6* expression increased in the brain following the LPS stimulation ($p < 0.001$ for both markers; electronic supplementary material, S1, tables S11–S13). By contrast, the expression of *CNR1* was independent of the LPS treatment ($p > 0.05$; electronic supplementary material, S1, tables S11 and S14). Considering the putative interspecific differences, we next compared changes in *IL1B* and *CNR1* gene expression on the mRNA levels following LPS stimulation in the six parrot species (E2). The results confirmed that the expression of *IL1B* in the brain changes in response to LPS stimulation, regardless of species ($p = 0.005$; electronic supplementary material, S1, tables S11 and S15). Again, we found no effect of the LPS stimulation on *CNR1* mRNA expression ($p > 0.050$; electronic supplementary material, S1, tables S11 and S16). By contrast, in the zebra finch, a species with a functional *CNR2* receptor, there was no significant effect of the LPS treatment on expression changes of any of these genes ($p > 0.050$; electronic supplementary material, S1, tables S11, S17–S20). There was no correlation between *CNR1* and *IL1B* ($p > 0.050$) or *IL6* ($p > 0.050$) expression in brain in any of the compared taxa (electronic supplementary material, S1, figures S3–S7). However, expression of *CNR2* in zebra finch was significantly positively correlated with expression of *IL1B* ($p = 0.009$; $r = 0.711$; electronic supplementary material, S1, figure S8), but not *IL6* ($p = 0.753$, $r = 0.315$, electronic supplementary material, S1, figure S9). Taken altogether, these results confirm no overall increase in expression of neuroinflammatory markers in the *CNR2*-intact




Figure 4. Expression of (a) *IL1B* and (b) *IL6* following peripheral stimulation with LPS in the brains of budgerigars and zebra finches. Gene expression assessed based on RT-qPCR is shown as centred standardized relative expression ($\log Q_{st}$) values, i.e. species population average is equal to 0. C = controls, LPS = LPS-stimulated treatments. Statistically significant differences ($p \ll 0.001$; electronic supplementary material, table S11 in SM1) are marked with asterisks. (Online version in colour.)

passerines following LPS stimulation, but a contrasting significant upregulation of these markers in the brains of LPS-stimulated *CNR2*-deficient parrots (figure 4*a,b*).

4. Discussion

Our search through negative regulators of inflammatory responses revealed that *CNR2* is the only gene consistently missing in parrots being a candidate for their susceptibility to neuroinflammation. We show that the *CNR2* gene has been lost during parrot evolution through chromosomal rearrangements. We found no evidence for compensatory evolution in *CNR1* after *CNR2* loss in parrots and the comparative experimental findings suggest that the gene loss events affect neuroimmune regulation. While in passerines possessing functional *CNR2* (represented by the zebra finch) peripheral inflammation induced by LPS failed to trigger any neuroinflammation (measured on transcriptomic level), in the *CNR2*-deficient parrots we recorded in the brain a significant proinflammatory cytokine upregulation.

LPS-induced activation of the immune system in the periphery can trigger systemic immune responses with neuroinflammatory outcomes [72–74] that in mammals can cause important alterations in behaviour and cognition [75]. This phenomenon has not been recorded in birds, where even high doses of LPS (more than 6 mg kg^{-1} of body mass) trigger only mild and non-lethal inflammation [60,76,77]. However, most immunological data for birds have so far only been generated in poultry (evolutionarily basal Galloanserae lineage), or, to a much lesser extent, in passerine birds. Thus, diversity in avian immune responses to peripheral stimulation remains largely unknown. Of particular relevance is the immune response regulation in species with highly rearranged genomes, such as the parrots [29,31].

Peripheral inflammation can modulate the expression of *CNRs* in both the periphery and the brain, thereby altering neuronal processes and behavioural and cognitive functions [78]. We confirmed *CNR1* expression in the nervous system of birds (both zebra finches and parrots), suggesting its similar regulatory effect on neuronal processes as in mammals. In mammals, leucocyte-modulating *CNR2*, an inhibitor of the proinflammatory cytokine secretion [79], is also expressed in both the brain (microglia) and periphery [45], providing an important anti-neuroinflammatory protection to the brain [18–21]. However, somewhat surprisingly, previous radiographic investigations have revealed no signs of its expression in the brain of budgerigars [80]. Our genome-database search indicated a complete absence of functional *CNR2* genes in all parrot species investigated, which contrasts with its conserved presence in all lineages closely related to parrots (i.e. the falcons, seriemas and passerines, including the zebra finch). We were able to identify putative remnants of the *CNR2* pseudogene in the budgerigar genome, indicating apparent *CNR2* pseudogenization following massive karyotype rearrangements early in parrot phylogeny [29,31]. Interestingly, a comparison of the karyotype localization of passerine *CNR2*-neighbouring genes in the budgerigar and kakapo genomes suggested two presumably independent karyotype rearrangement events in parrots resulting in the *CNR2* loss. The absence of *CNR2* was confirmed through negative results for (i) *CNR2*-targeted amplification attempts in budgerigar gDNA using conserved PCR primers, and (ii) searches through Illumina NextSeq-generated transcriptomes of small intestine in six different parrot species. We consider this as a conclusive support for the complete absence of the *CNR2* gene in parrots, although further research should aim to support this finding on the chromosomal level.

This finding raises the question as to whether a pseudogenization event could have affected the regulation of neuroimmune interactions in parrots. Our positive selection

analysis indicates that CNR2 is functionally conserved across the avian taxa. As we found no other *CNR* gene in the parrot genomes aside from *CNR1*, we tested for evolutionary changes in *CNR1* that could be linked to *CNR2* absence. Nevertheless, our selection analysis showed that *CNR1* is also highly conserved throughout vertebrates, with no compensatory selection linked to the *CNR2* loss in parrots. This suggests that *CNR2* pseudogenization could have functional significance. To test this hypothesis, we compared data on systemic inflammation in passerines and parrots, focusing on the putative difference in neuroinflammation-linked cytokine expression caused by the lack of the *CNR2* negative regulation in parrots [45,81]. By contrast to the zebra finch, in the *CNR2*-deficient budgerigars, we observed upregulation of expression in proinflammatory cytokines such as *IL1B* and *IL6* in the hyperpallial tissue. The same pattern has been detected across all investigated parrot species, suggesting that parrots in general may be more vulnerable to neuroinflammation than other birds. This is supported by the fact that parrots are exceptionally susceptible to bornavirus-related neuropathy [8,82–84] and also other parrot pathogens including bacteria, viruses and fungi are suspected to frequently cause behavioural disorders [85–87].

Our data, therefore, suggest that *CNR2* loss in parrots could impair regulation which dampens systemic proinflammatory signalling (for example, mediated by *IL1B* and *IL6*). Evidence from *CNR2*-knock-out mice showing pronounced immunopathology [88], appears to support our interpretation. Thus, our results promote the hypothesis of regulatory relevance of *CNR2* absence in sensitivity to neuroinflammation and also suggest that parrots could be prone to neurological syndromes.

5. Conclusion

In this study, we provide comprehensive evidence for *CNR2* absence in parrots and initial results documenting the possible impact of this loss on the regulation of neuroinflammation. Specifically, we observed upregulated proinflammatory cytokine expression in parrot brains, but no similar changes in zebra finches possessing fully functional *CNR2*. With no apparent compensatory evolution in *CNR1*, parrots lacking functional *CNR2* may be more

susceptible to systemic neuroinflammation (e.g. induced by dysbiosis) than other avian species. Our findings do not only provide important insights into variability in susceptibility to immunopathology between species but also offer relevant evolutionary evidence for the functional effects of gene loss events during chromosomal rearrangements. Further research is needed to illuminate possible compensatory mechanisms in parrot immunity, and links to parrot infection ecology and evolution.

Ethics. The research was approved by the Ethical Committee of Charles University, Faculty of Science (permits 13882/2011-30 and MSMT-30397/2019-5) and was carried out in accordance with the current laws of the Czech Republic and the European Union.

Data accessibility. All raw and processed sequencing data generated in this study have been submitted to the NCBI (<https://www.ncbi.nlm.nih.gov/>) under accession numbers SAMN23963146, SAMN23963147, SAMN23963148, SAMN23963149, SAMN23963150, SAMN23963151, PRJNA751848, PRJNA879979.

The data are provided in the electronic supplementary material [89].

Authors' contributions. D.D.: conceptualization, data curation, formal analysis, investigation, project administration, visualization, writing—original draft, writing—review and editing; M.G.S.: formal analysis, investigation, writing—review and editing; N.K.V.: data curation, formal analysis, investigation, writing—review and editing; E.V.: formal analysis, investigation, writing—review and editing; Z.Š.: investigation, writing—review and editing; T.K.: investigation, project administration, writing—review and editing; M.T.: data curation, formal analysis, investigation, writing—review and editing; V.B.: methodology, writing—review and editing; D.E.: methodology, writing—review and editing; O.B.: methodology, writing—review and editing; M.V.: conceptualization, funding acquisition, investigation, methodology, project administration, supervision, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. This study was supported by Grant Schemes at Charles University (grant nos. GAUK 646119, PRIMUS/17/SCI/12 and START/SCI/113 with reg. no. CZ.02.2.69/0.0/0.0/19_073/0016935), the Czech Science Foundation (grant no. P502/19-20152Y) and Institutional Research Support (grant no. 260571/2022). Computational resources were supplied by the project 'e-Infrastruktura CZ' (e-INFRA CZ LM2018140) supported by the Ministry of Education, Youth and Sports of the Czech Republic.

Acknowledgements. We are grateful to Balraj Melepat and Tao Li for their help in the laboratory, and to Kevin Roche for language correction.

References

- Ferrari AJ, Charlson FJ, Norman RE, Patten SB, Freedman G, Murray CJL, Vos T, Whiteford HA. 2013 Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS Med.* **10**, e1001547. (doi:10.1371/journal.pmed.1001547)
- Koopman M, El Aidy S. 2017 Depressed gut? The microbiota-diet-inflammation triad in depression. *Curr. Opin Psychiatry* **30**, 369–377. (doi:10.1097/YCO.0000000000000350)
- Gaskins LA, Bergman L. 2011 Surveys of avian practitioners and pet owners regarding common behavior problems in psittacine birds. *J. Avian Med. Surg.* **25**, 111–118. (doi:10.1647/2010-027.1)
- Kinkaid HYM, Mills DS, Nichols SG, Meagher RK, Mason GJ. 2013 Feather-damaging behaviour in companion parrots: an initial analysis of potential demographic risk factors. *Avian Biol. Res.* **6**, 289–296. (doi:10.3184/175815513X13803574144572)
- Ebisawa K, Nakayama S, Pai C, Kinoshita R, Koie H. 2021 Prevalence and risk factors for feather-damaging behavior in psittacine birds: analysis of a Japanese nationwide survey. *PLoS ONE* **16**, e0254610. (doi:10.1371/journal.pone.0254610)
- van Zeeland YRA, Spruit BM, Rodenburg TB, Riedstra B, van Hierden YM, Buitenhuis B, Korte SM, Lumeij JT. 2009 Feather damaging behaviour in parrots: a review with consideration of comparative aspects. *Appl. Anim. Behav. Sci.* **121**, 75–95. (doi:10.1016/j.applanim.2009.09.006)
- Péron F, Grosset C. 2014 The diet of adult psittacids: veterinarian and ethological approaches. *J. Anim. Physiol. Anim. Nutr.* **98**, 403–416. (doi:10.1111/jpn.12103)
- Chen J, Guo J, Tian Y, Tizard I. 2020 The pathogenesis of proventricular dilatation disease caused by parrot bornaviruses: a possible role for neuropeptide Y (Npy). In review (cited 2 November 2020). See <https://www.researchsquare.com/article/rs-49954/v1>.

9. Swardfager W, Rosenblat JD, Benlamri M, McIntyre RS. 2016 Mapping inflammation onto mood: inflammatory mediators of anhedonia. *Neurosci. Biobehav. Rev.* **64**, 148–166. (doi:10.1016/j.neubiorev.2016.02.017)
10. Kumar H, Kawai T, Akira S. 2011 Pathogen recognition by the innate immune system. *Int. Rev. Immunol.* **30**, 16–34. (doi:10.3109/08830185.2010.529976)
11. Aguilera M, Vergara P, Martínez V. 2013 Stress and antibiotics alter luminal and wall-adhered microbiota and enhance the local expression of visceral sensory-related systems in mice. *Neurogastroenterol. Motil.* **25**, e515–e529. (doi:10.1111/nmo.12154)
12. Vanner SJ, Greenwood-Van Meerveld B, Mawe GM, Shea-Donohue T, Verdu EF, Wood J, Grundy D. 2016 Fundamentals of neurogastroenterology: basic science. *Gastroenterology* **150**, 1280–1291. (doi:10.1053/j.gastro.2016.02.018)
13. Acharya N, Penukonda S, Shcheglova T, Hagymasi AT, Basu S, Srivastava PK. 2017 Endocannabinoid system acts as a regulator of immune homeostasis in the gut. *Proc. Natl Acad. Sci. USA* **114**, 5005–5010. (doi:10.1073/pnas.1612177114)
14. Rider P, Carmi Y, Guttman O, Braiman A, Cohen I, Voronov E, White MR, Dinarello CA, Apte RN. 2011 IL-1 α and IL-1 β recruit different myeloid cells and promote different stages of sterile inflammation. *J. Immunol.* **187**, 4835–4843. (doi:10.4049/jimmunol.1102048)
15. DiSabato DJ, Quan N, Godbout JP. 2016 Neuroinflammation: the devil is in the details. *J. Neurochem.* **139**, 136–153. (doi:10.1111/jnc.13607)
16. Bird S, Zou J, Wang T, Munday B, Cunningham C, Secombes CJ. 2002 Evolution of interleukin-1 β . *Cytokine Growth Factor Rev.* **13**, 483–502. (doi:10.1016/S1359-6101(02)00028-X)
17. Ren K, Torres R. 2009 Role of interleukin-1 β during pain and inflammation. *Brain Res. Rev.* **60**, 57–64. (doi:10.1016/j.brainresrev.2008.12.020)
18. Klegeris A, Bissonnette CJ, McGeer PL. 2003 Reduction of human monocytic cell neurotoxicity and cytokine secretion by ligands of the cannabinoid-type CB2 receptor. *Br. J. Pharmacol.* **139**, 775–786. (doi:10.1038/sj.bjp.0705304)
19. Domenici MR. 2006 Cannabinoid receptor type 1 located on presynaptic terminals of principal neurons in the forebrain controls glutamatergic synaptic transmission. *J. Neurosci.* **26**, 5794–5799. (doi:10.1523/JNEUROSCI.0372-06.2006)
20. Solas M, Francis PT, Franco R, Ramirez MJ. 2013 CB2 receptor and amyloid pathology in frontal cortex of Alzheimer's disease patients. *Neurobiol. Aging* **34**, 805–808. (doi:10.1016/j.neurobiolaging.2012.06.005)
21. Tao Y *et al.* 2016 Cannabinoid receptor-2 stimulation suppresses neuroinflammation by regulating microglial M1/M2 polarization through the cAMP/PKA pathway in an experimental GMH rat model. *Brain Behav. Immun.* **58**, 118–129. (doi:10.1016/j.bbi.2016.05.020)
22. Emery NJ. 2006 Cognitive ornithology: the evolution of avian intelligence. *Phil. Trans. R. Soc. B* **361**, 23–43. (doi:10.1098/rstb.2005.1736)
23. Olkowicz S, Kocourek M, Lučan RK, Porteš M, Fitch WT, Herculano-Houzel S, Némec P. 2016 Birds have primate-like numbers of neurons in the forebrain. *Proc. Natl Acad. Sci. USA* **113**, 7255–7260. (doi:10.1073/pnas.1517131113)
24. Wang X, Grus WE, Zhang J. 2006 Gene losses during human origins. *PLoS Biol.* **4**, e52. (doi:10.1371/journal.pbio.0040052)
25. Olson MV. 1999 When less is more: gene loss as an engine of evolutionary change. *Am. J. Hum. Genet.* **64**, 18–23. (doi:10.1086/302219)
26. Charlesworth B. 2012 The effects of deleterious mutations on evolution at linked sites. *Genetics* **190**, 5–22. (doi:10.1534/genetics.111.134288)
27. Damas J, Corbo M, Lewin HA. 2021 Vertebrate chromosome evolution. *Annu. Rev. Anim. Biosci.* **9**, 1–27. (doi:10.1146/annurev-animal-020518-114924)
28. Huang Z *et al.* 2022 Recurrent chromosome reshuffling and the evolution of neo-sex chromosomes in parrots. *Nat. Commun.* **13**, 944. (doi:10.1038/s41467-022-28585-1)
29. Nanda I, Karl E, Griffin DK, Scharlt M, Schmid M. 2007 Chromosome repatterning in three representative parrots (Psittaciformes) inferred from comparative chromosome painting. *Cytogenet Genome Res.* **117**, 43–53. (doi:10.1159/000103164)
30. Harewood L, Fraser P. 2014 The impact of chromosomal rearrangements on regulation of gene expression. *Hum. Mol. Genet.* **23**, R76–R82. (doi:10.1093/hmg/ddu278)
31. Furo IDO, Kretschmer R, O'Brien PCM, Pereira JC, Garnero ADV, Gunski RJ, Ferguson-Smith MA, De Oliveira EH. 2018 Chromosome painting in neotropical long- and short-tailed parrots (Aves, Psittaciformes): phylogeny and proposal for a putative ancestral karyotype for tribe arini. *Genes* **9**, 491. (doi:10.3390/genes9100491)
32. Temperley ND, Berlin S, Paton IR, Griffin DK, Burt DW. 2008 Evolution of the chicken Toll-like receptor gene family: a story of gene gain and gene loss. *BMC Genom.* **9**, 62. (doi:10.1186/1471-2164-9-62)
33. Bainová H, Králová T, Bryjová A, Albrecht T, Bryja J, Vinkler M. 2014 First evidence of independent pseudogenization of Toll-like receptor 5 in passerine birds. *Dev. Comp. Immunol.* **45**, 151–155. (doi:10.1016/j.dci.2014.02.010)
34. van der Loo W, Magalhaes MJ, de Matos AL, Abrantes J, Yamada F, Esteves PJ. 2016 Adaptive gene loss? Tracing back the pseudogenization of the rabbit CCL8 chemokine. *J. Mol. Evol.* **83**, 12–25. (doi:10.1007/s00239-016-9747-7)
35. Velová H, Gutowska-Ding MW, Burt DW, Vinkler M. 2018 Toll-like receptor evolution in birds: gene duplication, pseudogenization, and diversifying selection. *Mol. Biol. Evol.* **35**, 2170–2184. (doi:10.1093/molbev/msy119)
36. Lu HC, Mackie K. 2016 An introduction to the endogenous cannabinoid system. *Biol. Psychiatry* **79**, 516–525. (doi:10.1016/j.biopsych.2015.07.028)
37. Elphick MR. 2012 The evolution and comparative neurobiology of endocannabinoid signalling. *Phil. Trans. R. Soc. B* **367**, 3201–3215. (doi:10.1098/rstb.2011.0394)
38. Gallègue S *et al.* 1995 Expression of central and peripheral cannabinoid receptors in human immune tissues and leukocyte subpopulations. *Eur. J. Biochem.* **232**, 54–61. (doi:10.1111/j.1432-1033.1995.tb20780.x)
39. Carlisle SJ, Marciano-Cabral F, Staab A, Ludwick C, Cabral GA. 2002 Differential expression of the CB2 cannabinoid receptor by rodent macrophages and macrophage-like cells in relation to cell activation. *Int. Immunopharmacol.* **2**, 69–82. (doi:10.1016/S1567-5769(01)00147-3)
40. Soderstrom K, Johnson F. 2000 CB1 cannabinoid receptor expression in brain regions associated with zebra finch song control. *Brain Res.* **857**, 151–157. (doi:10.1016/S0006-8993(99)02393-8)
41. Soderstrom K, Johnson F. 2003 Cannabinoid exposure alters learning of zebra finch vocal patterns. *Brain Res. Dev. Brain Res.* **142**, 215–217. (doi:10.1016/S0165-3806(03)00061-0)
42. Cristino L, Becker T, Di Marzo V. 2014 Endocannabinoids and energy homeostasis: an update: regulatory role of endocannabinoids in obesity. *Biofactors* **40**, 389–397. (doi:10.1002/biof.1168)
43. Cani PD, Plovier H, Van Hul M, Geurts L, Delzenne NM, Druart C, Everard A. 2016 Endocannabinoids — at the crossroads between the gut microbiota and host metabolism. *Nat. Rev. Endocrinol.* **12**, 133–143. (doi:10.1038/nrendo.2015.211)
44. Greenwood-Van Meerveld B (ed.) 2017 *Gastrointestinal pharmacology. Handbook of experimental pharmacology, vol. 239*. Cham, Switzerland: Springer International Publishing (cited 13 February 2019). See <http://link.springer.com/10.1007/978-3-319-56360-2>.
45. Maresz K, Carrier EJ, Ponomarev ED, Hillard CJ, Dittel BN. 2005 Modulation of the cannabinoid CB2 receptor in microglial cells in response to inflammatory stimuli. *J. Neurochem.* **95**, 437–445. (doi:10.1111/j.1471-4159.2005.03380.x)
46. Vincent L, Vang D, Nguyen J, Benson B, Lei J, Gupta K. 2016 Cannabinoid receptor-specific mechanisms to alleviate pain in sickle cell anemia via inhibition of mast cell activation and neurogenic inflammation. *Haematologica* **101**, 566–577. (doi:10.3324/haematol.2015.136523)
47. Krustev E, Muley MM, McDougall JJ. 2017 Endocannabinoids inhibit neurogenic inflammation in murine joints by a non-canonical cannabinoid receptor mechanism. *Neuropeptides* **64**, 131–135. (doi:10.1016/j.npep.2016.08.007)
48. Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S, AmiGO Hub, Web Presence Working Group. 2009 AmiGO: online access to ontology and annotation data. *Bioinformatics* **25**, 288–289. (doi:10.1093/bioinformatics/btn615)

49. Smedley D, Haider S, Ballester B, Holland R, London D, Thorisson G, Kasprzyk A. 2009 BioMart—biological queries made easy. *BMC Genom.* **10**, 22. (doi:10.1186/1471-2164-10-22)
50. Mueller RC, Mallig N, Smith J, Eöry L, Kuo RI, Kraus RHS. 2020 Avian immunome DB: an example of a user-friendly interface for extracting genetic information. *BMC Bioinf.* **21**, 502. (doi:10.1186/s12859-020-03764-3)
51. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018 MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547–1549. (doi:10.1093/molbev/msy096)
52. Letunic I, Bork P. 2021 Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res.* **49**, W293–W296. (doi:10.1093/nar/gkab301)
53. Glaser F, Pupko T, Paz I, Bell RE, Bechor-Shental D, Martz E, Ben-Tal N. 2003 ConSurf: identification of functional regions in proteins by surface-mapping of phylogenetic information. *Bioinformatics* **19**, 163–164. (doi:10.1093/bioinformatics/19.1.163)
54. Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Kosakovsky Pond SL, Scheffler K. 2013 FUBAR: a fast, unconstrained Bayesian approximation for inferring selection. *Mol. Biol. Evol.* **30**, 1196–1205. (doi:10.1093/molbev/mst030)
55. Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Kosakovsky Pond SL. 2012 Detecting individual sites subject to episodic diversifying selection. *PLoS Genet.* **8**, e1002764. (doi:10.1371/journal.pgen.1002764)
56. Smith MD, Wertheim JO, Weaver S, Murrell B, Scheffler K, Kosakovsky Pond SL. 2015 Less is more: an adaptive branch-site random effects model for efficient detection of episodic diversifying selection. *Mol. Biol. Evol.* **32**, 1342–1353. (doi:10.1093/molbev/msv022)
57. Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. 2015 RELAX: detecting relaxed selection in a phylogenetic framework. *Mol. Biol. Evol.* **32**, 820–832. (doi:10.1093/molbev/msu400)
58. Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. 2012 Predicting the functional effect of amino acid substitutions and indels. *PLoS ONE* **7**, e46688. (doi:10.1371/journal.pone.0046688)
59. Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC. 2016 SIFT missense predictions for genomes. *Nat. Protoc.* **11**, 1–9. (doi:10.1038/nprot.2015.123)
60. Wegmann M, Voegeli B, Richner H. 2015 Parasites suppress immune-enhancing effect of methionine in nestling great tits. *Oecologia* **177**, 213–221. (doi:10.1007/s00442-014-3138-9)
61. Grabherr MG *et al.* 2011 Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* **29**, 644–652. (doi:10.1038/nbt.1883)
62. Haas BJ, Papanicolaou A. 2020 TransDecoder (cited 12 April 2021). See <https://github.com/jls943/TransDecoder>.
63. Fu L, Niu B, Zhu Z, Wu S, Li W. 2012 CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* **28**, 3150–3152. (doi:10.1093/bioinformatics/bts565)
64. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015 BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210–3212. (doi:10.1093/bioinformatics/btv351)
65. Zhang Z, Schwartz S, Wagner L, Miller W. 2000 A greedy algorithm for aligning DNA sequences. *J. Comput. Biol.* **7**, 203–214. (doi:10.1089/10665270050081478)
66. Moll P, Ante M, Seitz A, Reda T. 2014 QuantSeq 3' mRNA sequencing for RNA quantification. *Nat. Methods* **11**, i–iii. (doi:10.1038/nmeth.f.376)
67. Bean AGD, Lowenthal JW. 2022 Chapter 9 - avian cytokines and their receptors. In *Avian immunology* (eds B Kaspers, KA Schat, TW Göbel, L Vervelde), pp. 249–276, 3rd edn. Boston, MA: Academic Press. (Cited 28 April 2022). See <https://www.sciencedirect.com/science/article/pii/B9780128187081000245>.
68. Kearse M *et al.* 2012 Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647. (doi:10.1093/bioinformatics/bts199)
69. Vinkler M, Leon AE, Kirkpatrick L, Dalloul RA, Hawley DM. 2018 Differing house finch cytokine expression responses to original and evolved isolates of *Mycoplasma gallisepticum*. *Front. Immunol.* **9**, 13. (doi:10.3389/fimmu.2018.00013/full) [cited 2020 Mar 7]
70. Pfaffl MW. 2001 A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **29**, e45. (doi:10.1093/nar/29.9.e45)
71. RStudio. Open source & professional software for data science teams (Internet). (Cited 17 January 2022). See <https://rstudio.com>.
72. Ban E. 1992 Brain interleukin 1 gene expression induced by peripheral lipopolysaccharide administration. *Cytokine* **4**, 48–54. (doi:10.1016/1043-4666(92)90036-Q)
73. Sköld-Chiriac S, Nord A, Nilsson JÅ, Hasselquist D. 2014 Physiological and behavioral responses to an acute-phase response in zebra finches: immediate and short-term effects. *Physiol. Biochem. Zool.* **87**, 288–298. (doi:10.1086/674789)
74. Batista CRA, Gomes GF, Candelario-Jalil E, Fiebach BL, de Oliveira ACP. 2019 Lipopolysaccharide-induced neuroinflammation as a bridge to understand neurodegeneration. *Int. J. Mol. Sci.* **20**, 2293. (doi:10.3390/ijms20092293)
75. Jurgens HA, Amancherla K, Johnson RW. 2012 Influenza infection induces neuroinflammation, alters hippocampal neuron morphology, and impairs cognition in adult mice. *J. Neurosci.* **32**, 3958–3968. (doi:10.1523/JNEUROSCI.6389-11.2012)
76. Vermeulen A, Eens M, Zaid E, Müller W. 2016 Baseline innate immunity does not affect the response to an immune challenge in female great tits (*Parus major*). *Behav. Ecol. Sociobiol.* **70**, 585–592. (doi:10.1007/s00265-016-2077-3)
77. Armour EM, Bruner TL, Hines JK, Butler MW. 2020 Low-dose immune challenges result in detectable levels of oxidative damage. *J. Exp. Biol.* **223**, jeb220095. (doi:10.1242/jeb.220095)
78. Procaccini C, Pucino V, De Rosa V, Marone G, Matarese G. 2014 Neuro-endocrine networks controlling immune system in health and disease. *Front. Immunol.* **6**, 143. (doi:10.3389/fimmu.2014.00143/abstract) [cited 2019 Oct 21]
79. Hasenoehrl C, Taschler U, Storr M, Schicho R. 2016 The gastrointestinal tract - a central organ of cannabinoid signaling in health and disease. *Neurogastroenterol. Motil.* **28**, 1765–1780. (doi:10.1111/nmo.12931)
80. Alonso-Ferrero ME, Paniagua MA, Mostany R, Pilar-Cuellar F, Diez-Alarcia R, Pazos A, Fernández-López A. 2006 Cannabinoid system in the budgerigar brain. *Brain Res.* **1087**, 105–113. (doi:10.1016/j.brainres.2006.02.119)
81. Klein TW, Newton C, Larsen K, Lu L, Perkins I, Nong L, Friedman H. 2003 The cannabinoid system and immune modulation. *J. Leukoc. Biol.* **74**, 486–496. (doi:10.1189/jlb.0303101)
82. Rinder M, Ackermann A, Kempf H, Kaspers B, Korbel R, Staeheli P. 2009 Broad tissue and cell tropism of avian bornavirus in parrots with proventricular dilatation disease. *J. Virol.* **83**, 5401–5407. (doi:10.1128/JVI.00133-09)
83. Staeheli P, Rinder M, Kaspers B. 2010 Avian bornavirus associated with fatal disease in psittacine birds. *J. Virol.* **84**, 6269–6275. (doi:10.1128/JVI.02567-09)
84. Rubbenstroth D, Rinder M, Kaspers B, Staeheli P. 2012 Efficient isolation of avian bornaviruses (ABV) from naturally infected psittacine birds and identification of a new ABV genotype from a salmon-crested cockatoo (*Cacatua moluccensis*). *Vet. Microbiol.* **161**, 36–42. (doi:10.1016/j.vetmic.2012.07.004)
85. Rubinstein J, Lightfoot T. 2012 Feather loss and feather destructive behavior in pet birds. *J. Exot. Pet. Med.* **21**, 219–234. (doi:10.1053/j.jepm.2012.06.024)
86. Speer B. 2014 Normal and abnormal parrot behavior. *J. Exot. Pet. Med.* **23**, 230–233. (doi:10.1053/j.jepm.2014.06.009)
87. Dovč A *et al.* 2016 Occurrence of bacterial and viral pathogens in common and noninvasive diagnostic sampling from parrots and racing pigeons in Slovenia. *Avian Dis.* **60**, 487–492. (doi:10.1637/11373-011116-Reg)
88. Karmaus PWF, Chen W, Crawford R, Kaplan BLF, Kaminski NE. 2013 Δ 9-Tetrahydrocannabinol impairs the inflammatory response to influenza infection: role of antigen-presenting cells and the cannabinoid receptors 1 and 2. *Toxicol. Sci.* **131**, 419–433. (doi:10.1093/toxsci/kfs315)
89. Divin D *et al.* 2022 Cannabinoid receptor 2 evolutionary gene loss makes parrots more susceptible to neuroinflammation. Figshare. (doi:10.6084/m9.figshare.c.6307543)