



Published in final edited form as:

Sci Signal. ; 13(630): . doi:10.1126/scisignal.aaz1854.

Inhibition of Hsp90 in the spinal cord enhances the antinociceptive effects of morphine by activating an ERK-RSK pathway

David I. Duron¹, Wei Lei¹, Natalie K. Barker², Carrie Stine¹, Sanket Mishra³, Brian S. J. Blagg³, Paul R. Langlais², John M. Streicher^{1,*}

¹Department of Pharmacology, College of Medicine, University of Arizona, Tucson, AZ 85724, USA.

²Division of Endocrinology, Department of Medicine, College of Medicine, University of Arizona, Tucson, AZ 85724, USA.

³Department of Chemistry and Biochemistry, College of Science, University of Notre Dame, Notre Dame, IN 46556, USA.

Abstract

Morphine and other opioids are commonly used to treat pain despite their numerous adverse side effects. Modulating μ -opioid receptor (MOR) signaling is one way to potentially improve opioid therapy. In mice, the chaperone protein Hsp90 mediates MOR signaling within the brain. Here, we found that inhibiting Hsp90 specifically in the spinal cord enhanced the antinociceptive effects of morphine in mice. Intrathecal, but not systemic, administration of the Hsp90 inhibitors 17-AAG or KU-32 amplified the effects of morphine in suppressing sensitivity to both thermal and mechanical stimuli in mice. Hsp90 inhibition enabled opioid-induced phosphorylation of the kinase ERK and increased abundance of the kinase RSK in the dorsal horns of the spinal cord, which are heavily populated with primary afferent sensory neurons. The additive effects of Hsp90 inhibition were abolished upon intrathecal inhibition of ERK, RSK, or protein synthesis. This mechanism downstream of MOR, localized to the spinal cord and repressed by Hsp90, may potentially be used to enhance the efficacy and presumably decrease the side effects of opioid therapy.

INTRODUCTION

Currently available therapeutics for the treatment of chronic pain are largely limited by their efficacy and undesired side effects. With more than 100 million individuals affected and an

*Corresponding author. jstreicher@email.arizona.edu.

Author contributions: D.I.D. performed and analyzed most of the experiments, participated in project and experimental design, and cowrote the manuscript. W.L. performed and analyzed the systemic behavioral experiments. N.K.B. performed and analyzed the quantitative proteomic experiments. C.S. performed and analyzed some of the costaining IHC experiments. S.M. synthesized and purified KU-32. B.S.J.B. supervised S.M. in the performance of the chemical synthesis and participated in project design. P.R.L. supervised N.K.B. in the performance of the proteomic experiments and participated in experimental and project design. J.M.S. supervised D.I.D., W.L., and C.S. in the performance of their work, secured funding for the project, conceived the initial idea of the project, participated in experimental and project design, and cowrote the manuscript. All authors had editorial input into the manuscript.

Competing interests: The authors declare that they have no competing interests.

economic burden exceeding \$600 billion in the United States alone, chronic pain remains an area of critical and growing medical need (1, 2). One of the more efficacious treatment options are opioid analgesics, such as morphine. Although these drugs can be very effective acutely, their side effects such as tolerance, addiction, and respiratory depression make them a high-risk choice when dealing with long-term medication regimens (3, 4). Accompanying these negative side effects is a growing social awareness of the potential dangers of opioids that have begun to negatively stigmatize their use, abetted by a growing opioid abuse and addiction crisis (5).

Intensive decades-long research has revealed a complex signaling network evoked by opioid treatment downstream of the μ -opioid receptor (MOR) (6). Increased understanding of the complexity of MOR signal transduction has resulted in new efforts for drug discovery and development, such as biased agonism to reduce the side effects of opioids (7–9). These efforts have produced new biased ligands, as well as additional drugs targeting key proteins, such as the kinase mammalian target of rapamycin, or MOR signaling–relevant receptors, such as protease-activated receptor 2, to either augment or reduce key behavioral outputs, such as antinociception (10–14). These efforts illuminate the relevant MOR signaling cascades beyond the “classical” $G\alpha_i$ cascade and show the value in elucidating key downstream signaling regulators.

Heat shock protein 90 (Hsp90) is a molecular chaperone protein that is up-regulated in response to stress. It regulates its client proteins through several molecular mechanisms, including protein folding, kinase modulation, protein complex formation, and subcellular localization (15, 16). Hsp90 makes up roughly 2% of the total protein pool in a given cell, highlighting its centrality to cell biology. Its functions have primarily been investigated in the context of cancer (17–19), but Hsp90 has been shown to have a key role in regulating signal transduction at the receptor and downstream cascade levels in a number of different tissues and physiological contexts (20).

In earlier work, we tested the hypothesis that Hsp90 could play a key role in MOR signal transduction by selectively inhibiting Hsp90 in the brain using intracerebroventricular administration of 17-*N*-allylamino-17-demethoxygeldanamycin (17-AAG). In that study, we found that brain Hsp90 inhibition completely suppressed the antinociceptive effects of systemically administered morphine in various murine pain models (21). In addition, we showed that intracerebroventricular administration of 17-AAG blocked the phosphorylation of extracellular signal–regulated kinase (ERK) in response to the selective MOR agonist and synthetic opioid peptide DAMGO ([D-Ala², *N*-MePhe⁴, Gly-ol]-enkephalin) and that this loss of ERK phosphorylation was responsible for the loss of morphine antinociception. Those findings demonstrated that Hsp90 promotes MOR signaling in the brain and identified new signaling pathways to explore further. However, that work left many mechanistic details unknown, as well as the contribution of other regions of the central nervous system (CNS) like the spinal cord, which may be more applicable clinically because of the potential for intrathecal drug delivery.

Here, we investigated Hsp90 modulation of MOR signal transduction in the spinal cord in mice. Contrary to what was observed in our previous study in the brain, Hsp90 inhibition

in the spinal cord amplified the antinociceptive effects of morphine. We further identified a molecular mechanism for this effect within the spinal cord dorsal horn. Our findings suggest a potential opioid dose reduction strategy via spinal Hsp90 inhibition to minimize the negative side effects of opioids while maintaining their analgesic benefits.

RESULTS

Inhibition of Hsp90 in the spinal cord enhances morphine-induced antinociception

We previously showed that intracerebroventricularly administered Hsp90 inhibitors completely ablated morphine-induced antinociception in multiple pain models (21). In addition, Hsp90 has a considerable number of client proteins, which differ in various tissue, cellular, and environmental contexts (22–24). This suggests the potential for context-specific roles for Hsp90 within downstream MOR signaling. We sought to test this role for Hsp90 in MOR signaling by considering the contribution of Hsp90 to morphine-induced antinociception within the spinal cord. CD-1 mice were treated with intrathecally administered 17-AAG, a geldanamycin derivative that competitively binds the N-terminal adenosine triphosphate (ATP) binding domain of Hsp90. Twenty-four hours after injection, mice were then treated with morphine [3.2 mg/kg, subcutaneously], and behavioral pain assays were performed.

Contrary to our previous report with intracerebroventricularly administered 17-AAG, we found that spinally inhibited Hsp90 resulted in an increased antinociceptive response (meaning, less pain-evoked sensitivity) due to morphine in both thermal tail-flick and mechanical post-operative paw incision pain models (Fig. 1, A and B). To verify the Hsp90 selectivity of these results, we used a C-terminal inhibitor of Hsp90, KU-32, that binds to an alternate site than 17-AAG and thus is unlikely to share off-target interactions (25, 26). KU-32 was administered to the spinal cord, followed by subcutaneous morphine 24 hours later. Enhanced morphine-induced antinociception was also observed with KU-32 treatment, confirming the Hsp90 selectivity of our results (Fig. 1C).

We next confirmed that these findings were not due to off-target motor or sedative effects using the rotarod test. Spinal 17-AAG treatment had no impact on rotarod performance in the mice, suggesting that our findings reflect bona fide changes to the opioid pain modulatory system (Fig. 1D). Last, in our previous report, we tested the brain role of Hsp90 in tail-flick pain using intracerebroventricular DAMGO instead of morphine, showing that inhibition of brain Hsp90 had no impact on tail-flick pain (21). To confirm that our tail-flick results here were due to changes in Hsp90 location (brain versus spinal cord) rather than drug and route (intracerebroventricular DAMGO versus subcutaneous morphine), we tested intracerebroventricular 17-AAG combined with subcutaneous morphine. We found that at both 3.2 and 10 mg/kg (subcutaneously), morphine had no impact on tail-flick response (Fig. 1E), the same as for intracerebroventricular DAMGO in our earlier study (21).

Inhibition of Hsp90 in the brain overrides inhibition in the spinal cord

With behavioral differences identified in brain versus spinal cord Hsp90 inhibition, we next tested the interaction of these two regions using systemic Hsp90 inhibition, which

would affect both brain and spinal cord. To do this, we injected mice with intraperitoneal 17-AAG and assessed its effects on morphine-induced antinociception after a 24-hour period. We tested for increased Hsp70 expression levels as a marker of Hsp90 inhibition within the periaqueductal gray (PAG) brain region and spinal cord tissue after 24 hours with intraperitoneal 17-AAG using Western blot (21). We found that Hsp70 levels were increased in the PAG as expected, validating our treatment regimen as effective in inhibiting CNS Hsp90 (Fig. 2, A and B). Unexpectedly, we were unable to detect an increase in spinal cord Hsp70 (Fig. 2, A and B). 17-AAG was likely reaching the spinal cord because we have shown that it can reach the PAG; thus, this result may represent different molecular mechanisms for Hsp90 in the brain versus the spinal cord. Now validated, we tested the impact of systemic 17-AAG on morphine antinociception. We observed that systemic delivery of 17-AAG had no impact on antinociception in the tail-flick pain model and markedly reduced antinociception in the paw incision pain model (Fig. 2, C and D); these results are very similar to what was observed with intracerebroventricular 17-AAG treatment previously (21) and here (Fig. 1E).

The effects on morphine antinociception seen with systemic Hsp90 inhibition suggest that the signaling events within the brain may override that of the spinal cord. To directly test this hypothesis and rule out peripheral mechanisms, we performed dual intracerebroventricular and intrathecal injections of 17-AAG. We found that dual brain and spinal cord injections recapitulated systemic injection in both the tail-flick and paw incision pain models (Fig. 2, E and F). These results suggest that the signaling events regulated by Hsp90 within the brain override MOR signaling within the spinal cord, which would otherwise allow for amplified pain relief in these models.

ERK mitogen-activated protein kinase signaling is activated by spinal Hsp90 inhibition and mediates enhanced antinociception

Our previous study within the brain demonstrated that blocked activation of ERK mitogen-activated protein kinase (MAPK) in the PAG by 17-AAG treatment is a mechanism for the reduction in morphine-induced antinociception (21). We thus tested ERK signaling activation within the spinal cord after intrathecal 17-AAG and DAMGO (selective MOR agonist) treatment using Western blot. DAMGO was used as a high-efficacy selective agonist, increasing our ability to observe kinase changes in tissue versus the partial agonist morphine; our results above (Fig. 1E) and experiments below validate this choice. DAMGO alone showed no ERK activation relative to vehicle treatment; in contrast, 17-AAG induced an elevated ERK baseline with a further increase in ERK phosphorylation when combined with DAMGO (Fig. 3, A and B). As in our systemic inhibition studies above, we also sought to confirm Hsp90 inhibition by 17-AAG by testing for Hsp70 up-regulation (21, 27, 28). We again found no Hsp70 up-regulation, even with direct intrathecal injection of inhibitor, confirming our systemic results above and further suggesting that Hsp90 molecular mechanisms may differ in the spinal cord versus the brain (Fig. 3, A and C).

To localize the observed increases in ERK phosphorylation within the spinal cord, we performed immunohistochemical (IHC) analysis of spinal cord tissue from mice treated with intrathecal 17-AAG and DAMGO as for our Western studies. Our findings confirmed

the Western results, with very low phosphorylated ERK (pERK) signal observed in vehicle-only-treated mice (Vehicle/Vehicle; order of treatment denoted by/) and vehicle followed by DAMGO-treated mice (Vehicle/DAMGO); we observed some increase in signal in the 17-AAG/Vehicle group and a large increase in specific pERK signal in the 17-AAG/DAMGO group (Fig. 3D). We particularly noted an apparent increase in ERK phosphorylation in the lamina I/II region of the dorsal horn, a region rich in nociceptive input and opioid receptors (Fig. 3D, white arrows). We also performed colocalization studies with neuronal nuclei (NeuN), a marker for neuronal cell bodies, and microtubule-associated protein 2 (MAP2), a neuronal cytoskeletal protein enriched in dendrites; we found that the pERK signal colocalized with MAP2 but not NeuN, suggesting ERK activation in postsynaptic dendrites (Fig. 3E). This was confirmed using high-magnification imaging (Fig. 3F), in which substantial, but not complete, pERK/MAP2 overlap was detected. We also quantitated the pERK signal in the dorsal horn region, which confirmed a significant increase with 17-AAG and DAMGO cotreatment (Fig. 3G).

To investigate whether these differences in ERK signaling contribute to the enhanced morphine-induced antinociception observed, we performed behavioral analysis with cotreatment of intrathecal 17-AAG and intrathecal U0126, a MAPK kinase/ERK inhibitor. In both tail-flick and paw incision models, U0126 treatment brought the enhanced morphine-induced antinociceptive profile back to the baseline morphine response (Fig. 3, H and I). In addition, mice treated with U0126 alone without 17-AAG showed no difference in morphine-induced antinociception (Fig. 3, H and I). These results demonstrate that ERK phosphorylation within the spinal cord is necessary for increased morphine-induced antinociception via spinal cord Hsp90 inhibition. They also support our Western and IHC results, suggesting that ERK is not activated by opioids without Hsp90 inhibition.

Rapid protein translation after ERK activation mediates enhanced morphine antinociception after spinal inhibition of Hsp90

Hsp90 and ERK MAPK signaling pathways have been previously connected to translational initiation (29–32). To evaluate the possibility of these pathways altering translation and subsequently contributing to the behavioral differences observed here, we intrathecally administered the translational inhibitor cycloheximide (CX) in the context of our behavioral experiments. In a very similar pattern to the ERK inhibitor experiments above, we found that CX, 24 hours after 17AAG and 30 min before morphine, reduced the enhancement of morphine-induced antinociception back to baseline in the tail-flick model (Fig. 4A). CX alone without 17-AAG treatment did not change morphine-induced antinociception (Fig. 4A). These findings suggest that rapid translation within 30 min of opioid treatment is necessary for the enhanced morphine antinociception seen through spinally inhibited Hsp90.

To identify the position of translation within the Hsp90/ERK molecular cascade, we performed Western blot analysis on spinal cord tissues harvested from mice treated with 17-AAG and combinations of CX and DAMGO. 17-AAG paired with DAMGO treatment stimulated ERK phosphorylation as above; CX treatment 30 min before DAMGO had no effect on this stimulation (Fig. 4, B and C). These results suggest that translational initiation is a downstream event from ERK phosphorylation after Hsp90 inhibition.

Proteomic analysis reveals a protein network altered by spinal Hsp90 inhibition

Our results above suggest that protein translation is altered by spinal Hsp90 inhibition; these changes should thus, in principle, be measurable by quantitative proteomics. We treated mice with intrathecal 17-AAG or Vehicle control as above for 24 hours and removed their spinal cords for analysis. We followed a protocol of protein extraction, SDS–polyacrylamide gel electrophoresis (PAGE) gel separation with six equal bands excised, tryptic digest, and a tandem mass spectrometry (MS/MS) analysis workflow (Fig. 5A). We detected 116 proteins significantly down-regulated by 17-AAG treatment and 69 proteins significantly up-regulated; unbiased hierarchical clustering analysis showed that the individual mice in each sample group (Vehicle versus 17-AAG) clustered together, validating a consistent effect of 17-AAG treatment (Fig. 5B). The full datasets for the significantly altered proteins in the whole analysis and in the subanalyses shown in this figure are available in data files S1 and S2. Of the proteins in this dataset, we noted that the abundance of the kinase ribosomal protein S6 kinase 2 (RSK2) was significantly up-regulated by 17-AAG treatment (Fig. 5C). RSK2 has been shown to promote acute opioid antinociception, highlighting this protein as a potential mechanism for spinal Hsp90 inhibition affecting opioid antinociception (33).

We next performed additional analyses to validate the proteomic dataset and explore the network of protein changes evoked by Hsp90 inhibition. Principal components analysis showed that the individual mice in each treatment group (vehicle versus 17-AAG) clustered together; the groups were strongly separated from each other on component 1, accounting for 75.1% of the variance, whereas within the same treatment, the samples were much closer together along component 2, accounting for only 8.5% of the variance (Fig. 5D). We also represented our data in a volcano plot, permitting an overall visualization of significance and fold change (Fig. 5E). Together, these analyses further confirm the quality of our data and analysis.

Last, we performed Gene Ontology and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis of the statistically significantly changed proteins using Database for Annotation, Visualization, and Integrated Discovery (DAVID) to identify broad themes in functions and processes altered by Hsp90 inhibition. We identified proteins heavily represented in molecular functions such as kinase activity, protein kinase binding, and protein phosphatase binding; pathways including metabolic pathways and oxytocin signaling; processes such as phosphorylation, cell proliferation, lipid metabolism, and synaptic plasticity; and cell components including synapse, exosome, focal adhesion, and postsynaptic density (Fig. 5F and data file S2). This network analysis begins to identify an overall role for Hsp90 in regulating protein networks in the spinal cord, which has not been previously reported.

RSK signaling mediates the enhanced antinociceptive effects of spinally inhibited Hsp90

Cytosolic RSK1 and RSK2 have both been implicated in translational initiation through several substrates, suggesting a potential link to our translation findings above (34–40). RSK2 has also been implicated in acute morphine-induced analgesia within the medial habenula (33). Our proteomic analysis demonstrated altered expression levels of RSK2

within the spinal cord due to Hsp90 inhibition. Therefore, we aimed to probe both RSK1 and RSK2 as a potential mechanism within this molecular pathway.

To evaluate the necessity of RSK activation within our behavioral model, we used the irreversible RSK1/2 inhibitor 1-[4-amino-7-(3-hydroxypropyl)-5-(4-methylphenyl)-7H-pyrrolo[2,3-d] pyrimidin-6-yl]-2-fluoroethanone (Fmk). In a similar design to the U0126 and CX experiments above, 24-hour intrathecal 17-AAG was combined with intrathecal Fmk 30 min before morphine treatment in the tail-flick model. Fmk treatment returned the enhanced morphine antinociception caused by 17-AAG treatment back to baseline, whereas Fmk alone without 17-AAG treatment had no effect on morphine antinociception (Fig. 6A). These results show the same pattern as the U0126 and CX experiments above and strongly suggest that RSK promotes morphine antinociception after spinal Hsp90 inhibition. Fmk is nonselective between RSK1 and RSK2, so either or both isoforms could promote antinociception.

To confirm and extend these findings, we evaluated phosphorylation levels of both isoforms by Western blot in treated spinal cords as above. We found that both RSK1 and RSK2 demonstrate a similar phosphorylation pattern to that of ERK. 17-AAG treatment alone elicits increases in both RSK1 and RSK2 phosphorylation that rises to the level of significance for RSK2; 17-AAG and DAMGO cotreatment significantly increases phosphorylation of both proteins versus the Vehicle/Vehicle control group and over the 17-AAG/Vehicle group for RSK2 (Fig. 6, B to D). These results show that both RSK1 and RSK2 are activated by 17-AAG and DAMGO cotreatment and may both promote morphine antinociception after spinal cord Hsp90 inhibition.

DISCUSSION

In this study, we have identified a previously unknown molecular ERK-RSK signaling circuit in the spinal cord that can promote acute opioid-induced antinociception; this circuit is normally suppressed by Hsp90 and is only uncovered by spinal Hsp90 inhibition. Our results place rapid protein translation as a downstream event of ERK activation. Given the extensive literature that has shown an ERK-RSK-translation cascade (34–40), we propose a model by which Hsp90 inhibition relieves the repression of ERK activation by MOR, resulting in an ERK-RSK-translation-mediated cascade facilitating opioid-induced antinociception (Fig. 7).

Our results provide strong support that spinal ERK, RSK, and translation are not active at baseline for acute opioid antinociception. The inhibitors U0126, Fmk, and CX all had no effect on their own without 17-AAG treatment; we also showed that neither ERK nor RSK phosphorylation was stimulated by opioid treatment in vehicle-treated control mice using both Western blot and IHC methods. We could not find any literature reports showing acute activation of these kinases by opioids in the spinal cord. This is in sharp contrast to the brain, where our results and others show that ERK and RSK are phosphorylated by baseline opioid treatment and contribute to opioid antinociception (21, 33, 41–44). This is not to say that ERK can have no impact on the opioid system in the spinal cord. Spinal ERK has been shown to have a role in mediating chronic opioid treatment side effects,

particularly tolerance (45). ERK also has a well-established role in promoting chronic pain states after activation in the dorsal horn by strong and chronic pain stimuli (46). These contrasting findings show the importance of context in the function of signaling kinases. ERK is downstream of numerous receptor systems in the same cell and must be able to carry out diverse functions in the same cell when stimulated by these different systems. We propose that ERK is organized uniquely within the spinal cord so that it does not respond to acute MOR activation but is free to act in response to chronic MOR activation and in response to other receptor systems; our results suggest that Hsp90 could be this organizing factor preventing acute activation by the MOR. Removing this blockade enables ERK activation, leading to RSK activation, translation of new proteins, and enhanced antinociception. Uncovering these additional mechanisms will lend great insight into how MOR signaling is organized in the spinal cord.

One potential clue to the unique organization of Hsp90 in the spinal cord is that we found that spinal Hsp90 inhibition does not result in Hsp70 up-regulation, confirmed in multiple experiments. Hsp70 up-regulation in response to Hsp90 inhibition has long been considered a canonical response, caused by the release of heat shock factor-1 when Hsp90 is inhibited; we and many others have shown in this paper and elsewhere that Hsp70 is up-regulated in response to Hsp90 inhibition in numerous cell lines and brain tissue (21, 47). However, we cannot find any reports of Hsp70 up-regulation in the wild-type spinal cord *in vivo* after Hsp90 inhibitor treatment. Others have pointed out that Hsp90 inhibition does not always result in a heat shock response leading to Hsp70 up-regulation (48). It may be that Hsp90 in the spinal cord is organized differently at the molecular level than in the brain; perhaps it does not interact with heat shock factor-1 or similar proteins in the spinal cord. These differences may point to the mechanism by which Hsp90 has different signaling roles in brain versus spinal cord.

Our observations are consistent with the ERK/RSK cascade enhancing opioid activation via rapid protein translation. Hsp90 and ERK have both been linked to the initiation of protein translation (29–32). RSK phosphorylation by ERK has been shown to activate translation through a variety of substrates including eukaryotic translation initiation factor-4B, tuberous sclerosis complex-1/2, the 40S ribosomal subunit protein S6, glycogen synthase kinase-3, and elongation factor-2 kinase (34–40). These studies provide plausible targets linking ERK/RSK to protein translation but do not provide a potential mechanism for how protein translation enhances antinociception. Among the full list of proteins altered by 17-AAG treatment in our proteomic analysis were candidate proteins for this mechanism (data file S1). These include ion channels, such as potassium voltage-gated channel subfamily A member 4 and the calcium voltage-gated channel auxiliary subunit $\alpha 2\delta 1$ subunit of the voltage-gated calcium channel, and numerous signaling proteins and signaling protein regulators (such as phospholipase C $\delta 3$, protein phosphatase 1, regulator of G protein signaling 12, and G protein-coupled receptor 162); these provide plausible future candidates to be investigated that could link the protein translation that we observed to enhanced antinociception. One finding that will guide such a search is that any candidate protein must have a rapid turnover half-life, given our observation that inhibiting translation within 30 min of opioid treatment abolished the response, suggesting that the protein must be degraded sufficiently within that time frame.

We also observed interesting systemic interactions above the level of molecular circuitry when investigating how Hsp90 inhibition in the brain and spinal cord interact. We found that Hsp90 inhibition in the brain had a dominant effect over that in the spinal cord in terms of the overall behavioral output, with either systemic or combined intracerebroventricular/intrathecal inhibition. For example, in the tail-flick pain model, brain Hsp90 inhibition had no notable effect on opioid-induced anti-nociception but nonetheless repressed the effects of spinal inhibition. The brain has a well-established circuit of opioidergic descending modulation with cell bodies in the rostroventral medulla and other regions and synapsing on nociceptive modulatory circuits in the spinal cord (49). It may be that descending modulatory neurons in the brain can override the spinal circuits when Hsp90 is inhibited in the brain. Lending some support to this hypothesis is our finding that spinal Hsp90 inhibition leads to enhanced ERK phosphorylation in lamina I/II of the dorsal horn of the spinal cord, which is a key target region for these descending neurons (50). Investigating the circuit context in which Hsp90 regulates antinociception will provide key insights into how the molecular circuitry translates into a whole animal behavioral response.

In this study, we demonstrated a spinal cord-specific role for Hsp90 within MOR downstream signaling and, in doing so, have begun to elucidate MOR-dependent downstream mechanisms of ERK phosphorylation within the spinal cord that can affect systemic morphine-induced antinociception. We propose a mechanism in which Hsp90 serves as a brake on ERK phosphorylation within neurons in the spinal cord dorsal horn. Once the brake is removed by a spinal Hsp90 inhibitor, ERK phosphorylation is “unchained” and can contribute to MOR agonist-induced antinociception through RSK activation and rapid translation. This translation event must up-regulate proteins that contribute to either hyperpolarization or the prevention of neurotransmitter release in primary or secondary nociceptive afferents within the spinal cord, further preventing the transmission of pain signals. Not only is this mechanism important in the context of molecular signaling, but there is also the potential to capitalize on these findings clinically by developing an opioid dose reduction strategy. Hsp90 inhibitors could be used to amplify morphine analgesia through the spinal cord without altering unwanted morphine side effects, many of which are evoked through brain regions such as the striatum (reward) or through the gut (constipation) and would not be affected by spinal cord treatment.

MATERIALS AND METHODS

Reagents

17-AAG (#AAJ66960MC), DAMGO (#11711), Fmk (#46-901-0), CX (#AC357420010), and U0126 (#11-445) were all purchased from Fisher Scientific (Waltham, MA). Morphine sulfate pentahydrate was obtained through the National Institute on Drug Abuse Drug Supply Program and distributed through the Research Triangle Institute. KU-32 was synthesized using published protocols, and purity (>95%) and identity were confirmed by high-performance liquid chromatography (HPLC) and mass spectrometry (25). 17-AAG, U0126, Fmk, KU-32, and CX were prepared as stock solutions in dimethyl sulfoxide (DMSO), and DAMGO was prepared as a stock solution in water. Morphine was prepared fresh for each experiment in United States Pharmacopeia (USP) saline. Powders were

stored as recommended by the manufacturer, and stock solutions were stored at -20°C . Appropriate vehicle controls were used for each experiment: 10% DMSO in water for KU-32, Fmk, and CX intrathecal injections; water for DAMGO intrathecal injections; USP saline for systemic morphine injections; and 10% DMSO, 10% Tween 80, and 80% USP saline for the 17-AAG and U0126 intrathecal, intracerebroventricular, and intraperitoneal injections.

Animals

Male and female CD-1 mice in age-matched controlled cohorts from 4 to 8 weeks of age were used for all experiments and were obtained from Charles River Laboratories (Wilmington, MA). Male and female mice were used in about equal numbers in each experiment; no sex differences were observed, so the male and female cohorts were combined for all data shown. CD-1 [also known as Institute for Cancer Research (ICR)] mice are commonly used in opioid research and in our previous work as a line with a strong response to opioid drugs [as in (21, 51)]. Mice were allowed to recover for at least 5 days after shipment before being used in experiments. The mice were kept in an Association for Assessment and Accreditation of Laboratory Animal Care–accredited vivarium at the University of Arizona under temperature control and 12-hour light/dark cycles with food (standard laboratory chow) and water available ad libitum. No more than five mice were kept in a cage. The animals were monitored daily, including after surgical procedures, by trained veterinary staff. All experiments performed were in accordance with Institutional Animal Care and Use Committee–approved protocols at the University of Arizona and according to the guidelines of the National Institutes of Health (NIH) *Guide for the Care and Use of Laboratory Animals* handbook.

Behavioral experiments

Before any behavioral experiment or testing, the animals were brought to the testing room in their home cages for at least 1 hour for acclimation. Testing always occurred within the same approximate time of day between experiments, and environmental factors (noise, personnel, and scents) were minimized. All testing apparatus (cylinders, grid boxes, etc.) were cleaned between uses. The experimenter was blinded to treatment group by another laboratory member delivering coded drug vials, which were then decoded after collection of all data.

Paw incision and mechanical allodynia

Mechanical thresholds were determined before surgery using calibrated Von Frey filaments (Ugo Basile, Varese, Italy) with the up-down method and four measurements after the first response per mouse (21, 52). The mice were housed in a homemade apparatus with Plexiglas walls and ceiling and a wire mesh floor (3 inches by 4 inches by 3 inches with 0.25-inch wire mesh). The surgery was then performed by anesthesia with $\sim 2\%$ isoflurane in standard air, preparation of the left plantar hindpaw with iodine and 70% ethanol, and a 5-mm incision made through the skin and fascia with a no. 11 scalpel. The muscle was elevated with curved forceps, leaving the origin and insertion intact, and the muscle was split lengthwise using the scalpel. The wound was then closed with 5–0 polyglycolic acid sutures. All intracerebroventricular and intrathecal injections were performed as described in our

previous work (21). For the 17-AAG/KU-32 experiments, the mice were then intrathecally injected and left to recover for 24 hours. The next day, the mechanical threshold was again determined as described above, and intrathecal injections took place for the U0126 experiments with a 15-min treatment time. Both the 17-AAG and the U0126 mice were then injected with morphine (3.2 mg/kg, subcutaneously), and mechanical thresholds were determined over a 3-hour time course. No animals were excluded from these studies.

Tail-flick assay

Preinjection tail-flick baselines were determined in a 52°C warm water tail-flick assay with a 10-s cutoff time (21). The mice were then intrathecally injected with 17-AAG, KU-32, CX, Fmk, or U0126 with a 24-hour (17-AAG and KU-32), 30-min (CX and Fmk), or 15-min (U0126) treatment time. Twenty-four hours after injection, baselines were determined for the 17-AAG experiments. The mice were then injected with morphine (3.2 mg/kg, subcutaneously), and tail-flick latencies were determined over a 2-hour time course. No animals were excluded from these studies.

Rotarod test

Mice were subjected to three training trials of 3 min each on a rotarod device, with the machine off for trial 1, the machine on but not rotating for trial 2, and the machine rotating at 4 rpm for trial 3 (21). An automatic timer in the unit was used to record fall latencies with a 3-min maximum time. The mice were then intrathecally injected with 17-AAG or vehicle and allowed to recover for 24 hours, and another 3-min rotarod trial was performed without additional treatments or interventions. This trial was done with an accelerating 4- to 16-rpm task over the 3-min trial time. No mice were excluded from these studies.

Western blotting and analysis

Mouse spinal cord or PAG protein lysates were prepared using our previously published protocol (21) and quantified with a bicinchoninic acid protein quantitation assay using the manufacturer's protocol (Bio-Rad). The protein was run on precast 10% bistris Bolt gels (Fisher Scientific, #NW00100BOX) using the Bolt gel apparatus and following the manufacturer's instructions. The gels were transferred to nitrocellulose membrane (Bio-Rad) using a wet transfer system (30 V, minimum of 1 hour on ice). The blots were blocked with 5% nonfat dry milk in tris-buffered saline (TBS) and incubated with primary antibody in 5% bovine serum albumin (BSA) in TBS + 0.1% Tween 20 (TBST) overnight rocking at 4°C. The blots were then washed three times for 5 min in TBST, incubated with secondary antibody (see below) in 5% milk in TBST for 1 hour of rocking at room temperature, washed again, and imaged with a LI-COR Odyssey infrared imaging system (LI-COR, Lincoln, NE). Some blots were then stripped with 25 mM glycine-HCl and 1% SDS (pH 2.0) for 30 to 60 min of rocking at room temperature before being washed and reexposed to primary antibody. The resulting image bands were quantified using Scion Image (based on NIH Image). All images were quantified in the linear signal range, which is easier to ensure because the Odyssey imager is a dynamic imager that allows for fine control of exposure. The pERK signal was normalized to the total ERK (tERK) signal, and pRSK1 and pRSK2 were normalized to tRSK1 and tRSK2, respectively, with both measured from the same blot

as the primary target. The normalized intensities were further normalized to a vehicle control present on the same blot.

Immunohistochemistry

Perfusions were performed on drug-treated mice with cold phosphate-buffered saline (PBS), followed by cold 4% paraformaldehyde in PBS. Shortly after the perfusions were complete, fixed spinal cords were extracted and immediately placed in cold 4% paraformaldehyde for ~6 hours. Spinal cords were then placed in 15% sucrose in PBS overnight, followed by 30% sucrose in PBS overnight. Spinal cords were then flash-frozen in optimal cutting temperature compound using liquid nitrogen and sectioned with a Microm HM 525 cryostat at a thickness of 20 μm between the L5 and L6 vertebrae and mounted on Surgipath X-tra microscope slides. Spinal cord sections were rehydrated in PBS in preparation for free-float staining. Samples were incubated in an endogenous peroxidase blocking buffer consisting of 60% methanol and 0.3% H_2O_2 in PBS at room temperature for 30 min and then washed with PBS + 0.1% Tween 20 (PBST). They were then incubated in 5% goat serum and 1% BSA in PBST at room temperature for 1 hour. Samples were then incubated with 1:5000 primary pERK antibody in 1.5% goat serum and 1% BSA in PBST at 4°C overnight. Samples were then washed with PBST and then incubated with a 1:400 biotinylated secondary goat anti-rabbit immunoglobulin G (IgG) antibody in 1.5% goat serum and 1% BSA in PBST at room temperature for 1 hour. Samples were prepared as instructed using the VECTASTAIN Elite ABC horseradish peroxidase kit (#PK-6101) and TSA Plus Fluorescein evaluation kit (#NEL741E001KT), both from PerkinElmer. NeuN and MAP2 primary antibodies were used at 1:1000 and 1:500, respectively, during the pERK primary incubation. The secondary for NeuN and MAP2 was Alexa Fluor goat anti-mouse IgG 594, which was used at 1:500 for both which was added during the pERK secondary incubation mentioned above. Stained spinal cord sections were then mounted onto slides with Novus FluorEver. Sections were imaged at 4 \times , 10 \times , and 63 \times using an Olympus BX51 microscope equipped with a Hamamatsu C8484 digital camera. Images were analyzed using ImageJ. Dorsal horn regions were selected, and average mean intensities were measured and normalized to no primary antibody and vehicle controls within experimental groups.

Antibodies

The antibodies used were as follows: Hsp70 (1:1000; Cell Signaling Technology, 4872S, lot 4, rabbit), glyceraldehyde phosphate dehydrogenase (GAPDH) (1:1000; Thermo Fisher Scientific, MA5-15738, lot PI209504, mouse), pERK (1:1000 for Westerns and 1:5000 for IHC; Cell Signaling Technology, 4370S, lot 12, rabbit), tERK (1:1000; Cell Signaling Technology, 4696S, lot 16, mouse), pRSK1 (1:1000; Cell Signaling Technology, 11989S, lot 4, rabbit), tRSK1 (1:1000; Cell Signaling Technology, 8408S, lot 5, rabbit), pRSK2 (1:1000; Cell Signaling Technology, 3556S, lot 4, rabbit), tRSK2 (1:1000; Cell Signaling Technology, 5528S, lot 1, rabbit), MAP2 (1:500; Invitrogen, 13-1500, lot TJ275359, mouse), NeuN (1:1000; Abcam, ab104224, lot GR3247200-1, mouse), secondary GaM680 (1:10,000 to 1:20,000; LI-COR, 926-68020, lot C50721-02, goat), secondary GaR800 (1:10,000 to 1:20,000; LI-COR, 926-32211, lot C50602-05, goat), and secondary Alexa Fluor goat anti-mouse IgG 594 (1:500; Invitrogen, A11032, lot 1985396, mouse).

Proteomics: In-gel digestion

Mouse spinal cord protein lysates (100 µg) were prepared as for Western blot from animals that were treated with either 17-AAG or vehicle ($N=3$ each) and were separated on a 10% SDS-PAGE gel and stained with Bio-Safe Coomassie G-250 Stain. Each lane of the SDS-PAGE gel was cut into six slices, and the gel slices were subjected to trypsin digestion. The resulting peptides were purified by C18-based desalting exactly as previously described (53, 54).

Proteomics: Mass spectrometry and database search

HPLC–electrospray ionization–MS/MS was performed in positive ion mode on a Thermo Scientific Orbitrap Fusion Lumos Tribrid mass spectrometer fitted with an EASY-Spray Source (Thermo Scientific, San Jose, CA). NanoLC was performed exactly as previously described (53, 54). Tandem mass spectra were extracted from Xcalibur “RAW” files, and charge states were assigned using the ProteoWizard 3.0 msConvert script using the default parameters. The fragment mass spectra were searched against the *Mus musculus* SwissProt_2018_01 database (16,965 entries) using Mascot (version 2.6.0; Matrix Science, London, UK) using the default probability cutoff score. The search variables that were used were as follows: 10–parts per million mass tolerance for precursor ion masses and 0.5 Da for product ion masses; digestion with trypsin; a maximum of two missed tryptic cleavages; and variable modifications of oxidation of methionine and phosphorylation of serine, threonine, and tyrosine. Cross-correlation of Mascot search results with X! Tandem was accomplished with Scaffold (version Scaffold_4.8.7; Proteome Software, Portland, OR). Probability assessment of peptide assignments and protein identifications were made using Scaffold. Only peptides with 95% probability were considered. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the Proteomics Identifications Database (PRIDE) (55, 56) partner repository with the dataset identifier PXD015060 and 10.6019/PXD015060. The reviewer account details are as follows: username, reviewer97855@ebi.ac.uk and password, 8AM00kfd.

Label-free quantitative proteomics

Progenesis QI for proteomics software (version 2.4; Nonlinear Dynamics Ltd., Newcastle upon Tyne, UK) was used to perform ion intensity–based label-free quantification as previously described (54). In an automated format, .RAW files were imported and converted into two-dimensional maps (y axis, time and x axis, m/z), followed by selection of a reference run for alignment purposes. An aggregate dataset containing all peak information from all samples was created from the aligned runs, which was then further narrowed down by selecting only +2, +3, and +4 charged ions for further analysis. The samples were then grouped, and a peak list of fragment ion spectra from only the top eight most intense precursors of a feature was exported to a Mascot generic file (.MGF) format and searched using Mascot (version 2.4; Matrix Science, London, UK) with the same search variables as described above. The resulting Mascot.XML file was then imported into Progenesis, allowing for peptide/protein assignment, whereas peptides with a Mascot ion score of <25 were not considered for further analysis. Protein quantification was performed using only

nonconflicting peptides, and precursor ion abundance values were normalized in a run to those in a reference run (not necessarily the same as the alignment reference run).

Statistical analysis

All data were reported as means \pm SEM and normalized where appropriate as described above to total protein and/or Vehicle control groups. The behavioral data were reported raw without maximum possible effect or other normalization. Biological and technical replicates are described in the figure legends. Comparisons between two groups (Hsp70 protein expression) were performed by unpaired two-tailed *t* tests. Comparisons of more than two groups (ERK and RSK signaling, paw incision, tail flick, and rotarod) were performed by two-way analysis of variance (ANOVA) with Sidak's (behavior) or Tukey's (Western) post hoc tests. In all cases, significance was defined as $P < 0.05$. All graphing and statistical analyses were performed using GraphPad Prism 8.1 (San Diego, CA).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments:

We acknowledge T. Largent-Milnes for assistance in analyzing the spinal cord IHC results and T. Vanderah for sharing his rotarod device.

Funding:

This work was funded by an Arizona Biomedical Research Commission New Investigator Award #ADHS18-198875 and institutional funds from the University of Arizona, both to J.M.S.

Data and materials availability:

A material transfer agreement is required for the molecules used in this study, and quantity is limited (B.S.J.B.). The full results of the proteomic analysis are contained in the Supplementary Materials and have been uploaded to the ProteomeXchange Consortium via the PRIDE (55, 56) partner repository: PXD015060 and 10.6019/PXD015060. All other data needed to evaluate the conclusions in the paper are present in the paper or the Supplementary Materials.

REFERENCES AND NOTES

1. Gaskin DJ, Richard P, The economic costs of pain in the United States. *J. Pain* 13, 715–724 (2012). [PubMed: 22607834]
2. Johannes CB, Le TK, Zhou X, Johnston JA, Dworkin RH, The prevalence of chronic pain in United States adults: Results of an internet-based survey. *J. Pain* 11, 1230–1239 (2010). [PubMed: 20797916]
3. National Collaborating Centre for Cancer, National Institute for Health and Clinical Excellence: Guidance, in Opioids in Palliative Care: Safe and Effective Prescribing of Strong Opioids for Pain in Palliative Care of Adults (National Collaborating Centre for Cancer, 2012).
4. Fields HL, The doctor's dilemma: Opiate analgesics and chronic pain. *Neuron* 69, 591–594 (2011). [PubMed: 21338871]

5. Goodyear K, Haass-Koffler CL, Chavanne D, Opioid use and stigma: The role of gender, language and precipitating events. *Drug Alcohol Depend.* 185, 339–346 (2018). [PubMed: 29499554]
6. Al-Hasani R, Bruchas MR, Molecular mechanisms of opioid receptor-dependent signaling and behavior. *Anesthesiology* 115, 1363–1381 (2011). [PubMed: 22020140]
7. Madariaga-Mazón A, Marmolejo-Valencia AF, Li Y, Toll L, Houghten RA, Martinez-Mayorga K, μ -Opioid receptor biased ligands: A safer and painless discovery of analgesics? *Drug Discov. Today* 22, 1719–1729 (2017). [PubMed: 28743488]
8. Olson KM, Lei W, Keresztes A, Vigne JL, Streicher JM, Novel molecular strategies and targets for opioid drug discovery for the treatment of chronic pain. *Yale J. Biol. Med* 90, 97–110 (2017). [PubMed: 28356897]
9. Villanueva MT, Analgesia: Designing out opioid side effects. *Nat. Rev. Drug Discov* 16, 311 (2017). [PubMed: 28417988]
10. Liang D-Y, Li W-W, Nwaneshiudu C, Irvine K-A, Clark JD, Pharmacological characters of oliceridine, a μ -opioid receptor G-protein-biased ligand in mice. *Anesth. Analg* 129, 1414–1421 (2019). [PubMed: 30044299]
11. la Rochelle AD, Guillemyn K, Dumitrascuta M, Martin C, Utard V, Quillet R, Schneider S, Daubeuf F, Willemse T, Mampuy P, Maes BUW, Frossard N, Bihel F, Spetea M, Simonin F, Ballet S, A bifunctional-biased μ -opioid agonist-neuropeptide FF receptor antagonist as analgesic with improved acute and chronic side effects. *Pain* 159, 1705–1718 (2018). [PubMed: 29708942]
12. Manglik A, Lin H, Aryal DK, McCorvy JD, Dengler D, Corder G, Levit A, Kling RC, Bernat V, Hübner H, Huang X-P, Sassano MF, Giguère PM, Löber S, Duan D, Scherrer G, Kobilka BK, Gmeiner P, Roth BL, Shoichet BK, Structure-based discovery of opioid analgesics with reduced side effects. *Nature* 537, 185–190 (2016). [PubMed: 27533032]
13. Wang Y, Ge Y-H, Wang Y-X, Liu T, Law P-Y, Loh HH, Chen H-Z, Qiu Y, Modulation of mTOR activity by μ -opioid receptor is dependent upon the association of receptor and FK506-binding protein 12. *CNS Neurosci. Ther* 21, 591–598 (2015). [PubMed: 26096047]
14. Bao Y, Gao Y, Hou W, Yang L, Kong X, Zheng H, Li C, Hua B, Engagement of signaling pathways of protease-activated receptor 2 and μ -opioid receptor in bone cancer pain and morphine tolerance. *Int. J. Cancer* 137, 1475–1483 (2015). [PubMed: 25708385]
15. Li J, Buchner J, Structure, function and regulation of the hsp90 machinery. *Biom. J* 36, 106–117 (2013).
16. Eckl JM, Richter K, Functions of the Hsp90 chaperone system: Lifting client proteins to new heights. *Int. J. Biochem. Mol. Biol* 4, 157–165 (2013). [PubMed: 24380020]
17. Kaigorodova EV, Bogatyuk MV, Heat shock proteins as prognostic markers of cancer. *Curr. Cancer Drug Targets* 14, 713–726 (2014). [PubMed: 25258164]
18. Schulz R, Moll UM, Targeting the heat shock protein 90: A rational way to inhibit macrophage migration inhibitory factor function in cancer. *Curr. Opin. Oncol* 26, 108–113 (2014). [PubMed: 24225413]
19. Jhaveri K, Ochiana SO, Dunphy MPS, Gerecitano JF, Corben AD, Peter RI, Janjigian YY, Gomes-DaGama EM, Koren III J, Modi S, Chiosis G, Heat shock protein 90 inhibitors in the treatment of cancer: Current status and future directions. *Expert Opin. Investig. Drugs* 23, 611–628 (2014).
20. Streicher JM, The role of heat shock proteins in regulating receptor signal transduction. *Mol. Pharmacol* 95, 468–474 (2019). [PubMed: 30670482]
21. Lei W, Mullen N, Carthy SM, Brann C, Richard P, Cormier J, Edwards K, Bilsky EJ, Streicher JM, Heat-shock protein 90 (Hsp90) promotes opioid-induced anti-nociception by an ERK mitogen-activated protein kinase (MAPK) mechanism in mouse brain. *J. Biol. Chem* 292, 10414–10428 (2017). [PubMed: 28450396]
22. Chiosis G, Dickey CA, Johnson JL, A global view of Hsp90 functions. *Nat. Struct. Mol. Biol* 20, 1–4 (2013). [PubMed: 23288357]
23. McClellan AJ, Xia Y, Deutschbauer AM, Davis RW, Gerstein M, Frydman J, Diverse cellular functions of the Hsp90 molecular chaperone uncovered using systems approaches. *Cell* 131, 121–135 (2007). [PubMed: 17923092]

24. Thanomkitti K, Fong-Ngern K, Sueksakit K, Thuangtong R, Thongboonkerd V, Molecular functional analyses revealed essential roles of HSP90 and lamin A/C in growth, migration, and self-aggregation of dermal papilla cells. *Cell Death Discov.* 4, 53 (2018). [PubMed: 29760951]
25. Burlison JA, Neckers L, Smith AB, Maxwell A, Blagg BSJ, Novobiocin: Redesigning a DNA gyrase inhibitor for selective inhibition of hsp90. *J. Am. Chem. Soc.* 128, 15529–15536 (2006). [PubMed: 17132020]
26. Ansar S, Burlison JA, Hadden MK, Yu XM, Desino KE, Bean J, Neckers L, Audus KL, Michaelis ML, Blagg BSJ, A non-toxic Hsp90 inhibitor protects neurons from Abeta-induced toxicity. *Bioorg. Med. Chem. Lett.* 17, 1984–1990 (2007). [PubMed: 17276679]
27. Smith V, Sausville EA, Camalier RF, Fiebig H-H, Burger AM, Comparison of 17-dimethylaminoethylamino-17-demethoxy-geldanamycin (17DMAG) and 17-allylamino-17-demethoxygeldanamycin (17AAG) in vitro: Effects on Hsp90 and client proteins in melanoma models. *Cancer Chemother. Pharmacol.* 56, 126–137 (2005). [PubMed: 15841378]
28. Kwon H-M, Kim YH, Yang S-I, Kim Y-J, Lee S-H, Yoon B-W, Geldanamycin protects rat brain through overexpression of HSP70 and reducing brain edema after cerebral focal ischemia. *Neurol. Res.* 30, 740–745 (2008). [PubMed: 18534054]
29. Cargnello M, Roux PP, Activation and function of the MAPKs and their substrates, the MAPK-activated protein kinases. *Microbiol. Mol. Biol. Rev.* 75, 50–83 (2011). [PubMed: 21372320]
30. Gehart H, Kumpf S, Ittner A, Ricci R, MAPK signalling in cellular metabolism: Stress or wellness? *EMBO Rep.* 11, 834–840 (2010). [PubMed: 20930846]
31. Pal JK, Hsp90 regulates protein synthesis by activating the heme-regulated eukaryotic initiation factor 2a (eIF-2a) kinase in rabbit reticulocyte lysates. *J. Biosci.* 23, 353–360 (1998).
32. Uma S, Hartson SD, Chen J-J, Matts RL, Hsp90 is obligatory for the heme-regulated eIF-2a kinase to acquire and maintain an activable conformation. *J. Biol. Chem.* 272, 11648–11656 (1997). [PubMed: 9111082]
33. Darcq E, Befort K, Koebel P, Pannetier S, Mahoney MK, Gaveriaux-Ruff C, Hanauer A, Kieffer BL, RSK2 signaling in medial habenula contributes to acute morphine analgesia. *Neuropsychopharmacology* 37, 1288–1296 (2012). [PubMed: 22218090]
34. Roux PP, Ballif BA, Anjum R, Gygi SP, Blenis J, Tumor-promoting phorbol esters and activated Ras inactivate the tuberous sclerosis tumor suppressor complex via p90 ribosomal S6 kinase. *Proc. Natl. Acad. Sci. U.S.A.* 101, 13489–13494 (2004). [PubMed: 15342917]
35. Shahbazian D, Roux PP, Mieulet V, Cohen MS, Raught B, Taunton J, Hershey JWB, Blenis J, Pende M, Sonenberg N, The mTOR/PI3K and MAPK pathways converge on eIF4B to control its phosphorylation and activity. *EMBO J.* 25, 2781–2791 (2006). [PubMed: 16763566]
36. Roux PP, Shahbazian D, Vu H, Holz MK, Cohen MS, Taunton J, Sonenberg N, Blenis J, RAS/ERK signaling promotes site-specific ribosomal protein S6 phosphorylation via RSK and stimulates cap-dependent translation. *J. Biol. Chem.* 282, 14056–14064 (2007). [PubMed: 17360704]
37. Sutherland C, Leighton IA, Cohen P, Inactivation of glycogen synthase kinase-3 beta by phosphorylation: New kinase connections in insulin and growth-factor signalling. *Biochem. J.* 296 (Pt 1), 15–19 (1993). [PubMed: 8250835]
38. Cohen P, Frame S, The renaissance of GSK3. *Nat. Rev. Mol. Cell Biol.* 2, 769–776 (2001). [PubMed: 11584304]
39. Wang X, Li W, Williams M, Terada N, Alessi DR, Proud CG, Regulation of elongation factor 2 kinase by p90(RSK1) and p70 S6 kinase. *EMBO J.* 20, 4370–4379 (2001). [PubMed: 11500364]
40. Anjum R, Blenis J, The RSK family of kinases: Emerging roles in cellular signalling. *Nat. Rev. Mol. Cell Biol.* 9, 747–758 (2008). [PubMed: 18813292]
41. Macey TA, Lowe JD, Chavkin C, Mu opioid receptor activation of ERK1/2 is GRK3 and arrestin dependent in striatal neurons. *J. Biol. Chem.* 281, 34515–34524 (2006). [PubMed: 16982618]
42. Duraffourd C, Kumala E, Anselmi L, Brecha NC, Sternini C, Opioid-induced mitogen-activated protein kinase signaling in rat enteric neurons following chronic morphine treatment. *PLOS ONE* 9, e110230 (2014). [PubMed: 25302800]
43. Gutstein HB, Rubie EA, Mansour A, Akil H, Woodgett JR, Opioid effects on mitogen-activated protein kinase signaling cascades. *Anesthesiology* 87, 1118–1126 (1997). [PubMed: 9366464]

44. Schmidt H, Schulz S, Klutzny M, Koch T, Händel M, Höllt V, Involvement of mitogen-activated protein kinase in agonist-induced phosphorylation of the μ -opioid receptor in HEK 293 cells. *J. Neurochem* 74, 414–422 (2000). [PubMed: 10617147]
45. Chen Y, Geis C, Sommer C, Activation of TRPV1 contributes to morphine tolerance: Involvement of the mitogen-activated protein kinase signaling pathway. *J. Neurosci* 28, 5836–5845 (2008). [PubMed: 18509045]
46. Popiolek-Barczyk K, Makuch W, Rojewska E, Pilat D, Mika J, Inhibition of intracellular signaling pathways NF- κ B and MEK1/2 attenuates neuropathic pain development and enhances morphine analgesia. *Pharmacol. Rep* 66, 845–851 (2014). [PubMed: 25149989]
47. Kim N, Kim JY, Yenari MA, Pharmacological induction of the 70-kDa heat shock protein protects against brain injury. *Neuroscience* 284, 912–919 (2015). [PubMed: 25446362]
48. Wang Y, McAlpine SR, Heat-shock protein 90 inhibitors: Will they ever succeed as chemotherapeutics? *Future Med. Chem* 7, 87–90 (2015). [PubMed: 25685998]
49. Ossipov MH, Lai J, Malan TP Jr., Porreca F, Spinal and supraspinal mechanisms of neuropathic pain. *Ann. N. Y. Acad. Sci* 909, 12–24 (2000). [PubMed: 10911921]
50. Heinricher MM, Tavares I, Leith JL, Lumb BM, Descending control of nociception: Specificity, recruitment and plasticity. *Brain Res. Rev* 60, 214–225 (2009). [PubMed: 19146877]
51. Ananthan S, Saini SK, Dersch CM, Xu H, Glinchey NM, Giuvelis D, Bilsky EJ, Rothman RB, 14-Alkoxy- and 14-acyloxyppyridomorphinans: μ agonist/ κ antagonist opioid analgesics with diminished tolerance and dependence side effects. *J. Med. Chem* 55, 8350–8363 (2012). [PubMed: 23016952]
52. Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL, Quantitative assessment of tactile allodynia in the rat paw. *J. Neurosci. Methods* 53, 55–63 (1994). [PubMed: 7990513]
53. Kruse R, Krantz J, Barker N, Coletta RL, Rafikov R, Luo M, Højlund K, Mandarino LJ, Langlais PR, Characterization of the CLASP2 protein interaction network identifies SOGA1 as a microtubule-associated protein. *Mol. Cell. Proteomics* 16, 1718–1735 (2017). [PubMed: 28550165]
54. Parker SS, Krantz J, Kwak E-A, Barker NK, Deer CG, Lee NY, Mouneimne G, Langlais PR, Insulin induces microtubule stabilization and regulates the microtubule plus-end tracking protein network in adipocytes. *Mol. Cell. Proteomics* 18, 1363–1381 (2019). [PubMed: 31018989]
55. Deutsch EW, Csordas A, Sun Z, Jarnuczak A, Perez-Riverol Y, Ternent T, Campbell DS, Bernal-Llinares M, Okuda S, Kawano S, Moritz RL, Carver JJ, Wang M, Ishihama Y, Bandeira N, Hermjakob H, Vizcaíno JA, The ProteomeXchange consortium in 2017: Supporting the cultural change in proteomics public data deposition. *Nucleic Acids Res.* 45, D1100–D1106 (2017). [PubMed: 27924013]
56. Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, Inuganti J, Griss, Mayer G, Eisenacher M, Pérez E, Uszkoreit J, Pfeuffer J, Sachsenberg T, Yilmaz , Tiwary S, Cox J, Audain E, Walzer M, Jarnuczak AF, Ternent T, Brazma A, Vizcaíno JA, The PRIDE database and related tools and resources in 2019: Improving support for quantification data. *Nucleic Acids Res.* 47, D442–D450 (2019). [PubMed: 30395289]




Fig. 1. Spinal cord Hsp90 inhibition enhances morphine antinociception. (A to E) Male and female CD-1 mice were treated as indicated in the labels with either 17-AAG (0.5 nmol) or KU-32 [0.01 nmol (C)] or vehicle (Veh) injected by the intracerebroventricular (icv) or intrathecal (it) route, followed by a 24-hour recovery, then injected subcutaneously with or without morphine [Mor; 3.2 mg/kg (A to D); 3.2 or 10 mg/kg (E)], and subjected to behavioral testing. BL, baseline response. Data are means \pm SEM from N (number of mice per group noted on each graph); each experiment was performed with one (D), two (A to C), or three (E) independent technical replicates (meaning groups of mice performed on different days). * $P < 0.05$, *** $P < 0.001$, and **** $P < 0.0001$ versus same time point in Vehicle group, by two-way ANOVA with Sidak's post hoc test.




Fig. 2. Brain Hsp90 inhibition overrides spinal cord Hsp90 inhibition with respect to opioid antinociception.

(A and B) Representative images (A) and analysis (B) of Western blotting for Hsp70 in the periaqueductal gray (PAG) and spinal cord (SC) from male and female CD-1 mice that received intraperitoneal (ip) injection with 17-AAG (50 mg/kg) or vehicle with a 24-hour recovery. Hsp70 densitometry was normalized to that of GAPDH (loading control) from each sample and was further normalized to the vehicle group within each tissue. Data are means \pm SEM of $N = 9$ to 10 mice, performed with two technical replicates. ** $P < 0.01$

versus same tissue in vehicle group, by unpaired two-tailed *t* test. **(C and D)** Tail-flick (C) and paw incision (D) pain behavior tests in mice injected with 17-AAG (50 mg/kg) or vehicle intraperitoneally, followed by a 24-hour recovery and then a subcutaneous injection of morphine (3.2 mg/kg). Data are means \pm SEM of *N* (number of mice per group), noted on each graph, performed with four (C) or two (D) technical replicates. **P* < 0.05 and *****P* < 0.0001 versus same time point in the 17-AAG group, by two-way ANOVA with Sidak's post hoc test. **(E and F)** Tail-flick (E) and paw incision (F) pain behavior tests in mice that received both intracerebroventricular and intrathecal injections of 0.5 nmol of 17-AAG or vehicle, followed by a 24-hour recovery and then a subcutaneous injection of morphine (3.2 mg/kg). Data are means \pm SEM of *N* (number of mice per group), noted on each graph, performed with two technical replicates. **P* < 0.05 and *****P* < 0.0001 versus same time point in the 17-AAG group, by two-way ANOVA with Sidak's post hoc test.




Fig. 3. Spinal Hsp90 inhibition enables opioid activation of ERK MAPK signaling, leading to enhanced antinociception.

(A to C) Representative images (A) and analysis (B and C) of Western blotting for phosphorylated ERK (pERK) and Hsp70 in the spinal cord from male and female CD-1 mice intrathecally injected with 0.5 nmol of 17-AAG or vehicle followed by a 24-hour recovery and then 0.1 nmol of DAMGO or vehicle intrathecally for 10 min. Densitometry of pERK was normalized to that of total ERK (tERK) within each sample, and the densitometry of Hsp70 was normalized to that of GAPDH; each was further normalized to that in the Vehicle/Vehicle group. Data are means \pm SEM from *N* (the number of mice per

group noted on the graphs), each performed as four technical replicates. In (B), $*P < 0.05$ and $****P < 0.0001$ versus Vehicle/Vehicle group and $##P < 0.01$ versus 17-AAG/Vehicle group (both by two-way ANOVA with Tukey's post hoc test). In (C), $P > 0.05$ by unpaired two-tailed t test. (D) Immunohistochemistry (IHC) for pERK (green) performed on L4-L6 region spinal cord tissue from mice treated as described in (A). Representative images from $N = 9$ to 10 mice per group are shown. (E and F) Assessment of colocalization (yellow staining; white arrow) of pERK (green) with neuronal markers NeuN or MAP2 (red) by IHC of the dorsal horn region from 17-AAG/DAMGO-treated mice. Representative images from $N = 3$ individual spinal cords per target, each performed as two independent technical replicates. Higher-magnification images (63 \times) are shown in (F). (G) Quantitation of the pERK signal in the dorsal horn region from all four groups in (D). Intensity values were normalized to the Vehicle/Vehicle group. $N = 9$ to 10 mice per group, each performed in four independent technical replicates. $**P < 0.01$ versus Vehicle/Vehicle group by two-way ANOVA with Tukey's post hoc test. (H and I) Tail-flick (H) and paw incision (I) pain behavior tests in mice intrathecally treated with 0.5 nmol of 17-AAG or vehicle for 24 hours, followed by 5 μ g of U0126 or vehicle intrathecally for 15 min, followed by morphine (3.2 mg/kg, subcutaneously). Data are means \pm SEM from N (the number of mice per group noted on the graphs), each performed as four (H) or three (I) independent technical replicates. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$ versus same time point in the Veh/Veh group by two-way ANOVA with Sidak's post hoc test.




Fig. 4. Spinal Hsp90 inhibition evokes rapid protein translation that mediates enhanced morphine-evoked antinociception.

(A) Tail-flick assay on male and female CD-1 mice intrathecally injected with 0.5 nmol of 17-AAG or vehicle for 24 hours, then 85 nmol of cycloheximide (CX) or vehicle intrathecally for 30 min, and then morphine (3.2 mg/kg, subcutaneously). * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ versus corresponding Veh/Veh data; # $P < 0.05$ and ## $P < 0.01$ versus corresponding 17-AAG/CX data (by two-way ANOVA with Sidak's post hoc test). Data are means \pm SEM from N (the number of mice per group as noted in the graph), each performed as four technical replicates. (B and C) Western blotting and densitometry analysis of pERK abundance in the spinal cords of mice intrathecally treated with 17-AAG then CX or vehicle as in (A), followed by 0.1 nmol of DAMGO or vehicle for 10 min. pERK density was normalized to tERK density in each sample and further normalized to the 17-AAG/Vehicle/Vehicle group. * $P < 0.05$ versus 17-AAG/Vehicle/Vehicle and # $P < 0.05$ versus 17-AAG/CX/Vehicle (both by two-way ANOVA with Tukey's post hoc test). Data are means \pm SEM from N (the number of mice per group as noted in the graph), each performed as six technical replicates.




Fig. 5. Quantitative proteomic analysis reveals a protein network altered by spinal Hsp90 inhibition.

(A) Protein sample preparation and proteomic analysis workflow, as detailed in Materials and Methods. The samples were prepared using female CD-1 mice ($N = 3$ per group), which were intrathecally injected with 0.5 nmol of 17-AAG or vehicle for 24 hours. Spinal cords were removed for proteomic analysis, and protein was extracted as for Western blotting (detailed in Materials and Methods). These samples were used for all subsequent analysis in this figure. (B) Unbiased hierarchical clustering and heat map analysis of proteins significantly altered by 17-AAG treatment ($P < 0.05$). Red, increased; green, decreased; rows, individual proteins; columns, individual samples. Protein quantity traces for all proteins in each sample are shown (right insets). (C) Protein quantity data for the protein kinase RSK2, shown as means \pm SEM of $N = 3$ per group. * $P < 0.05$ by unpaired two-tailed t test. (D) Principal components analysis of the full proteomic dataset was performed. Both treatment groups cluster together and are well separated along component 1, accounting for 75.1% of the variance. Within-group variance only occurs along component 2, accounting for only 8.5% of the variance. (E) Volcano plot of all detected proteins from the full proteomic dataset, plotting P value versus fold change. Red, significantly down-regulated; blue, significantly up-regulated; gray, not significant. (F) Gene ontology (GO) and KEGG pathway analysis of significantly altered proteins from (B) (see Materials and Methods for

details). Data are plotted as significance versus fold enrichment. JNK, c-Jun N-terminal kinase.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 6. Spinal Hsp90 inhibition activates RSK1/2 phosphorylation, which mediates enhanced morphine-evoked antinociception.

(A) Tail-flick assay in male and female CD-1 mice intrathecally injected with 0.5 nmol of 17-AAG or vehicle for 24 hours, followed by 10 nmol of Fmk or vehicle intrathecally for 30 min and then by morphine (3.2 mg/kg, subcutaneously). Data are means \pm SEM from N (number of mice per group noted in the graph), each as three technical replicates. ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$ versus same time point Vehicle/Vehicle group by two-way ANOVA with Sidak's post hoc test. (B to D) Western blotting for phosphorylated (p) and total (t) RSK1 and RSK2 in the spinal cords from mice intrathecally injected with 0.5 nmol of 17-AAG or Vehicle for 24 hours, followed by 0.1 nmol of DAMGO or Vehicle intrathecally for 10 min. Densitometry of pRSK1 (C) and pRSK2 (D) was normalized to the corresponding tRSK within each sample and further normalized to the Vehicle/Vehicle group. Data are means \pm SEM from N (number of mice per group noted in the graph), each as three technical replicates. * $P < 0.05$, *** $P < 0.001$, and **** $P < 0.0001$ versus Vehicle/

Vehicle group and $##P < 0.01$ versus 17-AAG/Vehicle group (both by two-way ANOVA with Tukey's post hoc test).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 7. Proposed model of Hsp90 regulation of opioid signaling in the spinal cord.

Our data suggest that phosphorylation of ERK-MAPK proteins in the spinal cord by the MOR in response to opioids is blocked by Hsp90. Thus, Hsp90 inhibition (by 17-AAG or KU-32) enables ERK MAPK phosphorylation by the MOR with opioid treatment, leading to an ERK-RSK-translation cascade that promotes opioid antinociception.