

*erratum*

# Dosimetric study for spine stereotactic body radiation therapy: magnetic resonance guided linear accelerator versus volumetric modulated arc therapy

Poonam Yadav, Hima B. Musunuru, Jacob S. Witt, Michael Bassetti, John Bayouth, Andrew M. Baschnagel

Department of Human Oncology, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Radiol Oncol 2019; 53(3): 362-368.

Received 24 March 2019  
Accepted 22 July 2019

Correspondence to: Poonam Yadav, Ph.D., Department of Human Oncology, University of Wisconsin School of Medicine and Public Health, 600 Highland Avenue, K4/B74, Madison, WI, USA 53792-0600. Phone: 001 608-235-2594; Fax: 001 608-263-0990; E-mail: yadav@humonc.wisc.edu

Disclosure: No potential conflicts of interest were disclosed.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

doi: 10.2478/raon-2019-0042

---

The authors regret that the following financial relationship was not previously disclosed upon publication of this article. Please note that Dr. John E. Bayouth has reported being a member of the Advisory Board of ViewRay Inc, dating from April 4, 2018, and became a corporate manager of MR Guidance LLC dating from May 19, 2019, which provides consulting services for MRI-guided radiation therapy, and has business activity with a company that utilizes deformable image registration and motion management technology (ViewRay, Inc.).

The authors would like to apologise for any inconvenience caused.

---