



Review

Active compounds in kepok banana peel as anti-inflammatory in acne vulgaris: Review article



Dwiana Savitri^{a,b}, Khairuddin Djawad^c, Mochammad Hatta^{d,*}, Sitti Wahyuni^e, Agussalim Bukhari^f

^a Doctoral Program in Medical Science, Faculty of Medicine, Hasanuddin University, Makassar, South Sulawesi, Indonesia

^b Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, South Kalimantan, Indonesia

^c Department of Dermatology and Venereology, Faculty of Medicine, Hasanuddin University, Makassar, South Sulawesi, Indonesia

^d Department of Molecular Biology and Immunology, Faculty of Medicine, Hasanuddin University, Makassar, South Sulawesi, Indonesia

^e Department of Parasitology, Faculty of Medicine, Hasanuddin University, Makassar, South Sulawesi, Indonesia

^f Department of Nutrition, Faculty Medicine, Universitas Hasanuddin, Makassar, South Sulawesi, Indonesia

ARTICLE INFO

Keywords:

Kepok banana peel
Anti-inflammatory
Acne vulgaris

ABSTRACT

Background: Acne vulgaris (AV) is a chronic inflammatory skin condition affecting the pilosebaceous units characterized by recurrent comedones, erythematous papules and pustules. The disease is benign however may produce scarring, erythema, and hyperpigmentation resulting in physical and psychological problems. Conventional therapy may reduce the symptoms of AV nevertheless, has a possibility of resistance, unwanted side effects, and has high cost. Thus, utilizing natural remedies may be a useful.

Methods: The data in this study were collect by search the keyword combinations of medical subject heading (mesh) of "inhibition", "antimicrobial", "banana peel", "acne vulgaris" and "antiinflammation" and relevant reference lists were manually searched in PubMed, EMBRASE and Scopus database. All relevant articles in data base above were included and narratively discussed in this review article.

Objective: To discuss the bioactive potential of banana peel as an inflammatory modulator in acne vulgaris.

Results: Banana peel contains many bioactive compounds, particularly phenolic and non-phenolic antioxidants (ascorbic acid, carotene, and cyanidin) which are pivotal in removing inflammatory products by inhibiting reactive oxygen species (ROS), protecting protease inhibitors from oxidative damage, and preventing fibroblasts degradation. Banana peel also contains anti-inflammatory agents such as trigonelline which inhibits bacterial enzymes and nucleic acid synthesis; isovanillic acid which suppresses TNF- α production; and ferulic acid which inhibits the production of proinflammatory signaling and cytokines.

Conclusion: Banana peel contains many bioactive compounds which demonstrate anti-inflammatory properties through several processes of the inflammatory pathway. However further research is needed to confirm this finding.

1. Introduction

Acne vulgaris (AV) is a chronic inflammatory skin disease of the pilosebaceous units (sebaceous glands and hair follicles) on the face, neck, trunk, or proximal upper extremities. This disorder is characterized by the development of chronic or recurrent comedones, erythematous papules, and pustules [1]. The degree of severity is in the form of open or closed comedones (non-inflammatory) lesions followed by pustules and papules (inflammation), and residual pathology in the form

of nodules and cysts [2,3]. AV is a disease that is widespread in all races, ethnicities, and cultures. Although it is benign, self-limiting and not life-threatening, in chronic inflammation cases it will trigger symptoms of scars, erythema, and hyperpigmentation that trigger physical and psychological problems [1]. Psychological problems can be depression, anxiety [4], and unemployment [5].

Conventional topical therapy refers to the AV type and is associated with symptoms. Therapies applied include antibacterial agents, decreased sebum secretion, triggers of cellular turnover, anti-

* Corresponding author. Molecular Biology and Immunology Laboratory, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia.

E-mail addresses: vinnadwiana@gmail.com (D. Savitri), duddin@ymail.com (K. Djawad), hattaram@yahoo.com (M. Hatta), sittiwahyuni@gmail.com (S. Wahyuni), agussalimbukhari@yahoo.com (A. Bukhari).

<https://doi.org/10.1016/j.amsu.2022.104868>

Received 12 August 2022; Received in revised form 4 October 2022; Accepted 6 November 2022

Available online 12 November 2022

2049-0801/© 2022 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

inflammatory agents. In severe cases the clinician will give oral antibiotics, retinoids, and hormonal therapy [6]. Conventional therapy still has high costs and leaves side effects, in the form of resistance [7,8]. This leads to a new approach in the form of utilizing natural or herbal resources.

Previous study has shown that some herbs can be used as antimicrobial [9–12], antifungal [13], anti-inflammatory [10,11,14–18], antidiabetic [19], antihypertensive [20], anti-sedative [21,22], and reducing oxidative stress after traumatic brain injury [23–25].

Banana (*Musa spp.*) is one of the fruits that is widely distributed and consumed throughout the world. Bananas have an attractive nutritional value and taste and are a popular fruit in industrialized countries. This plant is widely grown in tropical and subtropical regions and is centered in its original area, namely Southeast Asia and the Western Pacific [26]. This plant is found in more than 130 countries as a food source and contributes to 17% of world fruit production [27,28]. Banana flesh is wrapped by a yellow skin. When ripe, bananas have a light yellow skin with brown specks. Until now, there have not been many studies in terms of nutrition, pharmacy and its bioactive potential for banana peels [29]. Several active compounds from ripe and unripe banana peels include fiber, pectin, phytosterols, phenolics, biogenic amines, and carotenoids. The latter four exhibit antioxidant properties, thus producing health benefits [30,31]. However, there are limited studies that demonstrate a direct association between banana peel and acne vulgaris. Therefore, this paper aims to discuss the bioactive potential of banana peel as an inflammatory modulator in acne vulgaris.

2. Methods

Then data in this study were collect by search the keyword of medical subject headings (MeSH) of “inhibition”, “antimicrobial”, “banana peel”, “acne vulgaris”, and “antiinflammation” and relevant reference list were manually search in PubMed, EMBRASE and Scopus database. All relevant articles in data base above were included and narratively discussed in this review articles [27,28].

3. Body of article

3.1. Epidemiology of acne vulgaris

There is little to no universal standard on the minimal criteria of acne vulgaris diagnosis. Different studies employ different criteria and therefore it becomes difficult to compare, compile, and extrapolate their results. The prevalence estimates are also influenced by various factors such as the country where the data was gathered from and the sample size. This creates stark variation across different studies [32]. A study conducted in Germany found the prevalence estimate of 26.8%, whilst a study in Brazil found the prevalence estimate to be 96% [33]. According to the Global Burden of Disease Study 2010, acne vulgaris is the eight most common skin disorder. The prevalence estimate reached 9.38% globally. The study also mentioned the estimated prevalence of 8.96% in males and 9.81% in females [34]. In contrast, a systematic review revealed that the pooled odds ratio for males is 1,07 (95%CI = 1,97–2, 83) with reference to females. The prevalence was also found to increase with age. Teenagers have the highest incidence and pre-pubertal children have the lowest. During the young adult years, the prevalence declines gradually as the age increases. Aside from sex and age, family history and BMI are also proven to bear significant association with acne vulgaris. There is a strong association between acne vulgaris and having parents with the history of acne vulgaris. The pooled odds ratio is 2,91 (95%CI = 2,58–3,28). Obesity is also proven to be a risk factor of acne vulgaris. According to the pooled odds ratio, An obese patient is estimated 2,36 (95%CI = 1,97–2,83) times to have acne vulgaris compared to normal/underweight patient. The association between acne vulgaris and smoking, dietary factors, and the severity of acne was found to be inconsistent across multiple studies [32].

3.2. Pathogenesis of acne vulgaris

Acne vulgaris most commonly affects pilosebaceous units of the face, chest, upper arms, and upper back due to their high density of sebaceous glands. There are four main sequential factors leading to acne vulgaris, namely increased sebum production, followed by irregular follicular desquamation, proliferation of *C. acnes*, and inflammation [35–37].

Sebum production is proportional to the severity of the acne lesions. Androgen hormones increase sebum production since the increase in androgens is not accompanied with an increase in the sensitivity of androgen receptors [38]. Sebum production is also influenced by growth hormone. IGF-1 reduces the intracellular level of nuclear metabolic forkhead box class O transcription factor 1 which leads to activation of the mammalian target of rapamycin complex 1 (mTORC1). mTORC1 is associated with cell proliferation and metabolism, in this case, the hyperproliferation of sebaceous gland and keratinocyte. IGF-1 also increases androgen levels and endogenous IGF-1, thereby creating a positive feedback loop [39].

Androgens also cause hyperproliferation of the keratinocytes which occlude the follicles and cause microcomedones to form. Comedogenesis requires hyperproliferation of the keratinocyte and subsequent gathering of desquamated corneocytes in the follicle. Microcomedone fills the follicle with lipids and monofilaments which can turn into non-inflammatory comedones or inflammatory acne lesions. There are open comedones or blackheads and closed comedones or whiteheads. Blackheads contain desquamated keratinocytes and sebum. Whiteheads appear as whitish bumps beneath the skin, which may rupture to the surrounding tissue [35].

Comedones are the suitable site for microbial proliferation. The three normal microflora of the sebaceous follicles are coagulase-negative staphylococci, anaerobic diphtheroids, and lipophilic yeasts. Yeasts rarely partake in the pathogenesis of acne vulgaris. Coagulase-negative staphylococci elicits harmless antibody responses in comparison to *C. acnes*. *S. epidermidis* is an aerobic organism which grows superficially instead of deeply in the anaerobic infra-infundibulum where inflammation takes place. On the contrary, *C. acnes* is an anaerobic organism which colonizes the sebaceous follicles to obtain nutritious lipids. The response of immune system towards the bacteria trigger the formation of papules, pustules, and even cysts and nodules in severe cases [32,40].

3.3. *C. acnes* as initiator

A human skin commensal bacterium, *Propionibacterium acnes* (*P. acnes*), is now called *Cutibacterium acnes* (*C. acnes*) based on a new taxonomy. *C. acnes* is a commensal and predominant Gram-positive bacteria (>60% of total bacteria) in the skin. Almost everyone is a host of this bacterium, with a density of 102–105-6/cm² accounting for half of the total microbiome [41]. *C. acnes* is believed to play an important role in the pathogenesis of the development of AV lesions [42]. *C. acnes* induces sebaceous triglyceride metabolism, glycerol consumption, and release of free fatty acids, neutrophils, and complement attractants. *C. acnes* contributes to inflammation through monocyte induction of pro-inflammatory cytokine secretion. In vitro, *C. acnes* also triggers the secretion of antimicrobial peptides and cytokines from keratinocytes [43].

Innate immunity is the first line of defense against pathogens (pathogen-associated molecular pattern) and danger (danger-associated molecular pattern). Responses to PAMPs or DAMPs are mediated by pattern recognition receptors, such as Nod like receptors (NLRP1, NLRP2, NLRC4, and AIM2). These receptors will respond in the form of formation of inflammasomes that control the activation and secretion of pro-inflammatory cytokines [44]. *C. acnes* can activate the NLRP3 inflammasome pathway leading to an increase in IL-1 β levels. Furthermore, *C. acnes* will be recognized by pattern recognition receptors, including TLR2 and TLR4 to elicit an immune response against bacteria. TLR2 activation triggers the release of cytokines and attracts

inflammatory cells toward the lesion. This event will activate infiltrating cells and release pro-inflammatory cytokines namely IL-6, TNF- α , IL-12, IL-8, and IL-1 β . In recognition of *C. acnes*, TLRs can activate cellular components including TNF- α and IL-1A, then activate the chemokines IL-8 and IL-10, finally activating downstream components of inflammation. IL1A activity is also modulated by IL1RN [45,46].

TNF- α will interact with its receptor, TNFR2 to activate the inflammatory signaling cascade. This cascade will activate MAPKs, including MAPK11 which upregulates the expression of pro-inflammatory mediators, and also activates matrix MMPs, including MMP-2 to trigger tissue remodeling. MMP activity is regulated by TIMPs, altering the balance of MMP and TIMP activity which also contributes to acne development. IL-6 (stimulated by IL17A, IL17F, and ACE) plays a role in modulating host defense and inflammation. On the other hand, TGF- β not only plays a role in the synthesis of lipid sebocytes, but also controls the immune response against *C. acnes* [45,46].

3.4. Inflammation

Inflammation is required for host defense as a response against invading pathogens, toxic compounds, or noxious endogenous signals. On the other hand, the body's failure to stop this process will trigger tissue or organ damage [47]. Inflammation in acne vulgaris is a consequence of the interaction between *C. acnes* and innate immunity. This inflammation is characterized by the production of cytokines, chemokines, and antimicrobial peptides from the epidermis and immune cells. The effects of *C. acnes* infection, hyperkeratinization, and activation of innate immunity are prolongation of inflammation that triggers the occurrence of severe acne lesions [48].

There are different strains of *C. acnes* with varying resistance profiles which colonize different pilosebaceous units. The bacteria induce inflammation after being detected with toll-like receptors in macrophages and keratinocytes [49]. As *C. acnes* is detected by the immune system, the inflammatory process begins. The immune system, particularly from the lymphocytes, neutrophils, and macrophages, releases chemostatic factors. These molecules trigger follicular damage, rupture, as well as leakage of not just bacteria but also lipids and fatty acids into the surrounding dermis. As a result, inflammatory lesions such as nodules, pustules, papules, and cyst emerge. There are also reactive oxygen species released by neutrophils which damage the follicular epithelium and aggravate the inflammatory process [50].

3.5. Antimicrobial peptides

Antimicrobial peptides are usually small (<20 kDa), positively charged amphiphilic molecules with a linear -helix or -sheet motif. Antimicrobial peptides are important effectors of the innate immune system and provide the first line of defense against microorganisms [51, 52]. In human skin, antimicrobial peptides are produced by keratinocytes, neutrophils, sebocytes, or sweat glands. There is a correlation between disease severity and the amount of antimicrobial peptide production [53].

Antimicrobial peptides are amphipathic and cationic so they can interact with bacterial anionic membranes to trigger membranolysis (Wimley et al., 2010). In addition to membranolysis, antimicrobial peptides are also anti-inflammatory, bind to lipopolysaccharide/and/or lipoteichoic acid, and trigger bacterial agglutination [42,54-56]. Compared to conventional antibiotics, antimicrobial peptides are able to modulate the host immune response and are less likely to induce short-term bacterial resistance due to different actions [57]. Thus, antimicrobial peptides can be candidates for AV through direct bacterial killing and inhibition of Toll-like receptors in NF- κ B activation [58].

Human β -defensins are found primarily in epithelial cells and a number of sites in the body. The human β -defensin gene was found at constitutively low levels in epithelial cells. This transcription is induced by various factors, namely microbes and cytokines. There is ample

evidence that pro-inflammatory cytokines (IL-1 β and TNF- α) and bacterial lipopolysaccharides can upregulate human β -defensins. Research on the Han population in China found that there was a significant difference in the DEF1 C-44G and G-20 A polymorphisms between acne vulgaris cases compared to controls ($p < 0.05$). Alleles C-44G and G-20 A are underexpressed in acne vulgaris, which correlates with a lower risk of acne vulgaris in Han Chinese patients [59].

Two antimicrobial peptides from frog skin, namely [D4k]ascaphin-8 and [T5k]temporin-Dra can inhibit the growth of *P. acnes* and the release of cytokines and even trigger the production of anti-inflammatory cytokines [60]. Antimicrobial peptide HPA3NT3 as a synthetic peptide derived from *Helicobacter pylori* is antimicrobial against *P. acnes* and can block *P. acnes*-induced inflammation [61]. The antimicrobial peptide CEN1Hc-Br derived from green sea urchin is antimicrobial and inhibits the induction of toll-like receptor-2 expression by *P. acnes*. In addition, CEN1Hc-Br is also able to downregulate inflammatory cytokines in vitro and in vivo [62]. Bombinin is an antimicrobial peptide from the skin secretions of *Bombina* species [63]. BLP-7 contains 27 amino acids, including 4 cationic amino acids, namely Lis, Arg and His and 18 hydrophobic amino acids. It appears that the antimicrobial peptide properties of BLP-7 are via hydrophobicity. BLP-7 inhibits the expression of IL-8 and GM-CSF in NHEKs [42].

3.6. Active compound in kepok banana peels as anti-inflammation in acne vulgaris

Despite occupying 40% of the total weight, banana peels are often discarded. Banana peel has only been used as animal feed, which is not economically and environmentally friendly [64]. In many Asian countries, banana peel is used for traditional medicine and cosmetic purposes such as astringent [65,66]. Banana peel contains plenty of bioactive compounds. According to a study on *Musa sapientum* Linn. (*Musaceae*), the fresh ripe peel contains NO inhibitory activity without any antioxidant effect. However, the decoction from the unripe peel demonstrated significant antioxidant activity and high phenolic content. The study further shows that the antioxidant activity is correlated to the total phenolic content [65].

Phenolic is one of the active compounds from banana peel. The phenolic content of banana peel compounds is 35 mg equivalent of gallic acid/g extract and 428 mg/g equivalent of quercetin [67]. Galocatechin is another phenolic antioxidant identified in banana peel. It has higher amount in the peel (158 mg/100 g of dry weight) than in the pulp (29.6 mg/100 g of dry weight) according to studies on *M. cavendishii* Lamb [68, 69]. Other phenolics found in banana peel include catecholamines, flavanones, flavonols, tocopherols [70].

Non-phenolic antioxidant compounds in banana peel are ascorbic acid, carotene, and cyanidin. Banana peel contains higher carotenoid values, particularly *trans*-beta-carotene, *trans*-alpha-carotene, and *cis*-beta-carotene which are respectively 174.87 ± 7.86 μ g/g dry weight, 164.87 ± 10.51 μ g/g dry weight, and 92.21 ± 5.37 μ g/g dry weight [68]. On another hand, sterol compounds from banana peels include stigmasterol, -sitosterol, and campesterol. The triphenic alcohols from banana peels include cycloartenol, cycloeucalenol, and 2,4-methylene cyanartanol. Sterols and triphenic alcohols are lipids without a phenolic ring and without antioxidant activity [71]. Stigmasterol can inhibit the production of pro-inflammatory mediators (TNF- α , IL-6, IL-1b, iNOS, and COX-2) and increase the expression of anti-inflammatory mediators (IL-10) [72]. Antioxidants play an important role in the removal of inflammatory products by working against proteases as well as reactive oxygen species (ROS). Antioxidants also protect protease inhibitors from oxidative damage. Antioxidants also prevent fibroblasts and other cells degradation caused by ROS [68].

An experimental study on rats proved that oral banana peel extract is associated with lower concentrations of hydroperoxides, peroxidation products (MDA), and conjugated dienes. Simultaneously, rats treated with banana peel showed increased catalase and superoxide dismutase

activities. This is also followed by reduced concentration of glutathione [68].

LC-HRMS analysis on the ethanolic extract of Kepok banana peel found compounds of trigonelline, isovanillic acid, vanillin, ferulic acid, 3-methoxyflavone, rutin, and salsolinol. Trigonelline is an alkaloid found in the ethanolite extract of the Kepok banana peel [73]. Although nothing has been proven directly on acne vulgaris, various studies have proven the role of trigonelline as an anti-inflammatory in other diseases. The anti-inflammatory trigonelline pathway is through depression of nuclear factor kappa beta (NF- κ B), depression of TLR4, and inhibition of pro-inflammatory cytokine production [73]. Trigonelline as an alkaloid is also antibacterial through the mechanism of inhibiting nucleic acid synthesis through dihydrofolate reductase or topoisomerase activity, inhibition of cell division protein (FtsZ), inhibition of bacterial enzymes, and disruption of the outer membrane and bacterial cytoplasmic membrane integrity [74]. Isovanillic acid can suppress the production of TNF- α due to induction by LPS [75]. Ferulic acid is an organic acid. Ferulic acid can restore the production of inflammatory cytokines, suppress the production of proinflammatory cytokine mRNA, inhibit I κ B degradation and phosphorylation of NF- κ B p65 due to LPS induction [76,77].

A study on 45 female Indonesian adolescents was carried out to determine the efficacy of banana peel on acne vulgaris in 2018. The research found that before the application, the prevalence of mild, moderate, and severe acne vulgaris was 62.2% (28), 33.3% (15), 4.4% (2) respectively among the subjects. The ripe banana peel was rubbed on the acne and left for 30 min to an hour. The process was repeated every day for 7 days. After the application of banana peel, the prevalence of mild, moderate, and severe acne vulgaris became 62.2% (28), 33.3% (15), 4.4% (2) respectively. Furthermore, the study also found that 57.9% of the samples who used to have moderate acne vulgaris recede to mild acne vulgaris. Wilcoxon signed rank proved significant difference (p value = 0.016) [66].

3.7. Implication

The implications of this potential point to the need for in vivo studies. The active compounds from banana peel have broad pharmacological potential, as anti-inflammatory and antibacterial that can be used to inhibit the development of acne vulgaris.

4. Conclusion

Banana peel contains many bioactive compounds, particularly phenolic and non-phenolic antioxidants such as ascorbic acid, carotene, and cyanidin. Antioxidants play an important role in the removal of inflammatory products by inhibiting reactive oxygen species (ROS), protecting protease inhibitors from oxidative damage, and preventing fibroblasts degradation [68]. Banana peel also contains anti-inflammatory agents called trigonelline, isovanillic acid, and ferulic acid [73]. Trigonelline inhibits bacterial enzymes and nucleic acid synthesis [74]. Isovanillic acid suppresses TNF- α production due to induction by LPS [75]. Ferulic acid suppresses proinflammatory signaling and cytokine production [76,77]. An experimental study on the application of banana peel on 45 female subjects found significant alteration of acne vulgaris severity [66]. However further research is needed to confirm this finding.

Ethical approval

None.

Sources of funding

None.

Author contribution

Concept or design: Dwiana Savitri, Khairuddin Djawad, Mochammad Hatta, Agussalim Bukhari.

Data collection: Dwiana Savitri, Sitti Wahyuni, Agussalim Bukhari.

Data analysis: Dwiana Savitri, Sitti Wahyuni, Agussalim Bukhari.

Writing the paper: Dwiana Savitri, Khairuddin Djawad, Mochammad Hatta, Agussalim Bukhari.

Registration of research studies

- 1 Name of the registry: Prof. Mochammad Hatta, MD, PhD
- 2 Unique Identifying number or registration ID: 0000-0002-84564203
- 3 Hyperlink to your specific registration (must be publicly accessible and will be checked):-

Guarantor

Guarantor: Khairuddin Djawad, Mochammad Hatta, Agussalim Bukhari.

Consent

None.

Declaration of competing interest

The authors declares there is no conflict of interest.

Abbreviations

AV	acne vulgaris
IGF-1	Insulin-like growth factor 1
mTORC1	mammalian target of rapamycin complex 1
IL	Interleukin
TNF	Tumor necrosis factor
NF- κ B	Nuclear factor kappa beta
PAMPs	Pathogen-associated molecular patterns
DAMPs	Danger-associated molecular patterns
NLRP	Nod like receptors
TLR	Toll-like receptors
MAPK	Mitogen-activated protein kinase
MMP	Matrix metalloproteinase
BLP-7	Bombinin-like peptide 7
COX-2	Cyclooxygenase-2
mRNA	Messenger ribonucleic acid
LPS	Lipopolysaccharide

References

- [1] A. Leung, B. Barankin, J. Lam, K. Leong, K. Hon, *Dermatology: how to manage acne vulgaris*, *Drugs Context (US)* 10 (2021) 1–18.
- [2] H. Williams, R. Dellavalle, S. Garner, *Acne vulgaris*, *Lancet* 379 (9813) (2012) 361–372.
- [3] A. Mahto, *Acne vulgaris*, *Medicine* 45 (6) (2017) 386–389.
- [4] R. Motley, A. Finlay, *How much disability is caused by acne?* *Clin. Exp. Dermatol.* 14 (3) (1989) 194–198.
- [5] W. Cunliffe, *Acne and unemployment*, *Br. J. Dermatol.* 115 (3) (1986), 386–386.
- [6] A. Zaenglein, A. Pathy, B. Schlosser, A. Alikhan, H. Baldwin, D. Berson, et al., *Guidelines of care for the management of acne vulgaris*, *J. Am. Acad. Dermatol.* 74 (2016) 945–973.
- [7] B. Van den Bergh, J. Michiels, T. Wenseleers, E. Windels, P. Boer, D. Kestemont, et al., *Frequency of antibiotic application drives rapid evolutionary adaptation of Escherichia coli persistence*, *Nature Microbiology* 1 (5) (2016).
- [8] F. Masoud, P. Badali, M. Isa, H. Alamdari, S. Asnaashari, J. Shokri, et al., *The novel topical herbal gel might be an alternative treatment in patients with acne vulgaris: a randomized, double-blind controlled study*, *Phytomedicine* 2 (2) (2022), 100232.
- [9] T. Wahyuni, M. Hatta, A. Bukhari, A. Santoso, M. Massi, *Increasing Natural Resistance Associated Macrophage Protein 1 serum level after Miana treatment in BALB/c induced Klebsiella pneumoniae experimental research*, *Ann. Med. Surg.* 65 (2021), 102262.

- [10] T.A. Yanto, Mochammad Hatta, A. Bukhari, R. Natzir, Molecular and immunological mechanisms of miana leaf (*Coleus scutellarioides* [L.] Benth) in infectious diseases, *Biomed. Pharmacol. J.* 13 (4) (2020) 1607–1618, <https://doi.org/10.13005/bpj/2036>, 13(4).
- [11] L.I. Syarif, A.R. Junita, M. Hatta, R. Dwiyantri, C. Kaelan, et al., A mini review: medical plants for typhoid fever in Indonesia, *Sys. Rev. Pharm.* 11 (6) (2020) 1171–1180, <https://doi.org/10.31838/srp.2020.6.170>. E-ISSN 0976-2779 P-ISSN 0975-8453.
- [12] A. Febriza, M. Hatta, R. Natzir, V.N.A. Kasim, H.H. Idrus, Activity of antimicrobial peptide; cathelicidin, on bacterial infection, *Open Biochem. J.* 13 (2019) 45–53, <https://doi.org/10.2174/1874091X01913010045>.
- [13] M. Karo, M. Hatta, T. Simanjuntak, T. Rina, E. Kamelia, et al., A mini review of Indonesian medicinal plants for vulvovaginal candidiasis, *Rasayan J. Chem.* 10 (4) (2017) 1280–1288, <https://doi.org/10.7324/RJC.2017.1041887>.
- [14] M. Karo, M. Hatta, I. Patellongi, R. Natzir, T. Tambaip, et al., IgM antibody and colony fungal load impacts of orally administered ethanol extract of *Plectranthus scutellarioides* on mice with systemic candidiasis, *J. Pharma. Pharmacog. Res.* 6 (1) (2018) 27–34. <http://jppres.com/jppres>.
- [15] U.K. Amsyah, M. Hatta, H. Tahir, G. Alam, A. Asmawati, Expression of IL-10 in A. actinomycetemcomitans induced rat treated by purple miana leaves, *Biomed. Pharmacol. J.* 12 (4) (2019), <https://doi.org/10.13005/bpj/1845>, 2099–21.
- [16] M.B. Karo, M. Hatta, W. Salma, I. Patellongi, R. Natzir, Effects of miana (*Coleus scutellarioides* (L.) Benth) to expression of mRNA IL-37 in Balb/c mice infected *Candida albicans*, *Phcog. J.* 10 (1) (2018) 16–19, <https://doi.org/10.5530/pj.2018.1.3>.
- [17] M. Hatta, T. Simanjuntak, A. Tahir, R. Sirait, M. Karo, T. Tambaip, et al., Analysis of anti-toxoplasma immunoglobulin G and immunoglobulin M antibody levels after intervention with *Curcuma Longa* extract on early pregnant mice with acute toxoplasmosis, *J. Global Infect. Dis.* 11 (1) (2019) 25. <http://www.jgid.org/text.asp?2019/11/1/25/251839>.
- [18] T. Simanjuntak, M. Hatta, S. Rauf, S. Prabdandari, C. Siagian, R. Dwiyantri, Tumor necrosis factor- α levels and histopathology finding after intervention with *Curcuma longa* extract, *J. Med. Sci.* 18 (2) (2018) 56–62, <https://doi.org/10.3923/jms.2018.56.62>.
- [19] W. El-Tantawy, A. Temraz, Management of diabetes using herbal extracts: review, *Arch. Physiol. Biochem.* 124 (5) (2017) 383–389.
- [20] I. Royani, S. As'ad, N. Mappaware, M. Hatta, Rabia, Effect of ajwa dates consumption to inhibit the progression of preeclampsia threats on mean arterial pressure and roll-over test, *BioMed Res. Int.* 2019 (2019) 1–5, <https://doi.org/10.1155/2019/2917895>.
- [21] E. Mulyawan, M. Ahmad, A. Islam, M. Massi, M. Hatta, S. Arif, Effect of valerian extract on GABRB3 gene mRNA expression and sedation in BALB/C mice, *Curr. Bioact. Compd.* 16 (8) (2020) 1249–1257, <https://doi.org/10.2174/15734072146661809>, 2020.
- [22] E. Mulyawan, M. Ahmad, A. Islam, M. Massi, M. Hatta, S. Arif, Analysis of GABRB3 protein level after administration of valerian extract (*valeriana officinalis*) in BALB/c mice, *Phcog. J.* 12 (4) (2020) 821–827, <https://doi.org/10.5530/pj.2020.12.118>.
- [23] R. Nasution, A. Islam, M. Hatta, Prihantono, Decreased neutrophil levels in mice with traumatic brain injury after cape administration, *Annal. Med. Surg.* 54 (2020) 89–92.
- [24] T. Tommy, A. Islam, M. Hatta, A. Bukhari, Immunomodulatory properties of high mobility group box 1 and its potential role in brain injury: review article, *Annal. Med. Surg.* 59 (2020) 106–109.
- [25] A. Wu, L. He, W. Long, Q. Zhou, S. Zhu, P. Wang, et al., Novel mechanisms of herbal therapies for inhibiting HMGb1 secretion or action, *Evid. base Compl. Alternative Med.* 2015 (2015) 1–11.
- [26] T.X. Dou, J.F. Shi, Y. Li, F.C. Bi, H.J. Gao, C.H. Hu, C.Y. Li, Q.S. Yang, G.M. Deng, O. Sheng, W.D. He, G.J. Yi, T. Dong, Influence of harvest season on volatile aroma constituents of two banana cultivars by electronic nose and HS-SPME coupled with GC-MS, *Scientia Horticulture* 265 (2020), 109214.
- [27] A. D'Hont, F. Denoed, J.M. Aury, The banana (*Musa acuminata*) genome and the evolution of monocotyledonous plants, *Nature* 488 (2012) 213–217.
- [28] Food and Agriculture Organization (Fao), FAOSTAT, Food and Agriculture Organization, Geneva: Switzerland, 2013.
- [29] K. Akamine, T. Koyama, K. Yazawa, Banana peel extract suppressed prostate gland enlargement in testosterone-treated mice, *Biosci., Biotechnol., Biochem.* 73 (9) (2009) 1911–1914.
- [30] T. Happi Emaga, C. Robert, S. Ronkart, B. Wathélet, M. Paquot, Dietary fibre components and pectin chemical features of peels during ripening in banana and plantain varieties, *Bioresour. Technol.* 99 (10) (2008) 4346–4354.
- [31] B. Singh, J. Singh, A. Kaur, N. Singh, Bioactive compounds in banana and their associated health benefits – a review, *Food Chem.* 206 (2016) 1–11.
- [32] A.H.S. Heng, F.T. Chew, Systematic review of the epidemiology of acne vulgaris, *Sci. Rep.* 10 (1) (2020) 5754.
- [33] E. Bagatin, D.L. Timpano, Lr Dos S. Guadanhim, V.M.A. Nogueira, L.R. Terzian, D. Steiner, et al., Acne vulgaris: prevalence and clinical forms in adolescents from São Paulo, Brazil, *An. Bras. Dermatol.* 89 (3) (2014) 428–435.
- [34] T. Vos, A.D. Flaxman, M. Naghavi, R. Lozano, C. Michaud, M. Ezzati, et al., Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010, *Lancet* 380 (9859) (2012) 2163–2196.
- [35] L. Fox, C. Csongradi, M. Aucamp, J. du Plessis, M. Gerber, Treatment modalities for acne, *Molecules* 21 (8) (2016) 1063.
- [36] L.K. Ogé, A. Broussard, M.D. Marshall, Acne vulgaris: diagnosis and treatment, *Am. Fam. Physician* 100 (8) (2019) 475–484.
- [37] B. Dréno, What is new in the pathophysiology of acne, an overview, *J. Eur. Acad. Dermatol. Venereol.* 31 (2017) 8–12.
- [38] A. Zaenglein, A. Pathy, B. Schlosser, A. Alikhan, H. Baldwin, D. Berson, et al., Guidelines of care for the management of acne vulgaris, *J. Am. Acad. Dermatol.* 74 (5) (2016) 945–973.
- [39] A. Kucharska, A. Szmurlo, B. Sińska, Significance of diet in treated and untreated acne vulgaris, *Adv. Dermatol. Allergol.* 23 (2) (2016) 81–86.
- [40] A.K.C. Leung, B. Barankin, J.L. Lam, K.F. Leong, K.L. Hon, Dermatology: how to manage acne vulgaris, *Drugs Context* 10 (2021), 2021.
- [41] Y. Wang, T.R. Hata, Y.L. Ton, M.S. Kao, C.C. Zouboulis, R.L. Gallo, C.M. Huang, The anti-inflammatory activities of *Propionibacterium acnes* CAMP factor-targeted acne vaccines, *J. Invest. Dermatol.* 138 (2018) 2355–2364.
- [42] T. Ye, J. Wu, Z. Xu, J. Chai, Q. Zeng, B. Zeng, et al., Esc-1GN shows therapeutic potentials for acne vulgaris and inflammatory pain, *J. Pept. Sci.* 26 (8) (2020).
- [43] C. Mayslich, P. Grange, N. Dupin, Cutibacterium acnes as an opportunistic pathogen: an update of its virulence-associated factors, *Microorganisms* 9 (2) (2021) 303.
- [44] J. Danis, M. Mellett, Nod-like receptors in host defence and disease at the epidermal barrier, *Int. J. Mol. Sci.* 22 (9) (2021) 4677.
- [45] J. Kim, Review of the innate immune response in acne vulgaris: activation of toll-like receptor 2 in acne triggers inflammatory cytokine responses, *Dermatology* 211 (3) (2005) 193–198.
- [46] M. Kistowska, S. Gehrke, D. Jankovic, K. Kerl, A. Fettelschoss, L. Feldmeyer, G. Fenini, A. Kolios, A. Navarini, R. Ganceviciene, J. Schaubert, E. Contassot, L. E. French, IL-1 β drives inflammatory responses to propionibacterium acnes in vitro and in vivo, *J. Invest. Dermatol.* 134 (2014) 677–685.
- [47] P. Martin, J.D. Goldstein, L. Mermoud, A. Diaz-Barreiro, G. Palmer, IL-1 family antagonists in mouse and human skin inflammation, *Front. Immunol.* 12 (2021), 652846.
- [48] S.B. Clarke, A.M. Nelson, R.E. George, D.M. Thiboutot, Pharmacologic modulation of sebaceous gland activity: mechanisms and clinical applications, *Dermatol. Clin.* 25 (2) (2007) 137–146.
- [49] B. Zhang, Y. Choi, J. Lee, I. An, L. Li, C. He, et al., Toll-like receptor 2 plays a critical role in pathogenesis of acne vulgaris, *Biomed. Dermatol.* 3 (1) (2019).
- [50] E.A. Tanghetti, The role of inflammation in the pathology of acne, *J. Clin. Aesthetic Dermatol.* 6 (9) (2013) 27–35.
- [51] S. Mansour, O. Pena, R. Hancock, Host defense peptides: front-line immunomodulators, *Trends Immunol.* 35 (9) (2014) 443–450.
- [52] K. Woodburn, J. Jaynes, L. Clemens, Evaluation of the antimicrobial peptide, RP557, for the broad-spectrum treatment of wound pathogens and biofilm, *Front. Microbiol.* 10 (2019), <https://doi.org/10.3389/fmicb.2019.01688>.
- [53] J. Kim, B.E. Kim, K. Ahn, D.Y.M. Leung, Interactions between atopic dermatitis and *Staphylococcus aureus* infection: clinical implications, *Allergy Asthma Immunol. Res.* 11 (2019) 593–603, <https://doi.org/10.4168/air.2019.11.5.593>.
- [54] L. Heinbockel, S. Sánchez-Gómez, G. Martínez de Tejada, S. Dömming, J. Brandenburg, Y. Kaconis, et al., Preclinical investigations reveal the broad-spectrum neutralizing activity of peptide pep19-2.5 on bacterial pathogenicity factors, *Antimicrob. Agents Chemother.* 57 (3) (2013) 1480–1487, <https://doi.org/10.1128/AAC.02066-12>.
- [55] X. Xu, R. Lai, The chemistry and biological activities of peptides from Amphibian skin secretions, *Chem. Rev.* 115 (4) (2015) 1760–1846, <https://doi.org/10.1021/cr4006704>.
- [56] B. Zeng, J. Chai, Z. Deng, T. Ye, W. Chen, D. Li, et al., Functional characterization of a novel lipopolysaccharide-binding antimicrobial and anti-inflammatory peptide in vitro and in vivo, *J. Med. Chem.* 61 (23) (2018) 10709–10723, <https://doi.org/10.1021/acs.jmedchem.8b01358>.
- [57] I. Zelezetsky, A. Pontillo, L. Puzzi, N. Antcheva, L. Segat, S. Pacor, et al., Evolution of the primate cathelicidin, *J. Biol. Chem.* 281 (29) (2006) 19861–19871, <https://doi.org/10.1074/jbc.M511108200>.
- [58] M. Marta Guarna, R. Coulson, E. Rubinchik, Anti-inflammatory activity of cationic peptides: application to the treatment of acne vulgaris, *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Lett.* 257 (1) (2006) 1–6, <https://doi.org/10.1111/j.1574-6968.2006.00156.x>.
- [59] L.M. Tian, D. Ke, Acne vulgaris is associated with the human β -Defensin 1-gene polymorphisms in han Chinese ethnic group patients, *Clin. Cosmet. Invest. Dermatol.* 14 (2021) 123–128.
- [60] S. Popovic, E. Urbán, M. Lukic, J. Conlon, Peptides with antimicrobial and anti-inflammatory activities that have therapeutic potential for treatment of acne vulgaris, *Peptides* 34 (2) (2012) 275–282.
- [61] S. Ryu, H. Han, P. Song, C. Armstrong, Y. Park, Suppression of propionibacterium acnes infection and the associated inflammatory response by the antimicrobial peptide P5 in mice, *PLoS One* 10 (7) (2015), e0132619.
- [62] R. Han, H. Blencke, H. Cheng, C. Li, The antimicrobial effect of CEN1HC-Br against *Propionibacterium acnes* and its therapeutic and anti-inflammatory effects on acne vulgaris, *Peptides* 99 (2018) 36–43.
- [63] M. Simmaco, G. Kreil, D. Barra, Bombinins, antimicrobial peptides from Bombina species, *Biochim. Biophys. Acta Biomembr.* 1788 (8) (2009) 1551–1555.
- [64] L. Qiu, G. Zhao, H. Wu, L. Jiang, Investigation of combined effects of independent variables on extraction of pectin from banana peel using response surface methodology, *Carbohydr. Polym.* 80 (2) (2010) 326–331.
- [65] P. Phuaklee, S. Ruangnoo, A. Itharut, Anti-inflammatory and antioxidant activities of extracts from *Musa sapientum* peel, *J. Med. Assoc. Thai.* 95 (Suppl 1) (2012) S142–S146.
- [66] A. Amalia, S. Sulistiyowati, The effect of banana skin on acne vulgaris, *J. Keperawatan* 10 (1) (2019) 1.

- [67] A.K. Anal, S. Jaisanti, A. Noomhorm, Enhanced yield of phenolic extracts from banana peels (*Musa acuminata* Colla AAA) and cinnamon barks (*Cinnamomum varum*) and their antioxidative potentials in fish oil, *J. Food Sci. Technol.* 51 (10) (2014) 2632–2639.
- [68] A. Pereira, M. Maraschin, Banana (*Musa spp*) from peel to pulp: ethnopharmacology, source of bioactive compounds and its relevance for human health, *J. Ethnopharmacol.* 160 (2015) 149–163.
- [69] Danc Von Atzingen, Ar Dos A. Mendonça, M.M. Filho, V.A. Alvarenga, V.A. Assis, A.E. Penazzo, et al., Repair of surgical wounds in rats using a 10% unripe *Musa sapientum* peel gel, *Acta Cir. Bras.* 30 (9) (2015) 586–592.
- [70] S. Someya, Y. Yoshiki, K. Okubo, Antioxidant compounds from bananas (*Musa cavendish*), *Food Chem.* 79 (2002) 351–354.
- [71] A. Subagio, N. Morita, S. Sawada, Carotenoids and their fatty acid esters in banana peel, *J. Nutr. Sci. Vitaminol.* 42 (1996) 553–566.
- [72] M.A. Khan, A.H.M.G. Sarwar, R. Rahat, R.S. Ahmed, S. Umar, Stigmasterol protects rats from collagen induced arthritis by inhibiting proinflammatory cytokines, *Int. Immunopharm.* 85 (2020), 106642.
- [73] M. Khalili, M. Alavi, E. Esmaeil-Jamat, T. Baluchnejadmojarad, M. Roghani, Trigonelline mitigates lipopolysaccharide-induced learning and memory impairment in the rat due to its anti-oxidative and anti-inflammatory effect, *Int. Immunopharm.* 61 (2018) 355–362.
- [74] T.P. Cushnie, B. Cushnie, A.J. Lamb, Alkaloids: an overview of their antibacterial, antibiotic-enhancing and antivirulence activities, *Int. J. Antimicrob. Agents* 44 (2014) 377–386.
- [75] J. Di Gesso, Investigating the Effects of Dietary Flavonoids and Their Metabolites on Biomarkers of Inflammation and Antioxidant Defence [internet], Norwich Medical School, Norwich, 2015 [cited 2022 Jul 22]. Available from: <https://www.researchgate.net/pub>.
- [76] Y.M. Liu, J.D. Shen, L.P. Xu, H.B. Li, Y.C. Li, L.T. Yi, Ferulic acid inhibits neuroinflammation in mice exposed to chronic unpredictable mild stress, *Int. Immunopharm.* 45 (2017) 128–134.
- [77] P. Yin, Z. Zhang, J. Li, Y. Shi, N. Jin, W. Zou, et al., Ferulic acid inhibits bovine endometrial epithelial cells against LPS-induced inflammation via suppressing NK-kB and MAPK pathway, *Res. Vet. Sci.* 126 (2019) 164–169.