
Correction

Correction to: Low-intensity cigarette smoking and mortality risks: a pooled analysis of prospective cohort studies in Japan

Maki Inoue-Choi  , Neal D Freedman, Eiko Saito, Shiori Tanaka  , Mayo Hirabayashi, Norie Sawada  , Shoichiro Tsugane  , Yoshiaki Usui, Hidemi Ito, Chaochen Wang  , Akiko Tamakoshi, Taro Takeuchi, Yuri Kitamura, Mai Utada  , Kotaro Ozasa  , Yumi Sugawara  , Ichiro Tsuji  , Keiko Wada  , Chisato Nagata, Taichi Shimazu  , Tetsuya Mizoue, Keitaro Matsuo  , Mariko Naito, Keitaro Tanaka, Kota Katanoda  and Manami Inoue  ; for the Research Group for the Development and Evaluation of Cancer Prevention Strategies in Japan

Int J Epidemiol 2022;51:1276–90.

In Figure 2, the plots for all-cause mortality in both men and women and the plot for smoking-related cancer mortality in men were incorrect.

Specifically:

- All-cause mortality in men was shown as the plot for all-cause mortality in women and the 95% confidence

intervals at 1–2, 3–5, and >40 cigarettes per day were also shown incorrectly.

- All-cause mortality in women was shown as the plot for smoking-related cancer mortality in men.
- Smoking-related cancer mortality in men was shown as the plot for all-cause mortality in men.

The Figure has been corrected.