



## Original Research Article

# Fermented mixed feed alters growth performance, carcass traits, meat quality and muscle fatty acid and amino acid profiles in finishing pigs

Shiqi Liu <sup>a, b, c, d</sup>, Man Du <sup>a, b, c, d</sup>, Yuang Tu <sup>a, b, c, d</sup>, Wenjing You <sup>a, b, c, d</sup>,  
Wentao Chen <sup>a, b, c, d</sup>, Guoliang Liu <sup>e</sup>, Junyue Li <sup>e</sup>, Yizhen Wang <sup>a, b, c, d</sup>,  
Zeqing Lu <sup>a, b, c, d</sup>, Tenghao Wang <sup>e, \*</sup>, Tizhong Shan <sup>a, b, c, d, \*</sup>

<sup>a</sup> College of Animal Sciences, Zhejiang University, Hangzhou, China

<sup>b</sup> Key Laboratory of Molecular Animal Nutrition (Zhejiang University), Ministry of Education, Hangzhou, China

<sup>c</sup> Key Laboratory of Animal Nutrition and Feed Science (Eastern of China), Ministry of Agriculture and Rural Affairs, Hangzhou, China

<sup>d</sup> Key Laboratory of Animal Feed and Nutrition of Zhejiang Province, Hangzhou, China

<sup>e</sup> Zhejiang Qinglian Food Co Ltd, Jiaxing, China

## ARTICLE INFO

## Article history:

Received 12 May 2022

Received in revised form

20 August 2022

Accepted 20 September 2022

Available online 27 September 2022

## Keywords:

Fermented mixed feed

Growth performance

Meat quality

Amino acid profile

Fatty acid composition

Finishing pig

## ABSTRACT

This study was conducted to investigate the effects of fermented mixed feed (FMF) on growth performance, carcass traits, meat quality, muscle amino acid and fatty acid composition and mRNA expression levels of genes related to lipid metabolism in finishing pigs. In the present study, 144 finishing pigs (Duroc × Berkshire × Jiaxing Black) were randomly allocated to 3 dietary treatments with 4 replicate pens per group and 12 pigs per pen. The dietary treatments included a basal diet (CON), a basal diet + 5% FMF and a basal diet + 10% FMF. The experiment lasted 38 d after 4 d of acclimation. The results showed that 5% and 10% FMF significantly increased the average daily gain (ADG) of the females but not the males ( $P < 0.05$ ), but FMF supplementation showed no impact on carcass traits. Moreover, 10% FMF supplementation increased the meat color<sub>45 min</sub> and meat color<sub>24 h</sub> values, while it decreased the shear force relative to CON ( $P < 0.05$ ). In addition, 10% FMF significantly increased the contents of flavor amino acids (FAA), total essential AA (EAA), total non-EAA (NEAA) and total AA relative to CON ( $P < 0.05$ ). Furthermore, the diet supplemented with 10% FMF significantly increased the concentration of n-3 polyunsaturated fatty acids (PUFA), n-6 PUFA and total PUFA, and the PUFA to saturated fatty acids ratio ( $P < 0.05$ ), suggesting that FMF supplementation increased meat quality. Moreover, compared with the CON, 10% FMF supplementation increased the mRNA expression of lipogenic genes, including *CEBPα*, *PPARγ*, *SREBP1* and *FABP4*, and upregulated the expression of unsaturated fatty acid synthesis (*ACAA1* and *FADS2*). Together, our results suggest that 10% FMF dietary supplementation improved the female pigs' growth performance, improved the meat quality and altered the profiles of muscle fatty acids and amino acids in finishing pigs. This study provides a reference for the production of high-quality pork.

© 2022 The Authors. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding authors.

E-mail addresses: [wangth85@126.com](mailto:wangth85@126.com) (T. Wang), [tzshan@zju.edu.cn](mailto:tzshan@zju.edu.cn) (T. Shan).

Peer review under responsibility of Chinese Association of Animal Science and Veterinary Medicine.



Production and Hosting by Elsevier on behalf of KeAi

<https://doi.org/10.1016/j.aninu.2022.09.003>

2405-6545/© 2022 The Authors. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Pork is one of the most widely consumed meats and among the most important sources of protein for humans. In recent years, consumer demand for high-quality and healthy pork has been increasing, hence improving the meat quality of pork is an effective practice in the development of animal husbandry. The evaluation indices of meat quality include shear force, drip loss, meat color, pH value and intramuscular fat (IMF) content (Frank et al., 2016; Lomiwes et al., 2014), among which the IMF content is closely related to the sensory characteristics of meat, including flavor,

juiciness and tenderness (Hao et al., 2020; Van Elswyk and McNeill, 2014). Additionally, the composition and content of amino acids, especially the FAA, greatly affect the flavor and taste of meat (Pereira and Vicente, 2013). Furthermore, the content and composition of fatty acids in muscle play a key role in meat quality, which determines the nutritional value and flavor of the meat. Previous studies have demonstrated that polyunsaturated fatty acids (PUFA) are tied to anti-obesity and anti-inflammatory properties, particularly n-3 PUFA (Buckley and Howe, 2009; Tortosa-Caparrós et al., 2017). Nutritional regulation is an effective means to improve pork quality. Indeed, some feed additives, such as lycopene,  $\beta$ -glucan and betaine, have been shown to increase the IMF content and improve the pork amino acid and fatty acid profiles in finishing pigs (Luo et al., 2019; Wen et al., 2022; Zhong et al., 2021).

Fermented mixed feed (FMF) is a product of microbial fermentation, which effectively degrades antinutritional factors in feed while producing probiotics and their metabolites (Cao et al., 2012; Shi et al., 2015). Our previous study showed that microbial fermentation improved feed nutritional value and bioavailability (Hao et al., 2020; Shi et al., 2017; Wang et al., 2018). Furthermore, dietary FMF supplementation contributes to improving growth performance and meat quality, enhancing gut health and reducing the diarrhea rate in animal production (Ding et al., 2020; Kiers et al., 2003; Missotten et al., 2013; Tang et al., 2021). Our group also found that growth performance and meat quality were improved in finishing pigs (Duroc  $\times$  Landrace  $\times$  Large White) fed FMF (Hao et al., 2020). Taken together, FMF shows great application prospects to improve the growth performance of pigs and the quality of meat produced.

However, the effects of dietary FMF supplementation on growth performance and meat quality in Duroc  $\times$  Berkshire  $\times$  Jiaying Black pigs have not been reported. Here, we used corn, soybean meal, and wheat bran as substrates and fermented them with a complex microbial combination (*Bacillus subtilis* and *Enterococcus faecalis*) to obtain FMF. Notably, the basal substrate and microbial combination were different from our previous studies (Hao et al., 2020). Therefore, this study aimed to explore the effects of FMF supplementation on the growth performance, carcass traits, meat quality, muscle fatty acid and amino acid profiles and mRNA expression levels of genes related to lipid metabolism in the longissimus dorsi muscle (LDM) in finishing Duroc  $\times$  Berkshire  $\times$  Jiaying Black pigs.

## 2. Materials and methods

### 2.1. Animal ethics statement

All the procedures were approved by the Institutional Animal Care and Use Committee at Zhejiang University.

### 2.2. Fermented mixed feed preparation and chemical analysis

The FMF preparation method was based on previous research (Wang et al., 2018). *Enterococcus faecium* (*E. faecium*) was obtained from Baolai-leelai Biotech Co., Ltd (Tai'an, China). The *B. subtilis* (*B. subtilis*) ZJU12 used in this study was isolated from traditional fermented food (pickled vegetables). Pilot production of FMF was carried out at the Kesheng Feed Co., Ltd, Zhejiang, China. The basal substrate contained 40% corn, 40% soybean meal and 20% wheat bran. Whilst stirring evenly, sterile water was added to make the total moisture in system 40%. One kilogram of *E. faecium* powder ( $10^8$  cfu/g) and 1.25 kg of *B. subtilis* powder ( $3 \times 10^8$  cfu/g) was inoculated into 1,000 kg of wet mixed substrate. Then, the wet mixed substrate was transferred to a plastic bag with a one-way valve (Rou Duoduo Biotechnology Co., Beijing, China), sealed, and fermented at room temperature for 72 h.

**Table 1**

Nutrient composition of mixed feed and fermented mixed feed (as-fed basis).

Item	MF	FMF
DM, %	92.77	93.01
CP, %	26.39	28.03
TCA-SP, %	2.58	5.05
TCA-SP:CP ratio, %	9.77	18.01
NDF, %	34.11	32.45
ADF, %	20.02	19.42
Hemicellulose, %	14.09	13.03
Lactic acid, mmol/kg	96.67	117.78
pH	6.62	3.84
Live BS cells, cfu/g	—	$1.3 \times 10^8$
Live EF cells, cfu/g	—	$8.0 \times 10^8$

MF = mixed feed; FMF = fermented mixed feed (40% corn, 40% soybean meal, 20% wheat bran); CP = crude protein; TCA-SP = trichloroacetic acid-soluble protein (small peptides); TCA-SP:CP ratio = TCA-SP to CP ratio; NDF = neutral detergent fiber; ADF = acid detergent fiber; Hemicellulose = NDF-ADF; BS = *Bacillus subtilis*; EF = *Enterococcus faecium*.

The contents of dry matter (DM), crude protein (CP), neutral detergent fiber (NDF) and acid detergent fiber (ADF) in mixed feed (MF) and fermented mixed feed (FMF) were analyzed according to the AOAC International guidelines (2005). The trichloroacetic acid-soluble protein (TCA-SP) was measured according to reported methods (Ovissipour et al., 2009). The lactic acid content was evaluated using a commercial assay kit (Nanjing Jiancheng Bio Co., Nanjing, China) according to the manufacturer's instructions. Chemical analysis of the MF and FMF is presented in Table 1.

### 2.3. Experimental design and diets

A total of 144 finishing pigs (body weight =  $77.21 \pm 8.31$  kg) were randomly divided into 3 dietary treatments with 4 replications (12 pigs per replication). Each treatment contained 2 male replicates and 2 female replicates. In addition, each group consisted of half males and half females. The dietary treatments included a basal diet (CON), a basal diet + 5% FMF and a basal diet + 10% FMF. All experimental diets were formulated to meet the nutritional requirements of finishing pigs according to the Chinese National Feeding Standard for swine and contained similar levels of crude protein (CP) and metabolizable energy (ME) (Table 2). All pigs were weighed after 4 d of prefeeding. The experiment lasted 38 days. All pigs had free access to water and food and were fed 3 times (at 07:30, 13:30 and 17:30) per day.

### 2.4. Sample collection and carcass characteristics measurements

All pigs were weighed on d 38 and fasted for 12 h before slaughter. Then, 18 pigs (6 pigs/treatment group) were selected and slaughtered humanely after electrical stunning in a commercial slaughterhouse (Qinglian Food Co., Ltd., Zhejiang, China). After slaughter, the hot carcass weight of each pig was measured and used to calculate carcass yield. The oblique length and straight length were measured using band tape. The backfat thickness at the thickest part of the shoulder, thoracolumbar junction and lumbar-sacral junction were recorded and used to calculate the average backfat value. Additionally, the LDM was removed for on-site testing and the rest of the muscle samples were quickly frozen in liquid nitrogen and stored at  $-80^\circ\text{C}$  to determine LDM amino acids, fatty acids and gene expression.

### 2.5. Meat quality measurements

The LDM samples stored at  $4^\circ\text{C}$  were used for pH, meat color, drip loss, shear force and marbling score analyses. Meat quality

**Table 2**  
Ingredients and nutrient levels of experimental diets (% of as-fed basis).<sup>1</sup>

Item	Diet		
	CON	5% FMF	10% FMF
<b>Ingredients</b>			
Corn	40.50	36.20	32.10
Soybean meal	14.00	13.30	12.60
Barley	15.00	15.00	15.00
Rice bran meal	4.00	3.00	1.80
Flour	8.00	8.00	8.00
Defatted rice bran	8.00	8.00	8.00
Wheat middlings	6.00	6.00	6.00
Soyabean oil	1.00	2.00	3.00
FMF	0.00	5.00	10.00
Premix <sup>2</sup>	3.50	3.50	3.50
<b>Total</b>	<b>100.00</b>	<b>100.00</b>	<b>100.00</b>
<b>Nutrient level</b>			
Digestible energy, MJ/kg	13.44	13.43	13.43
Crude protein	15.37	15.37	15.36
Crude fat	4.43	5.33	6.23
Crude fiber	3.38	3.34	3.28
Calcium	0.57	0.57	0.58
Total phosphorus	0.58	0.56	0.54
Available phosphorus	0.21	0.20	0.20
Lysine	0.95	0.95	0.96
Methionine	0.30	0.30	0.30
Methionine + Cystine	0.57	0.57	0.57
Threonine	0.65	0.65	0.65
Tryptophan	0.17	0.17	0.17
Valine	0.71	0.71	0.71

CON = basal diet; FMF = fermented mixed feed.

<sup>1</sup> Basal diet formulated according to the Chinese National Feeding Standard for swine.

<sup>2</sup> Provided the following per kilogram of complete diet: vitamin A, 7,500 IU; vitamin D<sub>3</sub>, 1,800 IU; vitamin E, 54 IU; vitamin K<sub>3</sub>, 4.8 mg; vitamin B<sub>12</sub>, 0.024 mg; vitamin B<sub>1</sub>, 2.7 mg; vitamin B<sub>2</sub>, 7.2 mg; vitamin B<sub>6</sub>, 5.4 mg; D-biotin, 0.18 mg; folic acid, 0.9 mg; nicotinamide, 36 mg; D-pantothenic acid, 24 mg; Cu, 15 mg; Fe, 50 mg; Zn, 45 mg; Mn, 20 mg; I, 0.5 mg; Se, 0.35 mg.

evaluation was carried out according to a previous report (Nong et al., 2020). After slaughter, the LDM pH value was measured at 45 min and 24 h with a portable pH meter (pH-Star Matthäus GmbH, Pöttmes, Germany). The meat color values, including lightness (L\*), redness (a\*) and yellowness (b\*), were also evaluated at 45 min and 24 h after slaughter using a Minolta CM-2002 spectrophotometer (Osaka, Japan). For drip loss, approximately 10 g of each LDM sample was hung in a special sealed plastic tube at 4 °C and weighed after 24 h. The shear force was measured with a C-LM tenderness tester (Tenovo International Co., Limited, Beijing, China) according to the manufacturer's instructions. The marbling score at 45 min after slaughter was evaluated according to an NPPC meat color chart (Nanjing Mingao Instrument Equipment Co., Ltd., Nanjing, China).

## 2.6. Amino acid composition

First, approximately 0.1 g LDM of each sample was weighed and digested with 5 mL of 6 mol/L HCl solution in 105 °C oven for 24 h. Then, the volume was brought up to 50 mL in a volumetric flask and the sample was filtered through a 0.22-µm water phase filter into a centrifuge tube. Next, 2 mL of the filtrate was placed in an evaporating dish in a 60 °C water bath for evaporation and 4 mL of 0.02 mol/L HCl solution was added. After dissolution, the sample was stored at 4 °C for detection with an ion-exchange AA analyzer (L8900, Hitachi, Tokyo, Japan).

## 2.7. Fatty acid composition

The fatty acid composition was determined by gas chromatography (GC) as described previously (Hao et al., 2020). First, the LDM

samples were extracted with a mixture of chloroform and methanol (2:1; vol/vol). Approximately 20 g of each LDM sample was weighed and dried in an oven at 105 °C for 1 h, and then 1 g of each dried sample was weighed and leached with petroleum ether for 3 h. A total of 60 mg of the extracted fat was placed in a test tube, 4 mL of iso-octane was added to fully dissolve the sample, and then 200 µL potassium hydroxide-methanol and 1 g of sodium bisulfate were added. After salt precipitation, the solution containing the methyl esters was drawn into the upper layer and stored in a refrigerator at 4 °C. Each sample was filtered through a 0.22-µm filter membrane before GC detection (Model 7890 A, Agilent Technologies, Palo Alto, CA, USA). Finally, the fatty acid concentration was analyzed by GC ChemStation software (Agilent Technologies, Palo Alto, CA, USA).

## 2.8. Hematoxylin-eosin staining

After slaughter, the LDM was isolated and fixed in fixative purchased from Servicebio (CAT: G1111-100ML) followed by paraffin sectioning to obtain tissue sections. Then, the LDM paraffin sections were stained using a hematoxylin-eosin staining kit (Servicebio, Wuhan, China). In brief, after deparaffinization with xylene, the sections were stained with hematoxylin solution for 5 min. After that, they were immersed in 1% acid ethanol (1% HCl in 70% ethanol) for 10 s and rinsed with running water. Finally, the sections were stained with eosin for 2 to 3 min, dehydrated with pure alcohol and rendered transparent with xylene. The morphological structure and size of the cells were observed with an Olympus BX61 fluorescence microscope (Japan).

## 2.9. Intramuscular fat and triglyceride measurement

The IMF content was evaluated by following the Chinese Agriculture Industry Standard NY/T 2793-2015. For triglyceride measurement, the samples of LDM were homogenized with normal saline solution (tissue weight:normal saline solution = 1:9) and centrifuged for 10 min. The concentrations of triglycerides in the LDM were measured with the corresponding assay kits (Nanjing Jiancheng Bioengineering Institute, China) according to the manufacturer's instructions. In addition, the concentration of protein in the homogenate was detected using a BCA Protein Assay Reagent Kit (Thermo Fisher Scientific Inc., Waltham, MA, USA).

## 2.10. Real-time quantitative PCR

RNA extraction, library construction and quantitative real-time PCR (qPCR) of the LDM samples were performed as described in our previous study (Shan et al., 2013; Xu et al., 2021). Briefly, total RNA was isolated from the LDM muscle using TRIzol reagent (Thermo Fisher, Waltham, MA, USA) according to the instructions of the manufacturer as previously described (Shan et al., 2013; Xu et al., 2021). RNA quality and concentration were measured with a NanoDrop 2000 instrument (Gene Company Limited, Hong Kong, China). A ReverAid First Strand cDNA Synthesis Kit (Thermo Fisher) and random primers were used to generate cDNA by reverse transcription. The primer sequences are listed in Table 3. Quantitative real-time PCR was performed with an Applied Biosystems StepOnePlus Real-Time PCR System with SYBR Green Master Mix (Roche, Indianapolis, IN, United States). Relative gene expression was analyzed by the  $2^{-\Delta\Delta CT}$  method.

## 2.11. Statistical analyses

All data in the current study were analyzed by one-way analysis of variance (ANOVA) with the statistical software SPSS 20.0

**Table 3**  
Primer sequences used in this study.

Gene	Primer	Sequence (5' to 3')	GenBank ID	Size, bp
GAPDH	Forward	AAGGAGTAAGAGCCCTGGA	NM_001206359.1	140
	Reverse	TCTGGGATGGAACCTGGAA		
CEBP $\alpha$	Forward	AACAACCTGAGCCGAACTGGA	XM_003127015.4	170
	Reverse	CTTGAGATCTGGAGACCCGAAACC		
PPAR $\gamma$	Forward	AAGACGGGGTCTCATCTCC	NM_214379.1	149
	Reverse	CGCCAGGTCGCTGTCATCT		
SREBP1	Forward	AAGCGGACGGCTCACAATG	NM_214157.1	122
	Reverse	CGCAAGACGGCGGATTTAT		
ACC $\alpha$	Forward	TTCAGGCACAGTCCITAGG	NM_001114269.1	161
	Reverse	TCATCCAACACGAGCTCAGT		
FABP4	Forward	TGGAAACTGTCTCCAGTG	NM_001002817.1	147
	Reverse	GGTACTTTCTGATCTAATGGTG		
ACAA1	Forward	CGAGCTTCTCTGCAGTCAT	XM_003132103.4	148
	Reverse	TGGGATGTCACCTAGAACTGG		
FADS2	Forward	GCTGGATTCCAACCTCATG	NM_0011171750.1	56
	Reverse	AGCCTGGCCCTGAGAGGTA		
SCD	Forward	GCCACCTTTCTCGTTACG	NM_213781.1	142
	Reverse	CCTCACCCACAGCTCCAAT		

GAPDH = glyceraldehyde-3-phosphate dehydrogenase; CEBP $\alpha$  = CCAAT enhancer-binding protein  $\alpha$ ; PPAR $\gamma$  = peroxisome proliferator-activated receptor  $\gamma$ ; SREBP1 = sterol regulatory element binding protein 1; ACC $\alpha$  = acetyl CoA carboxylase  $\alpha$ ; FABP4 = fatty acid-binding protein 4; ACAA1 = acetyl-CoA acyltransferase 1; FADS2 = fatty acid desaturase 2; SCD = stearyl Coenzyme A desaturase.

(Chicago, IL, USA) followed by Duncan's multiple range analysis.  $P < 0.05$  was considered to be statistically significant. All results are shown as the means and standard error of the mean (SEM).

### 3. Results

#### 3.1. Nutrient composition of the fermented mixed feed

The nutrient compositions of MF and FMF are shown in Table 1. The contents of DM, CP and lactic acid in FMF were higher than those in unfermented MF, while FMF contained less NDF, ADF and hemicellulose. Furthermore, as presented in the Supplementary Data (Tables S1 and S2), we also analyzed the amino acid and fatty acid compositions of MF and FMF. After fermentation, the contents of EAA, NEAA and total AA were higher in FMF than those in unfermented MF. Regarding fatty acid composition, microbial fermentation improved the contents of monounsaturated fatty acid (MUFA) and PUFA, while reducing the levels of saturated fatty acids (SFA).

#### 3.2. Growth performance

The growth performance of the finishing pigs is presented in Table 4. The ADG values of the females in the 5% and 10% FMF groups were significantly higher than those in the CON group ( $P = 0.026$ ), however, the ADG of the males showed no significant difference. In addition, there was no difference in the other indices of growth performance.

#### 3.3. Carcass characteristics

As shown in Table 5, the live weights and carcass traits were higher in the 10% FMF group than those in the CON group ( $P < 0.05$ ). No difference in carcass characteristics was found between the 5% FMF group and the CON group. In addition, compared with the 5% FMF group, 10% FMF supplementation significantly increased backfat thickness at the lumbar-sacral junction ( $P = 0.018$ ).

**Table 4**  
Effects of fermented mixed feeds on the growth performance of finishing pigs.

Item	Treatment			SEM	P-value
	CON	5% FMF	10% FMF		
<b>Males</b>					
Initial weight, kg	76.88	77.40	75.75	0.95	0.776
Final weight, kg	108.73	109.42	109.13	1.22	0.974
ADG, kg/d	0.84	0.84	0.88	0.01	0.459
ADFI, kg/d	3.32	3.28	3.31		
F:G	4.06	3.99	3.88		
<b>Females</b>					
Initial weight, kg	79.67	76.35	77.52	1.03	0.412
Final weight, kg	107.07	106.00	107.74	1.19	0.837
ADG, kg/d	0.72 <sup>b</sup>	0.78 <sup>a</sup>	0.79 <sup>a</sup>	0.01	0.026
ADFI, kg/d	3.19	2.90	3.22		
F:G	4.41	3.75	4.14		

CON = basal diet; FMF = fermented mixed feed; ADG = average daily gain; ADFI = average daily feed intake; F:G = the ratio of feed intake to body weight gain. <sup>a,b</sup> Within a row, values with different superscripts differ significantly at  $P < 0.05$ . Data are expressed as means and SEM,  $n = 24$ .

**Table 5**  
Effects of fermented mixed feeds on the carcass traits of finishing pigs.

Item	Treatment			SEM	P-value
	CON	5% FMF	10% FMF		
Live weight, kg	104.58 <sup>b</sup>	111.33 <sup>ab</sup>	116.08 <sup>a</sup>	1.88	0.032
Carcass weight, kg	77.35 <sup>b</sup>	81.33 <sup>ab</sup>	85.23 <sup>a</sup>	1.32	0.039
Carcass yield, %	72.34	73.04	74.56	0.46	0.129
Carcass oblique length, cm	84.17	88.08	88.17	0.98	0.167
Carcass straight length, cm	97.08	101.33	100.92	1.29	0.354
Skin thick, mm	3.35	3.30	2.93	0.13	0.377
Backfat thickness at the thickest part of the shoulder, mm	46.38	46.07	49.37	1.46	0.622
Backfat thickness at thoracolumbar junction, mm	29.48	33.15	34.18	1.14	0.218
Backfat thickness at lumbar-sacral junction, mm	22.83 <sup>ab</sup>	19.13 <sup>b</sup>	26.22 <sup>a</sup>	1.09	0.018
Backfat, mm	31.75	31.77	37.37	1.50	0.222
Loin-eye area, cm	45.40	48.74	51.07	1.27	0.191

CON = basal diet; FMF = fermented mixed feed. <sup>a,b</sup> Within a row, values with different superscripts differ significantly at  $P < 0.05$ . Data are expressed as means and SEM,  $n = 6$ .

#### 3.4. Meat quality

The effects of FMF on the meat quality of finishing pigs are presented in Table 6. On the one hand, compared with CON, dietary 5% FMF supplementation significantly increased the meat color<sub>45 min</sub> value while significantly reducing the shear force ( $P < 0.05$ ). On the other hand, 10% FMF supplementation significantly increased meat color<sub>45 min</sub> and meat color<sub>24 h</sub> values, and significantly decreased shear force relative to CON ( $P < 0.05$ ). In addition, the pH<sub>24 h</sub> of 10% FMF supplementation was higher than that in the CON group ( $P = 0.051$ ).

#### 3.5. Free amino acid profiles in the longissimus dorsi muscle

The free amino acid profiles in the LDM are shown in Table 7. Both 5% and 10% FMF supplementation significantly increased the contents of flavor AA (FAA), total EAA, total NEAA and total AA relative to CON ( $P < 0.05$ ). In detail, the concentrations of EAA (lysine, methionine, and threonine) and NEAA (alanine, aspartate, glutamate, arginine, serine, and tyrosine) were significantly increased with FMF supplementation ( $P < 0.05$ ). In addition, only 10% FMF supplementation improved the concentration of



**Table 6**  
Effects of fermented mixed feeds on the meat quality of finishing pigs.

Item	Treatment			SEM	P-value
	CON	5% FMF	10% FMF		
pH <sub>45 min</sub>	6.41	5.88	6.21	0.09	0.054
pH <sub>24 h</sub>	5.59	5.52	5.92	0.07	0.051
pH <sub>48 h</sub>	5.67	5.61	5.79	0.07	0.659
Meat color <sub>45 min</sub>	77.09 <sup>b</sup>	82.60 <sup>a</sup>	82.72 <sup>a</sup>	1.06	0.034
L* <sub>45 min</sub>	45.03	42.99	41.69	0.70	0.141
a* <sub>45 min</sub>	9.56	9.91	10.08	0.19	0.557
b* <sub>45 min</sub>	9.28	8.47	8.44	0.19	0.112
Meat color <sub>24 h</sub>	69.95 <sup>b</sup>	72.97 <sup>b</sup>	79.36 <sup>a</sup>	1.40	0.010
L* <sub>24 h</sub>	50.00	49.06	45.70	0.86	0.091
a* <sub>24 h</sub>	11.18	12.03	11.28	0.34	0.578
b* <sub>24 h</sub>	11.58	12.51	10.49	0.45	0.188
Drip loss, %	1.96	1.65	1.55	0.08	0.082
Shear force, N	149.33 <sup>a</sup>	93.62 <sup>b</sup>	96.50 <sup>b</sup>	9.31	0.012
Marbling scores	1.83	1.67	2.17	0.18	0.537

CON = basal diet; FMF = fermented mixed feed.

<sup>a,b</sup> Within a row, values with different superscripts differ significantly at  $P < 0.05$ . Data are expressed as means and SEM,  $n = 6$ .

**Table 7**  
Effects of fermented mixed feeds on the free amino acid profile of the *longissimus dorsi* muscle in finishing pigs (g/kg, as-fresh basis).

Item	Treatment			SEM	P-value
	CON	5% FMF	10% FMF		
<b>EAA</b>					
Lysine	18.45 <sup>b</sup>	19.58 <sup>a</sup>	19.51 <sup>a</sup>	0.17	0.003
Methionine	4.42 <sup>b</sup>	4.75 <sup>a</sup>	4.80 <sup>a</sup>	0.07	0.048
Valine	10.47	10.87	10.88	0.08	0.055
Isoleucine	9.88 <sup>b</sup>	10.23 <sup>ab</sup>	10.33 <sup>a</sup>	0.08	0.047
Leucine	16.12 <sup>b</sup>	16.80 <sup>a</sup>	16.65 <sup>ab</sup>	0.12	0.044
Phenylalanine	6.68 <sup>c</sup>	7.10 <sup>b</sup>	7.47 <sup>a</sup>	0.09	< 0.001
Histidine	9.35	9.65	9.50	0.09	0.395
Threonine	8.60 <sup>b</sup>	9.20 <sup>a</sup>	9.18 <sup>a</sup>	0.10	0.008
<b>NEAA</b>					
Alanine	11.12 <sup>b</sup>	11.73 <sup>a</sup>	11.63 <sup>a</sup>	0.10	0.009
Aspartate	17.80 <sup>b</sup>	19.08 <sup>a</sup>	18.97 <sup>a</sup>	0.20	0.005
Glutamate	28.45 <sup>b</sup>	30.48 <sup>a</sup>	30.25 <sup>a</sup>	0.35	0.024
Arginine	12.48 <sup>b</sup>	13.27 <sup>a</sup>	13.07 <sup>a</sup>	0.13	0.026
Glycine	9.15	9.17	9.23	0.08	0.921
Serine	7.10 <sup>b</sup>	7.60 <sup>a</sup>	7.60 <sup>a</sup>	0.09	0.016
Tyrosine	6.78 <sup>b</sup>	7.23 <sup>a</sup>	7.17 <sup>a</sup>	0.07	0.005
Proline	7.52	7.60	7.43	0.08	0.743
FAA <sup>1</sup>	79.00 <sup>b</sup>	83.73 <sup>a</sup>	83.15 <sup>a</sup>	0.80	0.018
Total EAA	83.97 <sup>b</sup>	88.18 <sup>a</sup>	88.32 <sup>a</sup>	0.74	0.012
Total NEAA	100.40 <sup>b</sup>	106.17 <sup>a</sup>	105.35 <sup>a</sup>	0.99	0.024
Total AA	184.50 <sup>b</sup>	194.67 <sup>a</sup>	193.67 <sup>a</sup>	1.71	0.016
Total protein	205.33 <sup>b</sup>	219.67 <sup>a</sup>	214.50 <sup>ab</sup>	2.29	0.023

CON = basal diet; FMF = fermented mixed feed; EAA = essential amino acids; NEAA = non-essential amino acids; FAA = flavor amino acids.

<sup>a,b</sup> Within a row, values with different superscripts differ significantly at  $P < 0.05$ . Data are expressed as means and SEM,  $n = 6$ .

<sup>1</sup>FAA = glutamate + aspartate + alanine + arginine + glycine.

isoleucine compared with CON ( $P = 0.047$ ). The content of leucine in the 5% FMF group was higher than that in the CON group ( $P = 0.044$ ). Moreover, the concentration of phenylalanine showed a linear increase in response to the FMF supplementation ratio, with a maximum observed in the 10% FMF group ( $P < 0.001$ ). Notably, 5% FMF supplementation markedly increased the total protein content relative to CON ( $P = 0.023$ ).

### 3.6. Fatty acid profiles in the *longissimus dorsi* muscle

Diets supplemented with FMF greatly altered the fatty acid profiles in the LDM (Table 8). Compared with CON, 10% FMF supplementation significantly increased the concentration of n-3

**Table 8**  
Effects of fermented mixed feeds on the fatty acid profile of the *longissimus dorsi* muscle in finishing pigs (g/kg, as-fresh basis).

Item	Treatment			SEM	P-value
	CON	5% FMF	10% FMF		
C10:0	0.11	0.09	0.10	0.00	0.228
C14:0	1.45 <sup>a</sup>	0.99 <sup>b</sup>	1.36 <sup>a</sup>	0.08	0.034
C16:0	26.27	23.45	24.67	1.06	0.580
C16:1	2.10	1.75	1.77	0.11	0.331
C17:0	0.26	0.23	0.26	0.02	0.617
C18:0	11.71	5.61	9.86	1.15	0.077
C18:1n9c	36.58	31.03	38.20	1.57	0.149
C18:2n6c	13.47 <sup>b</sup>	12.87 <sup>b</sup>	17.82 <sup>a</sup>	0.69	0.001
C18:3n3	0.83 <sup>b</sup>	0.90 <sup>b</sup>	1.39 <sup>a</sup>	0.09	0.010
C20:0	0.27 <sup>a</sup>	0.17 <sup>b</sup>	0.24 <sup>a</sup>	0.00	0.001
C20:1	0.25 <sup>ab</sup>	0.20 <sup>b</sup>	0.27 <sup>a</sup>	0.00	0.029
C20:2	0.75	0.65	0.83	0.03	0.118
C20:3n3	0.13 <sup>b</sup>	0.13 <sup>b</sup>	0.18 <sup>a</sup>	0.01	0.003
C20:3n6	0.13	0.14	0.15	0.00	0.242
C20:4n6	0.27	0.37	0.38	0.02	0.088
C22:1n9	0.09	0.08	0.09	0.01	0.796
C24:1	0.08 <sup>b</sup>	0.18 <sup>a</sup>	0.12 <sup>ab</sup>	0.02	0.022
SFA <sup>1</sup>	40.07	30.54	36.48	1.84	0.095
MUFA <sup>2</sup>	39.11	33.25	40.46	1.66	0.170
PUFA <sup>3</sup>	15.58 <sup>b</sup>	15.05 <sup>b</sup>	20.74 <sup>a</sup>	0.80	0.001
PUFA:SFA ratio	0.40 <sup>b</sup>	0.50 <sup>ab</sup>	0.59 <sup>a</sup>	0.29	0.022
n-3 PUFA <sup>4</sup>	0.96 <sup>b</sup>	1.02 <sup>b</sup>	1.57 <sup>a</sup>	0.10	0.008
n-6 PUFA <sup>5</sup>	13.86 <sup>b</sup>	13.37 <sup>b</sup>	18.34 <sup>a</sup>	0.70	0.001
n-6:n-3 PUFA ratio	14.50	13.13	12.21	0.94	0.131

CON = basal diet; FMF = fermented mixed feed; SFA = saturated fatty acid; MUFA = monounsaturated fatty acid; PUFA = polyunsaturated fatty acid.

<sup>a,b</sup> Within a row, values with different superscripts differ significantly at  $P < 0.05$ . Data are expressed as means and SEM,  $n = 6$ .

<sup>1</sup> SFA = C10:0 + C14:0 + C16:0 + C17:0 + C18:0 + C20:0.

<sup>2</sup> MUFA = C16:1 + C18:1n9c + C20:1 + C22:1n9 + C24:1.

<sup>3</sup> PUFA = C18:2n6c + C18:3n3 + C20:2 + C20:3n3 + C20:3n6 + C20:4n6.


<sup>4</sup> n-3 PUFA = C18:3n3 + C20:3n3.

<sup>5</sup> n-6 PUFA = C18:2n6c + C20:3n6 + C20:4n6.

PUFA, n-6 PUFA, total PUFA and the PUFA:SFA ratio ( $P < 0.05$ ). The concentrations of C18:2n6c, C18:3n3, and C20:3n3 were higher in the 10% FMF group than in the CON group ( $P < 0.05$ ). Additionally, 5% FMF supplementation significantly decreased the concentrations of C14:0 and C20:0 ( $P < 0.05$ ), whereas the concentration of C24:1 markedly increased relative to CON ( $P = 0.022$ ). However, the concentration of C20:1 in the 10% FMF group was higher than that in the 5% FMF supplementation group ( $P = 0.029$ ). There were no pronounced differences in the concentrations of MUFA or the n-6 to n-3 ratio ( $P > 0.05$ ).

### 3.7. The IMF contents and mRNA expression levels of genes related to lipid metabolism

As shown in Fig. 1A, FMF supplementation increased the number of adipocytes by H&E staining. Furthermore, 5% and 10% FMF supplementation significantly increased the IMF content in the LDM ( $P < 0.05$ ) (Fig. 1B). Among the three groups, the TG content in the 10% FMF group was significantly higher than that in CON ( $P < 0.05$ ) (Fig. 1C). To further explore how FMF improves the IMF content and fatty acid profile, we detected the mRNA expression levels of genes related to lipid metabolism, such as fatty acid synthesis and transport, in the LDM. As shown in Fig. 1D, the expression of *CEBPα*, *PPARγ* and *SREBP1*, which are key transcription factors for lipid metabolism, in the 10% FMF supplementation group was significantly higher than that in the CON group ( $P < 0.05$ ). Moreover, compared with CON, 10% FMF supplementation significantly upregulated the expression levels of genes related to fatty acid uptake and transport, such as *FABP4* ( $P < 0.05$ ). Furthermore, 10% FMF supplementation significantly increased the expression levels



**Fig. 1.** Effects of fermented mixed feeds on the intramuscular fat content and relative mRNA expression of genes related to lipid metabolism in longissimus dorsi muscle (LDM) of finishing pigs. (A) Hematoxylin and eosin staining analyzed the percentage of adipocytes and the morphological structure of LDM. (B) Intramuscular fat (IMF) content in LDM. (C) Triglyceride (TG) content in LDM. (D) Real-time quantitative PCR analyzed the mRNA expressions of CCAAT enhancer-binding protein  $\alpha$  (*CEBP $\alpha$* ), peroxisome proliferator-activated receptor  $\gamma$  (*PPAR $\gamma$* ), sterol regulatory element binding protein 1 (*SREBP1*), acetyl CoA carboxylase  $\alpha$  (*ACC $\alpha$* ), fatty acid-binding protein 4 (*FABP4*), acetyl-CoA acyltransferase 1 (*ACAA1*), fatty acid desaturase 2 (*FADS2*) and stearoyl Coenzyme A desaturase (*SCD*). Data are presented as the mean and SEM ( $n = 6$ ). <sup>a, b</sup> Bars with different letters indicate significant difference ( $P < 0.05$ ). CON = basal diet; FMF = fermented mixed feed.

of the unsaturated fatty acid synthesis genes *ACAA1* and *FADS2* ( $P < 0.05$ ). However, there was no dramatic difference in the mRNA expression levels of *ACC $\alpha$*  and *SCD*.

#### 4. Discussion

The present study was designed to examine the effects of FMF supplementation on growth performance, carcass characteristics, meat quality and muscle amino acid and fatty acid composition in finishing pigs (Duroc  $\times$  Berkshire  $\times$  Jiaxing Black). We found that FMF supplementation improved the growth performance, meat quality and muscle amino acid and fatty acid compositions. Furthermore, we also found that FMF supplementation upregulated the mRNA expression of genes related to adipogenesis.

Previous studies have shown that FMF provides many advantages to animal growth, such as increased feed intake, nutrient utilization and gut health maintenance (Canibe and Jensen, 2003; Ding et al., 2020; Mukherjee et al., 2016). These effects are attributed to the ability of microorganisms, such as *B. subtilis*, to degrade antinutritional factors and macromolecular nutrients in the feed (Chun-Hua et al., 2016). Our previous study found that *B. subtilis* and *E. faecium* co-fermentation decreased the contents of  $\beta$ -conglycinin and glycinin, two antinutritional factors, while increasing the contents of CP, TCA-SP, Ca and total P in the feed (Wang et al., 2018). Furthermore, FMF improved the performance of females and their progeny. Therefore, in this study, we focused on the effects of *B. subtilis* and *E. faecium* co-fermentation on feed nutrient levels, such as fatty acid and amino acid compositions. We observed that *B. subtilis* and *E. faecium* co-fermentation increased the concentrations of CP, total AA and FAA, and improved the contents of MUFA and PUFA. Intriguingly, the SFA content decreased after *B. subtilis* and *E. faecium* co-fermentation. In line with previous reports, microbial fermentation also increased the concentrations of TCA-SP, a class of easily absorbed small molecule nutrients, and lactic acid (Hao et al., 2020; Wang et al., 2018).

Previous studies have shown that diets supplemented with FMF improved growth performance and nutrient digestibility in finishing pigs (Tian et al., 2020; Xu et al., 2017; Yan et al., 2012). In the present study, we found that FMF supplementation increased the ADG of females, but did not affect the ADG of males. This may be due to gender differences in the effect of FMF on growth performance, but the specific mechanism is unclear. Notably, although the carcass weights were higher in the 10% FMF group than in the CON group, this could be attributed to the difference in live weight and did not account for the benefits of FMF. The slaughter weight was not consistent with the final weight, because we calculated final weights using 48 finishing pigs per group while the calculation of carcass weights only referred to 6 randomly selected pigs. Unlike previous reports (Hao et al., 2020), we did not observe an increase in ocular muscle area with dietary FMF addition. This difference may be attributed to the different breeds of pigs.

The evaluation indicators of meat quality included IMF contents, marbling score, meat color, shear force and drip loss. Previous studies have shown that FMF supplementation increased marbling scores and IMF contents in finishing pigs (Duroc  $\times$  Landrace  $\times$  Large White) (Hao et al., 2020). In the present study, the IMF contents were significantly increased in the FMF supplementation group compared to the CON group, but there was no significant difference in marbling scores between the three groups. In line with previous studies (Guo et al., 2020; Hao et al., 2020), we observed that FMF improved the meat color of the LDM in finishing pigs, as the meat color<sub>45 min</sub> and meat color<sub>24 h</sub> values were greater in the 10% group in comparison to CON, whereas 5% FMF administration also significantly increased the meat color<sub>45 min</sub> values. Muscle shear force represents the tenderness of meat. Others have shown that

FMF supplementation had no effect on muscle shear force (Lu et al., 2020; Tian et al., 2020). Contrary to previous reports, we demonstrated that FMF markedly decreased the shear force of the LDM in finishing pigs. Together, the present findings confirm that dietary FMF supplementation improved meat quality, especially in the 10% FMF group. A popular explanation is that *B. subtilis* and *E. faecium* in FMF play a key role because the antibiotic metabolites from these microbes have been shown to improve meat quality (Meng et al., 2010). Specifically, metabolites produced by microorganisms may be beneficial for improving meat quality, such as through promoting the deposition of IMF and the conversion of fast-twitch fibers to slow-twitch fibers. However, these explanations need further verification.

The amino acid profiles of muscles are closely related to the flavor and nutritional value of meat (Moeller et al., 2010). Recently, nutritional intervention has been recognized as an effective means to improve the quality of pork, including amino acid composition (Xu et al., 2019; Yu et al., 2020). In this study, we observed that dietary FMF supplementation increased the concentrations of total EAA, NEAA and AA, whereas 5% FMF significantly improved the total protein content. Specifically, our results showed that pigs fed FMF had increased concentrations of lysine, methionine, phenylalanine, threonine, alanine, aspartate, glutamate, arginine, serine and tyrosine. However, previous research has not found such significant benefits from FMF (Lu et al., 2020). We speculate that this might be due to differences in the experimental animals, fermentation substrates and strains. In addition, the content of FAA in muscles is closely related to meat flavor (Pereira and Vicente, 2013). Furthermore, in the current study, the contents of flavor amino acids (FAA), such as glutamate, aspartate, alanine and arginine, were increased with FMF supplement. The findings are directly in line with previous studies (Li et al., 2022; Tian et al., 2020). To date, the exact mechanism that regulates amino acid composition remains unclear. Notably, the contents of CP, total AA and FAA increased after microbial fermentation in this study, which may be the main reason for the change in the amino acid profiles (Table S1).

The fatty acid profiles in the IMF determine the nutritional value and oxidative stability of muscle and are particularly important for meat quality and meat product acceptability (Duan et al., 2016). A high intake of SFAs has been reported to increase the risk of type-2 diabetes (T2D) and heart disease (Lenighan et al., 2019), while PUFA, in particular EPA and DHA, have been found to possess numerous benefits, such as anti-inflammatory, glycolipid metabolism regulation and muscle development properties (Tachtsis et al., 2018; Tortosa-Caparrós et al., 2017; Vaidya and Cheema, 2014; Vissers et al., 2019). In the current study, only 10% FMF supplementation increased the content of PUFA and the PUFA:SFA ratio. These results tie in well with previous studies (Hao et al., 2020). Specifically, our results indicated that 10% FMF supplementation increased the concentrations of n-3 PUFA and n-6 PUFA, such as C18:2n6c, C18:3n3, and C20:3n3. Previous studies have shown that high n-6 and low n-3 PUFA intake (average ratio 15:1) contributes to the development of nonalcoholic fatty liver disease (NAFLD) (Toshimitsu et al., 2007). Our results suggested that the n-6 to n-3 PUFA ratio in the LDM was lower than 15:1 in all groups. Although FMF supplementation reduced the n-6 to n-3 ratio, no significant difference was found. However, there have been few published studies on how FMF affects the fatty acid profiles of pork, and it is difficult to explain the exact mechanism by which the fatty acid profiles of pigs fed FMF were altered. A plausible explanation may be that microbial fermentation optimized the fatty acid composition of the feed, as microbial fermentation increased the contents of MUFA and PUFA and decreased the content of SFAs in this study. As for why the effect of the 10% FMF group was more pronounced than that of the 5% FMF group, we speculated that the



main reason was that the addition of FMF was positively correlated with the nutritional levels of amino acids and fatty acids in the feed. As shown in Table S1 and Table S2, microbial fermentation increased the content of amino acids and fatty acids in the feed, resulting in a more significant effect on the composition of muscle amino acids and fatty acids in the 10% FMF group.

To date, few studies have been conducted on the role of FMF on lipid metabolism-related genes. As stated above, the IMF content increased with 5% and 10% FMF supplementation. However, it is unclear whether the expression of genes involved in the control of fat deposition was altered. *ACC $\alpha$*  and *FABP4* are two key genes involved in fatty acid synthesis and transport (Furuhashi and Hotamisligil, 2008; Munday, 2002). In the present study, 10% FMF supplementation upregulated the expression level of *FABP4*. These results are not entirely consistent with previous reports (Yu et al., 2020), as there was no difference in *ACC $\alpha$*  expression. The addition of FMF significantly increased the contents of unsaturated fatty acids and changed the fatty acid composition. We further examined the expression of unsaturated fatty acid synthesis-related genes, including *ACAA1*, *FADS2* and *SCD* (Xu et al., 2021). We observed that 10% FMF supplementation increased the expression levels of *ACAA1* and *FADS2*. *CEBP $\alpha$* , *PPAR $\gamma$*  and *SREBP* are key nuclear transcription factors related to lipid synthesis and metabolism that can regulate the expression of downstream target genes, including *ACC $\alpha$*  and *FABP4* (Oishi et al., 2017; Stoeckman and Towle, 2002; Wang et al., 2020). In the present study, the mRNA expression levels of *CEBP $\alpha$* , *PPAR $\gamma$*  and *SREBP1* were all markedly upregulated in the 10% FMF group. In summary, this study provides evidence that 10% FMF supplementation promoted fatty acid synthesis and transport, which was attributed to fatty acid profile alterations and IMF deposition.

## 5. Conclusion

In conclusion, the current study demonstrated that dietary 5% and 10% FMF supplementation improved the ADG in female finishing pigs. In addition, dietary FMF supplementation improved the meat quality and altered the amino acid and fatty acid compositions in the LDM. Furthermore, the upregulated expression of genes related to lipid metabolism might mediate these benefits. However, further research is needed to explore the mechanism by which FMF improves meat quality in finishing pigs. Taken together, this study suggests that FMF has great prospects for improving growth performance and meat quality in finishing pigs.

## Declaration of competing interest

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, and there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the content of this paper.

## Acknowledgments

We thank all the members of the Shan Laboratory for their support and constructive comments. This work was partially supported by the National Key R&D Program of China (2021YFC2103005) and the Zhejiang Provincial Key R&D Program of China (2021C02008).

## Appendix Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.aninu.2022.09.003>.

## References

- Buckley JD, Howe PR. Anti-obesity effects of long-chain omega-3 polyunsaturated fatty acids. *Obes Rev* 2009;10:648–59.
- Canibe N, Jensen BB. Fermented and nonfermented liquid feed to growing pigs: effect on aspects of gastrointestinal ecology and growth performance. *J Anim Sci* 2003;81:2019–31.
- Cao FL, Zhang XH, Yu WW, Zhao LG, Wang T. Effect of feeding fermented ginkgo biloba leaves on growth performance, meat quality, and lipid metabolism in broilers. *Poultry Sci* 2012;91:1210–21.
- Chun-Hua Chi, Cho Seong-Jun. Improvement of bioactivity of soybean meal by solid-state fermentation with *Bacillus amyloliquefaciens* versus *Lactobacillus* spp. and *Saccharomyces cerevisiae*. *LWT - Food Sci Technol (Lebensmittel-Wissenschaft -Technol)* 2016.
- Ding X, Li H, Wen Z, Hou Y, Wang G, Fan J, Qian L. Effects of fermented tea residue on fattening performance, meat quality, digestive performance, serum antioxidant capacity, and intestinal morphology in fatteners. *Animals (Basel)* 2020;10.
- Duan Y, Li F, Li Y, Guo Q, Ji Y, Tan B, Li T, Yin Y. Effects of supplementation with branched-chain amino acids to low-protein diets on expression of genes related to lipid metabolism in skeletal muscle of growing pigs. *Amino Acids* 2016;48:2131–44.
- Frank D, Joo ST, Warner R. Consumer acceptability of intramuscular fat. *Korean J Food Sci Anim Resour* 2016;36:699–708.
- Furuhashi M, Hotamisligil GS. Fatty acid-binding proteins: role in metabolic diseases and potential as drug targets. *Nat Rev Drug Discov* 2008;7:489–503.
- Guo S, Zhang Y, Xv J, Hou Y, Ding B. Partial substitution of fermented soybean meal for soybean meal influences the carcass traits and meat quality of broiler chickens. *Animals* 2020;10:225.
- Hao L, Su W, Zhang Y, Wang C, Lu Z. Effects of supplementing with fermented mixed feed on the performance and meat quality in finishing pigs. *Anim Feed Sci Technol* 2020;266:114501.
- Kiers JL, Meijer JC, Nout MJ, Rombouts FM, Nabuurs MJ, Van Der Meulen J. Effect of fermented soya beans on diarrhoea and feed efficiency in weaned piglets. *J Appl Microbiol* 2003;95:545–52.
- Lenighan YM, McNulty BA, Roche HM. Dietary fat composition: replacement of saturated fatty acids with pufa as a public health strategy, with an emphasis on  $\alpha$ -linolenic acid. *Proc Nutr Soc* 2019;78:234–45.
- Li H, Duan Y, Yin F, Zhu Q, Hu C, Wu L, Xie P, Li F, Cheng R, Kong X. Dietary addition of fermented sorghum distiller's dried grains with soluble improves carcass traits and meat quality in growing-finishing pigs. *Trop Anim Health Prod* 2022;54:97.
- Lomiwes D, Farouk MM, Wiklund E, Young OA. Small heat shock proteins and their role in meat tenderness: a review. *Meat Sci* 2014;96:26–40.
- Lu J, Han Q, Wang S, Wang Z, Shi X. Effect of fermented corn–soybean meal on carcass and meat quality of grower–finisher pigs. *J Anim Physiol Anim Nutr* 2020.
- Luo J, Zeng D, Cheng L, Mao X, Yu J, Yu B, Chen D. Dietary  $\beta$ -glucan supplementation improves growth performance, carcass traits and meat quality of finishing pigs. *Anim Nutr* 2019;5:380–5.
- Meng QW, Yan L, Ao X, Zhou TX, Kim IH. Influence of probiotics in different energy and nutrient density diets on growth performance, nutrient digestibility, meat quality, and blood characteristics in growing–finishing pigs. *J Anim Sci* 2010;88:3320.
- Missotten JA, Michiels J, Dierick N, Ovary A, Akbarian A, De Smet S. Effect of fermented moist feed on performance, gut bacteria and gut histo-morphology in broilers. *Br Poultry Sci* 2013;54:627–34.
- Moeller SJ, Miller RK, Edwards KK, Zerby HN, Logan KE, Aldredge TL, Stahl CA, Boggess M, Box-Steffensmeier JM. Consumer perceptions of pork eating quality as affected by pork quality attributes and end-point cooked temperature. *Meat Sci* 2010;84:14–22.
- Mukherjee R, Chakraborty R, Dutta A. Role of fermentation in improving nutritional quality of soybean meal - a review. *Asian-Australas J Anim Sci* 2016;29:1523–9.
- Munday MR. Regulation of mammalian acetyl-coa carboxylase. *Biochem Soc Trans* 2002;30:1059–64.
- Nong Q, Wang L, Zhou Y, Sun Y, Chen W, Xie J, Zhu X, Shan T. Low dietary n-6/n-3 pufa ratio regulates meat quality, reduces triglyceride content, and improves fatty acid composition of meat in heigai pigs. *Animals (Basel)* 2020;10.
- Oishi Y, Spann NJ, Link VM, Muse ED, Strid T, Edillor C, Kolar MJ, Matsuzaka T, Hayakawa S, Tao J, Kaikkonen MU, Carlin AF, Lam MT, Manabe I, Shimano H, Saghatelian A, Glass CK. Srebp1 contributes to resolution of pro-inflammatory tlr4 signaling by reprogramming fatty acid metabolism. *Cell Metabol* 2017;25:412–27.
- Ovissipour M, Abedian A, Motamedzadegan A, Rasco B, Safari R, Shahiri H. The effect of enzymatic hydrolysis time and temperature on the properties of protein hydrolysates from Persian sturgeon (*Acipenser persicus*) viscera. *Food Chem* 2009;115:238–42.
- Pereira PM, Vicente AF. Meat nutritional composition and nutritive role in the human diet. *Meat Sci* 2013;93:586–92.
- Shan T, Liang X, Bi P, Kuang S. Myostatin knockout drives browning of white adipose tissue through activating the *ampk-pgc1 $\alpha$ -fndc5* pathway in muscle. *Faseb J* 2013;27:1981–9.
- Shi C, He J, Yu J, Yu B, Huang Z, Mao X, Zheng P, Chen D. Solid state fermentation of rapeseed cake with *Aspergillus Niger* for degrading glucosinolates and upgrading nutritional value. *J Anim Sci Biotechnol* 2015;6:13.



- Shi C, Zhang Y, Lu Z, Wang Y. Solid-state fermentation of corn-soybean meal mixed feed with bacillus subtilis and enterococcus faecium for degrading antinutritional factors and enhancing nutritional value. *J Anim Sci Biotechnol* 2017;8:50.
- Stoekman AK, Towle HC. The role of srebp-1c in nutritional regulation of lipogenic enzyme gene expression. *J Biol Chem* 2002;277:27029–35.
- Tachtsis B, Camera D, Lacham-Kaplan O. Potential roles of n-3 PUFA during skeletal muscle growth and regeneration. *Nutrients* 2018;10.
- Tang X, Liu X, Zhang K. Effects of microbial fermented feed on serum biochemical profile, carcass traits, meat amino acid and fatty acid profile, and gut microbiome composition of finishing pigs. *Front Vet Sci* 2021;8:744630.
- Tian Z, Deng D, Cui Y, Chen W, Yu M, Ma X. Diet supplemented with fermented okara improved growth performance, meat quality, and amino acid profiles in growing pigs. *Food Sci Nutr* 2020;8:5650–9.
- Tortosa-Caparrós E, Navas-Carrillo D, Marín F, Orenes-Piñero E. Anti-inflammatory effects of omega 3 and omega 6 polyunsaturated fatty acids in cardiovascular disease and metabolic syndrome. *Crit Rev Food Sci Nutr* 2017;57:3421–9.
- Toshimitsu K, Matsuura B, Ohkubo I, Niiya T, Furukawa S, Hiasa Y, Kawamura M, Ebihara K, Onji M. Dietary habits and nutrient intake in non-alcoholic steatohepatitis. *Nutrition* 2007;23:46–52.
- Vaidya H, Cheema SK. Sea cucumber and blue mussel: new sources of phospholipid enriched omega-3 fatty acids with a potential role in 3t3-L1 adipocyte metabolism. *Food Funct* 2014;5:3287–95.
- Van Elswyk ME, McNeill SH. Impact of grass/forage feeding versus grain finishing on beef nutrients and sensory quality: the U.S. Experience. *Meat Sci* 2014;96:535–40.
- Vissers LET, Rijkse J, Boer JMA, Verschuren WMM, Van Der Schouw YT, Sluijs J. Fatty acids from dairy and meat and their association with risk of coronary heart disease. *Eur J Nutr* 2019;58:2639–47.
- Wang C, Lin C, Su W, Zhang Y, Wang F, Wang Y, Shi C, Lu Z. Effects of supplementing sow diets with fermented corn and soybean meal mixed feed during lactation on the performance of sows and progeny. *J Anim Sci* 2018;96:206–14.
- Wang G, Wu B, Zhang L, Cui Y, Zhang B, Wang H. Laquinimod prevents adipogenesis and obesity by down-regulating ppar- $\gamma$  and c/ebp $\alpha$  through activating ampk. *ACS Omega* 2020;5:22958–65.
- Wen W, Chen X, Huang Z, Chen D, Yu B, He J, Luo Y, Yan H, Chen H, Zheng P, Yu J. Dietary lycopene supplementation improves meat quality, antioxidant capacity and skeletal muscle fiber type transformation in finishing pigs. *Anim Nutr* 2022;8:256–64.
- Xu X, Chen X, Chen D, Yu B, Yin J, Huang Z. Effects of dietary apple polyphenol supplementation on carcass traits, meat quality, muscle amino acid and fatty acid composition in finishing pigs. *Food Funct* 2019;10:7426–34.
- Xu X, Li LM, Li B, Guo WJ, Ding XL, Xu FZ. Effect of fermented biogas residue on growth performance, serum biochemical parameters, and meat quality in pigs. *Asian-Australas J Anim Sci* 2017;30:1464–70.
- Xu Z, Chen W, Wang L, Zhou Y, Nong Q, Valencak TG, Wang Y, Xie J, Shan T. Cold exposure affects lipid metabolism, fatty acids composition and transcription in pig skeletal muscle. *Front Physiol* 2021;12:748801.
- Yan L, Meng QW, Kim IH. Effects of fermented garlic powder supplementation on growth performance, nutrient digestibility, blood characteristics and meat quality in growing-finishing pigs. *Anim Sci J* 2012;83:411–7.
- Yu M, Li Z, Rong T, Wang G, Liu Z, Chen W, Li J, Li J, Ma X. Different dietary starch sources alter the carcass traits, meat quality, and the profile of muscle amino acid and fatty acid in finishing pigs. *J Anim Sci Biotechnol* 2020;11:78.
- Zhong Y, Yan Z, Song B, Zheng C, Duan Y, Kong X, Deng J, Li F. Dietary supplementation with betaine or glycine improves the carcass trait, meat quality and lipid metabolism of finishing mini-pigs. *Anim Nutr* 2021;7:376–83.