

European Association of Neuro-Oncology (EANO) guidelines for treatment of primary central nervous system lymphoma (PCNSL)

Khê Hoang-Xuan, Martina Deckert, Andrés J.M. Ferreri, Julia Furtner, Jaime Gallego Perez-Larraya, Roger Henriksson, Andreas F. Hottinger, Benjamin Kasenda, Florence Lefranc, Alexander Lossos, Catherine McBain, Matthias Preusser^o, Patrick Roth, Roberta Rudà, Uwe Schlegel, Riccardo Soffietti^o, Carole Soussain, Martin J.B. Taphoorn, Valérie Tuitou, Michael Weller^o and Jacoline E.C. Bromberg

APHP, Department of Neurology, Groupe Hospitalier Pitié-Salpêtrière; Sorbonne Université; IHU; ICM. Paris, France (K.H.X.); Institute of Neuropathology, Faculty of Medicine and University Hospital Cologne, University of Cologne, Cologne, Germany (M.D.); Lymphoma Unit, IRCCS San Raffaele Scientific Institute, Milano, Italy (A.J.M.F.); Department of Biomedical and Imaging Image-guided Therapy Medical University of Vienna, Vienna, Austria (J.F.); Health Research Institute of Navarra (IdiSNA), Program in Solid Tumors, Foundation for the Applied Medical Research, Department of Neurology, Clínica Universidad de Navarra, Pamplona, Navarra, Spain (J.G.P.L.); Department of Radiation Sciences, Oncology, University of Umeå, S-901 85 Umeå, Sweden (R.H.); Departments of Oncology and Clinical Neurosciences, CHUV University Hospital Lausanne and University of Lausanne, Lausanne Switzerland (A.F.H.); Department of Hematology/Oncology and Palliative Care, Klinikum Stuttgart, Stuttgart, Germany (B.K.); Department of Neurosurgery, Hôpital Erasme, Université Libre de Bruxelles, Belgium (F.L.); Head, Leslie and Michael Gaffin Center for Neuro-Oncology; Departments of Oncology and Neurology; Hadassah-Hebrew University Medical Center; Jerusalem, Israel (A.L.); Department of Clinical Oncology, The Christie NHS FT; Manchester; United Kingdom (C.M.B.); Division of Oncology, Department of Medicine I, Medical University of Vienna, Vienna, Austria (M.P.); Department of Neurology & Brain Tumor Center, University Hospital and University of Zurich, Zurich, Switzerland (P.R., M.W.); Department of Neurology, Castelfranco Veneto/Treviso Hospital, Italy (R.R.); Universitätsklinikum Knappschaftskrankenhaus Bochum GmbH, Germany (U.S.); Division of Neuro-Oncology, Department of Neuroscience, University of Turin, and City of Health and Science University Hospital, Turin, Italy (R.R., R.S.); Department of Hematology, Institut Curie, Site Saint-Cloud, France and INSERM U932 Institut Curie, PSL Research University, Paris, France (C.S.); Department of Neurology, Leiden University Medical Center and Department of Neurology, Haaglanden Medical Center The Hague, The Netherlands (M.J.B.T.); APHP, Department of Ophthalmology, Groupe Hospitalier Pitié-Salpêtrière; Sorbonne Université. Paris, France (V.T.); Department of Neuro-Oncology, Erasmus MC University Medical Center Cancer Institute, Rotterdam. The Netherlands (J.E.C.B.)

Corresponding Author: Khê Hoang-Xuan, Department of Neurology Mazarin, Division Mazarin, Groupe Hospitalier Pitié-Salpêtrière, 47 boulevard de l'hôpital, Paris 75013, France (khe.hoang-xuan@aphp.fr).

Abstract

The management of primary central nervous system (PCNSL) is one of the most controversial topics in neuro-oncology because of the complexity of the disease and the limited number of controlled studies available. In 2021, given recent advances and the publication of practice-changing randomized trials, the European Association of Neuro-Oncology (EANO) created a multidisciplinary task force to update the previously published evidence-based guidelines for immunocompetent adult patients with PCNSL and added a section on immunosuppressed patients. The guideline provides consensus considerations and recommendations for the treatment of PCNSL, including intraocular manifestations and specific management of the elderly. The main changes from the previous guideline include strengthened evidence for the consolidation with ASCT in first-line treatment, prospectively assessed chemotherapy combinations for both young and elderly patients, clarification of the role of rituximab even though the

data remain inconclusive, of the role of new agents, and the incorporation of immunosuppressed patients and primary ocular lymphoma. The guideline should aid the clinicians in everyday practice and decision making and serve as a basis for future research in the field.

Keywords

chemotherapy | immunotherapy | primary CNS lymphoma | radiotherapy | treatment

Primary diffuse large B cell lymphoma (DLBCL) of the central nervous system (PCNSL) is an aggressive neoplasm confined to the brain, eyes, cranial nerves, leptomeninges, or spinal cord in the absence of disease outside the CNS. Currently, PCNSL is estimated to account for up to 1% of lymphomas, 4–6% of all extranodal lymphomas, and about 3% of all CNS tumors.¹ Immunodeficiency is the only well-known risk factor for the development of the disease, and the incidence of PCNSL rose dramatically during the peak of the acquired immune deficiency syndrome (AIDS) epidemic in the late 1980s.² Highly active antiretroviral therapy (HAART), which led to an immune recovery in many HIV patients, was introduced in 1996–1997, and since then many studies have indicated a dramatic decrease in the incidence of PCNSL in AIDS patients^{2,3}

In contrast, the incidence continues intriguingly to rise in the elderly who consequently represent the majority of patients in the immunocompetent population in some recent studies.^{4–6} In this guideline, PCNSL in immunocompetent and immunodeficient patients will be discussed separately. Although the prognosis of PCNSL remains poor, it has significantly improved over the past decades as a result of better treatment strategies with a curative aim. Treatment of PCNSL remains challenging. Despite high chemo- and radiosensitivity, remissions are frequently short-lasting. The blood brain barrier (BBB) limits the access of many drugs to the CNS and patients, especially the elderly, are at high risk of developing severe treatment related-neurotoxicity. The majority of current evidence supporting therapeutic choices still results from retrospective series or single-arm phase II studies, but the insights gained from the publication of several recent randomized trials including one phase III,⁷ four phase II studies^{8–13} and abstracts of 3 further randomized studies,^{14–16} prompted us to update our previous recommendations.¹⁷ The objective of this guideline is to provide clinicians with updated evidence-based recommendations and consensus expert opinions on the management of patients with PCNSL.

The task force set up in 2013 under the auspices of the EANO (European Association for Neuro-Oncology) and selected to be representative of European-based medical experts, wrote the first guideline published in 2015¹⁷ and was partially renewed to update the guidelines and enlarge the scope to PCNSL in immunodeficient patients. The panel with specialists from 11 countries covered all fields of expertise in the management of PCNSL, ie, neurologists, hematologists, medical oncologists, neurosurgeons, pathologists, neuroradiologists, ophthalmologists, and radiation oncologists. Based on the best available evidence from the literature review, all experts were assigned to update and/or rewrite the different sections of the guidelines

and grade the evidence. The revised guideline, taking into account the comments of the panelists, was resubmitted by the two chairmen (KHX, JB) to the whole task force for review and amendments three times. The final agreement was obtained in June 2022. References for this review were identified through searches of PubMed from January 1980 to May 2022 and through searches of the authors' own files. The final reference list was generated on the basis of originality and relevance to the broad scope of this review. Relevant abstracts presented at international meetings were mentioned by task force members during manuscript preparation but were not taken into account for evidence grading. As for the previous EANO guidelines, the scientific evidence of papers collected from the literature was evaluated and graded and recommendations are given according to [Table 1](#).^{17,18} The main changes from the previous guideline are the incorporation of immunosuppressed patients, a more solid basis for the consolidation with ASCT in first-line treatment and more data regarding rituximab, even though the data are still not conclusive, and development of new agents though none have become standard yet.

General Recommendations

For recommendations regarding the general approach to patients with PCNSL, including pathology and genetics, clinical presentation, diagnostic confirmation, staging, prognostic factors, and response criteria to treatment, we refer to the table in the [supplemental appendix](#) that has been built and updated on our previous guideline.¹⁷ Similarly, updated key recommendations for treatment are summarized in [Table 2](#).

Surgery

Resection has historically not been considered to be the standard of care for PCNSL because of i) the microscopically multifocal and infiltrative nature of PCNSL, which may extend beyond the visible border of the lesion, ii) the fact that lesions are often located deeply in the periventricular space, and iii) historical series that suggested no clear benefit in the outcome of resection when used as the only treatment compared either to supportive care (Class IIIb)⁶⁵ or with biopsy performed for patients having received post-operative chemo-and/or radiotherapy (Class IIIb).⁶⁶ In addition, early retrospective studies on combination treatment found a high complication rate of surgery without a

Table 1 Grading of evidence and recommendations^{17,18}

Grading of evidence	
Class I	Prospective, randomized phase III studies
Class IIa	Prospective, randomized phase II studies
Class IIb	Phase I and II studies
Class IIIa	Prospective studies, including observational studies, cohort studies and case-control studies
Class IIIb	Retrospective studies
Class IV	Uncontrolled case series, case reports and expert opinion
Recommendations	
Level A	At least one Class I study or two consistent Class IIa studies
Level B	At least one Class IIa study or overwhelming Class IIb and III evidence
Level C	At least two consistent Class III studies
Good practice point	If insufficient evidence for Level A–C and consensus among task force

Table 2 Treatment recommendations (adapted and updated from the previous *EANO guidelines*. Hoang-Xuan et al, *Lancet Oncol* 16:e322-e332, 2015¹⁷)

Surgery	Reference
Surgical resection may be considered in patients suffering from a large space occupying lesion with acute symptoms of brain herniation to reduce rapidly intracranial pressure (Good practice point).	
Limited and only retrospective data exist regarding surgical resection or biopsy in a unifocal and resectable lesion suspected of PCNSL. No consensus was met in the panel for a recommendation.	19–21
Induction chemotherapy, immunochemotherapy	
HD-MTX is the drug of choice in PCNSL and chemotherapy should include MTX at HD (≥ 3 g/m ²) both to cross the BBB and yield cytotoxic levels in the CSF. It should be delivered in 2–3 h iv infusions for a minimum of 4–6 injections and at intervals that should not exceed 2–3 weeks (Level B).	22
Combination of HD-MTX with other chemotherapeutic agents improves the response and progression-free survival rates with respect to HD-MTX alone (Level B).	22
Chemotherapeutic agents to combine with HD MTX should be selected among active drugs known to cross the blood-brain-barrier, such as HD cytarabine and combinations used in large and/ or randomized prospective trials have to be preferred (Level B).	7,9,11,14,23
HD-MTX-based chemotherapy is feasible in elderly patients with adequate performance status and renal function (Level B).	8,24–26
Most combinations addressed in large clinical trials include HD-MTX associated with an alkylating agent (procarbazine, carmustine, temozolomide, and thiopeta) (Level B).	7,9,11,14,23,27
The value of IT chemotherapy is unclear. IT chemotherapy (intralumbar or preferably intraventricular through an Ommaya reservoir) can be proposed in case of documented meningeal involvement with insufficient response to iv HD MTX (>3 g/m ²) based chemotherapy (Good Practice Point).	
Conflicting data exist regarding iv Rituximab combined with a chemotherapy regimen in PCNSL. No consensus was met in the panel for recommendation (Level B).	7,9,12,28
Consolidation treatment	
Radiotherapy	
WBRT and the combination of HD-MTX with WBRT expose patients to an increased risk of neurotoxicity (Level A).	29–32
The role of consolidation WBRT following HD-MTX based chemotherapy remains debated and, if considered, the optimal dose is not yet defined. Risk of neurotoxicity and alternative consolidation options (eg, ASCT) or omitting consolidation should be weighed in the individual patient (Level B).	10–13,33
Reduced dose WBRT consolidation (23.4–30 Gy in 1.8–2.0 Gy fractions) in CR patients is a therapeutic option that is currently being investigated in randomized trials. (Good Practice Point).	14
In patients with progressive or significant residual disease after primary chemotherapy, a total dose of 36–40 Gy with a 1.8–2 Gy dose/fraction appears advisable. With such doses, there is no evidence to add a focal boost on the enhancing lesions (Good Practice Point).	
In patients >60 years, the risk of delayed neurotoxicity, after WBRT especially if following HDMTX, is unacceptably high and WBRT should be avoided in these elderly patients (Level B).	29–32

Table 2 Continued

Consolidation treatment	
High dose chemotherapy with autologous stem cell transplantation (HDC/ASCT)	
HDC/ASCT as consolidation is an effective treatment for younger (age up to 65–70) patients with newly-diagnosed PCNSL, though risk of acute toxicity should be taken into account (Level B).	10–13
HDC/ASCT is an effective consolidation treatment with efficacy at least comparable to that of WBRT	10–13
High-dose thiotepa-based conditioning chemotherapy should be preferred over the BEAM regimen (Level B).	34,35
Non myeloablative chemotherapy	
The value of nonmyeloablative consolidation chemotherapy and maintenance chemotherapy is currently under investigation in clinical trials (Good Practice Point)	36,37 NCT02531841, NCT01511562 NCT03495960 NCT02313389
Primary vitreoretinal lymphoma (PVRL)	
HD-MTX-based chemotherapy seems to improve OS in PVRL (level C), though local relapses occur frequently. Whether the addition of local treatment reduces local relapses is uncertain.	38
Local treatment (intravitreal immuno/chemotherapy or ocular RT) is a valid approach for patients with systemic chemotherapy contraindications or for elderly patients with relapsing intraocular disease (Good Practice Point).	39,40
Patients with concurrent intraocular and CNS lymphoma should be treated no differently from other patients with PCNSL (Good Practice Point).	
If consolidation WBRT is proposed, it should include both eyes (Good Practice Point).	
Refractory and relapsed PVRL should be treated according to the patients' characteristics and prior treatments. Treatments include intravitreal injections of MTX, focal radiotherapy, WBRT, systemic chemotherapy, targeted treatment and HDC/ASCT (Good Practice Point).	
Salvage treatment	
Patients with relapsed/ refractory PCNSL should be enrolled into clinical trials (Good Practice Point).	
HDC/ASCT is a valid therapeutic option in patients aged <70 years with chemosensitive relapsing PCNSL especially in patients without prior ASCT (Level B).	41–43
Salvage WBRT may be proposed in radiotherapy-naïve patients; it may be preceded by induction chemotherapy (Good Practice Point).	44,45
Salvage chemotherapy can be delivered as induction therapy before WBRT or HDC/ASCT, or as exclusive treatment in patients not eligible for these therapies.	46,47
MTX re-challenge should be considered in recurrent PCNSL patients who previously responded to HD MTX (Level C).	48–50
Isolated extra-CNS relapses should be managed with anthracycline-based chemotherapy followed or not by HDC/ASCT (Good Practice Point).	
Bruton Tyrosine kinase inhibitors, imids, immune checkpoint inhibitors and CART have shown clinical activity as single agents in relapsing PCNSL and may be considered in salvage treatments.	51–57
HIV related patients	
Initiation, if not yet done, or modification of ART should be done in conjunction with the infectious disease specialist (Good Practice Point).	58,59
Patients with adequate performance status (arbitrarily defined as KPS \geq 60) and able to tolerate it (adequate renal function, absence of pleural or abdominal effusion) should be offered treatment with HD-MTX based chemotherapy. Polychemotherapy should be preferred to MTX monotherapy (Good Practice Point).	60–62
In HIV-related PCNSL patients the risk of delayed neurotoxicity from WBRT is significant and radiation should therefore be avoided (Good Practice Point).	
If HD-MTX based regimens cannot be considered, combination of ART with other chemotherapeutic agents or with palliative WBRT may be an alternative (Good Practice Point).	
Combinations of chemotherapy with antiviral treatments against EBV, rituximab, immune check point inhibitors or targeted therapies need further evaluation in this population (Good Practice Point).	
PCNS-post-transplant lymphoproliferative disorder (PTLD) patients	
Immunosuppression should be reduced to the lowest level possible, in close collaboration with transplant specialists (Good Practice Point). Further antitumor treatment needs to be tailored to patient's age and performance status, and transplanted organ functioning (Good Practice Point).	63,64
Extrapolated from its use in systemic PTLD, systemic chemotherapy including HD-MTX based regimens should be considered in order to increase response rates and reduce the high risk of relapse (Good Practice Point).	
On the basis of its efficacy in systemic PTLD, treatment with rituximab might also be considered when possible, especially in patients with underlying renal failure that precludes usage of HD-MTX (Good Practice Point)	
There is lack of evidence supporting treatment with antiviral agents, and other therapeutic strategies need further evaluation in this population (Good Practice Point).	

positive effect on survival (Class IIIb).¹⁹ In contrast, recent studies suggest that resection may provide a therapeutic benefit in selected patients.^{20,21,67}

In a post hoc analysis of the German PCNSL Study Group-1 trial, including 526 patients with PCNSL, 67 of whom underwent gross total resection, PFS and OS were significantly shorter in the biopsied group as compared with patients with resections even when controlled for age and KPS.²⁰ When controlled for the number of lesions, the difference remained statistically significant only for PFS (Class IIIa).²⁰ Smaller single institution retrospective analyses revealed that surgical resection for PCNSL patients is safe for patients with good performance status and a single, superficial lesion with complication rates comparable to rates for other intracranial tumors, though the clinical benefit to resection could not be concluded (Class IVb).^{68,69} Data from 132 PCNSL patients cross-validated using data from 8,936 patients from two national American databases suggested that craniotomy is associated with increased survival over biopsy by 8.5 months for patients categorized in a low surgical risk, which includes lesion location and number, age and frailty (Class IIIb).²¹ There have not yet been any published series prospectively assessing morbidity or survival in PCNSL patients treated with cytoreductive surgery vs biopsy; selection bias to include patients with better prognosis, that is patients with single, superficial, and small lesions in resected subgroups, cannot be excluded in these retrospective studies. No consensus was met to recommend either resection or biopsy for patients with a unifocal and resectable lesion suspected of PCNSL. In such cases, decision-making should be discussed in a multidisciplinary tumor-board.

Systemic Chemotherapy

Based on convergent results from numerous prospective and retrospective studies, high-dose intravenous (iv) methotrexate (HD-MTX), an antifolate and antimetabolite, is the most important and beneficial single agent.²² Penetration of MTX into the CNS depends both on the total dose and rate of infusion. The optimal dose of MTX has not been determined. It has been estimated that the iv MTX dose should range between 1 g/m² and 8 g/m² to achieve sufficient drug levels within the CNS. In the absence of clear evidence for a dose-response relationship,²² and since the rapid infusion of MTX ≥ 3 g/m² over 3 h achieves cytotoxic levels in the CSF, there is a growing consensus to deliver MTX according to this protocol (Class IV).⁷⁰ Since the efficacy of MTX may also depend on the duration of exposure, the MTX administration intervals in most treatment protocols range between 10 days and 3 weeks.²² The optimal number of MTX injections to deliver has not been formally established. A minimum of 4–6 infusions is delivered in most chemotherapy regimens, especially if no consolidation treatment (radiotherapy and/or intensive chemotherapy) is scheduled in the protocol. Infusions of HD-MTX require hyperhydration, urine alkalinization, leucovorin rescue, and MTX concentration monitoring. HD-MTX has been used as monotherapy in single-arm studies though with a varying responses.^{71,72} Currently, most treatment protocols combine HD-MTX with a variety of other chemotherapeutic

agents to improve response rate and outcome. In the only randomized, though phase II, a study comparing HD-MTX (3 g/m²) alone with a HD-MTX (3 g/m²) combination regimen a significantly higher complete response (CR) rate was found in the HD-MTX + cytarabine arm (46%), compared with the HD-MTX-only control arm (18%, $P = .006$)²³ (Class IIa). In the subsequent IELSG32 phase II trial, three different HD-MTX-based induction regimens were compared for patients up to the age of 65 (ECOG ≤ 3) or 70 (ECOG ≤ 2).⁹ On completion of induction chemotherapy, eligible patients underwent a second randomization and received either high-dose chemotherapy and autologous stem cell transplantation (HD/ASCT) or WBRT as consolidation, with CR rate as a primary endpoint for induction. Patients receiving the most intensive induction regimen, consisting of HD-MTX, Ara-C, rituximab, and thiotepa (the MATRix regimen), showed a significantly higher CR rate (49%) in comparison to 23% in the least intensive regimen, consisting of HD-MTX and Ara-C, ($P = .0007$) with a 2-year OS of 69% and 42% respectively.⁹ (Class IIa).

Another approach is BBB disruption (BBBD) by intra-arterial (IA) infusion of hypertonic mannitol followed by intra-arterial (IA) chemotherapy to increase the drug concentration in the CNS. BBBD with IA MTX administered in newly diagnosed PCNSL has demonstrated a good safety profile and neurocognitive tolerance and achieved comparable outcomes to those observed with iv HD-MTX-based chemotherapy regimens (Class IIIb).^{27,73,74}

In a recent meta-analysis including only prospective phase II and III trials, treatment results (CR rate, PFS, OS) were compared for HD-MTX alone with HD-MTX-based polychemotherapy including 1, 2, 3, 4, or ≥ 5 additional drugs (Class IIIa).²² Results of this analysis suggest improved CR rates with the addition of two or three drugs in addition to the backbone of HD-MTX, but an impact on OS remains to be demonstrated in adequately powered trials. A randomized comparison of the different combinations has not been published except in elderly patients and except in the addition of agents to the same backbone. In patients who are not eligible for HD-MTX, treatment should be chosen from therapeutic regimens that have shown efficacy in PCNSL.

In summary, HD-MTX is the drug of choice for PCNSL and should be the backbone of polychemotherapy regimens that include additional substances which have been investigated in large prospective phase II and III trials (level B recommendation). Despite important advances in the induction chemotherapy of PCNSL in the past years, the complete response rate remains lower compared to that obtained in systemic DLBCL, and improving induction treatment in PCNSL to fill this gap remains an important goal.

Immunotherapy

The anti-CD20 antibody rituximab is a standard component in the treatment of B-cell non-Hodgkin lymphomas including DLBCL occurring outside the CNS. However, the high molecular weight of rituximab limits its penetration through the BBB.⁷⁵ On the other hand, BBB breakdown associated with PCNSL has been assumed to potentially

increase the rituximab concentration in the CNS to levels required for sufficient anti-neoplastic activity.⁷⁶ The effect of rituximab when used as monotherapy in PCNSL was evaluated in a single study in which 12 patients with refractory or relapsed PCNSL were treated with a weekly iv dose of 375 mg/m² rituximab infusion for up to eight doses (Class IV).⁷⁷ MRI responses were observed in 36% of patients.

The role of rituximab in combination with HD-MTX-based chemotherapy as initial treatment for PCNSL has been evaluated in small uncontrolled studies^{36,46,51,78–83} and two prospective randomized clinical trials.⁷⁹ In the randomized phase II IELSG-32 study (see also above) patients with newly diagnosed PCNSL randomized for MTX–cytarabine alone had an ORR of 53% (95% CI 42–64); patients randomized for MTX–cytarabine and rituximab had an ORR of 74% (95% CI 64–84) ($P = .01$), with OS at 2 years of 42% vs 56% respectively.⁹ A 2022 update showed OS at 7-years of 21% (95% CI = 4–47%) vs 37% (95% CI = 26–48%).¹² The phase II design of this study should be taken into account when considering these data supporting the use of rituximab, as well as the superior results found in patients treated with the MTX–cytarabine combination without rituximab in a previous study: an ORR of 69% and 2 year OS of just under 60%.²³ (Class IIa). The HOVON-105/ALLG-NHL-24 study was a randomized phase III study in which patients with newly diagnosed PCNSL aged 18–70 years were randomized for treatment with MBVP chemotherapy (HD-MTX, carmustine, teniposide, and prednisone) with or without rituximab.⁷ Responding patients were subsequently treated with HD-cytarabine, and patients under 61 years old were additionally consolidated with low-dose (30 Gy) WBRT with an integrated boost to the tumor area if they did not achieve CR. The primary endpoint, event-free survival (EFS) at 1 year, was 49% without and 52% with rituximab, with a hazard ratio (HR) of 1.0 (95% CI 0.7–1.43), $P = .99$, thus showing no effect of rituximab on EFS. Similarly, 1-year PFS did not differ between the arms with 58% in the MBVP group and 65% in the R-MBVP group (HR 0.77, 95% CI 0.52–1.13, $P = .18$)⁷ (Class I). An unplanned subgroup analysis suggested a possible effect of rituximab in patients aged up to 60 years, but given the relatively small number of patients in this analysis and the nature of such an unplanned analysis, no definitive conclusions can be drawn. A systematic review and meta-analysis investigated the influence of rituximab on outcomes in randomized prospective trials (Class IIIa).²⁸ The hazard ratio (HR) for death in the pooled analysis was 0.76 (95% CI, 0.52–1.12) and the HR for PFS was 0.65 (95% CI, 0.45–0.95), thus showing no statistically significant evidence for an OS benefit and low certainty for a PFS benefit of rituximab, with no evidence of increased toxicity.

In conclusion, conflicting data exist regarding the efficacy of rituximab in PCNSL and no consensus was met in the panel for a recommendation.

Intrathecal Chemotherapy

The role of intrathecal (IT) chemotherapy remains unclear because prospective trials focused on this important issue do not exist. This strategy is not used in most recent and ongoing prospective trials. Indirect

evidence of the potential role of IT chemotherapy comes from long-term results of a single-arm phase II trial using the Bonn protocol which included MTX, prednisolone, and cytarabine applied via an Ommaya reservoir over six cycles in addition to HD-MTX-based polychemotherapy^{74,84} and a subsequent single-arm trial using the same systemic polychemotherapy regimen but without intraventricular therapy.⁸⁵ The first version of the protocol has been associated with a 10 year OS of 53% in patients younger than 60 years, whereas the version without intraventricular chemotherapy led to inferior results. Although this seems to support the use of intraventricular chemotherapy (Class IIIa), three retrospective studies did not show benefit from the addition of intrathecal drugs (MTX, cytarabine) in patients treated with HD-MTX dosed at 3 g/m² that theoretically reaches cytotoxic thresholds in the CSF (Class IIIb).^{86–88} Therefore, given the low level of evidence, we currently do not advocate IT chemotherapy in PCNSL patients without CSF dissemination. In the case of lymphomatous meningitis, pragmatically, IT chemotherapy may be proposed depending on the initial early assessment of the leptomeningeal response to systemic chemotherapy that should be evaluated at the latest before each scheduled intravenous HD-MTX injection.

Radiotherapy

Although recognized as an active treatment modality, discussion of the role of radiotherapy in PCNSL has become inextricably linked to concerns about neurotoxicity; in patients >60 years old this risk is considered unacceptably high and WBRT should be avoided if possible; see below.^{29–32} As outcomes from systemic therapies improve and survival rates rise, avoidance of late toxicity becomes ever more critical also in younger patients. Despite high initial response rates, radiotherapy (RT) used alone provides limited survival benefit, with a median OS of 10–18 months and a 5-y survival rate of 5%⁸⁹ (Class IIb). Although never formally compared in a randomized trial, the introduction of HD-MTX followed by RT is considered to be superior to RT alone, with reported 2–4-fold increases in OS (median: 30–72 months) and more long-term survivors (5-year survival of 20–50%) for many protocols^{19,90–93} (Class IIb, IIIa, and IIIb). Whether RT following induction chemotherapy could be safely omitted was investigated in the non-inferiority phase III G-PCNSL-SG 1 trial,³³ in which patients who achieved a CR received either consolidation WBRT (45 Gy in 30 × 1.5 Gy fractions) or observation. OS was similar in both arms (32.4 months in the WBRT arm, 37.4 months in the non-WBRT arm) with a non-significant trend to improved median PFS with WBRT (18.3 months vs 11.9 months) (Class I). Full interpretation has been hampered by methodological limitations, and by the fact that only 318 of 551 enrolled patients were treated per protocol. A few retrospective studies similarly suggested that omission of WBRT from first-line treatment results in shorter PFS but does not compromise OS (Class IIIb).^{94–96} Nevertheless, as a result of the above-mentioned limitations and the suggested effect on PFS, the role of consolidation WBRT remains

debated. Alternative consolidation options such as reduced dose WBRT (described below) or autologous stem cell transplant (ASCT) have been explored (see ASCT section).

Changes in radiation parameters have been proposed to reduce the risk of radiotherapy-related neuro-toxicity while maintaining efficacy. Aspects to be considered include total dose, fraction size, irradiated volume, hippocampal avoidance, and the use of neuroprotective agents. The most commonly employed WBRT total dose has been 36–40 Gy shielding the orbits after 30 Gy (after 36 Gy in the case of intraocular disease). Higher doses have not demonstrated any benefit⁹⁷ (Class IIIb) and, following a phase II study demonstrating encouraging results,⁷⁸ interest now centers around the role of reduced dose WBRT, which is being evaluated in the RTOG 1114 randomized phase II study.¹⁴ Initial data suggest that the addition of LD-WBRT (23.4 Gy in 13 × 1.8 Gy fractions) to R-MPV-A improves PFS in newly diagnosed PCNSL.¹⁴ Neurotoxicity rates at the time of reporting were not statistically significantly increased, but further neuropsychological testing and neuro-imaging analyses are ongoing; full results need to be awaited for final interpretation, and radiation-induced neurotoxicity in PCNSL may become overt years after administration.^{29–32}

The standardly employed fraction size remains 1.8 Gy/day. No trials have specifically addressed this question, but series using smaller fraction sizes and/or twice daily hyperfractionation have not demonstrated different outcomes.^{98,99} (Class IIb, Class III). The presence of multifocal disease, CSF and ocular involvement, and diffuse involvement of the brain in autopsy studies¹⁰⁰ imply that the traditional WBRT volume must still be advocated, with fields extending to the inferior border of C2, and covering the meninges, including the posterior 2/3 of the orbit^{101,102} (Class IIIb, Class IV). A single retrospective study addressing partial brain RT demonstrated significantly more out-of-field recurrences using margins of <4cm compared to margins of ≥ 4cm (83% vs 22%) (Class IIIb).¹⁰³

Delayed radiation neurotoxicity, concerning primarily impaired psychomotor speed, executive function, attention, and memory,¹⁰⁴ has been found to have an incidence of 25–35%, related mortality of 30%, and typically occurs months to years after successful treatment (Class IIIb).^{105–107} The risk is substantially higher in patients ≥60 years (Class IIIb).³² After reduced-dose WBRT cognitive functions remained stable at least for the first 2–3 years.^{108,109} Thereafter some cognitive deterioration was found in a small series although this was not significantly more than in patients consolidated with high-dose chemotherapy and stem cell transplantation.¹⁰⁹

In summary, the role of WBRT in PCNSL continues to be defined, except in older patients in whom the risk of delayed neurotoxicity is unacceptably high. Increasing response rates to induction chemotherapy may facilitate a move to the routine use of reduced dose WBRT or alternative consolidation strategies. In all settings, each patient's individualized situation and the role of RT must be considered with care, taking into account the risks and benefits of response versus late toxicity and its impact on functioning and quality of life.

Consolidation High-Dose Chemotherapy and Autologous Stem Cell Transplantation (HDC/ASCT)

Several studies have addressed the safety and efficacy of HDC/ASCT as consolidation in first-line treatment in patients with PCNSL. The first study with HDC/ASCT without WBRT used the BEAM regimen (carmustine (BCNU), etoposide, cytarabine, and melphalan) as conditioning and reported a disappointing median event-free survival of 9.3 months (Class IIIa).¹¹⁰ Subsequently, encouraging studies for which WBRT had been omitted at least in patients in CR after HDC/ASCT using HD thiotepa-based conditioning regimens have been reported (Class IIIb and IV).^{111–114} In these studies the selection bias—patients must meet strict criteria to be eligible for transplantation—should be taken into account. Two prospective multicentre randomized phases II studies, the IELSG-32 and the PRECIS trials have evaluated the role of consolidative HDC/ASCT as part of first-line treatment in patients with PCNSL, in parallel with a control arm with conventional, 36–40 Gy WBRT consolidation.^{10,11} Both studies showed the feasibility and efficacy of thiotepa-based HDC/ASCT in first-line treatment. The per-protocol 2-year PFS was identical in the WBRT and ASCT arms (75–76%) from the date of trial registration in the IELSG study (Class IIa). The intention-to-treat 2-year PFS was 80% in the WBRT arm and 69% in the ASCT arm. In the PRECIS trial, the exploratory analysis of the per-protocol population showed a significant difference of the 2-year PFS from the time of consolidation in favor of ASCT (2-year PFS = 69% after WBRT; and 87% after ASCT; $P = .03$) (Class IIa). Both trials have reported an excess of cognitive decline in the WBRT arm and early and late lymphoma-unrelated deaths in five patients after ASCT in each study (treatment-related death rate of 9% and 11% in the IELSG and PRECIS studies, respectively). Considering all studies, although the direct comparison between conditioning regimens applied is difficult, HD thiotepa-based conditioning regimens seem more efficient than BEAM-based regimens (Class IIIa).^{34,35,115} Because of its toxicity risks, the HDC/ASCT is likely to be proposed for younger patients (<60–65 years) with suitable organ functions. A retrospective European study has suggested that HDC/ASCT with a thiotepa-based conditioning regimen is also feasible and effective in PCNSL patients over the age of 65 (Class IIIb).¹¹⁶

In summary, consolidative HDC/ASCT, preferably conditioned by a thiotepa-based regimen, represents a relevant treatment option with efficacy at least comparable with WBRT despite increased acute toxicity but with less long-term deleterious neurocognitive side-effects recorded after WBRT, as confirmed in recently published long term results of the IELSG and PRECIS trials.^{12,13}

Consolidation Chemotherapy without ASCT

Single-arm phase II trials have suggested that a de-escalated treatment intensity with non-myeloablative consolidation treatment, for example with cytarabine/etoposide, is feasible and effective in patients with newly-diagnosed PCNSL after induction with HD-MTX-based treatment, although with considerable risk of prolonged

grade IV neutropenia (Class IIb).^{36,37} This approach is being investigated in two ongoing randomized trials that use HDC/ASCT as a control arm (NCT02531841, NCT01511562). Preliminary results of the latter, the CALGB 51101 Alliance study, have been reported as a meeting abstract. In this randomized phase II study, 113 patients with newly-diagnosed PCNSL were enrolled and randomly allocated between myeloablative (thiotepa/carmustine) or non-myeloablative (etoposide/cytarabine) consolidation after HD-MTX-based induction. Though PFS was longer after myeloablative treatment (6 years vs 2.4 years) a significant part of this difference was caused by treatment failure before initiation of consolidation and 3 year OS was similar at 83% and 72%, respectively. Initial data suggest no advantage in safety profile in patients treated with non-myeloblastic consolidation. However, conclusions on feasibility, tolerability, and efficacy can be drawn only once full data are published.¹⁶

Salvage Treatment

The treatment of refractory or relapsed PCNSL still remains a huge therapeutic challenge since presently used treatments are of limited benefit.^{48,117} About one-third of patients with PCNSL have disease refractory to the first-line treatment and half of the responders will relapse despite the high response rates seen with initial treatment.^{48,117,118} Many of these patients die early due to lymphoma progression despite the use of salvage therapy, and in patients with recurrent or progressive disease, especially if combined with severe comorbidity or contraindications for chemotherapy, palliative care may be an appropriate strategy. Selection of salvage treatment should be based on results of published phase II trials and depends on performance status, comorbidities, prior treatments, and time from last treatment. After remission MTX rechallenge given as a single agent or in combination may yield a high rate of new objective response and durable remission most likely in patients who previously achieved response lasting after at least several months with HD MTX-based chemotherapy, suggesting retained chemosensitivity to MTX (Class III).^{48–50,117} Alternatively, the ICE/D regimen (ifosfamide, carboplatin, etoposide, and dexamethasone),⁴⁷ the R-IE regimen (rituximab, ifosfamide, and etoposide)⁴⁶ and the ESHAP/DHAP regimen (cytarabine, cisplatin, etoposide, and methylprednisolone) have shown activity as reinduction before consolidation therapy.^{119,120} In patients who had not received any consolidating treatment after HD-MTX-based induction chemotherapy, WBRT or HDC/ASCT should be considered to reduce the risk of relapse. Two retrospective studies have evaluated WBRT delivered as a single option in patients with relapsed PCNSL and reported a high rate of objective responses and a median survival of 11–16 months—quite similar to what is expected with WBRT alone as initial treatment (Class IIIb).^{44,45} Delayed neurotoxicity occurred in 15%–22% and is more pronounced in patients older than 60 years of age. Therefore, if a reasonable systemic treatment-option is available it is plausible to delay or even avoid WBRT due to the high risk of significant neurotoxicity.

HDC/ASCT is an alternative option for consolidation in patients with a good performance status, preferentially offered for younger patients (aged <65 years) (Class IIb).^{41–43} It has been addressed in patients with relapsed or refractory PCNSL in two multicentric phase II trials, both using thiotepa-based conditioning.^{41,121} In the intention to treat the population in the French trial, median PFS and OS were 11 and 18 months.⁴¹ However, in patients who completed the full HDC/ASCT procedure 2 and 3 year OS of 55–69% were found (Class IIIa) with treatment-related mortality of 7–12%.^{41,121} Subsequent retrospective analyses have confirmed these results (Class IIIb).^{42,43} However, age may be less important than overall fitness regarding toxicity, and some promising results are seen also in patients above 65 years of age (Class IIIb).¹¹⁶

Thus, consolidation HDC/ASCT following salvage induction chemotherapy may be associated with prolonged remission in a subset of fit patients (Class IIIa).^{48,117,118} However, most studies were performed in patients who had not received ASCT in the first line.

Other treatment options, if the patient is not suitable for HDC/ASCT, include conventional chemotherapy. There is, however, only a limited number of prospective studies available for guidance and these have been exclusively single-arm phase II trials precluding comparison across studies (Class IIb, III, and IV for all studies in this section). Drugs used as a single agent or in combination, with or without rituximab, that have been evaluated and have demonstrated modest activity include temozolomide,^{51,122} topotecan,¹²³ pemetrexed,¹²⁴ bendamustine,¹²⁵ the PCV regimen,¹²⁶ ifosfamide-etoposide based regimens,^{46,127} HD cytarabine,¹²⁸ cisplatin-cytarabine based regimens,¹²⁰ gemcitabine-oxaliplatin based regimen,¹²⁹ rituximab.⁷⁷ See [table 2](#).

The activity of R-CHOP, not usually used in PCNSL due to low CNS bioavailability, seems to be considerably improved by enhancing the vascular permeability and CNS access using NGR-TNF (Class IIb)¹³⁰ Further studies are underway to evaluate this approach in PCNSL. New innovative approaches using physical methods or physiological transporters to facilitate the passage of targeted therapies across the BBB are of interest to evaluate in the treatment of relapsed PCNSL. However, none of these can be recommended as a standard treatment yet.

Relapses outside the CNS account for only 3% of failures, and some studies suggest that extra-CNS relapses are associated with a better prognosis than CNS-involving relapses.^{118,131} The optimal salvage treatment for this condition remains to be defined, but excellent results have been reported with anthracycline-based chemotherapy consolidated or not with HDC/ASCT.¹³²

Novel Agents

BTK Inhibitors

Promising data are now available for targeted therapies in relapsed disease. Comprehensive molecular analyses of PCNSL tissue revealed a complex architecture of signaling pathways in tumor cells which may be exploited for

therapeutic targeting. The network that comprises Bruton tyrosine kinase (BTK), which acts as a central mediator of B cell receptor (BCR) and Toll-like receptor (TLR) signaling leading to NF- κ B activation,¹³³ has been considered as of particular importance in PCNSL. Therefore, BTK inhibitors have gained increasing interest as novel drugs for the treatment of PCNSL. High response rates to ibrutinib were observed in patients with newly diagnosed PCNSL after treatment with the ibrutinib-based DA-TEDDI-R regimen.¹³⁴ In two phase II trials,^{52,135} single-agent ibrutinib at a dose of 560–840 mg/d, resulted in a response rate of 70–77%, with a complete remission rate of 23–38%, in patients with relapsed or refractory PCNSL. A response rate of 80% was reported for the combination of ibrutinib with HD-MTX, and rituximab⁵³ and signs of clinical activity of tirabrutinib, a second-generation BTK inhibitor, have also been found.¹³⁶ While these findings are encouraging (Class IIb), data from randomized clinical trials are not yet available.

Imids

The thalidomide derivatives lenalidomide and pomalidomide have been investigated in a few prospective trials. Their mechanism of action involves the induction of targeted degradation of disease-relevant proteins but may also include a modulation of the tumor microenvironment.¹³⁷ Lenalidomide was used as single-agent treatment or in combination with other drugs such as rituximab resulting in an overall response rate of 68% and a median PFS of 6 months in patients with recurrent or refractory PCNSL.⁵⁴ Encouraging results were also seen in a phase 2 trial of lenalidomide in combination with rituximab in patients with recurrent or refractory PCNSL or primary intravitreal lymphoma, with an overall response rate in PCNSL patients of 65% and a median PFS of 3.9 months⁵⁵ (Class IIb). The combination of rituximab-lenalidomide-ibrutinib seems feasible and active in heavily pretreated R/R PCNSL (Class IV)¹³⁸; a prospective study is ongoing (NCT03703167). Clinical activity was also noted with pomalidomide in patients with recurrent or refractory PCNSL.¹³⁹ (Class IIIa). The results of ongoing trials which investigate lenalidomide, pomalidomide or other targeted agents in the first line or recurrent setting need to be awaited to clarify their role in PCNSL patients.

PI3K/mTor Inhibitors

The clinical evaluation of drugs targeting the phosphoinositide 3 kinase/mammalian target of the rapamycin (PI3K/mTor) pathway is ongoing. Treatment with the mTOR inhibitor temsirolimus led to an overall response rate of 54% in a phase 2 study in patients with recurrent or refractory PCNSL but a disappointing median PFS of 2.1 months (Class IIb).¹⁴⁰

Immune Checkpoint Inhibitors

Increased expression of PD1, PD-L1, and PD-L2 in PCNSL, recurrent 9p24.1 genomic alterations, and objective response to anti PD1 antibody in a small series (nivolumab)

or case report (pembrolizumab) in relapsing PCNSL^{141,142} (Class IV) has led to some ongoing phase II trials to better specify the therapeutic role of anti PD1 blockade (eg, NCT03012620).¹⁴³

CAR-T

While CD19-targeted chimeric antigen receptor T-cell (CAR-T) therapy has been shown to be active in systemic DLBCL, data supporting CAR-T use in PCNSL are scarce, due to concerns related to possible severe immune effector cell-associated neurotoxicity (ICANS). Two retrospective series (class IIIb) and one phase I/II trial (class IIb), including 5, 9, and 12 highly refractory PCNSL patients respectively, converge to show that CAR-T therapy was well tolerated with manageable and reversible toxicities. In addition, 3 out of 5, 5 out of 9, and 6 out of 12 patients respectively achieved a complete response.^{56,57,144}

In conclusion, patients with relapsed or refractory PCNSL should be assessed for participation in prospective trials addressing new agents, combinations or strategies. When a prospective trial is not available reinduction with HD-MTX or one of the agents studied in phase 2 studies (see Table 2) can be given with or without consolidation with HDCT/ASCT or WBRT or, alternatively, WBRT alone.

Specific Situations

Elderly Patients

As in other entities, the definition of “elderly” is not uniform. However, in the studies available which have evaluated prognostic factors, age over 60 (used by most studies as cut off) was consistently correlated with worse outcomes and found to be highly prognostic for chemoradiation-induced neurotoxicity.³² Elderly patients represent a vulnerable treatment subgroup, due to the poor prognosis and the peculiar sensitivity to treatment side effects. Some prospective, mainly phase II, studies have been published on the treatment of elderly patients with PCNSL (Class IIb),^{8,24–26,80,145,146} and seven prospective studies on patients of all ages which reported specifically on older patients (Class IIIa),^{84,89,93,98,147–149} In the RTOG phase II trial evaluating radiotherapy alone (40 Gy + 20 Gy boost), the median survival was only 7.8 months.⁸⁹ After HD-MTX-based therapy, defined as a dose of MTX \geq 1 g/m² PFS in patients aged 60 or 65 and older is reported between 6 and 16 months and OS between 14 and 37 months (Class IIb and Class III) with OS in the majority of prospective studies under 2 years.^{25,80,84,93,98,112,149–156}

Outside of retrospective studies, no direct comparisons have been made between treatment with HD-MTX-based chemotherapy and radiotherapy in this age group.¹⁵² However, the impression from the single arm and population-based studies is that survival is at least as good and probably better after HD-MTX-based chemotherapy than after radiotherapy (Class IIIb).¹⁵⁷ Formal comparisons of different HD-MTX-based regimens have not been published except in a randomized phase II study, where toxicity was similar and CR rate was 53% with MPV-A (MTX,

procarbazine, vincristine, and cytarabine) vs 38% for MTX and temozolomide, although the difference was not statistically significant (Class IIa).⁸ Concerning toxicity in patients aged over 60, with the exception of one study, in which an intensive multi-drug regimen was used and toxicity was exceedingly high in older patients,⁹³ chemotherapy with MTX doses up to 3.5 g/m² was well tolerated with 2–7% treatment-related mortality, less than 10% grade 3–4 nephrotoxicity and 7–10% of patients discontinuing treatment due to chemotherapy-associated toxicity, though MTX dose was reduced because of decreased renal function in 26–44% of patients.^{8,24–26,42,80,145,158} Retrospective studies substantiate this view. Thus, in general, older patients tolerate treatment with HD-MTX well when adequate supportive measures are used and renal function is accurately monitored.¹⁵⁹

As discussed above, the risk of delayed leukoencephalopathy is particularly high in patients older than 60 years managed with chemoradiotherapy.³² For patients treated with HD-MTX-based chemotherapy without radiotherapy only one study reporting specifically on older patients is available,⁸ but reports including neuropsychological assessment of patients of all ages show little or no cognitive decline in post-treatment evaluations (Class IIIb).^{160,161} Overall, if elderly patients are considered eligible, they should receive HD-MTX-based chemotherapy including drugs that cross the BBB such as an oral alkylating agent.¹⁶² Selected elderly patients with the good clinical condition can be considered for more intensive approaches.¹⁶³ As in other elderly cancer patients, future studies should address the role of geriatric assessment tools adapted to PCNSL patients to identify vulnerabilities to drive treatment choice in this population.¹⁶⁴ Prognosis in old patients with poor general condition (i.e., severe comorbidity, poor performance status lacking of autonomy) and in the very old (over 80) patients is very poor.¹⁵⁶ Acute morbidities and frequent admissions to hospitals associated with HD-MTX chemotherapy need to be individually weighed against the more limited survival benefits in this markedly frail population.

Primary Vitreoretinal Lymphoma

Intraocular infiltration can be the exclusive site of disease at presentation, referred to as primary vitreoretinal lymphoma (PVRL) or as a part of PCNSL with concomitant brain or meningeal disease. The optimal treatment for vitreoretinal lymphoma remains debated. Data on therapy and outcome are scarce and limited to retrospective case reports or mostly small series with heterogeneous patient populations and treatments. Up to 90% of patients with PVRL subsequently develop brain involvement over the course of the disease and dissemination to the brain is the main cause of death.^{39,165} The overall median survival of isolated PVRL regardless of treatment in retrospective series is approximately 60 months.^{39,166} Treatment may be focal, including ocular RT (historically, a total dose of 35–40 Gy, 2 Gy per fraction using opposed lateral beams to include both globes) (Class IV) and intravitreal drug delivery.^{167–169} Uncontrolled series have reported clinical remission with repeated intravitreal MTX and more recently after

rituximab injections (Class IV).^{168,169} More extensive treatments, including systemic chemotherapy and WBRT have also been evaluated and intraocular responses have been reported with HD-MTX,¹⁷⁰ HD cytarabine,^{171,172} ifosfamide, trofosfamide,¹⁷³ and temozolomide¹⁷⁴ used as single agent,¹⁷³ with HD-MTX-based polychemotherapy, and after HDC/ASCT (Class IV).¹⁷⁵ Retrospective multicenter studies have shown conflicting results regarding focal and extensive therapy for isolated PVRL in terms of disease control and survival (Class IIIb).^{38–40} These and other studies failed to provide reliable predictors of brain dissemination in PVRL patients. Nevertheless, given the high risk of relapse in the CNS, and the improved OS after systemic treatment in the most recent series,³⁸ most experts consider that initial treatment of PVRL should not differ from that of PCNSL ie, high-dose MTX-based polychemotherapy followed, or not, by consolidation treatment to eradicate the possible concomitant microscopic disease in the brain and in the CSF in patients fit to receive such treatment. Local treatments would remain options for refractory or recurrent disease confined to the eyes and for older and frail patients. Systemic treatment with HD-MTX-based chemotherapy is associated with an increased ocular relapse rate, probably because of insufficient intraocular drug availability, which led some investigators to add local ocular treatments to improve disease control.³⁸ The management decision should take into account the individual risk of treatment toxicities (including those related to ocular treatment) and center expertise.^{165,176} When intraocular lymphoma is concurrent with brain lesions, it has not been identified as an independent prognostic factor and the prognosis is similar to that of PCNSL without the intraocular disease (Class IIIb).¹⁷⁷ Accordingly, patients with concomitant intraocular and cerebral disease should be treated no differently from PCNSL. In these cases, the value of additional local ocular treatment (ie, intravitreal chemotherapy or ocular radiotherapy if WBRT has not been delivered) to systemic chemotherapy remains a matter of debate, with conflicting results in two retrospective studies (Class IIIb).^{177,178} No studies, other than practice surveys, have been done to evaluate response assessment and follow-up. However, follow-up at least by fundoscopy and slit-lamp examination as well as brain MRI after treatment is recommended; the French LOC group suggests every 6 months during the first 2 years and yearly thereafter.¹⁷⁹

Immunodeficiency-Related PCNSL

The pathogenesis of PCNSL in HIV-infected people and transplant recipients is strongly related to Epstein-Barr virus (EBV) reactivation, which can lead to chronic antigenic B cell stimulation and cell activity transformation.¹⁸⁰ There are no well-defined standard treatment regimens for immunodeficiency-associated PCNSL, including for patients with human immunodeficiency virus (HIV) infection and those receiving immunosuppressants following solid organ transplantation or for chronic autoimmune disorders. In general, the therapeutic strategy is aimed at reducing immunosuppression and administering antitumor treatment based on existing evidence in immunocompetent patients.

HIV-Related PCNSL

PCNSL occurs generally in HIV patients with severe immunosuppression. Thus, if not yet done, initiating or modifying prior antiretroviral therapy (ART) in an attempt to reconstitute the immune system constitutes a major therapeutic intervention in these patients. The optimal ART regimen must be defined in conjunction with the infectious disease specialist, taking into account the patient's HIV genotype, previous treatments, comorbidities, and potential drug interactions. Evidence on the benefit of ART in the treatment of HIV-related PCNSL derives from observational studies that showed that early institution of ART after PCNSL diagnosis was independently associated with improved survival.^{58,59,181} In a few cases, rapid immune recovery and prolonged tumor regression have been achieved with this sole approach.¹⁸² (Class IV). In addition to enhancement of the immune system, the other mainstay of HIV-related PCNSL treatment consists of concurrent administration of systemic chemotherapy with agents with suitable CNS drug concentration, ie, HD-MTX-based regimens.¹⁸³ In general, immunocompromised patients can undergo aggressive treatments identically to immunocompetent patients, with acceptable tolerance and limited morbidity, precluded that the general indications for such therapies are respected. Regarding efficacy, OS varies widely among series and ranges from few months to up to nearly 6 years.⁶⁰⁻⁶² (Class IV) Whether the addition of rituximab to first-line chemotherapy confers a benefit in this population is not clear. Likewise, there is little or no evidence supporting the use of anti-viral agents against EBV,¹⁸⁴ immune checkpoint inhibitors, or other targeted therapies in HIV-related PCNSL patients, but almost no studies have investigated this. Due to historical reasons, WBRT might be still considered by some institutions as an upfront therapeutic option in HIV-associated PCNSL. However, no prospective data are available, and although radiological response rates of up to 30–50% have been described, the outcome of patients treated with WBRT alone remains poor with a median OS of around 3 months (range 1.3–7.8 months).^{58,59,181} On the basis of short response durability and poor survival data, and the high risk of severe neurotoxicity in longer survivors, it is suggested to avoid or reserve palliative WBRT for patients with poor functional status who are not a candidate for chemotherapy (Class IV). For patients with morbidities that preclude safe usage of HD-MTX such as moderate or severe renal insufficiency, treatment decisions must be individualized. In this setting, and extrapolated from its use in immunocompetent patients, other agents with a more favorable toxicity profile such as temozolomide might be considered. WBRT combined with ART might be an alternative, especially for those patients with poor functional status and poor prognosis.

Post-transplant or Immune-Suppression-Related PCNSL

Evidence on the specific treatment of PCNSL in patients receiving immunosuppressants after solid organ transplantation or for a chronic autoimmune disease is lacking

owing to its rarity. In consequence, patients are usually treated in a similar manner to the general population and according to the principles of management of systemic post-transplant lymphoproliferative disorders (PTLD). Few retrospective series describing the management and outcome of patients with primary CNS PTL (PCNS-PTLD) have been published.^{63,64,185-187} Patients were mostly treated with both reduction of immunosuppression and heterogeneous regimens of chemotherapy, including HD-MTX alone or in combination with other agents, or radiation therapy. Despite the unclear benefit of rituximab in PCNSL, treatment protocols based on this agent were frequently employed in these series. The high prevalence of kidney transplants and underlying renal failure in addition to the risk of nephrotoxicity related to HD-MTX might have favored this later strategy. Widely variable OS times ranging from a few months to up to 4 years were observed (Class IV). There is anecdotal evidence of the management of PCNS-PTLD by sole reduction of immunosuppression.¹⁸⁸ A recent series, presented as an abstract, of six patients with PCNS-PTLD treated with temozolomide revealed a complete response in three patients with an OS of 100% at 17 months of follow-up, thus illustrating the potential beneficial role of this drug particularly in patients who are not suitable for HD-MTX treatment (Class IV).¹⁸⁹ Single cases of response with other agents such as ibrutinib have also been recently reported.¹⁹⁰

Conclusions

Guidelines reflect the state of knowledge at a given timepoint. The EANO website will inform of future updates on this guideline (<https://www.eano.eu>).

Funding

None declared.

Authorship

All authors have been involved in the writing of the manuscript at draft and any revision stages, and have read and approved the final version. The authors have been listed in alphabetical order with the exception of the two coordinators who are first and last authors respectively.

Conflict of interest statement: KHX: reports consulting fee from BTG. AJMF: reports speaker fee from Adienne; research grants from BMS, Beigene, Pharmacyclics, Hutchison Medipharma, Amgen, Genmab, ADC Therapeutics, Gilead, Novartis, and Pfizer; advisory boards from Gilead, Novartis, Juno, and PletixaPharm; inventor of patents on NGR-hTNF/RCHOP in relapsed or refractory PCNSL and SNGR-hTNF in brain tumours. BK: reports consultancies from Roche, Riemer, Astellas. CMB: reports honoraria

for webinar from Bayer. MP: reports honoraria for lectures, consultation or advisory board participation from Bayer, Bristol-Myers Squibb, Novartis, Gerson Lehrman Group, CMC Contrast, GlaxoSmithKline, Mundipharma, Roche, BMJ Journals, MedMedia, Astra Zeneca, AbbVie, Lilly, Medahead, Daiichi Sankyo, Sanofi, Merck Sharp & Dome, Tocagen, Adastra, Gan & Lee Pharmaceuticals. PR: reports research grants from MSD and Novocure, and honoraria for advisory board participation or lectures from Bristol-Myers Squibb, Covagen, Debiopharm, Medac, MSD, Novartis, Novocure, QED Therapeutics, Roche, and Virometix. RR: reports honoraria for lectures or advisory boards from Bayer, Novocure, UCB. MW: reports research grants from Apogenix, Merck, Sharp & Dohme, Merck (EMD), Philogen and Quercis, and honoraria for lectures or advisory board participation or consulting from Adastra, Bayer, Bristol Meyer Squibb, Medac, Merck, Sharp & Dohme, Merck (EMD), Nerviano Medical Sciences, Novartis, Orbus, Philogen and y-Mabs. MD, JF, JGPL, RH, AFH, FL, AL, US, RS, CS, MT, VT, JB declare no conflict of interest.

References

- Deckert M BT, Ferry JA, et al. *Primary diffuse large B cell lymphoma of the CNS, World Health Organization Classification of Tumours of the Central Nervous System 2021*. WHO Classification of Tumours Series (ed 5). Lyon, International Agency for Research on Cancer, 2021, pp 351–355
- Diamond C, Taylor TH, Im T, et al: Highly active antiretroviral therapy is associated with improved survival among patients with AIDS-related primary central nervous system non-Hodgkin's lymphoma. *Curr HIV Res*. 2006;4:375–8.
- Ammassari A, Scoppettuolo G, Murri R, et al: Changing disease patterns in focal brain lesion-causing disorders in AIDS. *J Acquir Immune Defic Syndr Hum Retrovirol*. 1998;18:365–71.
- O'Neill BP, Decker PA, Tieu C, et al: The changing incidence of primary central nervous system lymphoma is driven primarily by the changing incidence in young and middle-aged men and differs from time trends in systemic diffuse large B-cell non-Hodgkin's lymphoma. *Am J Hematol*. 2013;88:997–1000.
- van der Meulen M, Dinmohamed AG, Visser O, et al: Improved survival in primary central nervous system lymphoma up to age 70 only: a population-based study on incidence, primary treatment and survival in the Netherlands, 1989–2015. *Leukemia*. 2017;31:1822–1825.
- Mendez JS, Ostrom QT, Gittleman H, et al: The elderly left behind—changes in survival trends of primary central nervous system lymphoma over the past 4 decades. *Neuro Oncol*. 2018;20:687–694.
- Bromberg JEC, Issa S, Bakunina K, et al: Rituximab in patients with primary CNS lymphoma (HOVON 105/ALLG NHL 24): a randomised, open-label, phase 3 intergroup study. *Lancet Oncol*. 2019;20:216–228.
- Omuro A, Chinot O, Taillandier L, et al: Methotrexate and temozolomide versus methotrexate, procarbazine, vincristine, and cytarabine for primary CNS lymphoma in an elderly population: an intergroup ANOCEF-GOELAMS randomised phase 2 trial. *Lancet Haematol*. 2015;2:e251–9.
- Ferreri AJ, Cwynarski K, Pulczynski E, et al: Chemoimmunotherapy with methotrexate, cytarabine, thiopeta, and rituximab (MATRix regimen) in patients with primary CNS lymphoma: results of the first randomisation of the International Extranodal Lymphoma Study Group-32 (IELSG32) phase 2 trial. *Lancet Haematol*. 2016;3:e217–27.
- Ferreri AJM, Cwynarski K, Pulczynski E, et al: Whole-brain radiotherapy or autologous stem-cell transplantation as consolidation strategies after high-dose methotrexate-based chemoimmunotherapy in patients with primary CNS lymphoma: results of the second randomisation of the International Extranodal Lymphoma Study Group-32 phase 2 trial. *Lancet Haematol*. 2017;4:e510–e523.
- Houillier C, Taillandier L, Dureau S, et al: Radiotherapy or autologous stem-cell transplantation for primary CNS lymphoma in patients 60 years of age and younger: results of the intergroup ANOCEF-GOELAMS randomized phase II PRECIS study. *J Clin Oncol*. 2019;37:823–833.
- Ferreri AJM, Cwynarski K, Pulczynski E, et al: Long-term efficacy, safety and neurotolerability of MATRix regimen followed by autologous transplant in primary CNS lymphoma: 7-year results of the IELSG32 randomized trial. *Leukemia*. 2022;36:1870–1878.
- Houillier C DS, Taillandier L, et al. Radiotherapy or autologous stem-cell transplantation for primary CNS lymphoma in patients 60 years of age and younger: long-term results of the randomized phase II PRECIS study. *J Clin Oncol*. 2022;CQ2200491. doi: 10.1200.
- Omuro A, DeAngelis L, Karrison T, et al: Randomized phase II study of rituximab, methotrexate (MTX), procarbazine, vincristine, and cytarabine (R-MPV-A) with and without low-dose whole-brain radiotherapy (LD-WBRT) for newly diagnosed primary CNS lymphoma (PCNSL). *J Clin Oncol*. 2020;38:2501–2501.
- Kazuhiko M, Ryo N, Yoshitaka N, et al: Randomized phase III study of high-dose methotrexate and whole brain radiotherapy with or without concomitant and adjuvant temozolomide in patients with newly diagnosed primary central nervous system lymphoma: JCOG1114C. *J Clin Oncol*. 2020;38:2500–2500.
- Batchelor T. GS, Uppert A.S., Bartlett N.I., Hsi E.D., Cheson B.D., Nayak L., Leonard J.P., Rubenstein J.L.: Myeloablative versus non-myeloablative consolidative chemotherapy for newly diagnosed primary central nervous system lymphoma: results of induction therapy in Alliance 51101. *J Clin Oncol*. 2021;39:7506–7506.
- Hoang-Xuan K, Bessell E, Bromberg J, et al: Diagnosis and treatment of primary CNS lymphoma in immunocompetent patients: guidelines from the European Association for Neuro-Oncology. *Lancet Oncol*. 2015;16:e322–e332.
- Brainin M, Barnes M, Baron JC, et al: Guidance for the preparation of neurological management guidelines by EFNS scientific task forces—revised recommendations 2004. *Eur. J. Neurol*. 2004;11:577–581.
- DeAngelis LM, Yahalom J, Thaler HT, et al: Combined modality therapy for primary CNS lymphoma. *J. Clin. Oncol*. 1992;10:635–643.
- Weller M, Martus P, Roth P, et al: Surgery for primary CNS lymphoma? Challenging a paradigm. *Neuro Oncol*. 2012;14:1481–4.
- Rae AI, Mehta A, Cloney M, et al: Craniotomy and survival for primary central nervous system lymphoma. *Neurosurgery*. 2019;84:935–944.
- Yu J, Du H, Ye X, et al: High-dose methotrexate-based regimens and post-remission consolidation for treatment of newly diagnosed primary CNS lymphoma: meta-analysis of clinical trials. *Sci Rep*. 2021;11:2125.
- Ferreri AJ, Reni M, Foppoli M, et al: High-dose cytarabine plus high-dose methotrexate versus high-dose methotrexate alone in patients with primary CNS lymphoma: a randomised phase 2 trial. *Lancet*. 2009; 374:1512–1520.
- Fritsch K, Kasenda B, Schorb E, et al: High-dose methotrexate-based immuno-chemotherapy for elderly primary CNS lymphoma patients (PRIMAIN study). *Leukemia*. 2017;31:846–852.
- Hoang-Xuan K, Taillandier L, Chinot O, et al: Chemotherapy alone as initial treatment for primary CNS lymphoma in patients older than 60 years: a multicenter phase II study (26952) of the European Organization for Research and Treatment of Cancer Brain Tumor Group. *J Clin Oncol*. 2003;21:2726–2731.
- Pulczynski EJ, Kuittinen O, Erlanson M, et al: Successful change of treatment strategy in elderly patients with primary central nervous system lymphoma by de-escalating induction and introducing temozolomide

- maintenance: results from a phase II study by the Nordic Lymphoma Group. *Haematologica*. 2015;100:534–40.
27. Angelov L, Doolittle ND, Kraemer DF, et al: Blood-brain barrier disruption and intra-arterial methotrexate-based therapy for newly diagnosed primary CNS lymphoma: a multi-institutional experience. *J Clin Oncol*. 2009;27:3503–9.
 28. Schmitt AM, Herbrand AK, Fox CP, et al: Rituximab in primary central nervous system lymphoma-A systematic review and meta-analysis. *Hematol Oncol*. 2019;37:548–557.
 29. Correa DD, DeAngelis LM, Shi W, et al: Cognitive functions in survivors of primary central nervous system lymphoma. *Neurology*. 2004;62:548–555.
 30. Harder H, Holtel H, Bromberg JE, et al: Cognitive status and quality of life after treatment for primary CNS lymphoma. *Neurology*. 2004;62:544–547.
 31. Doolittle ND, Korfel A, Lubow MA, et al: Long-term cognitive function, neuroimaging, and quality of life in primary CNS lymphoma. *Neurology*. 2013;81:84–92.
 32. Omuro AM, Ben Porat LS, Panageas KS, et al: Delayed neurotoxicity in primary central nervous system lymphoma. *Arch Neurol*. 2005;62:1595–1600.
 33. Thiel E, Korfel A, Martus P, et al: High-dose methotrexate with or without whole brain radiotherapy for primary CNS lymphoma (G-PCNSL-SG-1): a phase 3, randomised, non-inferiority trial. *Lancet Oncol*. 2010;11:1036–1047.
 34. Alnahhas I, Jawish M, Alsawas M, et al: Autologous stem-cell transplantation for primary central nervous system lymphoma: systematic review and meta-analysis. *Clin Lymphoma Myeloma Leuk*. 2019;19:e129–e141.
 35. Scordo M, Wang TP, Ahn KW, et al: Outcomes associated with thiotepa-based conditioning in patients with primary central nervous system lymphoma after autologous hematopoietic cell transplant. *JAMA Oncol*. 2021;7:993–1003.
 36. Rubenstein JL, Hsi ED, Johnson JL, et al: Intensive chemotherapy and immunotherapy in patients with newly diagnosed primary CNS lymphoma: CALGB 50202 (Alliance 50202). *J Clin Oncol*. 2013;31:3061–8.
 37. Birsan R, Willems L, Pallud J, et al: Efficacy and safety of high-dose etoposide cytarabine as consolidation following rituximab methotrexate temozolomide induction in newly diagnosed primary central nervous system lymphoma in immunocompetent patients. *Haematologica*. 2018;103:e296–e299.
 38. Lam M, Touitou V, Choquet S, et al: Intravenous high-dose methotrexate based systemic therapy in the treatment of isolated primary vitreoretinal lymphoma: An LOC network study. *Am J Hematol*. 2021;96:823–833.
 39. Grimm SA, Pulido JS, Jahnke K, et al: Primary intraocular lymphoma: an International Primary Central Nervous System Lymphoma Collaborative Group Report. *Ann Oncol*. 2007;18:1851–1855.
 40. Riemens A, Bromberg J, Touitou V, et al: Treatment strategies in primary vitreoretinal lymphoma: a 17-center European collaborative study. *JAMA Ophthalmol*. 2015;133:191–197.
 41. Soussain C, Hoang-Xuan K, Taillandier L, et al: Intensive chemotherapy followed by hematopoietic stem-cell rescue for refractory and recurrent primary CNS and intraocular lymphoma: Societe Francaise de Greffe de Moelle Osseuse-Therapie Cellulaire. *J Clin Oncol*. 2008;26:2512–2518.
 42. Soussain C, Choquet S, Fourme E, et al: Intensive chemotherapy with thiotepa, busulfan and cyclophosphamide and hematopoietic stem cell rescue in relapsed or refractory primary central nervous system lymphoma and intraocular lymphoma: a retrospective study of 79 cases. *Haematologica*. 2012;97:1751–6.
 43. Welch MR, Sauter CS, Matasar MJ, et al: Autologous stem cell transplant in recurrent or refractory primary or secondary central nervous system lymphoma using thiotepa, busulfan and cyclophosphamide. *Leuk Lymphoma*. 2015;56:361–7.
 44. Hottinger AF, DeAngelis LM, Yahalom J, et al: Salvage whole brain radiotherapy for recurrent or refractory primary CNS lymphoma. *Neurology*. 2007;69:1178–1182.
 45. Nguyen PL, Chakravarti A, Finkelstein DM, et al: Results of whole-brain radiation as salvage of methotrexate failure for immunocompetent patients with primary CNS lymphoma. *J Clin Oncol*. 2005;23:1507–1513.
 46. Mappa S, Marturano E, Licata G, et al: Salvage chemoimmunotherapy with rituximab, ifosfamide and etoposide (R-IE regimen) in patients with primary CNS lymphoma relapsed or refractory to high-dose methotrexate-based chemotherapy. *Hematol Oncol*. 2013;31:143–50.
 47. Choi MK, Kang ES, Kim DW, et al: Treatment outcome of relapsed/refractory primary central nervous system diffuse large B-cell lymphoma: a single-center experience of autologous stem cell transplantation. *Int J Hematol*. 2013;98:346–54.
 48. Holdhoff M, Wagner-Johnston N, Roschewski M. Systemic approach to recurrent primary CNS lymphoma: perspective on current and emerging treatment strategies. *Oncotargets Ther*. 2020;13:8323–8335
 49. Plotkin SR, Betensky RA, Hochberg FH, et al: Treatment of relapsed central nervous system lymphoma with high-dose methotrexate. *Clin Cancer Res*. 2004;10:5643–5646.
 50. Pentsova E, Deangelis LM, Omuro A. Methotrexate re-challenge for recurrent primary central nervous system lymphoma. *J Neurooncol*. 2014;117:161–5.
 51. Nayak L, Abrey LE, Drappatz J, et al: Multicenter phase II study of rituximab and temozolomide in recurrent primary central nervous system lymphoma. *Leuk Lymphoma*. 2013;54:58–61.
 52. Soussain C, Choquet S, Blonski M, et al: Ibrutinib monotherapy for relapse or refractory primary CNS lymphoma and primary vitreoretinal lymphoma: final analysis of the phase II “proof-of-concept” iLOC study by the Lymphoma study association (LYSA) and the French oculo-cerebral lymphoma (LOC) network. *Eur J Cancer*. 2019;117:121–130.
 53. Grommes C, Tang SS, Wolfe J, et al: Phase 1b trial of an ibrutinib-based combination therapy in recurrent/refractory CNS lymphoma. *Blood*. 2019;133:436–445.
 54. Rubenstein JL, Geng H, Fraser EJ, et al: Phase 1 investigation of lenalidomide/rituximab plus outcomes of lenalidomide maintenance in relapsed CNS lymphoma. *Blood Adv*. 2018;2:1595–1607.
 55. Ghesquieres H, Chevrier M, Laadhari M, et al: Lenalidomide in combination with intravenous rituximab (REVRI) in relapsed/refractory primary CNS lymphoma or primary intraocular lymphoma: a multicenter prospective “proof of concept” phase II study of the French Oculo-Cerebral lymphoma (LOC) Network and the Lymphoma Study Association (LYSA)†. *Ann Oncol*. 2019;30:621–628
 56. Alcantara M, Houillier C, Blonski M, et al: CAR T-cell therapy in primary central nervous system lymphoma: the clinical experience of the French LOC network. *Blood*. 2022;139:792–796.
 57. Frigault MJ, Dietrich J, Gallagher K, et al: Safety and efficacy of tisagenlecleucel in primary CNS lymphoma: a phase 1/2 clinical trial. *Blood*. 2022;139:2306–2315.
 58. Hoffmann C, Tabrizian S, Wolf E, et al: Survival of AIDS patients with primary central nervous system lymphoma is dramatically improved by HAART-induced immune recovery. *AIDS*. 2001;15:2119–27.
 59. Skiest DJ, Crosby C. Survival is prolonged by highly active antiretroviral therapy in AIDS patients with primary central nervous system lymphoma. *AIDS*. 2003;17:1787–93.
 60. Moulignier A, Lamirel C, Picard H, et al: Long-term AIDS-related PCNSL outcomes with HD-MTX and combined antiretroviral therapy. *Neurology*. 2017;89:796–804.
 61. Gupta NK, Nolan A, Omuro A, et al: Long-term survival in AIDS-related primary central nervous system lymphoma. *Neuro Oncol*. 2017;19:99–108.

62. Lurain K, Uldrick TS, Ramaswami R, et al: Treatment of HIV-associated primary CNS lymphoma with antiretroviral therapy, rituximab, and high-dose methotrexate. *Blood*. 2020;136:2229–2232.
63. Cavaliere R, Petroni G, Lopes MB, et al: Primary central nervous system post-transplantation lymphoproliferative disorder: an International Primary Central Nervous System Lymphoma Collaborative Group Report. *Cancer*. 2010;116:863–70
64. Evens AM, Choquet S, Kroll-Desrosiers AR, et al: Primary CNS posttransplant lymphoproliferative disease (PTLD): an international report of 84 cases in the modern era. *Am J Transplant*. 2013;13:1512–22.
65. Henry JM, Heffner RR, Jr., Dillard SH, et al: Primary malignant lymphomas of the central nervous system. *Cancer*. 1974;34:1293–302.
66. Bataille B, Delwail V, Menet E, et al: Primary intracerebral malignant lymphoma: report of 248 cases. *J Neurosurg*. 2000;92:261–266.
67. Wu S, Wang J, Liu W, et al: The role of surgical resection in primary central nervous system lymphoma: a single-center retrospective analysis of 70 patients. *BMC Neurol*. 2021;21:190
68. Cloney MB, Sonabend AM, Yun J, et al: The safety of resection for primary central nervous system lymphoma: a single institution retrospective analysis. *J Neurooncol*. 2017;132:189–197.
69. Jahr G, Da Broi M, Holte H, Jr., et al: The role of surgery in intracranial PCNSL. *Neurosurg Rev*. 2018;41:1037–1044.
70. Hiraga S, Arita N, Ohnishi T, et al: Rapid infusion of high-dose methotrexate resulting in enhanced penetration into cerebrospinal fluid and intensified tumor response in primary central nervous system lymphomas. *J Neurosurg*. 1999;91:221–30.
71. Herrlinger U, Schabet M, Brugger W, et al: German Cancer Society Neuro-Oncology Working Group NOA-03 multicenter trial of single-agent high-dose methotrexate for primary central nervous system lymphoma. *Ann Neurol*. 2002;51:247–52.
72. Batchelor T, Carson K, O'Neill A, et al: Treatment of primary CNS lymphoma with methotrexate and deferred radiotherapy: a report of NABTT 96-07. *J Clin Oncol*. 2003;21:1044–1049.
73. McAllister LD, Doolittle ND, Guastadisegni PE, et al: Cognitive outcomes and long-term follow-up results after enhanced chemotherapy delivery for primary central nervous system lymphoma. *Neurosurgery*. 2000;46:51–60; discussion 60-1.
74. Seidel S, Pels H, Schlömer S, et al: Twenty-year follow-up of a pilot/phase II trial on the Bonn protocol for primary CNS lymphoma. *Neurology*. 2020;95:e3138–e3144.
75. Rubenstein JL, Combs D, Rosenberg J, et al: Rituximab therapy for CNS lymphomas: targeting the leptomeningeal compartment. *Blood*. 2003;101:466–468.
76. Iwamoto FM, Schwartz J, Pandit-Taskar N, et al: Study of radiolabeled indium-111 and yttrium-90 ibritumomab tiuxetan in primary central nervous system lymphoma. *Cancer*. 2007;110:2528–2534.
77. Batchelor TT, Grossman SA, Mikkelsen T, et al: Rituximab monotherapy for patients with recurrent primary CNS lymphoma. *Neurology*. 2011;76:929–30.
78. Morris PG, Correa DD, Yahalom J, et al: Rituximab, methotrexate, procarbazine, and vincristine followed by consolidation reduced-dose whole-brain radiotherapy and cytarabine in newly diagnosed primary CNS lymphoma: final results and long-term outcome. *J Clin Oncol*. 2013;31:3971–3979.
79. Chamberlain MC, Johnston SK. High-dose methotrexate and rituximab with deferred radiotherapy for newly diagnosed primary B-cell CNS lymphoma. *Neuro Oncol*. 2010;12:736–44.
80. Fritsch K, Kasenda B, Hader C, et al: Immunochemotherapy with rituximab, methotrexate, procarbazine, and lomustine for primary CNS lymphoma (PCNSL) in the elderly. *Ann Oncol*. 2011;22:2080–2085.
81. Birnbaum T, Stadler EA, von Baumgarten L, et al: Rituximab significantly improves complete response rate in patients with primary CNS lymphoma. *J Neurooncol*. 2012;109:285–91.
82. Gregory G, Arumugaswamy A, Leung T, et al: Rituximab is associated with improved survival for aggressive B cell CNS lymphoma. *Neuro Oncol*. 2013;15:1068–73.
83. Holdhoff M, Ambady P, Abdelaziz A, et al: High-dose methotrexate with or without rituximab in newly diagnosed primary CNS lymphoma. *Neurology*. 2014;83:235–9.
84. Pels H, Schmidt-Wolf IG, Glasmacher A, et al: Primary central nervous system lymphoma: results of a pilot and phase II study of systemic and intraventricular chemotherapy with deferred radiotherapy. *J Clin Oncol*. 2003;21:4489–4495.
85. Pels H, Juergens A, Glasmacher A, et al: Early relapses in primary CNS lymphoma after response to polychemotherapy without intraventricular treatment: results of a phase II study. *J Neurooncol*. 2009;91:299–305.
86. Ferreri AJ, Reni M, Pasini F, et al: A multicenter study of treatment of primary CNS lymphoma. *Neurology*. 2002;58:1513–1520.
87. Khan RB, Shi W, Thaler HT, et al: Is intrathecal methotrexate necessary in the treatment of primary CNS lymphoma? *J Neurooncol*. 2002;58:175–178.
88. Sierra Del Rio M, Ricard D, Houillier C, et al: Prophylactic intrathecal chemotherapy in primary CNS lymphoma. *J Neurooncol*. 2012;106:143–6.
89. Nelson DF, Martz KL, Bonner H, et al: Non-Hodgkin's lymphoma of the brain: can high dose, large volume radiation therapy improve survival? Report on a prospective trial by the Radiation Therapy Oncology Group (RTOG): RTOG 8315. *Int J Radiat Oncol Biol Phys*. 1992;23:9–17.
90. Gavrilovic IT, Hormigo A, Yahalom J, et al: Long-term follow-up of high-dose methotrexate-based therapy with and without whole brain irradiation for newly diagnosed primary CNS lymphoma. *J Clin Oncol*. 2006;24:4570–4574.
91. Blay JY, Bouhour D, Carrie C, et al: The C5R protocol: a regimen of high-dose chemotherapy and radiotherapy in primary cerebral non-Hodgkin's lymphoma of patients with no known cause of immunosuppression. *Blood*. 1995;86:2922–9.
92. Poortmans PM, Kluin-Nelemans HC, Haaxma-Reiche H, et al: High-dose methotrexate-based chemotherapy followed by consolidating radiotherapy in non-AIDS-related primary central nervous system lymphoma: European Organization for Research and Treatment of Cancer Lymphoma Group Phase II Trial 20962. *J Clin Oncol*. 2003;21:4483–4488.
93. Ghesquieres H, Ferlay C, Sebban C, et al: Long-term follow-up of an age-adapted C5R protocol followed by radiotherapy in 99 newly diagnosed primary CNS lymphomas: a prospective multicentric phase II study of the Groupe d'Etude des Lymphomes de l'Adulte (GELA). *Ann Oncol*. 2010;21:842–850.
94. Ekenel M, Iwamoto FM, Ben-Porat LS, et al: Primary central nervous system lymphoma: the role of consolidation treatment after a complete response to high-dose methotrexate-based chemotherapy. *Cancer*. 2008;113:1025–1031.
95. Omuro A, Taillandier L, Chinot O, et al: Primary CNS lymphoma in patients younger than 60: can whole-brain radiotherapy be deferred? *J Neurooncol*. 2011;104:323–30.
96. Gerard LM, Imrie KR, Mangel J, et al: High-dose methotrexate based chemotherapy with deferred radiation for treatment of newly diagnosed primary central nervous system lymphoma. *Leuk Lymphoma*. 2011;52:1882–90.
97. Ferreri AJ, Verona C, Politi LS, et al: Consolidation radiotherapy in primary central nervous system lymphomas: impact on outcome of different fields and doses in patients in complete remission after upfront chemotherapy. *Int J Radiat Oncol Biol Phys*. 2011;80:169–75.
98. DeAngelis LM, Seiferheld W, Schold SC, et al: Combination chemotherapy and radiotherapy for primary central nervous system

- lymphoma: Radiation Therapy Oncology Group Study 93-10. *J Clin Oncol.* 2002;20:4643–4648.
99. Herrlinger U, Schäfer N, Fimmers R, et al: Early whole brain radiotherapy in primary CNS lymphoma: negative impact on quality of life in the randomized G-PCNSL-SG1 trial. *J Cancer Res Clin Oncol.* 2017;143:1815–1821.
 100. Lai R, Rosenblum MK, DeAngelis LM. Primary CNS lymphoma: a whole-brain disease? *Neurology.* 2002;59:1557–1562.
 101. Citterio G, Ferreri AJ, Reni M. Current uses of radiation therapy in patients with primary CNS lymphoma. *Expert Rev Anticancer Ther.* 2013;13:1327–37.
 102. Dabaja B, Milgrom S, Parikh R, et al: The challenges of applying radiation in primary central nervous system lymphoma. *Int J Radiat Oncol Biol Phys.* 2020;107:398–400.
 103. Shibamoto Y, Hayabuchi N, Hiratsuka J, et al: Is whole-brain irradiation necessary for primary central nervous system lymphoma? Patterns of recurrence after partial-brain irradiation. *Cancer.* 2003;97:128–133.
 104. Correa DD, Maron L, Harder H, et al: Cognitive functions in primary central nervous system lymphoma: literature review and assessment guidelines. *Ann Oncol.* 2007;18:1145–1151.
 105. Blay JY, Conroy T, Chevreau C, et al: High-dose methotrexate for the treatment of primary cerebral lymphomas: analysis of survival and late neurologic toxicity in a retrospective series. *J Clin Oncol.* 1998;16:864–871.
 106. Abrey LE, DeAngelis LM, Yahalom J. Long-term survival in primary CNS lymphoma. *J Clin Oncol.* 1998;16:859–863.
 107. van der Meulen M, Dirven L, Habets EJJ, et al: Cognitive functioning and health-related quality of life in patients with newly diagnosed primary CNS lymphoma: a systematic review. *Lancet Oncol.* 2018;19:e407–e418.
 108. van der Meulen M, Dirven L, Habets EJJ, et al: Neurocognitive functioning and radiologic changes in primary CNS lymphoma patients: results from the HOVON 105/ALLG NHL 24 randomized controlled trial. *Neuro Oncol.* 2021;23:1315–1326.
 109. Correa DD, Braun E, Kryza-Lacombe M, et al: Longitudinal cognitive assessment in patients with primary CNS lymphoma treated with induction chemotherapy followed by reduced-dose whole-brain radiotherapy or autologous stem cell transplantation. *J Neurooncol.* 2019;144:553–562.
 110. Abrey LE, Moskowitz CH, Mason WP, et al: Intensive methotrexate and cytarabine followed by high-dose chemotherapy with autologous stem-cell rescue in patients with newly diagnosed primary CNS lymphoma: an intent-to-treat analysis. *J Clin Oncol.* 2003;21:4151–6.
 111. Alimohamed N, Daly A, Owen C, et al: Upfront thiotepa, busulfan, cyclophosphamide, and autologous stem cell transplantation for primary CNS lymphoma: a single centre experience. *Leuk Lymphoma.* 2012;53:862–7.
 112. Illerhaus G, Muller F, Feuerhake F, et al: High-dose chemotherapy and autologous stem-cell transplantation without consolidating radiotherapy as first-line treatment for primary lymphoma of the central nervous system. *Haematologica.* 2008;93:147–148.
 113. Kiefer T, Hirt C, Späth C, et al: Long-term follow-up of high-dose chemotherapy with autologous stem-cell transplantation and response-adapted whole-brain radiotherapy for newly diagnosed primary CNS lymphoma: results of the multicenter Ostdeutsche Studiengruppe Hamatologie und Onkologie OSHO-53 phase II study. *Ann Oncol.* 2012;23:1809–12.
 114. Schorb E, Kasenda B, Atta J, et al: Prognosis of patients with primary central nervous system lymphoma after high-dose chemotherapy followed by autologous stem cell transplantation. *Haematologica.* 2013;98:765–70.
 115. Schenone L, Houillier C, Tanguy ML, et al: Intensive chemotherapy followed by autologous stem cell transplantation in primary central nervous system lymphomas (PCNSLs). Therapeutic outcomes in real life-experience of the French Network. *Bone Marrow Transpl.* 2022;57:966–974.
 116. Schorb E, Fox CP, Fritsch K, et al: High-dose thiotepa-based chemotherapy with autologous stem cell support in elderly patients with primary central nervous system lymphoma: a European retrospective study. *Bone Marrow Transpl.* 2017;52:1113–1119.
 117. Tao K, Wang X, Tian X. Relapsed primary central nervous system lymphoma: current advances. *Front Oncol.* 2021;11:649789.
 118. Langner-Lemercier S, Houillier C, Soussain C, et al: Primary CNS lymphoma at first relapse/progression: characteristics, management, and outcome of 256 patients from the French LOC network. *Neuro Oncol.* 2016;18:1297–303.
 119. Ungur R, Tempescul A, Berthou C, et al: ESHAP chemotherapy is efficient in refractory/relapsed primary central nervous system lymphoma: report of four cases. *Onco Targets Ther.* 2015;8:2771–3.
 120. del Rio MS, Choquet S, Hoang-Xuan K, et al: Platine and cytarabine-based salvage treatment for primary central nervous system lymphoma. *J Neurooncol.* 2011;105:409–14.
 121. Kasenda B, Ihorst G, Schroers R, et al: High-dose chemotherapy with autologous haematopoietic stem cell support for relapsed or refractory primary CNS lymphoma: a prospective multicentre trial by the German Cooperative PCNSL study group. *Leukemia.* 2017;31:2623–2629.
 122. Reni M, Mason W, Zaja F, et al: Salvage chemotherapy with temozolomide in primary CNS lymphomas: preliminary results of a phase II trial. *Eur J Cancer.* 2004;40:1682–1688.
 123. Fischer L, Thiel E, Klasen HA, et al: Prospective trial on topotecan salvage therapy in primary CNS lymphoma. *Ann Oncol.* 2006;17:1141–1145.
 124. Raizer JJ, Rademaker A, Evens AM, et al: Pemetrexed in the treatment of relapsed/refractory primary central nervous system lymphoma. *Cancer.* 2012;118:3743–8.
 125. Chamberlain MC. Salvage therapy with bendamustine for methotrexate refractory recurrent primary CNS lymphoma: a retrospective case series. *J Neurooncol.* 2014;118:155–62.
 126. Herrlinger U, Brugger W, Bamberg M, et al: PCV salvage chemotherapy for recurrent primary CNS lymphoma. *Neurology.* 2000;54:1707–1708.
 127. Arellano-Rodrigo E, Lopez-Guillermo A, Bessell EM, et al: Salvage treatment with etoposide (VP-16), ifosfamide and cytarabine (Ara-C) for patients with recurrent primary central nervous system lymphoma. *Eur J Haematol.* 2003;70:219–224.
 128. Chamberlain MC. High-dose cytarabine salvage therapy for recurrent primary CNS lymphoma. *J Neurooncol.* 2016;126:545–50.
 129. Collignon A, Houillier C, Ahle G, et al: (R)-GEMOX chemotherapy for unfit patients with refractory or recurrent primary central nervous system lymphoma: a LOC study. *Ann Hematol.* 2019;98:915–922.
 130. Ferreri AJM, Calimeri T, Ponzoni M, et al: Improving the antitumor activity of R-CHOP with NGR-hTNF in primary CNS lymphoma: final results of a phase 2 trial. *Blood Adv.* 2020;4:3648–3658.
 131. Provencher S, Ferlay C, Alaoui-Slimani K, et al: Clinical characteristics and outcome of isolated extracerebral relapses of primary central nervous system lymphoma: a case series. *Hematol Oncol.* 2011;29:10–6.
 132. Ferreri AJ, Ciceri F, Brandes AA, et al: MATILDE chemotherapy regimen for primary CNS lymphoma: results at a median follow-up of 12 years. *Neurology.* 2014;82:1370–3.
 133. Hendriks RW, Yuvaraj S, Kil LP. Targeting Bruton's tyrosine kinase in B cell malignancies. *Nat Rev Cancer.* 2014;14:219–32.
 134. Lionakis MS, Dunleavy K, Roschewski M, et al. Inhibition of B cell receptor signaling by ibrutinib in primary CNS lymphoma. *Cancer Cell.* 2017;31:833–843 e5.

135. Grommes C, Pastore A, Palaskas N, et al: Ibrutinib unmasks critical role of Bruton tyrosine kinase in primary CNS lymphoma. *Cancer Discov.* 2017;7:1018–1029.
136. Narita Y, Nagane M, Mishima K, et al: Phase I/II study of tirabrutinib, a second-generation Bruton's tyrosine kinase inhibitor, in relapsed/refractory primary central nervous system lymphoma. *Neuro Oncol.* 2021;23:122–133.
137. Jan M, Sperling AS, Ebert BL. Cancer therapies based on targeted protein degradation—lessons learned with lenalidomide. *Nat Rev Clin Oncol.* 2021;18:401–417.
138. Houillier C, Chabrot CM, Moles-Moreau MP, et al: Rituximab-lenalidomide-ibrutinib combination for relapsed/refractory primary CNS lymphoma: a case series of the LOC network. *Neurology.* 2021;97:628–631.
139. Tun HW, Johnston PB, DeAngelis LM, et al: Phase 1 study of pomalidomide and dexamethasone for relapsed/refractory primary CNS or vitreoretinal lymphoma. *Blood.* 2018;132:2240–2248.
140. Korfel A, Schlegel U, Herrlinger U, et al: Phase II trial of temsirolimus for relapsed/refractory primary CNS lymphoma. *J Clin Oncol.* 2016;34:1757–63.
141. Nayak L, Iwamoto FM, LaCasce A, et al: PD-1 blockade with nivolumab in relapsed/refractory primary central nervous system and testicular lymphoma. *Blood.* 2017;129:3071–3073.
142. Graber JJ, Plato B, Mawad R, et al: Pembrolizumab immunotherapy for relapsed CNS lymphoma. *Leuk Lymphoma.* 2020;61:1766–1768.
143. Hoang-Xuan K HR, Soussain C et al. First results of the Acsé pembrolizumab phase II in the primary CNS lymphoma (PCNSL) Cohort, *ASH.* 2020:15–16. doi:10.1182/blood-2020-141773.
144. Siddiqi T, Wang X, Blanchard MS, et al: CD19-directed CAR T-cell therapy for treatment of primary CNS lymphoma. *Blood Adv.* 2021;5:4059–4063.
145. Illerhaus G, Marks R, Muller F, et al: High-dose methotrexate combined with procarbazine and CCNU for primary CNS lymphoma in the elderly: results of a prospective pilot and phase II study. *Ann Oncol.* 2009;20:319–325
146. Laack NN, Ballman KV, Brown PB, et al: Whole-brain radiotherapy and high-dose methylprednisolone for elderly patients with primary central nervous system lymphoma: Results of North Central Cancer Treatment Group (NCCTG) 96-73-51. *Int J Radiat Oncol Biol Phys.* 2006;65:1429–1439.
147. O'Neill BP, O'Fallon JR, Earle JD, et al: Primary central nervous system non-Hodgkin's lymphoma: survival advantages with combined initial therapy? *Int J Radiat Oncol Biol Phys.* 1995;33:663–673.
148. Schultz C, Scott C, Sherman W, et al: Preirradiation chemotherapy with cyclophosphamide, doxorubicin, vincristine, and dexamethasone for primary CNS lymphomas: initial report of radiation therapy oncology group protocol 88-06. *J Clin Oncol.* 1996;14:556–564.
149. Roth P, Martus P, Kiewe P, et al: Outcome of elderly patients with primary CNS lymphoma in the G-PCNSL-SG-1 trial. *Neurology.* 2012;79:890–6.
150. Omuro AM, Taillandier L, Chinot O, et al: Temozolomide and methotrexate for primary central nervous system lymphoma in the elderly. *J Neurooncol.* 2007;85:207–211.
151. Ney DE, Reiner AS, Panageas KS, et al: Characteristics and outcomes of elderly patients with primary central nervous system lymphoma: the Memorial Sloan-Kettering Cancer Center experience. *Cancer.* 2010;116:4605–4612.
152. Schuurmans M, Bromberg JEC, Doorduijn J, et al: Primary central nervous system lymphoma in the elderly: a multicentre retrospective analysis. *Br J of Haematol.* 2010;151:179–184.
153. Taoka K, Okoshi Y, Sakamoto N, et al: A nonradiation-containing, intermediate-dose methotrexate regimen for elderly patients with primary central nervous system lymphoma. *Int J Hematol.* 2010;92:617–623.
154. Zhu JJ, Gerstner ER, Engler DA, et al: High-dose methotrexate for elderly patients with primary CNS lymphoma. *Neuro Oncol.* 2009;11:211–215.
155. Kurzwelly D, Glas M, Roth P, et al: Primary CNS lymphoma in the elderly: temozolomide therapy and MGMT status. *J Neurooncol.* 2010;97:389–392.
156. Welch MR, Omuro A, DeAngelis LM. Outcomes of the oldest patients with primary CNS lymphoma treated at Memorial Sloan-Kettering Cancer Center. *Neuro Oncol.* 2012;14:1304–1311.
157. van der Meulen M, Bromberg JEC, Nijland M, et al: Primary therapy and survival in patients aged over 70-years-old with primary central nervous system lymphoma: a contemporary, nationwide, population-based study in the Netherlands. *Haematologica.* 2021;106:597–600.
158. Jahnke K, Korfel A, Martus P, et al: High-dose methotrexate toxicity in elderly patients with primary central nervous system lymphoma. *Ann Oncol.* 2005;16:445–449.
159. Bessell EM, Dickinson P, Dickinson S, et al: Increasing age at diagnosis and worsening renal function in patients with primary central nervous system lymphoma. *J Neurooncol.* 2011;104:191–193.
160. Juergens A, Pels H, Rogowski S, et al: Long-term survival with favorable cognitive outcome after chemotherapy in primary central nervous system lymphoma. *Ann Neurol.* 2010;67:182–189.
161. Correa DD, Shi W, Abrey LE, et al: Cognitive functions in primary CNS lymphoma after single or combined modality regimens. *Neuro Oncol.* 2012;14:101–108.
162. Kasenda B, Ferreri AJ, Marturano E, et al: First-line treatment and outcome of elderly patients with primary central nervous system lymphoma (PCNSL)—a systematic review and individual patient data meta-analysis. *Ann Oncol.* 2015;26:1305–1313.
163. Schorb E, Kasenda B, Ihorst G, et al: High-dose chemotherapy and autologous stem cell transplant in elderly patients with primary CNS lymphoma: a pilot study. *Blood Adv.* 2020;4:3378–3381.
164. Mohile SG, Dale W, Somerfield MR, et al: Practical assessment and management of vulnerabilities in older patients receiving chemotherapy: ASCO guideline for geriatric oncology. *J Clin Oncol.* 2018;36:2326–2347.
165. Chan CC, Rubenstein JL, Coupland SE, et al: Primary vitreoretinal lymphoma: a report from an International Primary Central Nervous System Lymphoma Collaborative Group symposium. *Oncologist.* 2011;16:1589–99.
166. Hormigo A, Abrey L, Heinemann MH, et al: Ocular presentation of primary central nervous system lymphoma: diagnosis and treatment. *Br J Haematol.* 2004;126:202–8.
167. Berenbom A, Davila RM, Lin HS, et al: Treatment outcomes for primary intraocular lymphoma: implications for external beam radiotherapy. *Eye (Lond).* 2007;21:1198–201.
168. Frenkel S, Hendler K, Siegal T, et al: Intravitreal methotrexate for treating vitreoretinal lymphoma: 10 years of experience. *Br J Ophthalmol.* 2008;92:383–8.
169. Hashida N, Ohguro N, Nishida K. Efficacy and complications of intravitreal rituximab injection for treating primary vitreoretinal lymphoma. *Transl Vis Sci Technol.* 2012;1:1.
170. Batchelor TT, Kolak G, Giordia R, et al: High-dose methotrexate for intraocular lymphoma. *Clin Cancer Res.* 2003;9:711–5.
171. Siegel MJ, Dalton J, Friedman AH, et al: Ten-year experience with primary ocular "reticulum cell sarcoma" (large cell non-Hodgkin's lymphoma). *Br J Ophthalmol.* 1989;73:342–6.
172. Strauchen JA, Dalton J, Friedman AH. Chemotherapy in the management of intraocular lymphoma. *Cancer.* 1989;63:1918–21.

173. Jahnke K, Thiel E, Bechrakis NE, et al: Ifosfamide or trofosfamide in patients with intraocular lymphoma. *J Neurooncol.* 2009;93:213–7.
174. Baron M, Belin L, Cassoux N, et al: Temozolomide is effective and well tolerated in patients with primary vitreoretinal lymphoma. *Blood.* 2020;135:1811–1815.
175. Soussain C, Suzan F, Hoang-Xuan K, et al: Results of intensive chemotherapy followed by hematopoietic stem-cell rescue in 22 patients with refractory or recurrent primary CNS lymphoma or intraocular lymphoma. *J Clin Oncol.* 2001;19:742–749.
176. Rajagopal R, Harbour JW. Diagnostic testing and treatment choices in primary vitreoretinal lymphoma. *Retina.* 2011;31:435–40.
177. Grimm SA, McCannel CA, Omuro AM, et al: Primary CNS lymphoma with intraocular involvement: International PCNSL Collaborative Group Report. *Neurology.* 2008;71:1355–1360.
178. Ferreri AJ, Blay JY, Reni M, et al: Relevance of intraocular involvement in the management of primary central nervous system lymphomas. *Ann Oncol.* 2002;13:531–538.
179. Malaise D, Houillier C, Touitou V, et al: Primary vitreoretinal lymphoma: short review of the literature, results of a European survey and French guidelines of the LOC network for diagnosis, treatment and follow-up. *Curr Opin Oncol.* 2021;33:420–431.
180. Schaadt E, Baier B, Mautner J, et al: Epstein-Barr virus latent membrane protein 2A mimics B-cell receptor-dependent virus reactivation. *J Gen Virol.* 2005;86:551–559.
181. Newell ME, Hoy JF, Cooper SG, et al: Human immunodeficiency virus-related primary central nervous system lymphoma: factors influencing survival in 111 patients. *Cancer.* 2004;100:2627–36.
182. Travi G, Ferreri AJ, Cinque P, et al: Long-term remission of HIV-associated primary CNS lymphoma achieved with highly active antiretroviral therapy alone. *J Clin Oncol.* 2012;30:e119–21.
183. González-Aguilar A, Soto-Hernández JL. The management of primary central nervous system lymphoma related to AIDS in the HAART era. *Curr Opin Oncol.* 2011;23:648–53.
184. Bossolasco S, Falk KI, Ponzoni M, et al: Ganciclovir is associated with low or undetectable Epstein-Barr virus DNA load in cerebrospinal fluid of patients with HIV-related primary central nervous system lymphoma. *Clin Infect Dis.* 2006;42:e21–5.
185. Castellano-Sanchez AA, Li S, Qian J, et al: Primary central nervous system posttransplant lymphoproliferative disorders. *Am J Clin Pathol.* 2004;121:246–53.
186. Mahale P, Shiels MS, Lynch CF, et al: Incidence and outcomes of primary central nervous system lymphoma in solid organ transplant recipients. *Am J Transpl.* 2018;18:453–461
187. Kaulen LD, Galluzzo D, Hui P, et al: Prognostic markers for immunodeficiency-associated primary central nervous system lymphoma. *J Neurooncol.* 2019;144:107–115.
188. Valavoor SH, Ashraf Z, Narwal R, et al: Conservative management of post-transplant central nervous system lymphoma. *Int Urol Nephrol.* 2013;45:1219–22
189. Krishnamurthy J GT, Bhatt V, White M, Bierman P. Temozolomide in primary central nervous system post-transplant lymphoproliferative disorder (CNS-PTLD): a case series. *Neuro-Oncology.* 2014;16:v119
190. Kallam A, Hansen N, Bierman P. Central Nervous System Post-transplant lymphoproliferative disorder: response to ibrutinib. *Clin Lymphoma Myeloma Leuk.* 2020;20:e930–e933.