



HHS Public Access

Author manuscript

Arch Clin Biomed Res. Author manuscript; available in PMC 2023 January 12.

Published in final edited form as:

Arch Clin Biomed Res. 2022 ; 6(5): 771–780. doi:10.26502/acbr.50170290.

Diaphragm Ultrasound: A Valuable Predictor of the Outcome of Extubation. An Observational Pilot Study in Covid-19 Related ARDS

Veronica Gagliardi^{1,*}, Angelo Butturini², Gioconda Ferraro², Giuseppe Gagliardi^{1,2}

¹University of Padua, Italy

²Department of Anesthesia and Intensive Care, Ospedale Santa Maria della Misericordia, AULSS 5 Polesana, Rovigo, Italy

Abstract

Introduction: Because both early and delayed weaning are associated with increased mortality, longer stay in the ICU and higher economic costs, performing extubation once the patient can cope with the respiratory load is completely recommended. Ultrasound Sonography (US) is an available bedside tool that allows a rapid assessment and visualization of the different structures involved in spontaneous breath. M-mode ultrasonography can be useful for the assessment of diaphragm kinetics, providing valuable information about diaphragm dysfunction.

Aim of the Study: The aim of this study is to find a correlation between the value of the acceleration of the diaphragm detected with the US M-mode and the outcome of the weaning.

Materials and Methods: We have enrolled 19 patients admitted in our ICU. Each patient underwent the trial with the ultrasound M-mode to assess the acceleration of the diaphragm during the contraction. We have analyzed the results relating them to the outcome of the weaning.

Results: While 11 of our patients have had a successful weaning, 8 have failed it, and we can see that the outcome is associated to the values of acceleration.

*Corresponding author: Veronica Gagliardi, University of Padua, Italy.

Authors' Contributions

G.G. and V.G. conceived the original idea. V.G. developed the theory and performed the computations. V.G. verified the analytical methods. G.G., A.B. and G.F. carried out the experiment. G.G. supervised the findings of this work. All authors discussed the results and contributed to the final manuscript.

Competing Interests

I declare that the authors have no competing interests, or other interests that might be perceived to influence the results and/or discussion reported in this paper.

Declarations

Ethical Approval and Consent to Participate

Approved by the ethical committee (comitato Etico per la Sperimentazione Clinica delle Province di Verona e Rovigo), Project number: 3658CEESC. Written informed consent was obtained from all patients. All methods were carried out in accordance with relevant guidelines and regulations of ethical principles for medical research involving human, stated in the Declaration of Helsinki.

Consent for Publication

Not applicable.

Availability of supporting data

The data that support the findings of this study are openly available.

Discussion: Our study has demonstrated that an assessment of the diaphragm function using US could represent a usable and effective technique as the acceleration is related to the force generated by the diaphragm contraction.

Conclusions: In conclusion, the acceleration could be a useful parameter to consider when it comes to the prediction of the outcome of the weaning process.

Keywords

Diaphragm; ICU; Rapid Shallow Breathing Index; Ultrasound Sonography

Introduction

Even though mechanical ventilation is an established widespread supportive treatment for respiratory failure, it is not risk-free: when prolonged, it increases the risk of pneumonia, barotrauma, tracheal injuries, and musculoskeletal deconditioning [1]. Conversely, premature removal of mechanical ventilation entails a high risk of weaning failure, and prompting reintubation exposes the patient to unnecessary hemodynamic and respiratory stress [2,3]. Both early and delayed weaning are associated with increased mortality, longer stay in the ICU and higher economic costs: performing extubation once the patient is able to cope with the respiratory load is completely recommended [1,4]. Prolonged mechanical ventilation can lead to ventilator-induced diaphragmatic dysfunction, in a context where the diaphragm is also vulnerable to damage from hypotension, hypoxia and sepsis [4]. Disuse atrophy of the diaphragm plays a pivotal role in weaning failure and from the examination of diaphragm biopsies we can infer that the changes of the structure occur early after intubation [5]. To sum up, diaphragm dysfunction is defined as the inability of the diaphragm to generate reasonable levels of maximal force and it is an under-detected pathological condition in critically ill patients which could impair weaning from mechanical ventilation. Weaning failure, defined as the requirement of invasive or noninvasive mechanical ventilation within 48 hours after extubation, is extremely common: about 20% of mechanically ventilated patients require reintubation [6]. Although the implementation of spontaneous breathing trials is recommended by current guidelines, this practice is ineffective alone in predicting weaning failure and consequent reintubation [2]. The identification of the suitable conditions for weaning should be evaluated daily, assessing the improvement of the respiratory failure, of the ability to protect the airway and to start an inspiratory effort, of the fluid balance and the acid-base equilibrium, and of the level of consciousness [3]. In the last few years multiple indices and parameters have been proposed as predictors of weaning outcome, but none of them has shown more than a modest prognostic accuracy [2]: performing the weaning in the optimal moment is still a current challenge [3]. Considering the expiratory physiology research, the evaluation of esophageal pressure using esophageal manometry helps to estimate the trans-diaphragmatic pressure, the gold standard to evaluate the diaphragm force: when this is less than 11 cmH₂O diaphragm dysfunction occurs [8]. Many other weaning parameters have been used to predict weaning failure, including rapid shallow breathing index (RSBI), which is the ratio of respiratory rate to tidal volume: RSBI <105 correlates with weaning success, whereas a score >105 correlates with weaning failure [1], minute ventilation (VE),

maximum inspiratory pressure (P_Imax) [4], and the pressure developed in the occluded airway 100 ms after the onset of an inspiratory effort: P_{O.1} [7]. In this context, Ultrasound Sonography (US) is an available bedside tool that allows rapid assessment and visualization of the thoracic and abdominal structures. More recently, ultrasonography has been used to assess the diaphragm kinetics. The two-dimensional mode is initially used to obtain the best approach and select the exploration line; the M-mode is then used to display the motion of the anatomical structures along the selected line [9]. By providing a direct visualization of the diaphragm with both morphological and functional information in real time, [10] the use of ultrasonographic assessment of diaphragmatic dysfunction could well represent a valuable predictor of weaning failure [1]. To study the diaphragm function, the patient is placed in supine position with the trunk elevated of 10–15°, the function of each hemidiaphragm can be explored with a low frequency (3.5–5 MHz) ultrasound transducer (convex or phased array probe) placed along the midclavicular line or below the costal margin in the longitudinal plane. Technical difficulties in visualizing the left diaphragm have been described, leading most clinicians to perform right diaphragm measurements [11,12]. The M-mode is then used along the selected line to show movements and measure the excursion or displacement of a point of the hemidiaphragm during the respiratory phases. The ultrasound on M-mode trace appears as a wave shape: from the curve obtained the diaphragmatic excursion, the inspiratory and expiratory times. In addition, the M-mode can distinguish diaphragmatic weakness from paralysis: the first shows a reduced diaphragmatic caudal movement, whereas the second is characterized by a paradoxical motion. As far as the speed of diaphragmatic contraction is concerned, a study performed a diaphragmatic sonography with the M-mode technique, calculating the diaphragm contraction speed as the slope (Sc_{di}) of the curve provided by the diaphragm contraction during the inspiratory phase of the spontaneous breathing trials. The contraction speed represented a bedside, standardized and reproducible tool to predict the outcome of weaning [13]. Diaphragmatic ultrasound may identify patients at risk of weaning failure, it is mandatory to standardize the diagnostic criteria of diaphragm dysfunction and the diagnostic performance to predict weaning outcome [3], also because the technique is operator dependent and needs dedicated training. While studying the diaphragm dynamics with the US, we have detected a new parameter, the acceleration of the diaphragm, obtainable dividing the time of contraction to the speed of it, studied with the M-mode of the US.

Aim of the Study

Firstly, we want to study the correlation between the value of the acceleration of the diaphragm detected with the US M-mode and the outcome of the weaning. Secondly, we want to assess if this value could be predictive of success or failure in weaning.

Materials and Methods

We have enrolled 19 patients admitted in the ICU between March 2020 and April 2021. We have included only patients admitted in the ICU in the above-mentioned period in which neither the available recommended methods (SBT and RSBI, VE, P_Imax and P_{O.1}) nor clinical parameters were reliable to predict the outcome of extubation. Their mean age was 66 years old; 3 patients were female, with a mean age of 66 years old, and 16 patients were

male, with a mean age of 67 years old (Table 1). Each patient underwent the trial with the ultrasound M-mode to assess the acceleration of the diaphragm. The experimental protocol consists of an ultrasonographic analysis of the diaphragmatic activity, using the MyLab Esaote and the probe Convex 5–2 MHz. The probe has been located on the right middle axillary line, in plane along with the sagittal plan. An inclination of 15–20° has then been applied, to direct the ultrasound beam perpendicularly to the rear third of the homolateral hemidiaphragm. Because of the interindividual variability of the patients' built, the anatomic landmark for each has been identified either in the seventh, or in the eighth, or in the ninth intercostal space, choosing the one which allowed the best acoustic window to detect the diaphragmatic dome and its excursion both in M-mode and in B-mode. Focusing on the M-mode, the following parameters has been measured (Figure 1):

- Diaphragmatic excursion (displacement, cm)
- Time of inspiration (T_{insp}, s)
- Velocity of diaphragm contraction (slope, cm/s).

The outcome of the further extubation has been detected (Table 2). We define a successful weaning as the absence of ventilatory support for 48 hours after extubation, whereas weaning failure is defined as the need for reintubation within 48 h following extubation. Once we have obtained the measure of the time and the speed, we have calculated the value of the acceleration for each patient. Hence, we have analyzed our data and we found a cut-off predictive value.

Data Analysis

We have measured the space and the time of contraction from the graph obtained by M-mode, and we have applied the form of the uniformly accelerated movement to the curve of each patient:

$$x = x_0 + v_0t + 1/2at^2$$

where we consider x_0 to be 0 mm, and v_0 to be 0 mm/s, and we have obtained the following value, by explicating the acceleration:

$$a = 2 * s/t^2$$

We have calculated it for each patient, and we have analyzed the results relating them to the outcome of the weaning.

Statistical Analysis

We have applied Mann-Whitney U test for independent measures. We have calculated the rank sum, and we have obtained the U_{stat} value. Then, we have compared it with the U_{crit} value deriving from the table reported in Appendix A (critical values of the Mann-Whitney U test), obtaining our statistically significant result. Then we have used Chi-Square test

to find out the difference between the observed and the expected data to analyze if there is a relationship between the entity of the diaphragmatic acceleration and the outcome of extubation.

Results

At the end of our analysis, 11 of our patients have had a successful weaning, whereas 8 have failed it. This result does not seem to correlate neither with the sex and the age of the patients (Table 1), nor with the time of intubation, the time of ICU stay or the duration of positivity for SARS-CoV2 (Table 1, 2).

On the patients enrolled, we have conducted the analysis with the UltraSound: using the B-mode, we have identified a point belonging to the diaphragm, to study its kinetics using the M-mode. The curve detected in M-mode represents the oscillation of a point of the diaphragm, and, due the tension and elastic forces involved in the system, we can consider it as a uniformly accelerated motion, defined by a constant acceleration. Hence, the above-mentioned curve can be assimilated to a space-time graph showing the motion of a material point, where the x-axis shows the time (t), and the y-axis the distance covered by the material point: the space (s), which is the amplitude of contraction (Figure 1). We have then calculated the acceleration for each patient, using the form $a=2*s/t^2$: the obtained values are reported in Table 3. Our technique and the means through which we have obtained our results for each patient are shown in Figure 2.

We can see that both positive and negative outcomes are associated to different values of acceleration: in particular, patients with a negative outcome have a value of acceleration from 52 mm/s^2 to $128,9 \text{ mm/s}^2$, whereas patients who failed the extubation have values included between $7,4 \text{ mm/s}^2$ and 51 mm/s^2 , they are plotted in the graph of Figure 3. The median value of the population who failed extubation is far lower than the median value of the population with a successful extubation:

$$\text{MedianFailed} = 34,1 \text{ mm/s}^2 < \text{MedianSuccessful} = 101,4 \text{ mm/s}^2.$$

Furthermore, we can identify an estimated cut-off value, 52 mm/s^2 . A value of acceleration inferior to 52 mm/s^2 may well correlate with a negative outcome of weaning, whereas a value of acceleration superior to 52 mm/s^2 may well be predictable of successful weaning.

Statistical Analysis

We have applied Mann-Whitney U test for independent measures (Wilcoxon rank-sum test). We have calculated the rank sum, and we have obtained the U_{stat} value.

Null hypothesis H_0 : the distributions of both populations are equal.

Alternative hypothesis H_1 : the distributions are not equal.

We assigned each observation to a Rank (Table 4), and we calculated the rank-sum for each population:

$$\Sigma_{\text{Failed}} = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 = 36$$

$$\Sigma_{\text{Successful}} = 9 + 10 + 11 + 12 + 13 + 14 + 15 + 16 + 17 + 18 + 19 = 154$$

$$U_{\text{stat}} = \Sigma - n(n + 1)/2$$

$$U_{\text{statS}} = 154 - 11(11 + 1)/2 = 154 - 66 = 88$$

$$U_{\text{statF}} = 36 - 8(8 + 1)/2 = 36 - 36 = 0$$

$$U_{\text{Crit}} = 13 \text{ (Appendix A)} \quad \alpha = 0,01$$

$$U_{\text{statF}} < U_{\text{Crit}} \quad \text{with } \alpha = 0,01$$

We reject H₀, the difference between the two groups is statistically significant, it is not casual, so the distributions of populations are not equal (Figure 3).

Chi-Square Test

We have then calculated the χ^2 test, considering values reported in Table 4 and Table 5, and we have obtained:

$$\chi^2 = \sum (O_i - E_i)^2 / E_i = 19$$

where

O_i = observed value (actual value)

E_i = expected value.

From the table reported in Appendix B, we deduce that we have found a correlation between the value of the acceleration and the outcome of extubation of 99,5%, considering a level of confidence of 0,05.

Discussion

Defining the right time for extubation remains a debated issue because of the difficulty in predicting the outcome before the maneuver. Moreover, it results to be even more

challenging in patients who were affected by Covid-19 related ARDS (C-ARDS), because of the severity of the lung failure, the total amount, and the dosage of drugs in the ICU, and the aggressive prolonged mechanical ventilation. Therefore, a reliable method to make extubation as safe as possible is currently required. With regard to this subject, our study has demonstrated that an assessment of the diaphragm using US could well represent a usable and effective technique. It has been in fact broadly studied in literature, where the detection and the relationships between various parameters have been examined [4]. US is an available and easily usable bedside technique, cost effective and even though operator-dependent, repeatable, as we have focused on the detection of objective and measurable parameters. In this framework, our study highlights the use of B-mode to identify a point belonging to the diaphragm, and the use of M-mode to study the kinetics. The curve detected in M-mode represents the oscillation of a point of the diaphragm, and, due the tension and elastic forces involved in the system, we can consider it as a uniformly accelerated motion, defined by a constant acceleration. Hence, the above-mentioned curve can be assimilated to a space-time graph showing the motion of a material point, where the x-axis shows the time (t), and the y-axis the distance covered by the material point: the space (s) that is the amplitude of contraction. In particular, the uniform acceleration plays a pivotal role in our study, because it results to be directly proportional to the transdiaphragmatic pressure, the gold standard parameter considered when it comes to the prediction of success or failure in weaning.

We can demonstrate as follows:

P = resultant pressure acting on diaphragm, consisting of the trans-diaphragmatic pressure (P);

a = acceleration of the diaphragmatic contraction.

The resultant force acting on the diaphragm is the force of the diaphragmatic contraction, corresponding to $P \cdot S_d$, where S_d is the area of the diaphragmatic surface.

Moreover, for Newton's second law of motion, the resultant force results to be $m_d \cdot a$, where m_d is the diaphragm mass.

So we obtain the following system of equations:

$$F = m_d \cdot a$$

$$F = \Delta P \cdot S_d$$

$$\rightarrow \Delta P \cdot S_d = m_d \cdot a \rightarrow a = (S_d/m_d) \cdot \Delta P$$

The pressure and the acceleration results to be directly proportional, and we can consider the physical quantity S_d/m_d a constant quantity, typical of each system, so it is different from

one patient to another, but it is unnecessary to calculate, because our interest is on the fact that the two parameters are strictly related by a constant value that we can call k.

$$\rightarrow a = k * \Delta P.$$

Hence, we can demonstrate that the acceleration is related both to the force generated by the diaphragm contraction, and to the transdiaphragmatic pressure, a reliable measure rarely employed to decide the timing of weaning. Therefore, we can also consider our technique related to the measure of transdiaphragmatic pressure, but with the advantage of being non-invasive. To sum up, the outcome of extubation is not related to sex, age, mode of ventilation, or duration of ICU stay and of intubation. On the contrary, the calculated value of the acceleration is strongly correlated to the outcome of the weaning, so it represents a worth parameter which could play a pivotal role in the process of decision-making, detected with a bedside and cost-effective technique.

Conclusions

In conclusion, the acceleration, being directly proportional to the trans-diaphragmatic pressure, and to the force generated by diaphragm contraction, could well be a useful parameter to consider when it comes to the outcome of the weaning process. It is a reliable measure, bedside and cost effective. This is an observational pilot study, so more studies including a higher number of patients are needed to corroborate our findings.

Acknowledgements

All the material is owned by the authors and/or no permissions are required.

Availability of Data and Materials

All data generated or analysed during this study are included in this published article.

Appendix A

Critical Values of the Mann-Whitney U

(Two-Tailed Testing)

n ₂	α	n ₁																	
		3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
3	.05	--	0	0	1	1	2	2	3	3	4	4	5	5	6	6	7	7	8
	.01	--	0	0	0	0	0	0	0	0	1	1	1	2	2	2	2	3	3
4	.05	--	0	1	2	3	4	4	5	6	7	8	9	10	11	11	12	13	14
	.01	--	--	0	0	0	1	1	2	2	3	3	4	5	5	6	6	7	8
5	.05	0	1	2	3	5	6	7	8	9	11	12	13	14	15	17	18	19	20
	.01	--	--	0	1	1	2	3	4	5	6	7	7	8	9	10	11	12	13
6																			

n ₂	α	n ₁																	
		3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
	.05	1	2	3	5	6	8	10	11	13	14	16	17	19	21	22	24	25	27
	.01	--	0	1	2	3	4	5	6	7	9	10	11	12	13	15	16	17	18
7	.05	1	3	5	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34
	.01	--	0	1	3	4	6	7	9	10	12	13	15	16	18	19	21	22	24
8	.05	2	4	6	8	10	13	15	17	19	22	24	26	29	31	34	36	38	41
	.01	--	1	2	4	6	7	9	11	13	15	17	18	20	22	24	26	28	30
9	.05	2	4	7	10	12	15	17	20	23	26	28	31	34	37	39	42	45	48
	.01	0	1	3	5	7	9	11	13	16	18	20	22	24	27	29	31	33	36
10	.05	3	5	8	11	14	17	20	23	26	29	33	36	39	42	45	48	52	55
	.01	0	2	4	6	9	11	13	16	18	21	24	26	29	31	34	37	39	42
11	.05	3	6	9	13	16	19	23	26	30	33	37	40	44	47	51	55	58	62
	.01	0	2	5	7	10	13	16	18	21	24	27	30	33	36	39	42	45	48
12	.05	4	7	11	14	18	22	26	29	33	37	41	45	49	53	57	61	65	69
	.01	1	3	6	9	12	15	18	21	24	27	31	34	37	41	44	47	51	54
13	.05	4	8	12	16	20	24	28	33	37	41	45	50	54	59	63	67	72	76
	.01	1	3	7	10	13	17	20	24	27	31	34	38	42	45	49	53	56	60
14	.05	5	9	13	17	22	26	31	36	40	45	50	55	59	64	67	74	78	83
	.01	1	4	7	11	15	18	22	26	30	34	38	42	46	50	54	58	63	67
15	.05	5	10	14	19	24	29	34	39	44	49	54	59	64	70	75	80	85	90
	.01	2	5	8	12	16	20	24	29	33	37	42	46	51	55	60	64	69	73
16	.05	6	11	15	21	26	31	37	42	47	53	59	64	70	75	81	86	92	98
	.01	2	5	9	13	18	22	27	31	36	41	45	50	55	60	65	70	74	79
17	.05	6	11	17	22	28	34	39	45	51	57	63	67	75	81	87	93	99	105
	.01	2	6	10	15	19	24	29	34	39	44	49	54	60	65	70	75	81	86
18	.05	7	12	18	24	30	36	42	48	55	61	67	74	80	86	93	99	106	112
	.01	2	6	11	16	21	26	31	37	42	47	53	58	64	70	75	81	87	92
19	.05	7	13	19	25	32	38	45	52	58	65	72	78	85	92	99	106	113	119
	.01	3	7	12	17	22	28	33	39	45	51	56	63	69	74	81	87	93	99
20	.05	8	14	20	27	34	41	48	55	62	69	76	83	90	98	105	112	119	127
	.01	3	8	13	18	24	30	36	42	48	54	60	67	73	79	86	92	99	105

Appendix B

DF	Chi-Square Right-Tail Probability (χ^2)									
	0.995	0.99	0.975	0.95	0.9	0.1	0.05	0.025	0.01	0.005
1	---	---	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838

DF	Chi-Square Right-Tail Probability (χ^2)									
	0.995	0.99	0.975	0.95	0.9	0.1	0.05	0.025	0.01	0.005
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.070	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.300
13	3.565	4.107	5.009	5.892	7.042	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.408	7.564	8.672	10.085	24.769	27.587	30.191	33.409	35.718
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.844	7.633	8.907	10.117	11.651	27.204	30.144	32.852	36.191	38.582
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.034	8.897	10.283	11.591	13.240	29.615	32.671	35.479	38.932	41.401
22	8.643	9.542	10.982	12.338	14.041	30.813	33.924	36.781	40.289	42.796
23	9.260	10.196	11.689	13.091	14.848	32.007	35.172	38.076	41.638	44.181
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.559
25	10.520	11.524	13.120	14.611	16.473	34.382	37.652	40.646	44.314	46.928
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.808	12.879	14.573	16.151	18.114	36.741	40.113	43.195	46.963	49.645
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.993
29	13.121	14.256	16.047	17.708	19.768	39.087	42.557	45.722	49.588	52.336
30	13.787	14.953	16.791	18.493	20.599	40.256	43.773	46.979	50.892	53.672
40	20.707	22.164	24.433	26.509	29.051	51.805	55.758	59.342	63.691	66.766
50	27.991	29.707	32.357	34.764	37.689	63.167	67.505	71.420	76.154	79.490
60	35.534	37.485	40.482	43.188	46.459	74.397	79.082	83.298	88.379	91.952
70	43.275	45.442	48.758	51.739	55.329	85.527	90.531	95.023	100.425	104.215
80	51.172	53.540	57.153	60.391	64.278	96.578	101.879	106.629	112.329	116.321
90	59.196	61.754	65.647	69.126	73.291	107.565	113.145	118.136	124.116	128.299
100	67.328	70.065	74.222	77.929	82.358	118.498	124.342	129.561	135.807	140.169

References

1. Fadila M, Rajasurya V, Regunath H. Ventilator Weaning. 2020 Nov 28. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing (2021).

2. Llamas-Álvarez AM, Tenza-Lozano EM, Latour-Pérez J. Diaphragm and Lung Ultrasound to Predict Weaning Outcome: Systematic Review and Meta-Analysis. *Chest* 152 (2021): 1140–1150.
3. Qian Z, Yang M, Li L, et al. Ultrasound assessment of diaphragmatic dysfunction as a predictor of weaning outcome from mechanical ventilation: a systematic review and meta-analysis. *BMJ Open* 8 (2018): e021189.
4. Li C, Li X, Han H, et al. Diaphragmatic ultrasonography for predicting ventilator weaning: A meta-analysis. *Medicine (Baltimore)* 97 (2018): e10968. [PubMed: 29851847]
5. Pham T, Brochard LJ, Slutsky AS. Mechanical Ventilation: State of the Art. *Mayo Clin Proc* 92 (2017): 1382–1400. [PubMed: 28870355]
6. Baptistella AR, Sarmiento FJ, da Silva KR, et al. Predictive factors of weaning from mechanical ventilation and extubation outcome: A systematic review. *J Crit Care* 48 (2018): 56–62. [PubMed: 30172034]
7. Wawrzeniak IC, Regina Rios Vieira S, Almeida Victorino J. Weaning from Mechanical Ventilation in ARDS: Aspects to Think about for Better Understanding, Evaluation, and Management. *Biomed Res Int* 2018 (2018): 5423639. [PubMed: 30402484]
8. Vetrugno L, Guadagnin GM, Brussa A, et al. Mechanical ventilation weaning issues can be counted on the fingers of just one hand: part 1. *Ultrasound J* 12 (2020): 9. [PubMed: 32166566]
9. Matamis D, et al. Sonographic evaluation of the diaphragm in critically ill patients. Technique and clinical applications. *Intensive Care Med* 39 (2013): 801–810. [PubMed: 23344830]
10. de la Quintana Gordon FB, Nacarino Alcorta B, Fajardo Pérez M. Ultrasound evaluation of diaphragm function and its application in critical patients, mechanical ventilation, and brachial plexus block. *Rev Esp Anesthesiol Reanim* 64 (2017): 513–521. [PubMed: 28343681]
11. Kilaru D, Panebianco N, Baston C. Diaphragm Ultrasound in Weaning From Mechanical Ventilation. *Chest* 159 (2021): 1166–1172. [PubMed: 33309837]
12. Dres M, Goligher EC, Heunks LMA, et al. Critical illness-associated diaphragm weakness. *Intensive Care Med* 43 (2017): 1441–1452. [PubMed: 28917004]
13. Carbognin S, Pelacani L, Zannoni F, et al. P246 Diaphragm ultrasonography: a method for weaning patients from mechanical ventilation. 36th International Symposium on Intensive Care and Emergency Medicine: Brussels, Belgium. 15–18 March 2016. *Crit Care* (2016).




Figure 1:
Parameters Measured in the Diaphragm Kinetics.




Figure 2: US study, B-mode shows the diaphragm (1) next to the liver (2). Identification of a point of the diaphragm (3), of which the kinetics is studied in the figure below (4). The space has been measured (on the left) and identified in the figure (5), where we can see the time on the x-axis.




Figure 3:
Dot plot representing the ranks and the values of acceleration associated, failed in blue, successful in red.

Characteristics of the enrolled patients, including sex, age, date of the SARS-CoV2 test positivity, date of admission in the emergency department and date of admission in the ICU.

Table 1:

Patient n.	Sex	Date of birth	Date of the SARS-CoV2 test positivity	Date of admission in the emergency department	Date of admission in the ICU
1	M	01/01/1953	17/12/2020	17/12/2020	25/12/2020
2	M	22/07/1943	09/04/2020	07/04/2020	08/04/2020
3	M	18/07/1966	08/01/2021	13/01/2021	13/01/2021
4	M	30/11/1958	13/11/2020	24/11/2020	11/12/2020
5	M	24/11/1955	30/03/2020	30/03/2020	03/04/2020
6	M	18/01/1953	11/04/2020	09/04/2020	14/04/2020
7	M	27/02/1967	13/11/2020	13/11/2020	13/11/2020
8	M	06/02/1950	23/03/2020	13/03/2020	23/03/2020
9	M	11/09/1959	26/03/2021	26/03/2021	26/03/2021
10	M	22/03/1946	02/04/2021	02/04/2021	02/04/2021
11	M	22/03/1955	25/03/2021	27/03/2021	03/04/2021
12	M	26/01/1944	18/03/2020	21/03/2020	25/03/2020
13	M	04/04/1942	27/10/2020	05/11/2020	13/11/2020
14	F	22/08/1958	11/12/2020	18/12/2020	21/12/2020
15	F	12/08/1958	27/02/2021	27/02/2021	02/03/2021
16	M	29/11/1961	-	01/04/2020	04/04/2020
17	F	13/10/1946	23/03/2021	26/03/2021	28/03/2021
18	M	21/02/1950	19/03/2021	29/03/2021	06/04/2021
19	M	25/11/1957	-	20/03/2021	04/04/2021

Table 2:

Date of the US test with the applied ventilation mode, date of the extubation with the associated ventilation mode, and time of intubation.

Patient n.	Date of the US test	Ventilation mode during the test	Date of extubation	Ventilation mode at the extubation/death	Time of intubation (days)
1	27/01/2021	PSV 6+5; FiO ₂ 0,35	30/01/2021	PSV 8+6, FiO ₂ =0,35	38
2	24/04/2020	PSV 5+5; FiO ₂ 0,45	24/04/2020	PSV 5+5, FiO ₂ =0,45	25
3	04/02/2021	SIMV/P: P _{insp} =10, PEEP=6, FR=12/min FiO ₂ =0,65	05/02/2020	PSV 8+6, FiO ₂ = 0,5	32
4	08/12/2020	PSV 10+10, FiO ₂ =0,4	08/12/2020	PSV 10+5, FiO ₂ =0,4	14
5	17/04/2020	PSV 8+6, FiO ₂ =0,45	19/04/2021	PSV 10+5, FiO ₂ =0,4	16
6	30/04/2020	PSV 8+5, FiO ₂ =0,5	03/05/2020	PSV 8+8, FiO ₂ =0,4	19
7	21/11/2020	PSV 10+8, FiO ₂ =0,5	21/11/2020	PSV 10+8, FiO ₂ =0,5	12
8	20/04/2020	PSV 5+5, FiO ₂ =0,45	20/04/2020	C-PAP 5+5, FiO ₂ =0,35	27
9	13/04/2021	SIMV: (20+10), FiO ₂ =0,6	Dead	PCV: P _{insp} =20, RR=22/min, PEEP=10, FiO ₂ =0,5	29; dead
10	12/04/2201	SIMV-P: P _{insp} =14, RR= 15, PEEP=8, FiO ₂ =0,5	tracheostomy in SB from 21/04/2021	PSV 10+5, FiO ₂ =0,4	19
11	13/04/2021	PCV: FiO ₂ =0,7 P _{insp} =26 RR=22/ min, PEEP=10	Dead	PCV: P _{insp} =26, RR=26/min, PEEP=5, FiO ₂ =1	13; dead
12	30/04/2020	PSV 10+10, FiO ₂ = 0,4	Dead	PSV 15+5, FiO ₂ =0,5	46; dead
13	26/11/2020	PSV 10+8, FiO ₂ =0,4	26/11/2020	PSV 10+8, i FiO ₂ =0,4	13
14	16/01/2021	NAVA 2.5, FiO ₂ =0,8	Dead	PCV: P _{insp} =27, RR=26/min, PEEP=14, FiO ₂ =0,9	37; Dead
15	15/03/2021	SIMV-P: P _{insp} =14, PEEP=10, FiO ₂ =0,5	17/03/2021	C-PAP 8+8; FiO ₂ =0,4	15
16	18/04/2020	PSV 18+8, FiO ₂ =0,5	19/04/2020	PSV 18+8, FiO ₂ =0,5	15
17	17/04/2021	PCV: P _{insp} =28, RR=26/min, PEEP=10, FiO ₂ =0,95	Dead	PCV: P _{insp} =30, RR=20/min, PEEP 8, FiO ₂ =0,9	21; Dead
18	17/04/2021	PCV: P _{insp} =18 RR=20/min, PEEP=12, FiO ₂ =0,9	Dead	PCV: P _{insp} =20, RR=22/min, PEEP=10, FiO ₂ =0,9	24; Dead
19	17/04/2021	PCV: P _{insp} =26, RR=26/min, PEEP=10, FiO ₂ =0,9	Dead	PCV: P _{insp} =30, RR=20/min, PEEP=8, FiO ₂ =0,9	15; Dead

Table 3:

values of diaphragmatic acceleration of each patient, associated with the outcome of weaning.

Patient n.	Space (S) mm	Time (t)s	Acceleration (a) mm/s ²	Outcome
1	15,5	0,504	122,0	Successful
2	55,2	1,376	58,3	Successful
3	18,4	0,352	52,0	Successful
4	27,2	0,814	81,7	Successful
5	32,8	0,778	108,4	Successful
6	31,8	0,736	117,0	Successful
7	40,3	0,800	125,9	Successful
8	48,7	0,869	128,9	Successful
9	3,7	1,000	7,4	Failed
10	20,6	1,112	33,0	Failed
11	9,6	1,144	14,7	Failed
12	44,9	1,024	85,6	Successful
13	21,5	1,000	43,0	Failed
14	5,63	0,347	93,5	Successful
15	20,6	1,080	35,3	Failed
16	31,8	0,729	101,4	Successful
17	30,0	1,080	51,0	Failed
18	17,6	0,984	36,3	Failed
19	10,1	0,976	21,0	Failed

Table 4:

Ranks of Acceleration Values.

Failed	Successful	RANK
7,4		1
14,7		2
21,0		3
33,0		4
35,3		5
36,3		6
43,0		7
51,0		8
	52,0	9
	58,3	10
	81,0	11
	85,6	12
	93,5	13
	101,4	14
	108,4	15
	117,0	16
	122,0	17
	125,9	18
	128,9	19

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 5:

Observed Values.

	weaning failure	successful weaning		Total in the line
	0	11	a>52 mm/s ²	11
	8	0	a<52 mm/s ²	8
Total in the column	8	11		Total = 19

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 6:

Expected Values.

	weaning failure	successful weaning		Total in the line
	5	6	a>52 mm/s ²	11
	3	5	a<52 mm/s ²	8
Total in the column	8	11		Total = 19

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript