

LETTER

A pilot study of semiquantitative treatment evaluation following nonthermal atmospheric-pressure plasma administration for onychomycosis

Various methods have been used to treat onychomycosis, including application of topical and systemic antifungal agents and laser therapy. In current standard systemic treatment, the complete cure rate of toe nails is 38% with terbinafine and 14% with itraconazole.¹ In addition, recurrences (i.e., relapse [same infection after incomplete cure] or reinfection [same infection after complete cure]) occur at a rate of 20–25%.¹ Near-infrared diode lasers, 1064 nm neodymium-doped yttrium aluminum garnet (Nd:YAG) lasers, dual diode lasers, 1320 nm Nd:YAG lasers, and fractional CO₂ lasers have all been used for the treatment of onychomycosis.² A recent review reported that 1064 nm Nd:YAG laser treatment resulted in lower mycologic cure rates (11%) than oral and topical therapies approved by the Food and Drug Administration (29–61%).² Thus, an effective and safe local target therapy for onychomycosis is needed. Plasma lasers have antibacterial and antifungal effects; these lasers are generated in the air by a strong electric field pulse generated by the ionization of air molecules into ozone, hydroxyl radicals, and nitric oxide, which exert antifungal properties.³ In previous studies, nonthermal atmospheric-pressure plasma (NTAP) inhibited the growth of *Trichophyton rubrum* in vitro,⁴ achieving an overall clinical cure rate for infected toenails of 53.8%, and a mycological cure rate of 15.4% in an in vivo pilot study.⁵

This study aimed to evaluate the feasibility of NTAP as a treatment option for onychomycosis. A total of five participants (four females and one male, mean age 46.4) with toenail onychomycosis provided informed consent to participate. Reflectance confocal microscopy

(RCM, VivaScope® 1500; Lucid Inc., Rochester, NY, USA) was performed on the infected toenails before and immediately after NTAP (PLADUO; SHENB, Seoul, Korea), which generated plasma from argon and nitrogen gas sources, delivering pulses to the skin. The NTAP was applied to the affected nail plate with pulse energy 1.5 J, frequency 3 Hz of nitrogen gas and followed by 0.75 J, 3 Hz of argon gas, three passes, respectively.

The number of fungal spores and hyphae was counted in each 500 × 500 μm² RCM image. To evaluate fungal spore dispersion, the RCM images were divided in a grid pattern. Wilcoxon signed-rank test was then applied to compare the total number and dispersion effect of spores before and after NTAP treatment. Statistical significance was set at $p < 0.05$.

Before NTAP treatment, a spherical spore surrounded by a bright visible wall was observed in the RCM image; after NTAP treatment, cell wall disruption and fungal hyphae breakage were observed (Figure 1). The size of fungal spores is reported to range from 2 to 50 μm in diameter.⁶ We measured the size of the fungal spores to be 2–8 μm using Image J. The spores were clustered; however, immediately after treatment, the spores became more disperse ($p = 0.5$), and the total number of spores decreased from 216 to 138 ($p = 0.31$) (Figure 2), although this difference was not statistically significant. On the RCM image, crack-like changes in the nail plate were seen after treatment, which was thought to indicate a synergistic effect when combined with other treatments such as topical antifungal agent (Figure 3).




FIGURE 1 A hyphae with an intact cell wall was observed on reflectance confocal microscopy (RCM) image. After plasma treatment, disruption of the cell wall and broken hyphae (red arrow) were observed

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Skin Research and Technology* published by John Wiley & Sons Ltd.




FIGURE 2 An RCM image ($500 \times 500 \mu\text{m}^2$) with grid pattern showing that clusters of fungal spores (A) were dispersed (B) following plasma treatment. The mean number of fungal spores of all participants were counted, and the grid pattern showed a decreased density after plasma treatment (C and D)




FIGURE 3 On the RCM image, reflectance was increased and fine lines were clearly visible, reflecting the nail plate changes immediately after treatment

Biofilms are structured sessile microbial communities comprising microorganisms adhering to a surface and each other via an extracellular polymeric matrix.⁷ Similar to bacteria, fungi are also able to form biofilms; *T. rubrum* and *T. mentagrophyte* biofilms have been previously reported.⁸ The adhesion of fungal spores to surfaces occurs readily and is a prerequisite for biofilm formation.⁹ Recent studies have reported the positive effect of plasma on fungal biofilm inactivation.^{10,11} Our prior in vitro study showed destruction of colonies of *T. rubrum* with waxy surface after NTAP irradiation and scanning electron microscopy, confirming that the exopolymeric matrix of the biofilm⁸ disappeared (Figure S1). Despite the small sample size, this study revealed that NTAP treatment disrupted the cell wall and decreased the number and dispersed clusters of spores. These results suggest that NTAP treatment could exert preventive effects against fungal biofilms.

There were no serious side effects during NTAP irradiation, except for a minor heating sensation. To reduce this unpleasant sensation, it is necessary to keep moving the hand-piece, instead of fixing it

at one point. Although the effect may be weak as a single treatment, we suggest that irradiating onychomycosis with NTAP prior to topical antifungal or laser treatment could enhance the therapeutic effect.

ACKNOWLEDGMENT

This research was supported by a Paul Janssen grant from the Korean Dermatological Association, Republic of Korea, in 2022.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest.

ETHICS STATEMENT

Informed consents were obtained from all participants.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Hye-Jin Ahn¹

Tae-Eun Kim¹

Ye-Jin Lee¹

Su Jin Jeong²

Min Kyung Shin¹

Ki-Heon Jeong¹

¹Department of Dermatology, Kyung Hee University Medical Center, Seoul, South Korea

²Medical Science Research Institute, Kyung Hee University Medical Center, Seoul, South Korea

Correspondence

Ki-Heon Jeong, Department of Dermatology, College of Medicine,
Kyung Hee University, # Kyung HeeDae Ro 23, Dongdaemun-gu, Seoul
02447, South Korea.
Email: mdfamily@hanmail.net

Additional supporting information can be found online in the
Supporting Information section at the end of this article.

ORCID

Hye-Jin Ahn  <https://orcid.org/0000-0002-0085-9566>

Ye-Jin Lee  <https://orcid.org/0000-0003-0460-6272>

REFERENCES

1. Lipner SR, Scher RK. Onychomycosis: treatment and prevention of recurrence. *J Am Acad Dermatol*. 2019;80:853-67.
2. Gupta AK, Versteeg SG. A critical review of improvement rates for laser therapy used to treat toenail onychomycosis. *J Eur Acad Dermatol Venereol*. 2017;31:1111-8.
3. Ouf SA, El-Adly AA, Mohamed AH. Inhibitory effect of silver nanoparticles mediated by atmospheric pressure air cold plasma jet against dermatophyte fungi. *J Med Microbiol*. 2015;64:1151-61.
4. Heinlin J, Maisch T, Zimmermann JL, et al. Contact-free inactivation of *Trichophyton rubrum* and *Microsporum canis* by cold atmospheric plasma treatment. *Fut Microbiol*. 2013;8:1097-106.
5. Lipner SR, Friedman G, Scher RK. Pilot study to evaluate a plasma device for the treatment of onychomycosis. *Clin Exp Dermatol*. 2017;42:295-8.
6. Di Filippo PD, Pomata D, Riccardi C, Buiarelli F, Perrino C. Fungal contribution to size-segregated aerosol measured through biomarkers. *Atmos Environ*. 2013;64:132-40.
7. Heydorn A, Ersbøll BK, Hentzer M, Parsek MR, Givskov M, Molin S. Experimental reproducibility in flow-chamber biofilms. *Microbiology (Reading)*. 2000;146:2409-15.
8. Costa-Orlandi CB, Sardi JC, Santos CT, Fusco-Almeida AM, Mendes-Giannini MJ. In vitro characterization of *Trichophyton rubrum* and *T. mentagrophytes* biofilms. *Biofouling*. 2014;30:719-27.
9. Liauw CM, Slate AJ, Butler JA et al. The effect of surface hydrophobicity on the attachment of fungal conidia to substrates of polyvinyl acetate and polyvinyl alcohol. *J Polym Environ*. 2020;28:1450-64.
10. Los A, Ziuzina D, Boehm D, Cullen PJ, Bourke P. Inactivation efficacies and mechanisms of gas plasma and plasma-activated water against *Aspergillus flavus* spores and biofilms: a comparative study. *Appl Environ Microbiol*. 2020;86:e02619-19.
11. He M, Duan J, Xu J, et al. *Candida albicans* biofilm inactivated by cold plasma treatment in vitro and in vivo. *Plasma Process Polym*. 2020;17:1900068.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.