




Artificial Tears: A Systematic Review

David A Semp , Danielle Beeson , Amy L Sheppard, Debarun Dutta, James S Wolffsohn 

School of Optometry, College of Health and Life Sciences, Aston University, Birmingham, UK

Correspondence: James S Wolffsohn, Tel +44 121 2044140, Email j.s.wolffsohn@aston.ac.uk

Abstract: Artificial tears are the mainstay of dry eye disease management, but also have a role in corneal abrasion and wound healing, pain and inflammation management, conjunctivitis, keratitis, contact lens rewetting and removal, and foreign body removal. A systematic review of randomized controlled trials (PROSPERO registration CRD42022369619) comparing the efficacy of artificial tears in patients with dry eye to inform prescribing choices using Web of Science, PubMed and Medline databases identified 64 relevant articles. There is good evidence that artificial tears improve symptoms of dry eye disease within a month of regular use, applied about four times a day, but signs generally take several months to improve. Not all patients with dry eye disease benefit from artificial tears, so if there is no benefit over a month, alternative management should be considered. Combination formulations are more effective than single active ingredient artificial tears. Artificial tears containing polyethylene glycol are more effective than those containing carboxymethylcellulose/carmellose sodium and hydroxypropyl methylcellulose. Those classified as having evaporative dry eye disease, benefit from artificial tears with liposomes, especially of higher concentration. The data available is limited by the definition of dry eye disease applied in published studies being variable, as well as the disease severity examined and compliance with artificial tears being rarely quantified.

Keywords: artificial tears, dry eye, comfort, contact lenses

Artificial tear drops are most commonly associated with the management of dry eye disease (DED). Artificial tears are typically included in first-line management options for dry eye, as they are easy to use, accessible in a wide range of formulations, and have a low risk-profile.¹ Most artificial tear preparations have been found to be effective in reducing the symptoms and signs of DED, however the Tear Film and Ocular Surface Society (TFOS) dry eye workshop in 2017 (DEWS II) concluded there had been relatively few high quality randomized controlled trials comparing different formulations with each other.^{1,2} Furthermore, few clinical trials have compared the efficacy of different artificial tear products, and attempted to correlate this with patient characteristics, in order to aid management decisions for an individual.^{3,4} The issue with this is that both practitioners and patients are faced with a bewildering array of different products with varying ingredients, and little or no clear way of knowing which is the most effective. Practitioners will often be asked “which is the best drop for dry eye”, but with no scientific evidence to base their answer on. In addition, other aspects that influence practitioner and patient choices are:

- formulation
 - percentage concentration⁵
 - molecular weight⁵
 - preservative used⁶
- storage bottle design.^{7–10}

Patients may therefore face a trial-and-error approach to product selection, incurring mounting costs and frustration in the process. This will be felt even more keenly by patients who are highly price sensitive, since over-the-counter products are no longer easily available via National Health Service (NHS) subsidised prescriptions¹¹ in the UK. A recent study¹² on the reported experience of dry eye management across four continents identified that on average, DED still caused a

moderate impact on an individual's quality of life (median impact 3/10); less than half of the individuals in any country had undergone a consultation with an eye or health-care practitioner about their dry eye; about half had tried dry eye treatment, with artificial tears being the most common treatment, followed by warm compresses, and both therapies were rated as reasonably effective (median 5–7/10).

Formulation

The majority of artificial tear products are aqueous-based and contain viscosity-enhancing agents, such as carbomer 940, carboxymethyl cellulose (CMC), dextran, hyaluronic acid, sodium hyaluronate (which has a smaller molecular size), hydroxypropyl guar (HP-guar), hydroxypropyl methylcellulose (HPMC hypromellose), polyvinyl alcohol, polyvinylpyrrolidone and polyethylene glycol, which aid lubrication and increase on-eye retention time.¹ Other ingredients may include osmotic agents, osmoprotectants, antioxidants, preservatives and inactives such as pH buffers, excipients and electrolytes.¹ Aqueous-based artificial tears target principally the muco-aqueous phase of the tear film, but have been shown to improve dry eye symptoms related to all subtypes of DED.²

In recent years, there has been an increase in the popularity and availability of lipid-based drops, which target the superficial tear lipid layer^{13,14} as the emphasis on meibomian gland dysfunction and its role in evaporative dry eye continues to increase.¹ It has been demonstrated in randomised controlled trials that lipid-based drops are more effective at managing DED classified as evaporative.^{3,4} These can take the form of nano-emulsion drops or liposomal sprays, which are applied to the closed eye and may be easier for those who struggle to instil drops, for example those with reduced manual dexterity or hand tremor. A completely water-free drop comprised of 100% lipid (perfluorohexyloctane) is available now, with the added benefit of being preservative-free.¹⁵

Preservatives

Multidose eye drops, including artificial and medicated topical ocular drops, commonly contain preservatives to maintain sterility and prolong shelf life, however, these are also known to produce toxicity. Benzalkonium chloride, commonly found in multidose drops, can produce toxic, proinflammatory and detergent effects, which may actually lead to or exacerbate DED.¹⁶ For this reason, there has been a move towards preservative-free and unit dose formulations, due to the risk of toxic and allergic reactions, especially when frequent instillation is required. Newer preparations may contain less damaging preservatives such as polyquaternium, or “vanishing” preservatives such as sodium perborate and purite, or feature specially designed bottles, which prevent the entry of microorganisms.¹⁷ Preservative-free formulations are recommended for all types of dry eye, however this is even more important for severe dry eye or sensitive individuals, and more details can be found in the TFOS DEWS II iatrogenic report.⁶

Ideal Properties

It is important that artificial tear drops behave in a similar way to natural tears. One aspect of this is the physical property of rheology, which refers to the way fluids and soft solids flow. The viscosity of human tears is high between blinks, but reduces during each blink cycle in order to protect the ocular surface from damage due to fluid turbulence.¹ Hence, they do not display Newtonian behaviours and are referred to as having non-Newtonian properties. Hyaluronic acid has been the subject of a significant amount of research and has been shown to exhibit these non-Newtonian shear-thinning properties,¹⁸ making it more like the tear film and hence suitable for use in artificial tears.¹⁹ Hyaluronic acid, a common constituent of artificial tears, is a naturally occurring glycosaminoglycan, which is found in and around body cells and tissues, for example in synovial fluid, and vitreous and aqueous humour.²⁰ Its use in ophthalmology was pioneered by Andre Balazs in the late 1960s,²¹ with Polack and McNiece²² being the first to report its use in dry eye. Hyaluronic acid is water soluble and is capable of binding large quantities of water, compared to its own weight, but its physical properties vary depending upon its molecular weight.²³ There is evidence to suggest that high molecular weight hyaluronic acid (HMWHA) is clinically superior in the treatment of DED compared to its low molecular weight counterpart.²⁴ Furthermore, HMWHA has been found to be protective against corneal cell apoptosis due to benzalkonium chloride toxicity, ultraviolet light radiation and chemical burns,^{25–27} as well as being anti-inflammatory and having a role in reducing pain sensation.^{24,28}

Artificial Tears for Dry Eye Disease

There have been several systematic reviews^{2,29–31} conducted over the past decade, concluding that artificial tears are a safe and effective way of treating DED. A meta-analysis concluded that the effectiveness of sodium hyaluronate did not differ based on its preparation³⁰ and another³² suggested that CMC appeared to be better than hyaluronic acid in treating DED, but the results were not statistically significant. Two recent reviews^{5,33} both identified that while hyaluronic acid was effective in reducing the symptoms of DED, the ideal drop frequency and formulation (both concentration and molecular weight) for different ages and severities were yet to be investigated.

To date, there has been no review of studies which compared different artificial tears to identify whether certain formulations are more effective. Hence, with the objective to better understand the evidence for the effect of different artificial tears in managing dry eye, a search was made of the Web of Sciences databases (Clarivate Analytics, Philadelphia, USA) which includes the Science Citation Index Expanded covering over 9200 of the world's most impactful journals from 1900 to the present day along with PubMed (including MEDLINE) from its inception. The systematic review was prospectively registered on PROSPERO (CRD42022369619) and was conducted in the format prescribed by PRISMA (2020).³⁴ A search for “artificial tear*” AND “randomi?ed” identified 481 unique results which were screened independently by two researchers (DB and DS) and verified by a third (JSW). Studies were eligible to be accepted if they were in full paper form (not abstracts or book chapters), compared two or more artificial tears against each other (not just with a placebo) and involved randomisation to avoid bias. This resulted in 64 papers being accepted (Figure 1) and the full text scrutinized for the key factors, which were tabulated in a spreadsheet and are summarised in Table 1. The study design, artificial tears compared, number and age profile of participants completing the trial, duration of use and dosing, tests conducted which showed a significant difference/did not differentiate between the products or change from baseline and general comments (dyes used for ocular surface staining, adverse events when reported and subanalyses) were extracted. Missing information is highlighted in the table and risk of bias analysis performed with the Cochrane Tool reported.³⁵ No data synthesis was attempted due to heterogeneity particularly in drop duration.

All studies are prospective (as expected) and involve parallel groups (unless stated otherwise) of dry eye patients (diagnosed using National Eye Institute, arbitrary or recently TFOS DEWS II criteria). However, less than half (20 out of 42) are registered with a clinical trials database and even those that are have high risk of bias characteristics,³⁵ hence the certainty of the result is generally low. The lack of a definitive severity classification has been identified as a factor in differentiating the effectiveness of the available artificial tears,³¹ but previous attempts at a severity matrix table in TFOS DEWS I³⁶ led to patients being graded at different levels of severity by different tests and was abandoned in TFOS DEWS II,³⁷ severity to a dry eye patient is based on symptoms whereas it is more likely to be based on signs on the ocular surface to a cataract surgeon for example. While the intention of many of the analysed studies is to demonstrate non-inferiority compared to an established treatment, some are underpowered (see TFOS sample size recommendations)-³⁷ and/or both eyes included without accounting for the correlation between the two eyes³⁸ of an individual.^{39–43} In most studies, fluorescein sodium is used for assessing corneal staining (although an appropriate blue light with a peak around 395nm [not cobalt blue whose peak is ~450nm] and yellow filter with a cut off around 500nm is often not stated).⁴⁴ Most studies use lissamine green for conjunctival staining (unless otherwise stated in Table 1) which is the recommended practice,³⁷ but few state the brand which can dramatically affect the staining observed.⁴⁵ Some studies^{46,47} report differences even when they do not meet the standard criteria of $p < 0.05$ and therefore any “difference” should be considered as noise in the data. While many trials comparing artificial tears are manufacturer initiated or sponsored, unless the research was conducted by the company or not conducted by a reputable research organisation, this should not lead to concerns regarding bias.

From the studies summarised to date (with the caveat that the effects might be affected by dry eye severity and full artificial tear formulation as well as the patient demographic) it would appear from direct comparisons between artificial tears that:

- Combination formulations are more effective than single active ingredient artificial tears.
 - The combination of CMC with hyaluronic acid is more effective than either in isolation.^{48,49}




Figure 1 PRISMA 2020 flow diagram of the systematic review search results.

Notes: PRISMA figure adapted from Liberati A, Altman D, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Journal of clinical epidemiology*. 2009;62(10). Creative Commons.¹²⁷

- Hyaluronic acid⁵⁰ and sodium hyaluronate⁵¹ benefit from the addition of trehalose.
- CMC is enhanced by the addition of glycerine.⁵²
- CoQ10 enhances the effectiveness of hyaluronic acid.⁵³
- Newer versions of Systane (Complete and Balance) outperform earlier versions with less complexity (Ultra).^{54,55}
- Some studies suggest sodium hyaluronate could be more effective than CMC⁴⁰ and carbomers,⁵⁶ while others find no difference,^{57,58} the optimal percentage is not clear.^{59,60}
- PEG containing artificial tears are more effective than those containing CMC^{61–65} and HPMC.^{66,67}
- Cationic formulations are more effective than sodium hyaluronate (for objective signs)⁶⁸ and polyvinyl alcohol.⁶⁹
- Hyaluronic acid containing artificial tears might be better than those with HPMC,⁷⁰ but worse than those with CMC.³⁹
- Carbomer containing artificial tears might be more effective than those based on PVA⁷¹ or CMC⁷² or cellulose/mineral oils,⁷³ but less^{56,74} or as effective⁴³ as sodium hyaluronate.
- Most studies recommend 4x/day use, but reported/measured use is generally less than that advised.⁴²

Table I Randomised Clinical Trials That Have Used Artificial Tears for the Treatment of Dry Eye Disease

Paper	Design	Comparators	Participants Completing	Age (Years)	Duration (Dosing)	Tests Showing Significant Difference	Tests Not Differentiating Products	Tests Showing Significant Difference	Tests Showing no Change	General Comments
						Cross Comparator		Compared to Baseline		
Amran et al 2014 ⁶⁹	Randomized, open-label, multi-center study	Cationic Emulsion - Cationorm PVA-Povidone - Refresh	N = 44 N = 35	61.3 ± 15.4 61.9 ± 12.5	4 weeks (4x/day)	Symptoms, TBUT, eyelid erythema, Conjunctival staining with Cationic	Schirmer's Corneal staining			Sub-analysis with MGD participants CLINICAL TRIAL NOT REGISTERED
Aragona et al 2020 ⁴⁸	Randomized, double-masked, multi-center study	CMC + HA - Optive Fusion UD CMC - Refresh Optive Sensitive/Optive UD	N = 180 N = 184	59.4 ± 13.8 57.5 ± 13.7	90 days (2x/day)	Lower ocular pain/discomfort CMC-HA	OSDI TBUT Ocular surface staining Schirmer's II	OSDI Symptoms (VAS) TBUT Ocular surface staining	Schirmer's II	10% minor AE CLINICAL TRIAL NOT REGISTERED
Baeyens et al 2012 ⁷⁴	Randomized, double-masked, multi-center study	SH 0.18% - Vismed Carbomer 0.3% NaCL	N = 100 N = 96 N = 96	59.3 ± 15.0 (across groups)	84 days	Symptoms & Corneal staining with SH vs saline	Symptoms, corneal and conjunctival staining, Schirmer's, TBUT SH vs carbomer	Symptoms Corneal staining	Conjunctival staining, Schirmer's, TBUT	CLINICAL TRIAL NOT REGISTERED
Barabino et al 2014 ¹⁰⁴	Randomized, double-masked, multi-center study	CMC 0.5% / glycerin 0.9% - Optive HA 0.2% / tamarind seed polysaccharide 0.2% - Xilolial	N = 25 N = 23	57.1 ± 17.4 52.2 ± 14.9	3 months (4x/day)	Symptoms with HA+TS	TBUT Ocular surface staining Schirmer's	OSDI TBUT Ocular surface staining	Schirmer's	CLINICAL TRIAL NOT REGISTERED
Baudouin et al 2012 ⁵⁷	Randomized, investigator-masked, multi-center study	CMC 0.5% and Osmoprotectant - Optive SH 0.18% - Vismed Multi	N = 37 N = 29	58.1 ± 14.2 55.4 ± 13.4	3 months	None	Osmolarity, Schirmer's-I, OSDI, staining	Osmolarity, Schirmer's-I, OSDI, staining	None	Clinical Trial NCT00987727 - only symptom primary and secondary outcomes and day 35 data missing Cochrane Risk of Bias R? C?M-O?I-S-B?
Benelli et al 2010 ⁶²	Randomized, investigator-masked, single-center study	CMC 0.5% - Cellufresh PEG 400 2.5% - Blink Intensive HP-guar 0.18%/PEG 400/PG - Systane	N = 20 N = 20 N = 20	Not stated	30 days (up to 4x/day)	Osmolarity with PEG400	VA Aberrometry Staining TBUT Schirmer's	Aberrometry	Osmolarity VA Staining TBUT Schirmer's	CLINICAL TRIAL NOT REGISTERED
Brignole et al 2005 ⁴⁰	Randomized, masked-observer, single-center study	CMC 1% - Celluvisc SH 0.18% - Vismed	N = 11 N = 10	69 ± 2 57 ± 2	2 months (3x/day)	CD44, comfort (only at day 7), keratitis recovery with SH	All other inflammatory markers, cornea and conjunctival staining, TBUT, corneal topography, tear meniscus height	Symptoms and ocular surface staining	None	Moderate dry eye and keratitis patients CLINICAL TRIAL NOT REGISTERED
Brodwall et al 1997 ¹⁰⁵	Randomized, investigator-masked, single-center study	Polyacrylic acid 0.2% - Visco Tears PVA 1.4%	N = 38 N = 41	60.2 61.8	4 weeks (Drops/day a study variable; avg 3-5)	Symptoms (16/27 study days), hyperaemia, Rose Bengal staining, compliance with polyacrylic acid	TBUT Schirmer's	Symptoms & signs (unspecified)	TBUT Schirmer's	CLINICAL TRIAL NOT REGISTERED
Bron et al 1998 ¹⁰⁶	Randomized, double masked, multi-center study	Carbomer 940 0.2% - Lacrinorm/GelTears, Laboratoire Chauvin Carbomer 940 0.2% - Viscotears/Vidiscil/ Lacrigel	N = 92 N = 87	58.6 ± 16.2 64.0 ± 14.0	4 weeks (4x/day)	None	Symptoms TBUT, Schirmer's, corneal and conjunctival staining	Symptoms TBUT, Schirmer's, fluorescein/ lissamine green staining	None	AEs in n=21 Lacrinorm group and 17 Viscotears group CLINICAL TRIAL NOT REGISTERED

(Continued)

Table I (Continued).

Paper	Design	Comparators	Participants Completing	Age (Years)	Duration (Dosing)	Tests Showing Significant Difference	Tests Not Differentiating Products	Tests Showing Significant Difference	Tests Showing no Change	General Comments
						Cross Comparator		Compared to Baseline		
Calvao-Santos et al 2011 ⁴¹	Randomized, open-label, single-center study	Tears Again [lipidic] Opticol [aqueous] Optive [mucin] No treatment	N = 7 N = 6 N = 7 N = 7	24 to 53 years	30 days (not stated)	None	OSDI TBUT Schirmer's	Symptoms, Schirmer's for tears again	TBUT	Patients with digital eye strain. Compared drops primarily acting on one tear layer CLINICAL TRIAL NOT REGISTERED
Chiambaretta et al 2017 ⁵⁰	Randomized, investigator-masked, multi-center study	HA-trehalose HA	N = 52 N = 49	60.0 ± 12.2 58.5 ± 13.4	84 days (3–6x/day; average 4)	Symptoms with HA-trehalose	Cornea & conjunctival staining	OSDI [Schirmer's, TBUT, staining, hyperaemia, no statistics presented]	None	AEs: 3 with HA-trehalose vs 24 with HA Clinical Trial NCT02023268 - only staining as primary outcome and day 35 data missing Cochrane Risk of Bias R? C-M-O!!S-B?
Christensen et al 2004 ⁵¹	Randomized, double-masked, multi-center study	PEG 400 0.4% / PG/HP-guar 0.3% - Systane CMC 0.5% - Refresh Tears	N = 42 N = 45	58.5 59.5	6 weeks (4x/day)	Lissamine green staining, dryness, refreshed and FB symptoms with 0.5% PEG	Fluorescein staining, use ratings, ocular signs or symptom frequency	Corneal & conjunctival staining only with 0.4% PEG	Conjunctival staining with CMC	CLINICAL TRIAL NOT REGISTERED
Cohen et al 2014 ⁵³	Randomized, double-masked, multi-center study	CMC 1% - Refresh LiquiGel PEG 400 0.4% / PG/HP-Guar 0.3% - Systane Gel	N = 70 N = 67	57.5 ± 16.6 56.5 ± 15.0	6 weeks (4x/day)	Corneal staining with PEG	Conjunctival staining, TBUT, symptoms	Corneal staining	Lissamine green staining, TBUT, symptoms	CLINICAL TRIAL NOT REGISTERED
Comez et al 2013 ¹⁰⁷	Randomized, patient-masked, 2 group contralateral, single-center study	PG 0.3% and PEG 0.4% - Systane SH 15% - Eystil HPMC - Tears Naturale CMC 0.5% - Refresh Tears	N = 17 N = 13	47.4 ± 14.5 46.3 ± 15.5	12 weeks (5x/day)	None	OSDI, Osmolarity, Schirmer's, TBUT	OSDI, osmolarity, Schirmer's, TBUT	None	~30% drop-out CLINICAL TRIAL NOT REGISTERED
Craig et al 2021 ⁴	Randomized, double-masked, multi-center study	Aminomethylpropanol, HP-guar - Systane Ultra Dimyristoyl phosphatidylglycerol, HP-guar, mineral oil, polyoxl 40 stearate - Systane Complete	N = 49 N = 50	43 ± 17 45 ± 16	6 months (4x/day +)	Lipid thickness	Symptoms, TMH, lipid, osmolarity, hyperaemia, expressibility, blinking	Symptoms (OSDI, DEQ-5, SANDE) NIBUT, LWVE Cornea & conjunctival staining	TMH, osmolarity, hyperaemia, expressibility, blinking	Symptoms improved @ 1+ month, LWVE @ 2+ months, lipid @ 3+ months staining @ 4+ months. 1 in 3 had no benefit in signs or symptoms. Those with lipid layer grade ≤ 3 benefit more from lipid-based drop Clinical Trial ACTRN12619000390189 - additional questionnaire, acuity and lid data presented Cochrane Risk of Bias R +C+M+O+I+S-B+
Dausch et al 2006 ⁷⁶	Randomized, investigator-masked, cross-over, multi-center study	Liposomes - Tears Again Carbomer triglycerides - Liposic	N = 74 with deficient lipid layer	n=1 <25 years n=9 25–45 years n=16 46–60 years n=49 >60 years	6 weeks (3x/day)	Symptoms, LIPCOF, TBUT, Schirmer's, lid margin inflammation with Tears Again		Symptoms, LIPCOF, TBUT, Schirmer's, lid margin inflammation	-	Photo sequence of phospholipid liposomes sprayed on eyelid reaching ocular surface CLINICAL TRIAL NOT REGISTERED

Davitt et al 2010 ⁶⁴	Randomized, double-masked, single-center study	PEG 400/PG/HP-guar CMC 0.5% - Optive	N = 52 N = 53	33 x 18–64 years, 19 x ≥65 years 41 x 18–64 years, 12 x ≥65 years	6 weeks (4x/day)	Cornea & conjunctival staining with PEG 400/PG/HP-guar group	Symptoms, TBUT	Symptoms	TBUT	CLINICAL TRIAL NOT REGISTERED
Diaz-Llopis et al 2019 ¹⁰⁸	Randomized, investigator-masked, multi-center study	Seawater spray - Quinton CMC 0.5% -Viscofresh	N = 60 N = 60	68.1 ± 6.3 66.8 ± 8.4	12 weeks (5x/day)	OSDI, IL-1 β and IL-6 with seawater spray	Cornea & conjunctival staining, Schirmer I, osmolarity, TBUT, TMH	OSDI, Cornea & conjunctival staining	Schirmer I, osmolarity, TBUT, TMH	CLINICAL TRIAL NOT REGISTERED
Downie et al 2020 ¹⁰⁹	Randomized, double-masked, multi-center study	CMC, glycerin, flaxseed oil and castor oil and osmoprotectants (levocarnitine, Erythritol & trehalose) (OM3) Refresh Optive Advanced	N = 120 N = 122	54.3 ± 17.3 52.8 ± 16.7	90 days (2x/day +)	Combined corneal / conjunctival staining with OM3	OSDI TBUT	OSDI TBUT Combined corneal / conjunctival staining	None	AEs (OM3 0% vs ROA 4.1%) Clinical Trial NCT02553772 Cochrane Risk of Bias R +C+M+O+I+S+B?
Dumbleton et al 2009 ¹¹⁰	Randomized, double-masked, single-center study	PEG 400 0.25% - Blink gel tears CMC 1% - Refresh Liquigel	N = 56 N = 54	46.3 ± 19.3 47.2 ± 19.1	30 days (3x/day)	Symptoms with PEG	Phenol red test, TMH, NIBUT, hyperaemia, corneal and conjunctival staining		Hyperaemia, corneal and conjunctival staining	No notable AE's CLINICAL TRIAL NOT REGISTERED
Essa et al 2018 ³	Randomized, investigator-masked, crossover, single-center study	SH 0.4% - Clinitas Soothe SH 0.15% - Hyabak Phospholipid liposomes -Tears Again CMC - TheraTears	N = 50 (for all treatments)	60.8 ± 14.2	4 weeks (drops/day a study variable; average 2–3)	None	OSDI NIBUT FBUT TMH Phenol Red LIPCOF Ocular surface staining Lipid layer grading Osmolarity (baseline visit only)	OSDI LIPCOF Conjunctival staining	NIBUT FBUT TMH Phenol Red Lipid layer grading	Artificial tears performed similarly. However, osmolarity balanced preferred in those with low baseline tear volume and liposomal spray for those with lipid layer deficiency. Clinical Trial NCT02420834 Cochrane Risk of Bias R? C?M-O?!+S+B?
Fogt et al 2019 ¹¹¹	Randomized, observer-masked, crossover, non-dispensing, single-center study	Omega 3 - Refresh Optive MEGA-3 Refresh Optive	N = 19 with thin lipid	46.5 ± 8.7	60 minutes (Single application)	Lipid layer thickness (overall), Symptoms with MEGA-3	None	Lipid layer thickness Symptoms	Symptoms Schirmer's	Clinical Trial NCT03380624 - 15 min data missing Cochrane Risk of Bias R? C?M-O?!+S-B?
Fondi et al 2018 ¹¹²	Randomized, patient-masked, crossover single-center study	SH and trehalose - Thealoz Duo HA, trehalose and carbomer - Thealoz Duo Gel	N = 40 (for both treatment)	43.7 ± 12.3	1 week (actual 3.2 ± 2.6x/day HT & 1.9 ± 2.2x/day HTC-gel)	None	Corneal / conjunctival staining TBUT Sleep quality	Corneal / conjunctival staining TBUT Sleep quality	None	Clinical Trial NCT02980913 Cochrane Risk of Bias R? C?M-O?!+S+B?
Garcia-Lazaro et al 2011 ⁶⁷	Randomized, investigator masked, cross-over, single-center study	PEG 400 2.5% - Blink Intensive Tears HPMC 0.3% - Artific Tears	N = 20	57.5 ± 8.4	1 month (3x/day)	Tear meniscus volume with PEG	None	Tear meniscus volume	None	CLINICAL TRIAL NOT REGISTERED
Gensheimer et al 2012 ¹¹³	Randomized, double-masked, contralateral, non-dispensing, single-center study	Glycerin 1% with PLL-g-PEG - Eyeon PG 0.3% and PEG 0.4% - Systane	N = 16	44.5	120 mins (single application)	NIBUT, TBUT with glycerine	None	NIBUT with glycerine	TBUT	CLINICAL TRIAL NOT REGISTERED

(Continued)

Table I (Continued).

Paper	Design	Comparators	Participants Completing	Age (Years)	Duration (Dosing)	Tests Showing Significant Difference	Tests Not Differentiating Products	Tests Showing Significant Difference	Tests Showing no Change	General Comments
						Cross Comparator		Compared to Baseline		
Gokul et al 2018 ⁵⁴	Randomized, double-masked, contralateral, non-dispensing, single-center study	Systane Balance Systane Ultra	N = 30	27 ± 9	30 mins (following 2.5 mins in adverse conditions)	Lipid thickness with liposomal Systane Balance	NIBUT	Lipid thickness, NIBUT	Glare acuity, temperature variation, TMH	CLINICAL TRIAL NOT REGISTERED
Greene et al 1992 ⁶⁶	Randomized, double-masked, single-center study	CMC 1.0% - Celluvisc Lubricant HPMC 0.3% - Tears Naturale 2	N=28? N=28? severe	??	2 months (8x/day)	Symptoms, corneal erosions and impression cytology grades with CMC	Schirmer's Corneal & conjunctival staining Lid & conjunctival swelling	Corneal staining, Symptoms, impression cytology grade (CMC only)	Schirmer's	CLINICAL TRIAL NOT REGISTERED
Iester et al 2000 ⁷⁰	Randomized, open-label?, multi-center, study	HPMC 0.3% HA 0.4%	N = 55 N = 58	56.4 ± 12.8 52.2 ± 10.6	2–3 months (6x/day)	Symptoms, Tear ferning Osmolarity, impression cytology With HA	TBUT Staining Schirmer's I	TBUT, staining, Schirmer's I, symptoms Impression cytology	-	Ferning, osmolarity and impression cytology only measured in ~33% of sample each CLINICAL TRIAL NOT REGISTERED
Jacobi et al 2012 ¹¹⁴	Randomized, open-label? Single-center study	HP-Guar - Systane UD Tamarindus indica seed polysaccharide 1% - VISINE INTENSIV	N=14 N=14	44 ± 8 overall	3 months (5x/day)	TBUT with HP-Guar	OSDI Schirmer's II LIPCOF Corneal & conjunctival (rose Bengal)	TBUT LIPCOF OSDI with HP-Guar	Schirmer's II LIPCOF Corneal & conjunctival (rose Bengal)	CLINICAL TRIAL NOT REGISTERED
Jenkins et al 2020 ¹¹⁵	Randomized, double-masked, multi-center study	Systane Balance Refresh Optive advanced	N = 117 N = 114	56.7 ± 14.7 55.6 ± 16.4	35 days (4x/day)	TBUT with Systane	Symptoms	Symptoms TBUT	None	2 lipid based drops Clinical Trial NCT02776670 - exploratory lid wiper epitheliopathy and questionnaire additionally reported Cochrane Risk of Bias R +C+M!O!-S+B?
Johnson et al 2006 ⁶⁰	Randomized, double-masked, contralateral, single-center study	SH 0.1% SH 0.3% NaCL 0.9%	N = 13 (for all treatments)	Range 21–34	6 hours (single application)	NIBUT (0.3% SH performed better than 0.1% SH)	Symptoms	Symptoms NIBUT	None	CLINICAL TRIAL NOT REGISTERED
Johnson et al 2008 ⁵⁶	Randomized, double-masked study, single-center study	Carbomer 934 0.3% - Lacryvisc SH 0.18% - Vismed	N = 33 N = 32	Median 36 Median 39 Range 21–64	1 month (drops/day a study variable; median 2.1–2.3)	Corneal & conjunctival staining with SH	Symptoms NIBUT TBUT	Symptoms Corneal & conjunctival staining	NIBUT TBUT	CLINICAL TRIAL NOT REGISTERED
Khaireddin and Schmidt, 2010 ¹¹⁶	Randomized, multi-center study	HA - Vismed light Phospholipid - Tears Again	N = 103 N=113 Evaporative	n=9 <25 years, n=26 25–45 years, n=42 46–60 years, n=139 >60 years	3 months 3x/day +	LIPCOF, lid inflammation NIBUT with Tears Again	Schirmer's	LIPCOF, lid Inflammation NIBUT	Schirmer's	CLINICAL TRIAL NOT REGISTERED
Khanal et al 2007 ¹¹⁷	Randomized, investigator-masked, single-center study	Castor oil 0.1.25% HPMC 0.32% - Artelac Single Dose Unit	N = 27 N = 26	Unclear from text	1 month (3x/day)	Tear evaporation with HPMC	Schirmer's, osmolarity	Tear evaporation; Lipid layer with castor oil	Schirmer's, osmolality	CLINICAL TRIAL NOT REGISTERED

Labetoulle et al 2018 ¹¹⁸	Randomized, double-masked, multi-center study	HP-Guar - HA dual-polymer – Systane Hydration SH 0.15% - Hyabak	N = 50 N = 49	61.7 ± 12.3 56.7 ± 14.3	6 weeks (4x/day)	None	Symptoms, TBUT, ocular surface staining	Ocular surface staining	Symptoms, TBUT	Fluorescein dye only Clinical Trial NCT02470429 - exploratory end points additionally reported in n=30 Cochrane Risk of Bias R +C+M?O?I+S+B?
Lahia et al 2020 ¹¹⁹	Randomized, double-masked, single-center study	Sacha inchi microemulsion (SIME) HA 0.2%	N = 26 N = 26	53.3 ± 12.6 overall	1 month (3x/day)	Ocular protection index with SIME	Symptoms, Corneal & conjunctival staining, TBUT	Symptoms, osmolarity in hyperosmolar subgroup. Corneal and conjunctival (nasal) staining, TBUT & lid redness only with SIME	Osmolarity, Conjunctival temporal staining	Fluorescein dye only Clinical Trial NCT03569202 Cochrane Risk of Bias R +C+M+O+I+S+B+
Lee et al 2011 ⁵⁸	Randomized, observer-masked, single-center study	CMC 0.5% - Refresh Plus SH 0.1% - Hynex	N = 33 N = 32	39 ± 14.6 37 ± 13.4	2 months (6x/day)	None	Corneal & conjunctival staining TBUT Symptoms	Cornea & conjunctival staining TBUT Symptoms	None	Fluorescein staining only CLINICAL TRIAL NOT REGISTERED
Lievens et al 2019 ⁵²	Randomized, double-masked, multi-center study	CMC 1.0% and glycerin 0.9% CMC 1.0%	N = 94 N = 94	≥ 18 years of age	1 month (2x/day +)	Symptoms Corneal staining TBUT With CMC-GLY at day 7 only	Symptoms Corneal staining TBUT at all other time points	Symptoms, corneal staining, and TBUT	None	Clinical Trial NCT02280473 Cochrane Risk of Bias R +C+M+O+I+S+B+
Marner et al 1996 ⁷¹	Randomized, open-label, crossover, multi-center study	Carbomer gel - Lubrithal PVA 1.4% - Lacril/ Liquifilm	N=54 (for all treatment)	64.3, range 38–89	2 weeks (drops/day a study variable (carbomer 3.9 vs PVA 4.6x)	Symptoms, TBUT. Instillation frequency with carbomer	Schirmer's I Ocular surface staining, Corneal sensitivity	Schirmer's I, TBUT, ocular surface staining, symptoms	None	Rose Bengal only used AEs 33% with carbomer, 8% with PVA CLINICAL TRIAL NOT REGISTERED
Mihaltz et al 2018 ⁴³	Randomized, investigator-masked, single-center study	Carbomer, triglycerides - Artelac Lipids UD SH - Artelac Splash Edo UD	N=10 N=13	55.5 ± 11.3 53.8 ± 17.9	3 months (4x/day +)	None	Schirmer's, TBUT, Ocular surface staining Symptoms MG dropout aberrations	Schirmer's, TBUT, Ocular surface staining	None	Lipid drops better for those with >50% MG dropout improving Schirmer's & aberrations CLINICAL TRIAL NOT REGISTERED
Muntz et al 2020 ⁵⁵	Randomized, double-masked, contralateral crossover, single-center study	Lipid, PG, HP-guar and mineral oil - Systane Complete PEG 400, PG and HP-guar - Systane Ultra	N = 28 (for all treatments)	29 ± 9	Single application – adverse environment	Symptoms, lipid layer quality, NIBUT with Systane complete	TMH Hyperaemia	Symptoms, Lipid layer quality only with Systane Complete	TMH Hyperaemia	Clinical Trial ACTRN12619000361101 Cochrane Risk of Bias R +C+M+O+I+S+B?
Nelson and Farris, 1998 ¹²⁰	Randomized, double-masked, multi-center study	PVA 1.4% - Liquifilm SH 0.1%	N = 16 N = 20	52.3 ± 16.4 64.8 ± 10.8	8 weeks 8x/day +	-	Symptoms, Osmolality, TBUT, rose bengal staining, Schirmer's I, impression cytology	Symptoms, Osmolality, TBUT, rose bengal staining, Schirmer's I	Impression cytology	CLINICAL TRIAL NOT REGISTERED

(Continued)

Table I (Continued).

Paper	Design	Comparators	Participants Completing	Age (Years)	Duration (Dosing)	Tests Showing Significant Difference	Tests Not Differentiating Products	Tests Showing Significant Difference	Tests Showing no Change	General Comments
						Cross Comparator		Compared to Baseline		
Ousler et al 2007 ⁶⁵	Randomized, double-masked crossover, single-center study	PEG & HP-Guar - Systane CMC - Refresh Tears CMC - Refresh Endura	N = 50	62.7	Single application	TBUT, Ocular protection index with Systane	Blink rate	No comparison presented		No difference between CMC products CLINICAL TRIAL NOT REGISTERED
Park et al 2017 ⁵⁹	Randomized, investigator-masked, multi-center study	SH 0.1% SH 0.15% SH 0.3% Cyclosporine 0.05%	N = 43 N = 41 N = 47 N = 45	44.1 ± 13.9 46.2 ± 14.0 44.8 ± 16.2 45.2 ± 15.4	12 weeks (5–6x/day)	Schirmer's (0.15% SH group)	Corneal & conjunctival staining TBUT	Corneal & conjunctival staining TBUT	Schirmer's	AEs 13% 0.1% SH, 20% 0.15% SH, 13% 0.3% SH, 31% 0.05% CS group. Clinical Trial KCT0001796 Cochrane Risk of Bias R +C!M-O?!+S+B?
Perez-Balbuena et al 2016 ⁴⁷	Randomized, double-masked, multi-center study	Xanthan gum 0.09% and chondroitin sulfate 0.1% PEG 400 0.4% and PG 0.3%	N = 76 N = 72	49.9 ± 16.0 45.5 ± 12.7	2 months (4x/day)	None	Schirmer's, TBUT, Symptoms, Corneal & conjunctival staining	Schirmer's, TBUT, Symptoms	Corneal & conjunctival staining	Clinical Trial NCT01657253 Cochrane Risk of Bias R +C!M-O?!+S+B+
Pinto-Bonilla et al 2015 ⁴²	Randomized, open-label, crossover, single-center study	Trehalose and SH 1.5mg/mL -Thealoz Duo PEG & HP-guar - Systane	N = 9 N = 8	45.3 ± 11.8 53.8 ± 14.6	1 week (5x/day) (Actual 3.7±0.9 / 3.5 ±0.9)	None	Symptoms, Corneal & conjunctival staining, Schirmer's, TBUT	Symptoms	Schirmer's TBUT, Corneal & conjunctival staining	CLINICAL TRIAL NOT REGISTERED
Postorino et al 2018 ⁵³	Randomized, investigator-masked, single-center study	HA crosslinked + CoQ10 HA 0.15% crosslinked	N = 20 N = 20	60.2 ± 13.6 60.9 ± 12.5	3 months (4x/day)	Symptoms, MGD assessment, corneal / conjunctival staining, epithelial hyperreflectivity and keratocytes with HA + CoQ10	Symptoms, corneal aesthesiometry TBUT	OSDI MGD assessment, corneal / conjunctival staining, epithelial hyperreflectivity and keratocytes with HA + CoQ10 only	Corneal aesthesiometry TBUT	Fluorescein staining only Clinical Trial NCT03074344 - meibomian gland assessment and confocal additionally reported Cochrane Risk of Bias R +C+M-O?!+S-B?
Pult et al 2021 ⁷⁷	Randomized, double-masked, crossover, multi-center study	Phospholipid 0.98% - Tears Again Phospholipid 0.12% - Ocuvors	N=30 (all treatments)	33.2±1.8	Single application	Symptoms, NIBUT with high concentration lipid	None	Symptoms, NIBUT with high concentration lipid only	None	CLINICAL TRIAL NOT REGISTERED
Robert et al 2016 ⁶⁸	Randomized, investigator masked, multi-center study	Cationic Emulsion (Cation Norm) SH 0.18% - Vismed	N = 37 N = 37 Moderate to severe	60.0 ± 14.6 65.3 ± 11.1	3 months (4x/day)	Symptoms at 1 month with SH	TBUT, Schirmer's, Corneal & conjunctival staining, Osmolarity, Impression cytology	Symptoms, Corneal & conjunctival staining	Schirmer's, TBUT, Osmolarity, Impression cytology	AEs 18% CE, 27% HS >10% drop-out Clinical Trial EudraCT 2011-A00955-36 Cochrane Risk of Bias R +C!M-O?!+S+B?
Safarzadeh et al 2017 ¹²¹	Randomized patient-masked, single-center study	Dextran 70, 1 mg/mL and HPMC – Tears Naturale Dextran 70, 0.1 mg/mL and 0.3 g HPMC – Tearlose	N = 41 N = 47	44.1 ± 6.3 45.8 ± 8.4	4 weeks (2x/day)	None	Symptoms, TBUT, Schirmer's Corneal & conjunctival staining	Symptoms, TBUT, Corneal & conjunctival staining	Schirmer's	Fluorescein staining only CLINICAL TRIAL NOT REGISTERED

Sanchez et al 2017 ³⁹	Randomized, investigator-masked, single-center study	CMC 0.5% (Viscofresh) HA 0.15% (Lubristil)	N = 7 N = 8	51.8 ± 14.1 71.8 ± 12.2	1 month (4x/day)	TBUT, corneal staining, and HLA-DR with CMC	Schirmer's Other inflammatory markers	HLA-DR, TBUT & corneal staining with CMC	Schirmer's, Tear clearance,	No Aes CLINICAL TRIAL NOT REGISTERED
Schmidl et al 2015 ⁵¹	Randomized, double-masked, single-center study	Trehalose and SH 1.5mg/mL -Thealoz Duo SH, 0.15% - Hyabak NaCL 0.9% - Hydrabak	N = 20 N = 20 N = 20	43.6 ± 13.3 42.9 ± 12.0 41.8 ± 9.9	240 minutes Single application	Tear film thickness (SH +trehalose to 240min and SH to 40min only)	TBUT, Schirmer's	Tear film thickness (both SH products)	TBUT, Schirmer's	CLINICAL TRIAL NOT REGISTERED
Simmons and Vehige, 2007 ⁷⁸	Randomized, double-masked, crossover and parallel groups, multi-center studies	CMC 1.0% CMC 0.5% (Refresh Tears)	N = 43 single application Parallel N = 53 N = 50	Mean 62 Not stated	60 minute (single-application) 1 month (4x/day)	Ocular protection index (low viscosity to 20min, high viscosity to 30min). Corneal & conjunctival staining with higher viscosity	Symptoms	Symptoms, Corneal and conjunctival staining	None	Fluorescein staining only. More AEs with high viscosity – visual disturbance 23vs4%; discharge 13vs2% CLINICAL TRIAL NOT REGISTERED
Simmons et al 2015 ⁷⁹	Randomized, investigator-masked, multi-center study	CMC (Refresh Optive Advanced Sensitive), unit dose CMC (Refresh Optive Sensitive), unit dose CMC (Refresh Optive Advanced Sensitive), multi-dose CMC (Refresh Optive Sensitive), multi-dose	N = 105 N = 103 N = 51 N = 56	54.4 ± 14.8 55.8 ± 14.1 55.2 ± 14.5 53.5 ± 13.9	30 days (2x/day +)	None	Symptoms, TBUT, Corneal & conjunctival staining Schirmer's	OSDI TBUT	Corneal & conjunctival staining, Schirmer's	No clinically significant differences in safety, effectiveness, and acceptability between lipid and aqueous artificial tears Clinical Trial NCT01459588 Cochrane Risk of Bias R +C?M-O?I+S+B?
Simmons et al 2015 ⁴⁹	Randomized, double-masked, multi-center study	CMC 0.5% + 0.1% HA (Optive Fusion) CMC 0.5% + 0.15HA CMC 0.5% (Refresh Tears)	N = 87 N = 87 N = 90	59.6 ± 14.5 59.2 ± 16.3 60.0 ± 13.3	3 months (2x/day +) (actual 4.3, 3.9, 3.8x/day)	Some symptoms with Fusion Corneal staining with Fusion vs Refresh	Conjunctival staining	Symptoms, Corneal & Conjunctival staining	None	Investigational formulations Clinical Trial NCT01294384 - visual disturbance questionnaire additionally reported Cochrane Risk of Bias R +C+M+O?I+S-B?
Szegedi et al 2018 ¹²²	Randomized, patient-masked, single-center study	SH 0.18% + triglycerides, and phospholipids SH 0.18% - Vismed sodium chloride 0.9% - Hydrabak	N = 20? N = 20? N = 20?	34.6 ± 11.7 40.5 ± 9.9 39.2 ± 12.8	40 minutes Single-application	Tear film thickness 40min vs 20min vs 0min with phospholipids	TBUT, Corneal staining, Lipid thickness	Tear film thickness, TBUT, Corneal staining, Lipid thickness	None	Clinical Trial NCT03161080 Cochrane Risk of Bias R? C?M-O-I+S+B?
Tomlison et al 2013 ¹²³	Randomized, double-masked, crossover, single-center study	CMC 0.5% - Refresh Tears CMC 0.5%/castor oil - Optive Plus Glycerin 1%/castor oil - Refresh Ultra	N = 18 with dry eye N = 19 controls For all treatments	41 ± 14 30 ± 12	2 weeks 3x/day	Evaporation for both CMCs	Symptoms, TBUT, NIBUT (except for controls), osmolality	Symptoms, evaporation, TBUT, NIBUT (except for controls), osmolality	Lipid thickness	Measures taken after adaptation to environmental centre CLINICAL TRIAL NOT REGISTERED

(Continued)

Table I (Continued).

Paper	Design	Comparators	Participants Completing	Age (Years)	Duration (Dosing)	Tests Showing Significant Difference	Tests Not Differentiating Products	Tests Showing Significant Difference	Tests Showing no Change	General Comments
Troiano and Monaco, 2008 ⁴⁶	Randomized, patient-masked, crossover, single-center study	HA 0.4% 300mOsm/L HA 0.4% 150mOsm/L	N = 28 For all treatments	55.5 ± 7.3	7 days 4x/day	Foreign body and dryness symptoms and ocular surface staining with 150mOsm/L	None	Symptoms, hyperaemia, ocular surface staining	None	Reducing osmolarity effective Rose Bengal staining only CLINICAL TRIAL NOT REGISTERED
van Setten et al 2020 ¹²⁴	Substitution, open-label, multi-center study	High molecular weight HA 0.15% - Comfort Shield Over habitual controls	N = 44 N = 40	57.7 ± 14.4 59.5 ± 12.5	8 weeks Actual 8.2 vs 6.5			Symptoms, Visual acuity, nerve fibre length with high molecular weight HA	Corneal staining, TBUT, Schirmer's, Lid wiper epitheliopathy, mucotaneous junction, osmolarity	Change from habitual optimal artificial tears. No change with controls CLINICAL TRIAL NOT REGISTERED
Waduthantri et al 2012 ¹²⁵	Randomized, double-masked, single-center study	CMC 0.5% - Refresh Tears PEG 400 0.4% / PG/HP-guar 0.3% - Systane Ultra	N = 15 N = 15	54.7 ± 5.8 55.9 ± 5.3	6 weeks 4x/day	None	Symptoms Schirmer's TBUT, Corneal staining	Symptoms	Schirmer's TBUT, Corneal staining	Clinical Trial NCT00796926 - meibography, osmolarity and tear meniscus height not reported Cochrane Risk of Bias R +C/M+O+I+S-B+
Wang et al 2007 ⁷³	Randomized, open label, single-center study	Carbomer - Vidisic Ophthalmic Gel Cellulose - Artelac Ophthalmic Solution Mineral oil (lanolin) -Duratears Ointment	N = 22 N = 23 N = 22	55.9 ± 15.7 50.1 ± 14.3 60.3 ± 11.2	4 weeks (4x/day for Carbomer and Cellulose) (1x/day before sleep for mineral oil)	Schirmer's with Carbomer and Cellulose & TBUT with Carbomer	Schirmer's	Symptoms, TBUT, Schirmer's		Fluorescein staining only, but not reported in results CLINICAL TRIAL NOT REGISTERED
Wang et al 2010 ¹²⁶	Randomized, open label, single-center study	Carbomer + lipid gel - Liposic Ophthalmic Liquid Gel HP-guar gel - Systane Lubricant Eye Drops	N = 15 N = 15	40.4 ± 15.0 49.5 ± 12.2	2 months (4x/day)	Symptoms & Schirmer's with Carbomer + lipid	TBUT	Symptoms Schirmer's TBUT	None	Fluorescein staining only, but not analysed in results CLINICAL TRIAL NOT REGISTERED
Xiao et al 2008 ⁷²	Randomized, investigator-masked, single-center study	Carbomer-based 0.4% gel CMC 1.0%	N = 30 N = 30	46.7 ± 2.3 46.6 ± 2.1	3 months 3x/day +	Symptoms, TBUT, Schirmer's, corneal staining, ocular residence time with carbomer gel	None	Symptoms, TBUT, Schirmer's corneal staining (but no statistics presented)	None	Method relating to precorneal residence time missing. Fluorescein staining only. CLINICAL TRIAL NOT REGISTERED

Notes: Grey box = no statistical comparison made; ? = not certain from paper. Cochrane Risk of Bias Tool Rating:³⁵ "+": low risk; "-": high risk; "?": unclear risk; for Random sequence generation (selection bias), allocation Concealment (selection bias); Masking of participants/researchers (performance bias), masking of Outcome assessment (detection bias), Incomplete outcome data (attrition bias), Selective reporting (reporting bias), other Biases (respectively). CLINICAL TRIAL NOT REGISTERED - based on a search of the main registries and paper. © Aston University.

Abbreviations: CoQ10, coenzyme Q10; CMC, carboxymethylcellulose/ carmellose sodium; HA, hyaluronic acid/ hyaluronan; HPMC, Hydroxypropyl methylcellulose; HP-Guar, Hydroxypropyl guar; NaCl, sodium chloride; NIBUT, non-invasive tear breakup time; PEG, polyethylene glycol; PG, propylene glycol; PVA, polyvinyl alcohol; SH, sodium hyaluronate; TBUT, fluorescein tear breakup time; TMH, tear meniscus height; LIPCOF, Lid parallel conjunctival folds.

- Long-term compliance is needed to improve ocular surface signs rather than just symptoms⁴ and symptoms benefit from 4x/day compared to “as needed” dosing.⁷⁵
- Higher liposomal concentration increases effectiveness.^{76,77}
- Lower osmolarity drops increase the effectiveness of an artificial tear drop.⁴⁶
- Higher concentration (viscosity) CMC is more effective in reducing corneal and conjunctival staining, but caused more reports of visual disturbance.⁷⁸
- While drops targeting individual layers of the tear film seem equally effective,^{41,79} studies have shown that the most effective drop for an individual can be predicted from their baseline classification; drops containing phospholipids are more effective in those with evaporative dry eye^{3,4} and osmoprotectants benefit those with high tear film osmolarity.³
- Artificial tears may not be effective for as much as one-third of patients, but this can be predicted by one month of compliant use.⁴

These findings can inform clinical dry eye practice; in summary: non-preserved or soft preserved artificial tears being appropriate to prescribe to patients, regardless of the severity of their DED; patients with evaporative dry eye should be prescribed artificial tears containing a high concentration of liposomes; one month’s compliant use 4x/day is recommended to determine whether an artificial tear can manage the patients’ symptoms in the longer-term; signs of ocular surface disease typically take up to 4 months to start improving so patience is needed; artificial tears with multiple active ingredients (especially with PEG) seem to outperform more basic previous generation drops; ability to use different types of artificial tear bottles/sprays varies⁹ and should be part of the prescribing consideration. While the efficacy of artificial tears is well established for managing DED, its use in ocular surface disease without symptoms to improve post-surgical symptomology and to reduce refractive ‘surprises’ from poor ocular biometry⁸⁰ is less well established. The data available as reviewed in this study is limited by the definition of dry eye disease applied in published studies being variable as well as the disease severity examined and compliance with artificial tears being rarely quantified.

Other Therapeutic Functions of Artificial Tears

As well as being a management option for dry eye disease and the ocular surface, artificial tears can also be utilised for a wide range of therapeutic functions such as in the treatment of anterior eye trauma, infection, inflammation and disease as well as contact lens management.

Corneal Abrasion and Wound Healing

Corneal abrasions can be caused by foreign bodies, trauma, and trichiasis, and may result in pain, redness, lacrimation, and photophobia. Artificial tears improve epithelial healing.⁸¹ Ideally, preservative free drops are used as they tend to be associated with better ocular surface health and tolerability.⁸² The most common treatment for perioperative corneal abrasions is artificial tears followed by a combination of artificial tears and antibiotic ointment.⁸³ Most artificial tears contain hydrogels; these are known to activate the epidermal growth factor (EGF) receptor which promotes the healing of corneal epithelial wounds.⁸⁴

Pain and Inflammation Management

Artificial tears are commonly used in the management of ocular pain and inflammation. In the treatment of episcleritis, the combination of artificial tears and cold compresses provide symptomatic relief.⁸⁵ No significant differences have been observed in the signs or symptoms of idiopathic episcleritis when either artificial tears or topical ketorolac (NSAID) is used.⁸⁶ Following photorefractive keratectomy (PRK) surgery, the application of preservative-free artificial tears reduces postoperative ocular discomfort and increases visual recovery.⁸⁷ Cooled artificial tears have been shown to reduce corneal and conjunctival sensation, with 4°C being the most comfortable temperature.⁸⁸ In contrast to this, Bitton et al found no improvement in perceived patient comfort when refrigerated Systane Ultra artificial tears were used for mild to moderate dry eye sufferers.⁸⁹ It is also worth noting that pain complaints can be associated with contrasting subjective responses,⁹⁰ and in some patients artificial tears are not effective in relieving uncomfortable symptoms.⁹¹

Conjunctivitis

Allergic conjunctivitis causes ocular itching, watery discharge, lid oedema and conjunctival chemosis. Bilkhu et al exposed 18 participants (who had a known allergy to grass pollen) to grass pollen, and found that artificial tears and cold compresses improved the signs of allergic conjunctivitis and provided symptomatic relief.⁹² However, if symptoms are persistent, short-term use of topical antihistamines and mast cell stabiliser drops is recommended.⁹³

Viral (non-herpetic) conjunctivitis causes redness, discomfort, and watering. Follicles on the palpebral conjunctiva and punctate epithelial lesions on the cornea may also be observed. It has been shown that 0.5% topical ketorolac,⁹⁴ 0.45% ketorolac tromethamine,⁹⁵ and 1% prednisolone acetate⁹⁶ are no better in relieving signs or symptoms of viral conjunctivitis compared to artificial tears.

Bacterial conjunctivitis causes redness, discomfort, and produces a sticky discharge with crusting of the eyelids. Bacterial conjunctivitis usually self-resolves, but the application of artificial tears and eye bathing aids ocular comfort and hygiene. If bacterial conjunctivitis persists after 3–4 days, the application of topical antibiotics is usually recommended.⁹⁷

Keratitis

Keratitis is an inflammation of the cornea and has several different aetiologies including viral (Herpes Simplex), bacterial (marginal keratitis), fungal, contact-lens associated and unprotected exposure to ultraviolet radiation (photokeratitis). In dry eye and photokeratitis,⁹⁸ the application of artificial tears has been recommended. In herpetic keratitis, marginal keratitis, fungal keratitis, and contact-lens associated keratitis, artificial tears are advised (for lubrication and symptomatic relief) alongside additional treatment such as topical antivirals, topical and/or oral antibiotics, and antifungals.

Contact Lens Rewetting and Removal

Contact lens wearers commonly use preservative free artificial tears for ocular lubrication, comfort and contact lens rehydration.^{99–101} Towards the end of wear, contact lenses become drier and fit tighter. The application of artificial tears reduces friction against the cornea and can facilitate safe lens removal.

Foreign Body Removal

Corneal foreign bodies can cause irritation, lacrimation, blurred vision, and redness. Loose foreign bodies can be irrigated away with normal saline or artificial tears. Upon successful removal of a foreign body, prophylactic antibiotics,¹⁰² analgesia and artificial tears are advised.¹⁰³

Summary

Artificial tears are the mainstay of DED management, but also have a role in corneal abrasion and wound healing, pain and inflammation management, conjunctivitis, keratitis, contact lens rewetting and removal, and foreign body removal. A review of randomized controlled trials comparing artificial tears identified 64 papers. There is good evidence that artificial tears improve symptoms of DED within a month of regular use, applied ~4x a day, but signs generally take several months. Not all patients with DED benefit from artificial tears, so if there is no benefit over a month, alternative management should be considered. Combination formulations are more effective than single active ingredient artificial tears. PEG containing artificial tears are more effective than those containing CMC and HPMC. Those classified as having evaporative DED, benefit from artificial tears with liposomes, especially of higher concentration.

Disclosure

JSW is on the executive of the Tear Film and Ocular Surface Society and the Aston University Optometry Research Group have received research funding from Alcon, the Eye Doctor, Scope Ophthalmic and Thea Pharmaceuticals. No funding was received to conduct this review. The authors report no other conflicts of interest in this work.

References

1. Jones L, Downie LE, Korb D, et al. TFOS DEWS II management and therapy report. *Ocular Surface*. 2017;15(3):575–628. doi:10.1016/j.jtos.2017.05.006
2. Pucker AD, Ng SM, Nichols JJ. Over the counter (OTC) artificial tear drops for dry eye syndrome. *Cochrane Database Syst Rev*. 2016;2016(2):Cd009729. doi:10.1002/14651858.CD009729.pub2
3. Essa L, Laughton D, Wolffsohn JS. Can the optimum artificial tear treatment for dry eye disease be predicted from presenting signs and symptoms? *Contact Lens Anterior Eye*. 2018;41(1):60–68. doi:10.1016/j.clae.2017.07.007
4. Craig JP, Muntz A, Wang MTM, et al. Developing evidence-based guidance for the treatment of dry eye disease with artificial tear supplements: a six-month multicentre, double-masked randomised controlled trial. *Ocular Surface*. 2021;20:62–69. doi:10.1016/j.jtos.2020.12.006
5. Hynneklev L, Magno M, Vernhardsdottir RR, et al. Hyaluronic acid in the treatment of dry eye disease. *Acta Ophthalmol*. 2022;100:844–860. doi:10.1111/aos.15159
6. Gomes JAP, Azar DT, Baudouin C, et al. TFOS DEWS II iatrogenic report. *Ocular Surface*. 2017;15(3):511–538. doi:10.1016/j.jtos.2017.05.004
7. Dietlein TS, Jordan JF, Lüke C, Schild A, Dinslage S, Krieglstein GK. Self-application of single-use eyedrop containers in an elderly population: comparisons with standard eyedrop bottle and with younger patients. *Acta Ophthalmologica*. 2008;86(8):856–859. doi:10.1111/j.1755-3768.2007.01155.x
8. Connor A, Severn P. Force requirements in topical medicine use—the squeezability factor. *Eye*. 2011;25(4):466–469. doi:10.1038/eye.2011.5
9. Drew T, Wolffsohn JS. Usability of prostaglandin monotherapy eye droppers. *Br J Ophthalmol*. 2015;99(9):1251–1254. doi:10.1136/bjophthalmol-2014-306291
10. Kashiwagi K. Wide variation of squeezing force and dispensing time interval among eyedropper bottles. *J Ophthalmol*. 2019;2019:115.
11. NHS. Why can't I get a prescription for an over-the-counter medicine? NHS. Available from: <https://www.nhs.uk/common-health-questions/medicines/why-cant-i-get-prescription-over-counter-medicine/>. Accessed September 6, 2022.
12. Bilkhu P, Sivardeen Z, Chen C, et al. Patient-reported experience of dry eye management: an international multicentre survey. *Cont Lens Anterior Eye*. 2022;45(1):101450. doi:10.1016/j.clae.2021.101450
13. Lee S-Y, Tong L. Lipid-containing lubricants for dry eye: a systematic review. *Optomet Vision Sci*. 2012;89(11):1654–1661. doi:10.1097/OPX.0b013e31826f32e0
14. Moshirfar M, Pierson K, Hanamaikai K, Santiago-Caban L, Muthappan V, Passi SF. Artificial tears potpourri: a literature review. *Clin Ophthalmol*. 2014;8:1419. doi:10.2147/OPTH.S65263
15. Agarwal P, Khun D, Krösser S, et al. Preclinical studies evaluating the effect of semifluorinated alkanes on ocular surface and tear fluid dynamics. *Ocul Surf*. 2019;17(2):241–249. doi:10.1016/j.jtos.2019.02.010
16. Baudouin C, Labbé A, Liang H, Pauly A, Brignole-Baudouin F. Preservatives in eyedrops: the good, the bad and the ugly. *Prog Retinal Eye Res*. 2010;29(4):312–334. doi:10.1016/j.preteyeres.2010.03.001
17. Kathuria A, Shamloo K, Jhanji V, Sharma A. Categorization of marketed artificial tear formulations based on their ingredients: a rational approach for their use. *J Clin Med*. 2021;10(6):1289. doi:10.3390/jcm10061289
18. Pisárčik M, Bakoš D, Čepčan MJC, et al. Non-Newtonian properties of hyaluronic acid aqueous solution. *Am J Phys Anthropol*. 1995;97(3):197–202. doi:10.1002/ajpa.1330970209
19. Arshinoff SA, Hofmann I, Nae H, Surgery R. Role of rheology in tears and artificial tears. *J Cataract Refract Surg*. 2021;47(5):655–661.
20. Rah MJ. A review of hyaluronan and its ophthalmic applications. *Optomet-J Am Optometr Assoc*. 2011;82(1):38–43. doi:10.1016/j.optm.2010.08.003
21. Balazs E, Freeman M, Klöti R, Meyer-Schwickerath G, Regnault F, Sweeney D. Hyaluronic acid and replacement of vitreous and aqueous humor. *Mod Probl Ophthalmol*. 1972;10:3–21.
22. Polack FM, McNiece M. The treatment of dry eyes with Na hyaluronate (Healon®). *Cornea*. 1982;1(2):133–136. doi:10.1097/00003226-198201020-00007
23. Müller-Lierheim WG. Why chain length of hyaluronan in eye drops matters. *Diagnostics*. 2020;10(8):511. doi:10.3390/diagnostics10080511
24. Kojima T, Nagata T, Kudo H, et al. The effects of high molecular weight hyaluronic acid eye drop application in environmental dry eye stress model mice. *Int J Mol Sci*. 2020;21(10):3516. doi:10.3390/ijms21103516
25. Pauloin T, Dutot M, Warnet J-M, Rat P. In vitro modulation of preservative toxicity: high molecular weight hyaluronan decreases apoptosis and oxidative stress induced by benzalkonium chloride. *Eur J Pharma Sci*. 2008;34(4–5):263–273. doi:10.1016/j.ejps.2008.04.006
26. Pauloin T, Dutot M, Joly F, Warnet J-M, Rat P. High molecular weight hyaluronan decreases UVB-induced apoptosis and inflammation in human epithelial corneal cells. *Molecular Vision*. 2009;15:577.
27. Wu CL, Chou HC, Li JM, et al. Hyaluronic acid-dependent protection against alkali-burned human corneal cells. *Electrophoresis*. 2013;34(3):388–396. doi:10.1002/elps.201200342
28. Gomis A, Pawlak M, Balazs EA, Schmidt RF, Belmonte CJA. Effects of different molecular weight elastoviscous hyaluronan solutions on articular nociceptive afferents. *Arthritis Rheumat*. 2004;50(1):314–326. doi:10.1002/art.11421
29. Song JK, Lee K, Park HY, et al. Efficacy of carboxymethylcellulose and hyaluronate in dry eye disease: a systematic review and meta-analysis. *Korean J Fam Med*. 2017;38(1):2–7. doi:10.4082/kjfm.2017.38.1.2
30. Ang BCH, Sng JJ, Wang PXH, Htoon HM, Tong LHT. Sodium hyaluronate in the treatment of dry eye syndrome: a systematic review and meta-analysis. *Sci Rep*. 2017;79013. doi:10.1038/s41598-017-08534-5
31. Alves M, Fonseca EC, Alves MF, et al. Dry eye disease treatment: a systematic review of published trials and a critical appraisal of therapeutic strategies. *Ocul Surf*. 2013;11(3):181–192. doi:10.1016/j.jtos.2013.02.002
32. Liu R, Rong B, Tu P, et al. Analysis of cytokine levels in tears and clinical correlations after intense pulsed light treating meibomian gland dysfunction. *Am J Ophthalmol*. 2017;183:81–90. doi:10.1016/j.ajo.2017.08.021
33. Yang YJ, Lee WY, Kim YJ, Hong YP. A meta-analysis of the efficacy of hyaluronic acid eye drops for the treatment of dry eye syndrome. *Int J Environ Res Public Health*. 2021;18(5). doi:10.3390/ijerph18052383

34. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Rev Esp Cardiol.* 2021;74(9):790–799. doi:10.1016/j.rec.2021.07.010
35. Higgins JPT, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* 2011;343:d5928. doi:10.1136/bmj.d5928
36. Lemp MA, Foulks GN. The definition and classification of dry eye disease: report of the Definition and Classification Subcommittee of the International Dry Eye WorkShop (2007). *Ocul Surf.* 2007;5(2):75–92. doi:10.1016/s1542-0124(12)70081-2
37. Wolffsohn JS, Arita R, Chalmers R, et al. TFOS DEWS II diagnostic methodology report. *Ocul Surf.* 2017;15(3):539–574. doi:10.1016/j.jtos.2017.05.001
38. Armstrong RA. Statistical guidelines for the analysis of data obtained from one or both eyes. *Ophthalmic Physiol Opt.* 2013;33(1):7–14. doi:10.1111/opo.12009
39. Sanchez MA, Torralbo-Jimenez P, Giron N, et al. Comparative analysis of carmellose 0.5% versus hyaluronate 0.15% in dry eye: a flow cytometric study. *Cornea.* 2010;29(2):167–171. doi:10.1097/ICO.0b013e3181b11648
40. Brignole F, Pisella PJ, Dupas B, Baeyens V, Baudouin C. Efficacy and safety of 0.18% sodium hyaluronate in patients with moderate dry eye syndrome and superficial keratitis. *Graefes Arch Clin Exp Ophthalmol.* 2005;243(6):531–538. doi:10.1007/s00417-004-1040-6
41. Calvao-Santos G, Borges C, Nunes S, Salgado-Borges J, Duarte L. Efficacy of 3 different artificial tears for the treatment of dry eye in frequent computer users and/or contact lens users. *Eur J Ophthalmol.* 2011;21(5):538–544. doi:10.5301/ejo.2011.6324
42. Pinto-Bonilla JC, Del Olmo-Jimeno A, Llovet-Osuna F, Hernandez-Galilea E. A randomized crossover study comparing trehalose/hyaluronate eyedrops and standard treatment: patient satisfaction in the treatment of dry eye syndrome. *Ther Clin Risk Manag.* 2015;11:595–603. doi:10.2147/TCRM.S77091
43. Mihaltz K, Faschinger EM, Vecsei-Marlovits PV. Effects of lipid- versus sodium hyaluronate-containing eye drops on optical quality and ocular surface parameters as a function of the meibomian gland dropout rate. *Cornea.* 2018;37(7):886–892. doi:10.1097/ico.0000000000001523
44. Peterson RC, Wolffsohn JS, Fowler CW. Optimization of anterior eye fluorescein viewing. *Am J Ophthalmol.* 2006;142(4):572–575. doi:10.1016/j.ajo.2006.04.062
45. Delaveris A, Stahl U, Madigan M, Jalbert I. Comparative performance of lissamine green stains. *Cont Lens Anterior Eye.* 2018;41(1):23–27. doi:10.1016/j.clae.2017.11.002
46. Troiano P, Monaco G. Effect of hypotonic 0.4% hyaluronic acid drops in dry eye patients: a cross-over study. *Cornea.* 2008;27(10):1126–1130. doi:10.1097/ICO.0b013e318180e55c
47. Perez-Balbuena AL, Ochoa-Tabares JC, Belalcazar-Rey S, et al. Efficacy of a fixed combination of 0.09 % xanthan gum/0.1 % chondroitin sulfate preservative free vs polyethylene glycol/propylene glycol in subjects with dry eye disease: a multicenter randomized controlled trial. *BMC Ophthalmol.* 2016;16:164. doi:10.1186/s12886-016-0343-9
48. Aragona P, Benitez-del-Castillo JM, Coroneo MT, et al. Safety and efficacy of a preservative-free artificial tear containing carboxymethylcellulose and hyaluronic acid for dry eye disease: a randomized, controlled, multicenter 3-month study. *Clin Ophthalmol.* 2020;14:2951–2963. doi:10.2147/oph.S256480
49. Simmons PA, Liu H, Carlisle-Wilcox C, Vehige JG. Efficacy and safety of two new formulations of artificial tears in subjects with dry eye disease: a 3-month, multicenter, active-controlled, randomized trial. *Clin Ophthalmol.* 2015;9:665–675. doi:10.2147/oph.S78184
50. Chiambaretta F, Doan S, Labetoulle M, et al. A randomized, controlled study of the efficacy and safety of a new eyedrop formulation for moderate to severe dry eye syndrome. *Eur J Ophthalmol.* 2017;27(1):1–9. doi:10.5301/ejo.5000836
51. Schmidl D, Schmetterer L, Witkowska KJ, et al. Tear film thickness after treatment with artificial tears in patients with moderate dry eye disease. *Cornea.* 2015;34(4):421–426. doi:10.1097/ICO.0000000000000358
52. Lievens C, Berdy G, Douglass D, et al. Evaluation of an enhanced viscosity artificial tear for moderate to severe dry eye disease: a multicenter, double-masked, randomized 30-day study. *Contact Lens Anterior Eye.* 2019;42(4):443–449. doi:10.1016/j.clae.2018.12.003
53. Postorino EI, Rania L, Aragona E, et al. Efficacy of eyedrops containing cross-linked hyaluronic acid and coenzyme Q10 in treating patients with mild to moderate dry eye. *Eur J Ophthalmol.* 2018;28(1):25–31. doi:10.5301/ejo.5001011
54. Gokul A, Wang MTM, Craig JP. Tear lipid supplement prophylaxis against dry eye in adverse environments. *Contact Lens Anterior Eye.* 2018;41(1):97–100. doi:10.1016/j.clae.2017.09.013
55. Muntz A, Marasini S, Wang MTM, Craig JP. Prophylactic action of lipid and non-lipid tear supplements in adverse environmental conditions: a randomised crossover trial. *Ocular Surface.* 2020;18(4):920–925. doi:10.1016/j.jtos.2020.08.004
56. Johnson ME, Murphy PJ, Boulton M. Carbomer and sodium hyaluronate eyedrops for moderate dry eye treatment. *Optomet Vision Sci.* 2008;85(8):750–757. doi:10.1097/OPX.0b013e318182476c
57. Baudouin C, Cochener B, Pisella PJ, et al. Randomized, Phase III study comparing osmoprotective carboxymethylcellulose with sodium hyaluronate in dry eye disease. *Eur J Ophthalmol.* 2012;22(5):751–761. doi:10.5301/ejo.5000117
58. Lee JH, Ahn HS, Kim EK, Kim T-I. Efficacy of sodium hyaluronate and carboxymethylcellulose in treating mild to moderate dry eye disease. *Cornea.* 2011;30(2):175–179. doi:10.1097/ICO.0b013e3181e9adcc
59. Park Y, Song JS, Choi CY, et al. A randomized multicenter study comparing 0.1%, 0.15%, and 0.3% sodium hyaluronate with 0.05% cyclosporine in the treatment of dry eye. *J Ocular Pharmacol Therap.* 2017;33(2):66–72.
60. Johnson ME, Murphy PJ, Boulton M. Effectiveness of sodium hyaluronate eyedrops in the treatment of dry eye. *Graefes Arch Clin Exp Ophthalmol.* 2006;244(1):109–112. doi:10.1007/s00417-005-0028-1
61. Christensen MT, Cohen S, Rinehart J, et al. Clinical evaluation of an HP-guar gellable lubricant eye drop for the relief of dryness of the eye. *Curr Eye Res.* 2004;28(1):55–62. doi:10.1076/ceyr.28.1.55.23495
62. Benelli U, Nardi M, Posarelli C, Albert TG. Tear osmolarity measurement using the TearLab™ Osmolarity System in the assessment of dry eye treatment effectiveness. *Contact Lens Anterior Eye.* 2010;33(2):61–67. doi:10.1016/j.clae.2010.01.003
63. Cohen S, Martin A, Sall K. Evaluation of clinical outcomes in patients with dry eye disease using lubricant eye drops containing polyethylene glycol or carboxymethylcellulose. *Clin Ophthalmol.* 2014;8:157–164. doi:10.2147/oph.S53822
64. Davitt WF, Bloomstein M, Christensen M, Martin AE. Efficacy in patients with dry eye after treatment with a new lubricant eye drop formulation. *J Ocular Pharmacol Therap.* 2010;26(4):347–353. doi:10.1089/jop.2010.0025

65. Ousler GW, Michaelson C, Christensen MT. An evaluation of tear film breakup time extension and ocular protection index scores among three marketed lubricant eye drops. *Cornea*. 2007;26(8):949–952. doi:10.1097/ICO.0b013e3180de1c38
66. Grene RB, Lankston P, Mordaunt J, Harrold M, Gwon A, Jones R. Unpreserved carboxymethylcellulose artificial tears evaluated in patients with keratoconjunctivitis sicca. *Cornea*. 1992;11(4):294–301. doi:10.1097/00003226-199207000-00004
67. Garcia-Lazaro S, Belda-Salmeron L, Ferrer-Blasco T, Cervino A, Montes-Mico R. Comparison of two artificial tear formulations for dry eye through high-resolution optical coherence tomography. *Clin Exp Optomet*. 2011;94(6):549–556. doi:10.1111/j.1444-0938.2011.00632.x
68. Robert PY, Cochener B, Amrane M, et al. Efficacy and safety of a cationic emulsion in the treatment of moderate to severe dry eye disease: a randomized controlled study. *Eur J Ophthalmol*. 2016;26(6):546–555. doi:10.5301/ejo.5000830
69. Amrane M, Creuzot-Garcher C, Robert PY, et al. Ocular tolerability and efficacy of a cationic emulsion in patients with mild to moderate dry eye disease - A randomised comparative study. *J Francais D Ophthalmologie*. 2014;37(8):589–598. doi:10.1016/j.jfo.2014.05.001
70. Lester M, Orsoni GJ, Gamba G, et al. Improvement of the ocular surface using hypotonic 0.4% hyaluronic acid drops in keratoconjunctivitis sicca. *Eye*. 2000;14(6):892–898. doi:10.1038/eye.2000.244
71. Marner K, Moller PM, Dillon M, RaskPedersen E. Viscous carbomer eye drops in patients with dry eyes - Efficacy and safety. A randomized, open, cross-over, multicentre study. *Acta Ophthalmologica Scandinavica*. 1996;74(3):249–252. doi:10.1111/j.1600-0420.1996.tb00086.x
72. Xiao Q, Hu Y, Chen F, Chen X. A comparative assessment of the efficacy of carbomer gel and carboxymethyl cellulose containing artificial tears in dry eyes. *J Huazhong Univ Sci Technol*. 2008;28(5):592–595. doi:10.1007/s11596-008-0523-9
73. Wang IJ, Lin IC, Hou YC, Hu FR. A comparison of the effect of carbomer-, cellulose-, and mineral oil-based artificial tear formulations. *Eur J Ophthalmol*. 2007;17(2):151–159. doi:10.1177/112067210701700202
74. Baeyens V, Bron A, Baudouin C; Vismed Hylovis Study Group. Efficacy of 0.18% hypotonic sodium hyaluronate ophthalmic solution in the treatment of signs and symptoms of dry eye disease. *J Francais D Ophthalmologie*. 2012;35(6):412–419. doi:10.1016/j.jfo.2011.07.017
75. Asbell P, Vingrys AJ, Tan J, et al. Clinical outcomes of fixed versus as-needed use of artificial tears in dry eye disease: a 6-week, observer-masked phase 4 clinical trial. *Invest Ophthalmol Vis Sci*. 2018;59(6):2275–2280. doi:10.1167/iovs.17-23733
76. Dausch D, Lee S, Dausch S, Kim JC, Schwert G, Michelson W. Comparative study of treatment of the dry eye syndrome due to disturbances of the tear film lipid layer with lipid-containing tear substitutes - efficacy of lipid-containing tear substitutes. *Klin Monbl Augenheilkd*. 2006;223(12):974–983. doi:10.1055/s-2006-927266
77. Pult H, Khatum FS, Trave-Huarte S, Wolffsohn JS. Effect of eye spray phospholipid concentration on the tear film and ocular comfort. *Eye Contact Lens-Sci Clin Pract*. 2021;47(8):445–448. doi:10.1097/icl.0000000000000788
78. Simmons PA, Vehige JG. Clinical performance of a mid-viscosity artificial tear for dry eye treatment. *Cornea*. 2007;26(3):294–302. doi:10.1097/ICO.0b013e31802e1e04
79. Simmons PA, Carlisle-Wilcox C, Vehige JG. Comparison of novel lipid-based eye drops with aqueous eye drops for dry eye: a multicenter, randomized controlled trial. *Clin Ophthalmol*. 2015;9:657–664. doi:10.2147/oph.S74849
80. Röggl V, Leydolt C, Schartmüller D, et al. Influence of artificial tears on keratometric measurements in cataract patients. *Am J Ophthalmol*. 2021;221:1–8. doi:10.1016/j.ajo.2020.08.024
81. Prinz J, Mehta JS, Walter P, Fuest M. Simple limbal epithelial transplantation (SLET) A simple technique for the treatment of unilateral complete limbal stem cell deficiency. Video article. *Ophthalmologie*. 2021;118(4):404–412. doi:10.1007/s00347-021-01346-z
82. Walsh K, Jones L. The use of preservatives in dry eye drops. *Clin Ophthalmol*. 2019;13:1409–1425. doi:10.2147/oph.S211611
83. Segal KL, Fleischut P, Kim C, et al. Evaluation and treatment of perioperative corneal abrasions. *J Ophthalmol*. 2014;2014:901901. doi:10.1155/2014/901901
84. Lozano JS, Chay EY, Healey J, Sullenberger R, Klarlund JK. Activation of the epidermal growth factor receptor by hydrogels in artificial tears. *Exp Eye Res*. 2008;86(3):500–505. doi:10.1016/j.exer.2007.12.003
85. Salama A, Elsheikh A, Alweis R. Is this a worrisome red eye? Episcleritis in the primary care setting. *J Commun Hosp Intern Med Perspect*. 2018;8(1):46–48. doi:10.1080/20009666.2017.1418110
86. Williams CPR, Browning AC, Sleep TJ, Webber SK, McGill JI. A randomized, double-blind trial of topical ketorolac vs artificial tears for the treatment of episcleritis. *EYE*. 2005;19(7):739–742. doi:10.1038/sj.eye.6701632
87. Mohammadpour M, Khorrami-Nejad M, Shakoor D. Role of artificial tears with and without hyaluronic acid in controlling ocular discomfort following PRK: a randomized clinical trial. *Int J Ophthalmol*. 2021;14(8):1225–1230. doi:10.18240/ijo.2021.08.14
88. Fujishima H, Yagi Y, Shimazaki J, Tsubota K. Effects of artificial tear temperature on corneal sensation and subjective comfort. *Cornea*. 1997;16(6):630–634. doi:10.1097/00003226-199711000-00005
89. Bitton E, Crncich V, Brunet N. Does the temperature of an artificial tear affect its comfort? *Clin Exp Optomet*. 2018;101(5):641–647. doi:10.1111/coo.12664
90. Galor A, Batawi H, Felix ER, et al. Incomplete response to artificial tears is associated with features of neuropathic ocular pain. *Br J Ophthalmol*. 2016;100(6):745–749. doi:10.1136/bjophthalmol-2015-307094
91. Kim M, Lee Y, Mehra D, Sabater AL, Galor A. Dry eye: why artificial tears are not always the answer. *BMJ Open Ophthalmol*. 2021;6(1):e000697. doi:10.1136/bmjophth-2020-000697
92. Bilkhu PS, Wolffsohn JS, Naroo SA, Robertson L, Kennedy R. Effectiveness of nonpharmacologic treatments for acute seasonal allergic conjunctivitis. *Ophthalmology*. 2014;121(1):72–78. doi:10.1016/j.ophtha.2013.08.007
93. Castillo M, Scott NW, Mustafa MZ, Mustafa MS, Azuara-Blanco A. Topical antihistamines and mast cell stabilisers for treating seasonal and perennial allergic conjunctivitis. *Cochrane Database Syst Rev*. 2015;(6):Cd009566. doi:10.1002/14651858.CD009566.pub2
94. Shiuey Y, Ambati BK, Adamis AP; Viral Conjunctivitis Study Group. A randomized, double-masked trial of topical ketorolac versus artificial tears for treatment of viral conjunctivitis. *Ophthalmology*. 2000;107(8):1512–1517. doi:10.1016/S0161-6420(00)00177-9
95. Lyra AFV, Bastos LC, Lima RCD, Maranhao LDL, Arantes TE. Artificial tears alone versus 0.45% ketorolac tromethamine with artificial tears for the treatment of acute viral conjunctivitis. *Arq Bras Oftalmol*. 2014;77(2):99–102. doi:10.5935/0004-2749.20140025
96. Santiago LA, da Silva JMR, de Azevedo OGR, de Vasconcelos PRL. Comparative study on the efficacy of non-steroidal, steroid and non-use of anti-inflammatory in the treatment of acute epidemic conjunctivitis. *Acta Cirurgica Brasileira*. 2019;34(12):e201901206. doi:10.1590/s0102-865020190120000006

97. Messmer EM. [Bacterial conjunctivitis–diagnosis and therapy update] Bakterielle Konjunktivitis–Update zu Diagnose und Therapie. *Klin Monbl Augenheilkd.* 2012;229(5):529–533. doi:10.1055/s-0031-1299523
98. Sengillo JD, Kunkler AL, Medert C, et al. UV-photokeratitis associated with germicidal lamps purchased during the COVID-19 pandemic. *Ocul Immunol Inflamm.* 2021;29(1):76–80. doi:10.1080/09273948.2020.1834587
99. Choy CKM, Cho P, Boost MV. Cytotoxicity of rigid gas-permeable lens care solutions. *Clin Exp Optomet.* 2013;96(5):467–471. doi:10.1111/cxo.12039
100. Pucker AD. A review of the compatibility of topical artificial tears and rewetting drops with contact lenses. *Contact Lens Anterior Eye.* 2020;43(5):426–432. doi:10.1016/j.clae.2020.04.013
101. Pucker AD, McGwin G, Franklin QX, Dubey J, Nattis A, Lievens C. Application of systane complete for the treatment of contact lens discomfort. *Contact Lens Anterior Eye.* 2021;44(4):101399. doi:10.1016/j.clae.2020.12.004
102. Chou H-D, Chen K-J, Kang EY-C, et al. Eye irrigation as a first-line treatment and diagnostic method for emergency department patients who complain of ocular foreign bodies. *Sci Rep.* 2021;11(1):23386. doi:10.1038/s41598-021-02989-3
103. NICE. Corneal superficial injury. Available from: <https://cks.nice.org.uk/topics/corneal-superficial-injury/>. Accessed July 20, 2022.
104. Barabino S, Rolando M, Nardi M, Bonini S, Aragona P, Traverso CE. The effect of an artificial tear combining hyaluronic acid and tamarind seeds polysaccharide in patients with moderate dry eye syndrome: a new treatment for dry eye. *Eur J Ophthalmol.* 2014;24(2):173–178. doi:10.5301/ejo.5000355
105. Brodwall J, Alme G, Gedde-Dahl S, et al. A comparative study of polyacrylic acid (Viscotears®) liquid gel versus polyvinylalcohol in the treatment of dry eyes. *Acta Ophthalmologica Scandinavica.* 1997;75(4):457–461. doi:10.1111/j.1600-0420.1997.tb00413.x
106. Bron A, Daubas P, Siou-Mermet R, Trinquand C. Comparison of the efficacy and safety of two eye gels in the treatment of dry eyes: lacrinorm and viscotears. *Eye.* 1998;12(5):839–847. doi:10.1038/eye.1998.215
107. Comez AT, Tufan HA, Kocabiyik O, Gencer B. Effects of lubricating agents with different osmolalities on tear osmolarity and other tear function tests in patients with dry eye. *Curr Eye Res.* 2013;38(11):1095–1103. doi:10.3109/02713683.2013.806670
108. Diaz-Llopis M, Dolores Pinazo-Duran M, Diaz-Guionon L, et al. A randomized multicenter study comparing seawater washes and carmellose artificial tears eyedrops in the treatment of dry eye syndrome. *Clin Ophthalmol.* 2019;13:483–490. doi:10.2147/opth.S185409
109. Downie LE, Hom MM, Berdy GJ, et al. An artificial tear containing flaxseed oil for treating dry eye disease: a randomized controlled trial. *Ocul Surf.* 2020;18(1):148–157. doi:10.1016/j.jtos.2019.11.004
110. Dumbleton K, Woods C, Fonn D. An investigation of the efficacy of a novel ocular lubricant. *Eye Contact Lens-Sci Clin Pract.* 2009;35(3):149–155. doi:10.1097/ICL.0b013e3181a2c986
111. Fogt JS, Fogt N, King-Smith PE, Liu HX, Barr JT. Changes in tear lipid layer thickness and symptoms following the use of artificial tears with and without omega-3 fatty acids: a randomized, double-masked, crossover study. *Clin Ophthalmol.* 2019;13:2553–2561. doi:10.2147/OPHT.S228261
112. Fondi K, Wozniak PA, Schmidl D, et al. Effect of hyaluronic acid/trehalose in two different formulations on signs and symptoms in patients with moderate to severe dry eye disease. *J Ophthalmol.* 2018;2018:20184691417. doi:10.1155/2018/4691417
113. Gensheimer WG, Kleinman DM, Gonzalez MO, et al. Novel formulation of glycerin 1% artificial tears extends tear film break-up time compared with systane lubricant eye drops. *J Ocular Pharmacol Therap.* 2012;28(5):473–478. doi:10.1089/jop.2011.0053
114. Jacobi C, Kruse FE, Cursiefen C. Prospective, randomized, controlled comparison of SYSTANE UD eye drops versus VISINE INTENSIV 1% EDO eye drops for the treatment of moderate dry eye. *J Ocular Pharmacol Therap.* 2012;28(6):598–603. doi:10.1089/jop.2012.0066
115. Jerkins G, Greiner JV, Tong L, et al. A comparison of efficacy and safety of two lipid-based lubricant eye drops for the management of evaporative dry eye disease. *Clin Ophthalmol.* 2020;14:1665–1673. doi:10.2147/opth.S256351
116. Khairuddin R, Schmidt KG. Comparative investigation of treatments for evaporative dry eye. *Klin Monbl Augenheilkd.* 2010;227(2):128–134. doi:10.1055/s-0028-1109686
117. Khanal S, Tomlinson A, Pearce EI, Simmons PA. Effect of an oil-in-water emulsion on the tear physiology of patients with mild to moderate dry eye. *Cornea.* 2007;26(2):175–181. doi:10.1097/ICO.0b013e31802b492d
118. Labetoulle M, Schmickler S, Galarreta D, et al. Efficacy and safety of dual-polymer hydroxypropyl guar- and hyaluronic acid-containing lubricant eyedrops for the management of dry-eye disease: a randomized double-masked clinical study. *Clin Ophthalmol.* 2018;12:2499–2508. doi:10.2147/opth.S177176
119. Laihia J, Jarvinen R, Wylegala E, Kaarniranta K. Disease aetiology-based design of multifunctional microemulsion eye drops for moderate or severe dry eye: a randomized, quadruple-masked and active-controlled clinical trial. *Acta Ophthalmol.* 2020;98(3):244–254. doi:10.1111/aos.14252
120. Nelson JD, Farris RL. Sodium hyaluronate and polyvinyl-alcohol artificial tear preparations - a comparison in patients with keratoconjunctivitis sicca. *Arch Ophthalmol.* 1988;106(4):484–487. doi:10.1001/archoph.1988.01060130530029
121. Safarzadeh M, Azizzadeh P, Akbarshahi P. Comparison of the clinical efficacy of preserved and preservative-free hydroxypropyl methylcellulose-dextran-containing eyedrops. *J Optom.* 2017;10(4):258–264. doi:10.1016/j.optom.2016.11.002
122. Szegedi S, Scheschy U, Schmidl D, et al. Effect of single instillation of two hyaluronic acid-based topical lubricants on tear film thickness in patients with dry eye syndrome. *J Ocular Pharmacol Therap.* 2018;34(9):605–611. doi:10.1089/jop.2018.0069
123. Tomlinson A, Madden LC, Simmons PA. Effectiveness of dry eye therapy under conditions of environmental stress. *Curr Eye Res.* 2013;38(2):229–236. doi:10.3109/02713683.2012.757323
124. van Setten GB, Baudouin C, Horwath-Winter J, et al. The HYLAN M study: efficacy of 0.15% high molecular weight hyaluronan fluid in the treatment of severe dry eye disease in a multicenter randomized trial. *J Clin Med.* 2020;9(11):3536. doi:10.3390/jcm9113536
125. Waduthantri S, Yong SS, Tan CH, Htoon HM, Tong L. Lubricant with gelling agent in treating dry eye in adult Chinese patients. *Optomet Vision Sci.* 2012;89(11):1647–1653. doi:10.1097/OPX.0b013e31826cfc41
126. Wang TJ, Wang JJ, Ho JD, Chou HC, Lin SY, Huang MC. Comparison of the clinical effects of carbomer-based lipid-containing gel and hydroxypropyl-guar gel artificial tear formulations in patients with dry eye syndrome: a 4-week, prospective, open-label, randomized, parallel-group, noninferiority study. *Clin Ther.* 2010;32(1):44–52. doi:10.1016/j.clinthera.2010.01.024
127. Liberati A, Altman DG and Tetzlaff J, et al. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*, 339(jul21 1), b2700–b2700. 10.1136/bmj.b2700

Clinical Optometry

Dovepress

Publish your work in this journal

Clinical Optometry is an international, peer-reviewed, open access journal publishing original research, basic science, clinical and epidemiological studies, reviews and evaluations on clinical optometry. All aspects of patient care are addressed within the journal as well as the practice of optometry including economic and business analyses. Basic and clinical research papers are published that cover all aspects of optics, refraction and its application to the theory and practice of optometry. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-optometry-journal>