



# The definition of spine bone mineral density (BMD)-classified osteoporosis and the much inflated prevalence of spine osteoporosis in older Chinese women when using the conventional cutpoint T-score of $-2.5$

Yì Xiáng J. Wáng<sup>^</sup>

Department of Imaging and Interventional Radiology, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong SAR, China

Correspondence to: Dr. Yì Xiáng J. Wáng. Department of Imaging and Interventional Radiology, The Chinese University of Hong Kong, Hong Kong SAR, China. Email: yixiang\_wang@cuhk.edu.hk.

**Keywords:** Bone mineral density; osteoporosis; diagnostic criteria; Chinese women; prevalence

Submitted Sep 15, 2022. Accepted for publication Dec 15, 2022.

doi: 10.21037/atm-22-4559

View this article at: <https://dx.doi.org/10.21037/atm-22-4559>

Osteoporosis is a systemic skeletal disease characterised by a reduction in bone mass and qualitative skeletal changes that cause an increase in bone fragility and a higher fracture risk. The clinical significance of osteoporosis lies in the fractures that occur, and the most important fracture is hip fracture. According to the 1994 World Health Organization (WHO) criteria, the T-score is defined as:  $(BMD_{\text{patient}} - BMD_{\text{young normal mean}}) / SD_{\text{young normal population}}$ , where BMD is bone mineral density and SD is the standard deviation. When the femoral neck (FN) is measured in adult women, a cutpoint value of patient BMD of 2.5 SD below the  $BMD_{\text{young normal mean}}$  satisfies that the prevalence of osteoporosis for those aged  $\geq 50$  years is about 16.2%, the same as the lifetime risk of hip fragility fracture (FF) (1,2). If other sites are also considered, this cutpoint value identifies approximately 30% of postmenopausal women as having osteoporosis, which is approximately equivalent to the lifetime risk of FF at the spine, hip, or forearm. It is widely assumed that this osteoporotic subset of the population loses bone mass at a faster rate, and interventions should ideally begin before an FF occurs.

While some clinics measure only FN BMD, others measure anteroposterior lumbar spine BMD and total hip BMD as well, with the site presenting the lowest T-score


considered for the diagnosis of osteoporosis. However, the 1994 WHO document did not provide a specific definition of spine osteoporosis (1). Though primary osteoporosis is a systematic disease that affects the whole skeleton, it has been demonstrated that FN BMD reduction best predicts hip FF, and spine BMD reduction best predicts vertebral FF (3,4). Therefore, when both FN BMD and spine BMD are measured, it may be reasonably assumed that the clinical endpoint for spine osteoporosis is clinical vertebral FF. It is widely believed that the ratio of clinical vertebral FF to hip FF ratio is roughly 1 (5-7). In the study of Lam *et al.* (6) using data from Manitoba, Canada (2000 to 2007), for the female age group of 55–84 years, there were 2,313 clinical vertebral FFs and 2,395 hip FFs. The younger subjects ( $<70$  years) had a higher prevalence of vertebral FF and older subjects ( $>70$  years) had a higher prevalence of hip FF. The ratio of clinical vertebral FF to hip FF is close to 1 around the age of 70 years (Figure 1) (6,7). However, the reported prevalence of clinical vertebral FF is likely affected by the ease of access to medical care, with populations with easier access reporting a higher prevalence. If we use a T-score of  $-2.5$  as the cutpoint for defining spine osteoporosis in the National Health and Nutrition Examination Survey (NHANES) 2005–2008 data (8), the prevalence of spine

<sup>^</sup> ORCID: 0000-0001-5697-0717.


osteoporosis for older USA Caucasian women  $\geq 50$  years is 15.8% (9), which is a reasonable value because it is similar to the osteoporosis prevalence of the FN BMD definition (considering the ratio of clinical vertebral FF to hip FF is close to 1). Therefore, for Caucasian women, using the same cutpoint osteoporosis T-score (i.e.,  $\leq -2.5$ ) to define both FN and spine osteoporosis is justifiable, although it may not be ideal. Figure 2 shows that, if lumbar spine T-score  $\leq -2.5$  is taken as the cutpoint for defining osteoporosis, in addition to USA Caucasians and blacks (8,9), the data from Canada (10), Australia (11), Southern Italy (12), and Sweden (13) all demonstrate that FN BMD-

classified osteoporosis (FN osteoporosis) prevalence and spine BMD-classified osteoporosis (spine osteoporosis) prevalence are similar. Löfman *et al.* (13) studied women aged 70 years and reported that FN osteoporosis prevalence and spine osteoporosis prevalence were 36% and 32%, respectively. Mautalen *et al.* (14) reported higher spine osteoporosis prevalence for the ‘younger’ group of their study participants and higher FN osteoporosis prevalence for the ‘older’ group of their study participants, consistent with the pattern that the clinical vertebral FF to hip FF ratio is higher than 1 prior to the age of 70 years and lower than 1 after the age of 70 years.


The FF prevalence of older Chinese women is slightly less than half that of Caucasians (9,15). This is the case for hip FF, radiographic vertebral FF (Figure 3) (16), clinical vertebral FF (Figure 4) (17,18), and many other FF sites (9,15). Since the prevalences of vertebral FF and hip FF of older Chinese women are both slightly less than half of those of Caucasians, the clinical vertebral FF to hip FF ratio is also likely to be around 1 at the age of 70 for Chinese women (i.e., the same as for Caucasians), and this is demonstrated in Figure 4. Moreover, in a Korean study, Lim *et al.* (19) analysed female participants from the National Health Insurance Service senior cohort (mean age: approximately 68 years), and reported clinical vertebral FF and hip FF to be 10.7% and 9.6%, respectively (derived from Tab. 3 in reference 19). In a Japanese study of 1,342 postmenopausal women aged 50 years or more with a median follow-up of 15.2 years, Kamiya *et al.* (20) recorded 38 and 35 cases of clinical vertebral FF and hip




**Figure 1** The clinical vertebral FF to hip FF ratio at different age groups. Data from Sweden and Canada women (6,7). FF, fragility fracture.




**Figure 2** Graphic presentation of the ratios of areal DXA FN BMD-classified osteoporosis (FN osteoporosis) prevalence to spine BMD-classified osteoporosis (spine osteoporosis) prevalence, Caucasian and US blacks women results. X-axis: prevalence, with 0.1 denoting a prevalence of 10%. Data include US Caucasians [1], US blacks [2], Canadian [3], Australian [4], Southern Italians [5], and Swedish [6] (8-13). po: community population data. FN, femoral neck; BMD, bone mineral density; NHANES, National Health and Nutrition Examination Survey; DXA, dual-energy X-ray absorptiometry.



**Figure 3** Older Chinese women have a much lower radiographic osteoporotic vertebral fracture prevalence than that of Italian older women. (A) Total spinal deformity index score of four different age groups of Chinese women and Italian women (population-based and age-matched data). The spinal deformity index was calculated with each vertebra assigned a score of 0, 0.5, 1, 1.5, 2, 2.5, and 3 for no ROVD or ROVDs of <20%, 20~25%, ≥25%~1/3, ≥1/3~40%, ≥40%~2/3, and ≥2/3 vertebral height loss, respectively. The spinal deformity index for each subject was calculated by summing the scores of all vertebrae from T4 to L5. The red points indicate Hong Kong subjects, and the green points indicate Italian subjects. Lines denote the linear fit of the four values of the total spinal deformity index for the four age groups, with the slope being steeper for the Italian than for the Chinese data. N=49+49 means there are 49 Chinese subjects and 49 Italian subjects in this age group. (B) A comparison of the distribution of ECF sign positive ROVD among different vertebral levels (103 pairs of age-matched Chinese and Italian older women, population-based data, mean age: 72.4 years). The data in (B) is in addition to the data in (A). (C) A comparison of the distribution of apparent ROVD (i.e., with ≥20% vertebral height loss) among different vertebral levels. There were 122 population-based Chinese subjects and 61 Italian back pain patients (ROVD number for Chinese data was divided by 2 for presentation). X-axis in (B,C): vertebral level of the spine. Data are from Wang *et al.* Arch Osteoporos 2021;16:174, and Wang *et al.*, Arch Osteoporos 2022;17:13. ECF, endplate end/or cortex fracture; HK, Hong Kong; IL, Italian; ROVD/OVD, radiological osteoporotic vertebral deformity (equivalent to radiological vertebral fracture).



**Figure 4** The clinical vertebral FF to hip FF ratio is around 1 at the age of 77 years for Chinese women. In this graph, the values of *a1* and *a2* at age 77 years are similar. (A) Age-specific hip FF incidence among Hong Kong and USA older women according to the study of Ho *et al.* (Am J Public Health 1993;83:694-7). (B) Clinical vertebral FF incidence of older Asian women and Caucasian women. Data are from MsOS (Hong Kong) study; Sakuma *et al.*, J Bone Miner Metab 2008;26:373-8 (Japan Sado 2004), Imai *et al.*, J Bone Miner Metab 2019;37:484-90 (Japan Sado 2015); Kamiya *et al.*, Maturitas 2019;130:13-20 (JPOCS); Sanders *et al.* Osteoporos Int 1999;10:240-7 (Geelong study); Cooper *et al.* J Bone Miner Res 1992;7:221-7 (Rochester study); Fink *et al.* J Bone Miner Res 2005;20:1216-22 (FIT-USA study), and Papaioannou *et al.* Osteoporos Int 2005;16:568-78 (CaMos study). The dotted black and orange lines indicate linear fits for simplicity (ideally it should be an exponential growth fit). FF, fragility fracture.



**Figure 5** Graphical presentation of the ratios of areal DXA FN BMD-classified osteoporosis (FN osteoporosis) prevalence to spine BMD-classified osteoporosis (spine osteoporosis) prevalence, East Asia women results. Data from references (21-34). X-axis: prevalence, with 0.1 denoting a prevalence of 10%. [1]: meta-analysis results; [2]: Hologic device with local BMD reference; [3]: Hologic or GE Lunar devices, with local BMD reference; [4]: GE Lunar device with local BMD reference; [5]: Hologic or GE Lunar devices, with local BMD reference; [6]: Hologic device and its BMD reference; [7]: GE Lunar device with local BMD reference; [8]: Norland device with local BMD reference; [9]: Hologic device and its BMD reference; [10]: Hologic device and its BMD reference (specified for Caucasians); [11]: Lunar DPX or Norland devices and instrument BMD references; [12]: GE Lunar device with local BMD reference; [13]: Hologic device with local BMD reference; [14]: Hologic device with local BMD reference. [9]: rheumatic patients; [10]: total knee arthroplasty patients; [11]: type 2 diabetes mellitus patients. The prevalence ratio of spine osteoporosis to FN osteoporosis is: 1.89 [1], 3.6 [2], 1.91 [3], 2.64 [4], 3.84 [5], 4.8 [6], 1.54 [7], 1.61 [8], 1.84 [9], 3.0 [10], 3.94 [11], 4.66 [12], 2.1 [13], 3.28 [14], respectively. po: community population data; pt: patient data. FN, femoral neck; BMD, bone mineral density; DXA, dual-energy X-ray absorptiometry.

FF respectively. We anticipate that the dual-energy X-ray absorptiometry (DXA) based prevalence of FN osteoporosis in Chinese women will be approximately half of that of Caucasians, and that the prevalence of FN osteoporosis and spine osteoporosis will be similar in Chinese women. However, the reported data show a quite different picture. *Figure 5* shows one result from a meta-analysis of different Chinese female populations, four population-based studies from mainland China, two population-based studies from Taiwan and two population-based studies from Korea, and one each from Hong Kong and Japan (21-34). Except for the meta-analysis result, all other results were randomly selected from the literature. Overall, population-based studies from China, Korea, and Japan tend to report an FN osteoporosis prevalence of >8%, which is more than half of older Caucasian women's rate. Moreover, for all studies in *Figure 5*, the prevalence of spine osteoporosis was substantially higher than that of FN osteoporosis, with spine osteoporosis prevalence exceeding what would be expected

from the variation of the clinical vertebral FF to hip FF ratio across different age bands (5-7,9-14,19,20,35,36).

Based on statistical modeling, we recently proposed that the cutpoint T-score for defining FN osteoporosis should be revised from  $\leq -2.5$  to  $\leq -2.75$  for Hong Kong older women when a local BMD reference range is being applied (9). This lowers the prevalence of FN osteoporosis in women  $\geq 60$  years from 14.7% to 10%. More notably, if we use the Caucasian cutpoint T-score of  $\leq -2.5$  to define spine osteoporosis, the prevalence of spine osteoporosis for Hong Kong women  $\geq 60$  years is as high as 34.8%, which is unreasonable (also see *Figure 2* results for Caucasians). We also proposed that (9) the cutpoint T-score for defining spine osteoporosis should be revised from  $-2.5$  to  $-3.74$  in order to arrive at a prevalence of spine osteoporosis of 10% for Hong Kong women  $\geq 60$  years old using the local BMD reference of Lynn *et al.* (37). In an empirical study on women with radiographic vertebral FF, we recently demonstrated that, at the mean age of 73.5 years and

considering vertebral FF status, an FN T-score of  $-2.56$  for Italian Caucasians is equivalent to  $-2.61$  for Hong Kong Chinese, while a lumbar spine T-score of  $-2.7$  for Italian Caucasians is equivalent to  $-3.54$  for Hong Kong Chinese [n=310 Chinese and Italian pairs (38)]. For the 512 Chinese cases [mean age: 74.0 years, from the MsOS (Hong Kong) study] analysed in that study, 20.5% of the cases had FN osteoporosis, while 41.0% had spine osteoporosis when the conventional cutpoint T-score of  $-2.5$  was used (38).

In conclusion, while a cutpoint T-score  $\leq -2.5$  for defining spine osteoporosis is justified for Caucasian women, for Chinese women the same cutpoint T-score much inflates the estimated prevalence of spine osteoporosis. Despite DXA being an imperfect method for measuring bone strength, in addition to the use of a local BMD reference database, an adjusted cutpoint T-score for defining osteoporosis among elderly Chinese women should be applied, particularly for spine BMD measurements (9,38). Moreover, although this article presents female data, our earlier analysis showed a similarly inflated prevalence of spine osteoporosis in older Chinese men if the cutpoint T-score  $\leq -2.5$  is used (9). We expect that the principles presented here may also apply to osteoporosis estimation in other East Asian populations. For example, it has been suggested that older Korean women and Japanese women have FF risk profiles similar to those of older Chinese women (Figure 4) (5,39-41).

### Acknowledgments

The author thanks Professor Glen Mervyn Blake, King's College London, UK, for his constructive comments for this article.

*Funding:* None.

### Footnote

*Provenance and Peer Review:* This article was a standard submission to the journal. The article has undergone external peer review.

*Conflicts of Interest:* The author has completed the ICMJE uniform disclosure form (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-4559/coif>). The author has no conflicts of interest to declare.

*Ethical Statement:* The author is accountable for all aspects of the work in ensuring that questions related

to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

### References

1. World Health Organization. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis: report of a WHO study group [meeting held in Rome from 22 to 25 June 1992]. 1994. Available online: <https://apps.who.int/iris/handle/10665/39142>
2. Jain RK, Vokes T. Dual-energy X-ray Absorptiometry. *J Clin Densitom* 2017;20:291-303.
3. Fink HA, Harrison SL, Taylor BC, et al. Differences in site-specific fracture risk among older women with discordant results for osteoporosis at hip and spine: study of osteoporotic fractures. *J Clin Densitom* 2008;11:250-9.
4. Alarkawi D, Bliuc D, Nguyen TV, et al. Contribution of Lumbar Spine BMD to Fracture Risk in Individuals With T-Score Discordance. *J Bone Miner Res* 2016;31:274-80.
5. Bow CH, Cheung E, Cheung CL, et al. Ethnic difference of clinical vertebral fracture risk. *Osteoporos Int* 2012;23:879-85.
6. Lam A, Leslie WD, Lix LM, et al. Major osteoporotic to hip fracture ratios in canadian men and women with Swedish comparisons: a population-based analysis. *J Bone Miner Res* 2014;29:1067-73.
7. Kanis JA, Johnell O, Oden A, et al. Long-term risk of osteoporotic fracture in Malmö. *Osteoporos Int* 2000;11:669-74.
8. Looker AC, Borrud LG, Hughes JP, et al. Lumbar spine and proximal femur bone mineral density, bone mineral content, and bone area: United States, 2005-2008. *Vital Health Stat* 11 2012;(251):1-132.
9. Wáng YXJ, Xiao BH. Estimations of bone mineral density defined osteoporosis prevalence and cutpoint T-score for defining osteoporosis among older Chinese population: a framework based on relative fragility fracture risks. *Quant Imaging Med Surg* 2022;12:4346-60.

10. Tenenhouse A, Joseph L, Kreiger N, et al. Estimation of the prevalence of low bone density in Canadian women and men using a population-specific DXA reference standard: the Canadian Multicentre Osteoporosis Study (CaMos). *Osteoporos Int* 2000;11:897-904.
11. Henry MJ, Pasco JA, Nicholson GC, et al. Prevalence of osteoporosis in Australian women: Geelong Osteoporosis Study. *J Clin Densitom* 2000;3:261-8.
12. del Puente A, Heyse SP, Mandes MG, et al. Epidemiology of osteoporosis in women in southern Italy. *Aging (Milano)* 1998;10:53-8.
13. Löfman O, Larsson L, Toss G. Bone mineral density in diagnosis of osteoporosis: reference population, definition of peak bone mass, and measured site determine prevalence. *J Clin Densitom* 2000;3:177-86.
14. Mautalen C, Schianchi A, Sigal D, et al. Prevalence of Osteoporosis in Women in Buenos Aires Based on Bone Mineral Density at the Lumbar Spine and Femur. *J Clin Densitom* 2016;19:471-6.
15. Wáng YXJ. Fragility fracture prevalence among elderly Chinese is no more than half of that of elderly Caucasians. *Quant Imaging Med Surg* 2022;12:874-81.
16. Wáng YXJ, Diacinti D, Leung JCS, et al. Much lower prevalence and severity of radiographic osteoporotic vertebral fracture in elderly Hong Kong Chinese women than in age-matched Rome Caucasian women: a cross-sectional study. *Arch Osteoporos* 2021;16:174.
17. Wáng YXJ, Deng M, Griffith JF, et al. 'Healthier Chinese spine': an update of osteoporotic fractures in men (MrOS) and in women (MsOS) Hong Kong spine radiograph studies. *Quant Imaging Med Surg* 2022;12:2090-105.
18. Hayhoe RPG, Chan R, Skinner J, et al. Fracture Incidence and the Relevance of Dietary and Lifestyle Factors Differ in the United Kingdom and Hong Kong: An International Comparison of Longitudinal Cohort Study Data. *Calcif Tissue Int* 2021;109:563-76.
19. Lim JY, Yoo JI, Kim RB, et al. Comparison of the incidence rates of hip and vertebral fragility fractures according to cataract surgery in elderly population: a nationwide cohort study. *Arch Osteoporos* 2022;17:30.
20. Kamiya K, Kajita E, Tachiki T, et al. Association between hand-grip strength and site-specific risks of major osteoporotic fracture: Results from the Japanese Population-based Osteoporosis Cohort Study. *Maturitas* 2019;130:13-20.
21. Zhang ZQ, Ho SC, Chen ZQ, et al. Reference values of bone mineral density and prevalence of osteoporosis in Chinese adults. *Osteoporos Int* 2014;25:497-507.
22. Wu XP, Liao EY, Huang G, et al. A comparison study of the reference curves of bone mineral density at different skeletal sites in native Chinese, Japanese, and American Caucasian women. *Calcif Tissue Int* 2003;73:122-32.
23. Zhu H, Fang J, Luo X, et al. A survey of bone mineral density of healthy Han adults in China. *Osteoporos Int* 2010;21:765-72.
24. Wang L, Yu W, Yin X, et al. Prevalence of Osteoporosis and Fracture in China: The China Osteoporosis Prevalence Study. *JAMA Netw Open* 2021;4:e2121106.
25. Zeng Q, Li N, Wang Q, et al. The Prevalence of Osteoporosis in China, a Nationwide, Multicenter DXA Survey. *J Bone Miner Res* 2019;34:1789-97.
26. Woo J, Li M, Lau E. Population bone mineral density measurements for Chinese women and men in Hong Kong. *Osteoporos Int* 2001;12:289-95.
27. Lin YC, Pan WH. Bone mineral density in adults in Taiwan: results of the Nutrition and Health Survey in Taiwan 2005-2008 (NAHSIT 2005-2008). *Asia Pac J Clin Nutr* 2011;20:283-91.
28. Chan WP, Liu JF, Chi WL. Evaluation of bone mineral density of the lumbar spine and proximal femur in population-based routine health examinations of healthy Asians. *Acta Radiol* 2004;45:59-64.
29. Zhang X, Dai Z, Lau EHY, et al. Prevalence of bone mineral density loss and potential risk factors for osteopenia and osteoporosis in rheumatic patients in China: logistic regression and random forest analysis. *Ann Transl Med* 2020;8:226.
30. Tao Y, Tang S, Huang X, et al. Prevalence and Risk Factors of Osteoporosis in Chinese Postmenopausal Women Awaiting Total Knee Arthroplasty. *Clin Interv Aging* 2021;16:379-87.
31. Jiajue R, Qi X, Jiang Y, et al. Incident Fracture Risk in Type 2 Diabetic Postmenopausal Women in Mainland China: Peking Vertebral Fracture Study. *Calcif Tissue Int* 2019;105:466-75.
32. Cui LH, Choi JS, Shin MH, et al. Prevalence of osteoporosis and reference data for lumbar spine and hip bone mineral density in a Korean population. *J Bone Miner Metab* 2008;26:609-17.
33. Lee KS, Bae SH, Lee SH, et al. New reference data on bone mineral density and the prevalence of osteoporosis in Korean adults aged 50 years or older: the Korea National Health and Nutrition Examination Survey 2008-2010. *J Korean Med Sci* 2014;29:1514-22.
34. Iki M, Kagamimori S, Kagawa Y, et al. Bone mineral density of the spine, hip and distal forearm in

- representative samples of the Japanese female population: Japanese Population-Based Osteoporosis (JPOS) Study. *Osteoporos Int* 2001;12:529-37.
35. Pedrazzoni M, Girasole G, Bertoldo F, et al. Definition of a population-specific DXA reference standard in Italian women: the Densitometric Italian Normative Study (DINS). *Osteoporos Int* 2003;14:978-82.
  36. Legakis I, Papadopoulos V, Tataridas V, et al. Prevalence of osteoporosis-osteopenia at lumbar spine and femur in an urban female population. An epidemiological study in the Athens Metropolitan area, Greece. *Endocrinology Studies* 2012;2:e1.
  37. Lynn HS, Lau EM, Au B, et al. Bone mineral density reference norms for Hong Kong Chinese. *Osteoporos Int* 2005;16:1663-8.
  38. Wáng YXJ, Diacinti D, Leung JCS, et al. Conversion of osteoporotic vertebral fracture severity score to osteoporosis T-score equivalent status: a framework and a comparative study of Hong Kong Chinese and Rome Caucasian older women. *Arch Osteoporos* 2022;18:1.
  39. Kwok AW, Gong JS, Wang YX, et al. Prevalence and risk factors of radiographic vertebral fractures in elderly Chinese men and women: results of Mr. OS (Hong Kong) and Ms. OS (Hong Kong) studies. *Osteoporos Int* 2013;24:877-85.
  40. Patel MC, Chandra M, Lo JC. Mortality following hip fracture in Chinese, Japanese, and Filipina women. *Am J Manag Care* 2016;22:e358-9.
  41. Lo JC, Chandra M, Lee C, et al. Bone Mineral Density in Older U.S. Filipino, Chinese, Japanese, and White Women. *J Am Geriatr Soc* 2020;68:2656-61.

**Cite this article as:** Wáng YXJ. The definition of spine bone mineral density (BMD)-classified osteoporosis and the much inflated prevalence of spine osteoporosis in older Chinese women when using the conventional cutpoint T-score of -2.5. *Ann Transl Med* 2022;10(24):1421. doi: 10.21037/atm-22-4559