









Paraneoplastic arthritides: an up-to-date case-based systematic review

Suhel Gabriele Al Khayyat¹ , Edoardo Conticini¹ , Paolo Falsetti¹ , Stefano Gentileschi¹ , Antonio Vitale¹, Salvatore Massimo Stella² , Miriana D'Alessandro³ , Elena Bargagli³, Luca Cantarini¹ , Bruno Frediani¹ 

¹Rheumatology Unit, Department of Medicine, Surgery and Neurosciences, University of Siena, Italy

²Advanced Musculoskeletal Ultrasound, SIUMB School of Pisa, Italy

³Respiratory Diseases and Lung Transplantation Unit, Department of Medicine, Surgery and Neurosciences, University of Siena, Italy

Abstract

Among the rheumatic diseases whose symptoms are more often associated with the possibility of cancer and other malignancies are systemic sclerosis, dermatomyositis and rheumatic polymyalgia. However, a differential diagnosis should be performed in each case of non-typical rheumatic disease and/or other neoplastic disease risk factors. The article's aim was based on a literature review of this subject and presentation own a case description and discussion about arthritis as a paraneoplastic syndrome.

The conclusions of our analysis were as follows: more often paraneoplastic arthritis occurs in men, in ages higher than 50 years old, in patients who poorly respond to treatment of arthritis with polyarticular symmetrical involvement of the limbs, seronegative type of inflammatory joint disease. In this group of patients, complete remission after treatment of the primary tumor and recurrence of the symptoms in the presence of metastasis was observed.

Key words: rheumatoid arthritis, psoriatic arthritis, paraneoplastic arthritides, neoplasia.

Introduction

Paraneoplastic syndromes are heterogeneous and difficult to diagnose entities, especially in rheumatological practice where they act as mimickers of commoner clinical conditions. The inflammatory joint manifestations are the result of complex immunological mechanisms given the presence of the neoplasm in the absence of evident tumoral localization in the affected joint.

Herein we present the case of oligo- and seronegative arthritis with hand and atlantoaxial joint involvement in a patient with cutaneous psoriasis subsequently diagnosed with a neuroendocrine tumor of the ileocecal valve.

Furthermore, we perform a systematic review on paraneoplastic arthritis mimicking other rheumatological conditions pointing out the most relevant clinical

criteria to distinguish them from non-paraneoplastic disorders.

Material and methods

We extensively searched via PubMed articles using as key words “arthritis”, “rheumatoid arthritis”, “psoriatic arthritis”, “spondylarthritis” each combined with “tumor”, “cancer”, “paraneoplastic”. Given the rarity of the clinical manifestation, we decided to also include case reports in this review as well.

Studies that were not written in the English language and that were present only in the form of abstracts were not included. We set as reference the latest classification criteria for rheumatoid arthritis (RA) [1], psoriatic arthritis (PSA) [2] and axial spondylarthritis (AxSPA) [3].

Address for correspondence:

Suhel Gabriele Al Khayyat, Rheumatology Unit, Department of Medicine, Surgery and Neurosciences, University of Siena, Viale Mario Bracci, 16 53100 Siena, Italy, e-mail: alkhayatsuhel@gmail.com

Submitted: 04.10.2022; Accepted: 03.11.2022

We included studies with clinical and imaging proven arthritis. We excluded RS3PE, palmar fasciitis and digital clubbing from this review since in the literature are reported numerous evidences that correlate these pathologies to neoplasms [4]. Moreover, papers reporting cases of patients in treatment with checkpoint inhibitors were excluded beforehand. Informed written consent was obtained from the patient.

Results

We found 35 articles, for a total of 134 patients affected by a neoplasm presenting with articular inflammatory manifestations, most frequent type of tumor concomitant to an articular paraneoplastic manifestation was lung adenocarcinoma, followed by hematological malignancies and urinary tract cancers. Patients mean age was 53.6 ±17.23 years old, 60% males and 40% females, mean anti-citrullinated protein antibodies (ACPA) levels 134.8 ±70 IU/ml (6), mean rheumatoid factor (RF) 378.9 ±395.6 IU/ml (6), mean anti-nuclear antibody (ANA) titres 1 : 160 (only 3 positive), mean C-reactive protein (CRP) 8.72 ±7.01 mg/dl, mean erythrocyte sedimentation rate (ESR) 73.06 ±34.43 mm/h.

Moreover, 6 patients had an asymmetrical polyarthritis, 72 a symmetrical polyarthritis, 18 monoarthritis and 35 oligoarthritis, 76.4% displayed a poor or partial response to non-steroidal anti-inflammatory drugs (NSAIDs), glucocorticosteroids (GCs) or conventional synthetic disease-modifying antirheumatic drugs (csDMARDs), 17.6% displayed a good response.

Conversely, 39.47% of patients experienced a full remission after chemotherapy or surgery or both, 21% displayed an improvement of symptoms while 18.4% had a poor response to tumor treatment. In the examined papers only 26 patients had a definitive diagnosis of rheumatic disease: 20 RA, 3 defined as a pseudogout, 2 AxSPA, 1 reactive arthritis. Detailed results are displayed in Tables I-IV.

Case description

The patient was a Caucasian 67-year-old white man referred to our outpatient clinic in March 2021 for stiffness and pain of the right hand. At presentation the wrist appeared to be swollen, fovea sign was elicitable at palpation and the extremity was painful at extension and dorsiflexion. The patient had history of cutaneous psoriasis, hypertension and, recently, had undergone Bentall's surgery.

The following laboratory investigations were required: ESR, CRP, ANA, ACPA, RF, hepato-renal function, uric acid, serum protein electrophoresis, anti-HBV antibodies, anti-HCV, HIV test, QuantiFERON and Mantoux.

Table I. Epidemiological features of paraneoplastic patients' phenotype

Overall patients	Male/ female	Mean age	Mimicker	ACPA	RF	ANA	CRP	ESR	Involvement	Response to AT (NSAIDs/GCs/csDMARDs)	Response to TT
134	81 M (60%) /47 F (35%) 6 unknown (5%)	53.6 ±17.23	20 RA 3 pseudogout 2 AxSPA 1 reactive arthritis	134.8 ±70 IU/ml (6)	378.9 ±395.6 IU/ml (6)	1 : 160 (3)	8.72 ±7.01* mg/dl	73.06 ±34.43*	72 symmetrical polyarthritis 35 oligoarthritis 18 monoarthritis 6 asymmetrical polyarthritis	76.4% partial/ poor 17.6% good 8.8% n.a.	39.47% remission 21% improvement 18.4% poor 18% n.a.

*Approximately.

ACPA – anti-citrullinated protein antibodies, ANA – anti-nuclear antibodies, AxSPA – axial spondylarthritis, AT – arthritis treatment, CRP – C-reactive protein, csDMARDs – conventional synthetic disease-modifying antirheumatic drugs, ESR – erythrocyte sedimentation rate, GCs – glucocorticosteroids, NSAIDs – non-steroidal anti-inflammatory drugs, RA – rheumatoid arthritis, RF – rheumatoid factor, TT – tetanus toxoid.

Table II. Frequency of primary tumors and histotypes associated to an inflammatory articular involvement

Primary tumors	Frequency	Histotypes
Lung cancers	44	NSCLC (26), SCLC (3), epidermoidis (6), mesothelioma (1), bronchogenic (1), unknown (7)
Hematological malignancies	45	Chronic myeloproliferative disease (4), lymphocytic lymphoma (1), multiple myeloma (4), non-Hodgkin lymphoma (8), Hodgkin lymphoma (3), acute myeloblastic leukemia (8), myelodysplastic syndrome (6), acute immunoblastic leukemia (1), lymphoid granulomatosis (1), T-ALL (1), acute lymphoblastic leukemia (4), chronic lymphoblastic leukemia (1), leukemia n.o.s. (1)
Urinary tract cancers	9	Bladder cancer (4), prostatic cancer (3), renal cancer (2)
Gastrointestinal tumors	16	Pancreatic adenocarcinoma (3), gastric cancer (5), cilindric epithelioma (1), lingual squamous cell carcinoma (1), coledoc cancer (1), liver sarcoma (1), colon cancers (3), HCC (1)
Breast cancers	11	Galactophoric adenocarcinoma (1), papillary breast adenocarcinoma (1), breast ductal carcinoma (1), unknown (8)
Reproductive apparatus cancers	4	Penis squamous cell carcinoma (1), ovarian teratoma (1), endometrial carcinoma (1), ovarian cystoadenocarcinoma (1)
Thyroid cancers	1	Papillary carcinoma (1)
Bone cancers	1	Sternal condrosarcoma (1)
Cancer of unknown origin	3	n.a.

HCC – hepatocellular carcinoma, n.a. – not assessed, NSCLC – non-small cell lung cancer, SCLC – small-cell lung cancer, T-ALL – T cell acute lymphoblastic leukemia.




Fig. 1. Magnetic resonance T2-weighted sequences revealing mid-carpal and radio-carpal joint effusion as well as extensor tendons sheaths tenosynovitis of the right wrist.

Moreover, during the first visit, a musculoskeletal ultrasound was performed using a linear probe of 6–18 MHz which highlighted the presence of synovial hypertrophy of the radio-carpal and mid-carpal joints, associated with moderate joint effusion, marked inflammatory pattern at power Doppler (PD) and tenosynovitis of common extensor tendons with an inflammatory pattern at PD. Flexor tendon sheaths appeared within normal

limits and hyperechoic deposits suggestive for crystal arthropathy were absent.

During the first visit an intra-articular injection of the affected wrist was practiced with 20 mg of triamcinolone acetonide and the patient was prescribed with hand X-rays and a therapeutic cycle of methylprednisolone 16 mg/day to be tapered in 1 month.

The laboratory investigations were negative for RF, ACPA and ANA, exception made for ESR and CRP which appeared elevated. In the meantime, the intra-articular injection carried out at the wrist did not produce any improvement, nor did oral GCs therapy.

Hands X-rays were unremarkable for erosions, joint space narrowing or any sign of juxta-articular demineralization, therefore it was decided to subject the patient's limb to a nuclear magnetic resonance (MRI). The magnetic resonance confirmed the presence of severe synovitis of the radiocarpal and midcarpal joints, as well as common extensor tendon sheaths tenosynovitis (Fig. 1).

Psoriatic arthropathy was therefore diagnosed, and given the negativity of screening for occult infectious diseases, methotrexate 10 mg subcutaneously once a week, followed by 5 mg of folic acid within 24–48 hours was started.

No sooner did the patient performed the first methotrexate's syringe, that was admitted to the emergency department for gastrointestinal bleeding and anemia. An abdominal computed tomography (CT) scan was per-

Table III. Clinical and immunological features of paraneoplastic inflammatory manifestations

Author	Study type	Tumor(s)	N	TNM	Arthritis	Involved joints	ACPA	RF	ANA	ENA	Neo-plastic markers	Other signs	Outcome/median survival
Longley et al. 1986 [26]	CS	Myeloproliferative disease	2	n.a.	Asymmetrical polyarthritis	Wrist, MCP, IFJ, knee, ankle	n.a.	Neg.	Neg.	Neg.	n.a.	Cutaneous vasculitis	Death
Lambert, Nuki 1992 [39]	CR	Penis squamous carcinoma	1	n.a.	Symmetrical polyarthritis	Hands, wrist, knees	n.a.	Neg.	n.a.	n.a.	n.a.	Multicentric reticulohistiocytosis	Death
Cohen et al. 1993 [27]	CR	Lymphocytic lymphoma	1	IV	Axial arthritis	Bilateral SIJ, hips, lumbar and cervical spine, sternoclavicular	n.a.	Neg.	Neg.	n.a.	n.a.	Laterocervical lymphadenopathy, nephrolithiasis	n.a.
Drenth et al. 1995 [38]	CR	Breast ductal carcinoma	1	pT2N1M0	Polyarthritis	Knee, ankle, wrist, elbow, and shoulders	n.a.	Neg.	n.a.	n.a.	n.a.	Fever, erythematous rash	Survived
Stummvoll et al. 2001 [5]	CR	SCLC (1), colon adenocarcinoma (2)	2	pT2pN0 (1), SCLC), pT3pN1, G2 (2, colon)	Symmetrical polyarthritis	MCP and IFJ, shoulder, knee (1), joints hands (2)	n.a.	Neg (1)	1:80 (1)	n.a.	n.a.	n.a.	Survived (1)
Lima et al. 2002 [31]	CR	CML	1	n.a.	Polyarthritis	n.a.	n.a.	n.a.	Neg.	n.a.	n.a.	Weak ness, weight loss, diarrhea, myalgias	Death
Glinkov et al. 2003 [6]	CR	HCC	1	n.a.	Polyarthritis	n.a.	n.a.	n.a.	n.a.	n.a.	AFP	Erythema nodosum	Survived
Mok, Kwan 2003 [7]	CR	Adenocarcinoma of unknown origin	1	n.a.	Symmetrical polyarthritis	Shoulders, elbows, wrists, MCPs, PIPs, knee and ankle	n.a.	+	Neg.	n.a.	n.a.	Peritoneal carcinosis, ascites	Death
Wiese et al. 2004 [40]	CR	Ovarian teratoma	1	n.a.	Polyarthritis	Knees, wrists	n.a.	Neg.	Neg.	n.a.	n.a.	Hypersomnolence, fatigability, night sweats, and periodic mouth ulcers	Survived
Ardalan, Shoja 2007 [28]	CR	MM	1	n.a.	Symmetrical polyarthritis	Hands and feet	n.a.	Neg.	Neg.	n.a.	n.a.	AKI	Survived

Table III. Cont.

Author	Study type	Tumor(s)	N	TNM	Arthritis	Involved joints	ACPA	RF	ANA	ENA	Neo-plastic markers	Other signs	Outcome/median survival
Bivalacqua et al. 2007 [21]	CR	NSCLC + bladder cancer	1	T2N0 (Mx?)	Symmetrical polyarthritis	Ankles, knees, hands, elbow, shoulder	n.a.	n.a.	n.a.	n.a.	n.a.	Shortness of breath and wheezing	Survived
Cantini et al. 2007 [16]	CS	NSCLC	5	n.a.	Mononarthrits	Knee	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	Survived
Morel et al. 2008 [15]	CS	solid cancer (20 [13, 60% lung adenocarcinoma]), haematological disease (6)	26	T2 (50%) N1 (46.1%) M0 (80%)	Symmetric polyarthritis	Wrists, hands, knees	n.a.	Neg. (78.3%)	Neg. (70.8%)	n.a.	n.a.	Weight loss (42%), fatigue (46%), fever (27%)	1.2 years
Tedeschi et al. 2017 [29]	CR	MDS	1	n.a.	Asymmetrical oligoarthritis	Left ankle, atlanto-epistropheal, right knee	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	Death
Zupancic et al. 2008 [17]	CR	SCLC	1	n.a.	Asymmetrical polyarthritis	Left ankle, left wrist, shoulder	Neg.	n.a.	Neg.	Neg.	n.a.	n.a.	Death
Bahat et al. 2009 [9]	CS	Ail (1), colon adenocarcinoma (2)	2	n.a. (1) n.a.	Symmetrical polyarthritis	Hands, wrists (1 + 2)	Neg. (1 + 2)	n.a.	n.a.	n.a.	n.a.	Inguinal and axillary lymphadenopathies, splenomegaly (1)	Death (1), survived (2)
Kumar et al. 2009 [8]	CR	Pancreatic adenocarcinoma	1	n.a.	Asymmetrical polyarthritis	Wrist, hand, elbow, shoulder	154.6 IU/ml	419.5 IU/ml	Neg.	n.a.	373 U/ml CA 19.9	Nausea, vomiting, diarrhea, weight loss	Death
Larson et al. 2011 [19]	CR	Lung adenocarcinoma	1	IV	Symmetrical polyarthritis	Elbows, PIPs, MCP, knees	86 IU/ml	199 IU/ml	Neg.	n.a.	n.a.	Cough, shortness of breath, pulmonary embolism	Death
Kobak 2013 [25]	CR	Prostatic cancer	1	n.a.	Mononarthrits	Left ankle	n.a.	Neg.	Neg.	n.a.	PSA	Foot drop	Survived
Raja et al. 2010 [37]	CR	Lymphoid granulomatosis	1	IV	Symmetrical polyarthritis	Knees, ankles, wrists	133 IU/ml	> 250 IU/ml	n.a.	n.a.	n.a.	Maculopapular rash, fever, weight loss	Death

Table III. Cont.

Author	Study type	Tumor(s)	N	TNM	Arthritis	Involved joints	ACPA	RF	ANA	ENA	Neo-plastic markers	Other signs	Outcome/ median survival
Han et al. 2012 [18]	CR	Lung adenocarcinoma	1	n.a.	Symmetrical polyarthritis	Wrists, shoulder, knee, elbows, MCPs	n.a.	Neg.	Neg.	n.a.	CEA 11.9 ng/ml	Cough, fever, anorexia, weight loss	n.a.
Aruch, Mims 2013 [32]	CR	Hodgkin lymphoma	1	3A	Oligoarthritis	Ankle, knee	n.a.	n.a.	n.a.	n.a.	n.a.	Lymphadenopathies, nephrotic syndrome	Survived
Ochi et al. 2012 [10]	CR	Gastric cancer	1	n.a.	Symmetrical polyarthritis	Shoulders, knees, wrists	Neg.	Neg.	n.a.	n.a.	CA 19.9 197.4 U/ml	Weight loss	n.a.
Prashanth et al. 2013 [33]	CR	Leukemia	1	n.a.	Asymmetrical polyarthritis	Left knee, ankles	Neg.	n.a.	Neg.	n.a.	n.a.	n.a.	n.a.
Handy et al. 2015 [34]	CR	T-ALL	1	n.a.	Symmetrical Polyarthritis	Wrists, MCPs, knees, left ankle	242 IU/ml	1:148 IU/ml	1:80	n.a.	n.a.	Fever, tachycardia, tachypnea, adenopathies	Death
Kisacik et al. 2014 [22]	CS	Solid tumors (39), haematological malignancies (26)	65	n.a.	Polyarthritis (22), oligoarthritis (31), monoarthritis (12)	Hands and wrists (28), ankle (35), knee (39)	7/65 ml	15/65	10/65	n.a.	n.a.	n.a.	n.a.
Watson et al. 2015 [20]	CR	Papillary breast carcinoma (1), lung adenocarcinoma (2)	2	n.a. (1), n.a. (2)	Polyarthritis (1+2)	Knee (1), shoulder (1), wrist (1), hands, feets (2)	36 IU/ml (1), neg. (2)	42 IU/ml (1), neg. (2)	n.a. (1+2)	n.a. (1+2)	n.a. (1+2)	n.a. (1+2)	Survived (1+2)
Erlj et al. 2016 [35]	CR	Hodking lymphoma	1	n.a.	Symmetrical polyarthritis	MCP, PJJ	Neg.	Neg.	Neg.	n.a.	n.a.	Membranoproliferative glomerulonephritis + AKI + B symptoms	Survived
Levi Sandri et al. 2016 [13]	CR	Follicular dendritic cell liver sarcoma	1	n.a.	Polyarthritis	Wrists and knees	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.

Table III. Cont.

Author	Study type	Tumor(s)	N	TNM	Arthritis	Involved joints	ACPA	RF	ANA	ENA	Neo-plastic markers	Other signs	Outcome/median survival
Gamage et al. 2018 [36]	CR	DLBCL	1	IV	Polyarthritis	n.a.	Neg.	Neg.	Neg.	Neg.	n.a.	Fever, night sweats, weight loss, generalized lymphadenopathy	Survived
Iqbal et al. 2018 [30]	CR	MDS	1	n.a.	Polyarthritis	Left wrist, knee	Neg.	Neg.	Neg.	n.a.	n.a.	n.a.	Survived
Eidenschink et al. 2019 [24]	CR	Mesothelioma	1	IV	Oligoarthritis	Wrist, shoulder, knee	Neg.	Neg.	Neg.	Neg.	n.a.	Fever, hypertension, tachypnoea	n.a.
Briones-Figueroa et al. 2019 [14]	CR	Lingual squamous cell carcinoma	1	n.a.	Symmetrical polyarthritis	Wrists, proximal interphalangeal joints, knees and elbows	Neg.	Neg.	Neg.	n.a.	Neg.	Asthenia, hyporexia, and weight loss	Survived
Rabah et al. 2020 [11]	CR	Gastric cancer	1	T3N1Mx	Symmetrical polyarthritis	Hands, elbows, feet	Neg.	Neg.	n.a.	n.a.	n.a.	Weight loss, fatigue	n.a.
Sachdev Manjit Singh et al. 2021 [23]	CR	Lung adenocarcinoma (1), ovarian cystadenocarcinoma (2)	2	n.a. (1), IIc (2)	Asymmetrical oligoarthritis (1+2)	Ankles (1), wrist (2, 3), MCP (1-5) PIP (2)	n.a. (1)	Neg. (1+2)	1:320 (1), 1:160 (2)	n.a. (1)	n.a. (1)	Weight loss and anorexia(2)	n.a. (1), death (2)
Silvêrio-Antonio et al. 2021 [12]	CR	Gastric cancer	1	T3N2M0, G3	Symmetric polyarthritis	PIP, MCP, wrists, hands	156.9 IU/ml	215 IU/ml	n.a.	n.a.	n.a.	Vomiting, progressive fatigue, and weight loss	18 months

ACPA – anti-citrullinated protein antibodies, AFP – α -fetoprotein, AKI – acute kidney injury, AIL – aintegumenta-like, ANA – anti-nuclear antibody, CAE – caprine arthritis-encephalitis syndrome, CML – chronic myeloid leukemia, DLBCL – diffuse large B-cell lymphoma, ENA – 2'-O-4' C-Ethylene-bridged nucleic acid, HCC – hepatocellular carcinoma, n.a. – not applicable, MCP – metacarpophalangeal, MDS – myelodysplastic syndrome, Neg. – negative, NSCLC – non-small cell lung cancer, PJI – proximal interphalangeal joint, PIP – proximal interphalangeal, RF – rheumatoid factor, SCLC – small-cell lung cancer, T-ALL – T cell acute lymphoblastic leukemia, TNM – tumor-node-metastasis.

Table IV. Clinical, hematological and anthropological features of paraneoplastic arthritides

Author	CRP	ESR	Mimicker	Age	Arthritis treatment	Response to AT	Tumor medical treatment	Tumor surgical treatment	Response of articular manifestations to tumor treatments
Longley et al. 1986 [26]	n.a.	n.a.	n.a.	49 (g.u.)	NSAIDs, GCs	Partial	n.a.	n.a.	n.a.
Lambert, Nuki 1992 [39]	n.a.	43	RA	63 M	NSAIDs, cyclophosphamide	Partial	CHT (nos)	Penis amputation	n.a.
Cohen et al. 1993 [27]	n.a.	73	AxSPA	61 F	Piroxicam, SSZ, prednisone, AZT	n.a.	n.a.	n.a.	n.a.
Drenth et al. 1995 [38]	n.a.	135	RA	49 F	Naproxen, indometacin	Poor	Oral cyclophosphamide (100 mg/m ² days 1–14), 5-fluorouracil (600 mg/m ² days 1 and 8), methotrexate (40 mg/m ² days 1 and 8), and prednisone (30 mg/day)	n.p.	Remission after 3 months of CHT
Stummvoll et al. 2001 [5]	4.6 (1), 2.9 (2)	53 (1), 35 (2)	RA (1)	60 M	Diclofenac (1), Diclofenac (2)	Poor (1) Partial (2)	Cisplatin and etoposide soldesam, 5HT3 inhibitors (1), 5-fluorouracil and leucovorin (2)	Lobectomy and lymphadenectomy (1), surgery (2)	Response after chemotherapy (1), response after surgery (2)
Lima et al. 2002 [31]	n.a.	n.a.	n.a.	35 F	GCs	Poor	Cyclophosphamide, doxorubicin, vincristine, and prednisone	n.ap.	Improvement
Glinkov et al. 2003 [6]	n.a.	n.a.	n.a.	23 F	GCs, NSAIDs	Poor	n.a.	Resectomy	Full remission
Mok, Kwan 2003 [7]	2.2	80	RA	69 F	NSAIDs, SSZ, GCs	Poor to NSAIDs, partial to SSZ and GCs	n.p.	n.p.	Partial improvement
Wiese et al. 2004 [40]	n.a.	26	Reactive arthritis	34 F	NSAIDs	Poor	n.p.	Resectomy	Full remission
Ardalan, Shoja 2007 [28]	n.a.	53	RA	47 M	MPPT, PDN	Poor	VAD	n.a.	Full remission

Table IV. Cont.

Author	CRP	ESR	Mimicker	Age	Arthritis treatment	Response to AT	Tumor medical treatment	Tumor surgical treatment	Response of articular manifestations to tumor treatments
Bivalacqua et al. 2007 [21]	n.a.	n.a.	RA	52 M	n.a.	n.a.	n.a.	Lobectomy + cystectomy	Full remission
Cantini et al. 2007 [16]	n.a.	n.a.	n.a.	57 M	NSAIDs	Poor	n.a.	Resectomy	Full remission
Morel et al. 2008 [15]	↑	n.a.	RA	57 (16 M/10 F)	NSAIDs, GCs, csDMARDs	NSAIDs – good (45%) GCs – excellent (91%) csDMARDs (poor)	n.a.	n.a.	n.a.
Tedeschi et al. 2017 [29]	19.1	115	Pseudo-gout	75 M	Naproxen, MPPT colchicine, GCs injections	Partial	Azacitidine	n.a.	Poor
Zupancic et al. 2008 [17]	1.5	106	n.a.	43 M	Naproxen, prednisone	Poor	Cisplatin + etoposide	n.p.	Good
Bahat et al. 2009 [9]	n.a.	n.a.	RA	72 M (1), 63 F (2)	NSAIDs, PDN, MTX, plaquenil (1)	Good (1), poor (2)	COP (1), 5-FU (2)	n.a. (1), palliative (2)	Partial (1), good (2)
Kumar et al. 2009 [8]	19.1	115	RA	58 M	Naproxen, MPPT, colchicine, GCs injections	Partial	Azacitidine	n.a.	Poor
Larson et al. 2011 [19]	3.4	35	RA	45 F	Ibuprofen, GCs	Poor	n.a.	n.a.	n.a.
Kobak 2013 [25]	1.5	106	n.a.	43 M	Naproxen, prednisone	Poor	Cisplatin + etoposide	n.p.	Good
Raja et al. 2010 [37]	n.a.	n.a.	RA	40 F	NSAIDs, PDN, MTX, plaquenil (1)	Good (1), poor (2)	COP (1), 5-FU (2)	n.a. (1), palliative (2)	Partial (1), good (2)
Han et al. 2012 [18]	Normal	44	RA	55 F	n.a.	n.a.	Gefitinib	n.a.	Good
Aruch et al. 2013 [32]	n.a.	n.a.	n.a.	38 M	Dexamethasone	Poor	ABVD	n.a.	Full remission
Ochi et al. 2012 [10]	15.6	104	RA	71 M	n.a.	n.a.	n.a.	Total gastrectomy	Full remission after surgery
Prashanth et al. 2013 [33]	Normal	Normal	n.c.	10 M	Aspirin, naproxen, MPPT	Poor	n.a.	n.a.	Full remission with therapy

Table IV. Cont.

Author	CRP	ESR	Mimicker	Age	Arthritis treatment	Response to AT	Tumor medical treatment	Tumor surgical treatment	Response of articular manifestations to tumor treatments
Handy et al. 2015 [34]	7.1	122	RA	61 F	NSAIDs, GCs	Poor	CYC, vincristine, doxorubicin, and dexamethasone	n.ap.	Poor
Kisacik et al. 2014 [22]	65.1 ±85.5	58.5 ±31.9	RA	58.02 ±15.3	n.a.	n.a.	n.a.	n.a.	n.a.
Watson et al. 2015 [20]	11.8 (1), 1.2 (2)	21 (1), n.a. (2)	RA (1 + 2)	80 F (1), 71 F (2)	NSAIDs, GCs	Poor to NSAIDs, good to MPPT (1), poor to NSAIDs and GCs (2)	RT and CHT nos (1)	n.a. (1), lobectomy (2)	Remission (1), Remission after surgery (2)
Erlj et al. 2016 [35]		110	RA	46 M	MPPT, prednisolone	Partial	Doxorubicin, bleomycin, vinblastine and dacarbazine	n.ap.	Full remission
Levi Sandri et al. 2016 [13]	n.a.	n.a.	RA	19 M	n.a.	n.a.	n.a.	n.a.	n.a.
Gamage et al. 2018 [36]	↑ (n.r.)	↑ (n.r.)	RA	45 M	NSAIDs, colchicine	Poor	RCHOP	n.i.	Full remission
Iqbal et al. 2018 [30]	↑ (n.r.)	↑ (n.r.)	Pseudo-gout	83 M	GCs	Good	Azacitidine	n.ap.	Full remission
Eidenschink et al. 2019 [24]	15.6	69	n.a.	73 M	Naproxen	Partial	n.a.	n.a.	Improvement
Briones-Figueroa et al. 2019 [14]	9.5	51	n.a.	69 M	GCs (prednisone)	Poor	RT and CHT	n.p.	Improvement
Rabah et al. 2020 [11]	10.43	120	RA	83 M	NSAIDs, PDN	Poor	n.a.	n.a.	n.a.
Sachdev Manjith Singh et al. 2021 [23]	n.a. (1)	61 (1), raised (2)	SPA (1), RA (2)	65 F (1), 64 M (2)	PDN, SSZ (1 + 2)	Poor (1)	Gefitinib (1), carboplatin + paclitaxel + gemcitabine (2)	n.p.	Improvement (1), n.a. (2)
Silvério-Antônio et al. 2021 [12]	17.3	94	RA	64 M	Dexamethasone, HCQ, PDN	Good	Cisplatin, 5-FU, trastuzumab	n.p.	Poor

ABVD – doxorubicin, bleomycin, vinblastine, and dacarbazine, CHT – choline transporter, CYC – cyclophosphamide, GCs – glucocorticosteroids, F – female, HCQ – hydroxychloroquine, M – male, MPPT – methylprednisolone, MTX – methotrexate, n.a. – not assessed, n.p. – not performed, NSAIDs – non-steroidal anti-inflammatory drugs, PDN – prednisone, RA – rheumatoid arthritis, RCHOP – rituximab, cyclophosphamide, vincristine, doxorubicin, and prednisolone, SPA – spondyloarthritis, SSZ – salazopyrin, 5-FU – fluoropyrimidine 5-fluorouracil.

formed and revealed the presence of a solid, space-occupying lesion at the ileocecal valve with suspicious hepatic metastasis.

Therefore immunosuppressive therapy was interrupted, the patient underwent intestinal resection, from the subsequent and the histopathological analysis of the lesion the specimens revealed the presence of a neuroendocrine tumor (NET) with glandular pattern expressing chromogranin, synaptophysin, neuron-specific enolase (NSE), with a Ki-67 of 2% (G1 stage according to WHO) and a pathological staging of pT3pN2aM0. Serological chromogranin and NSE at time of the surgery were respectively 592 ng/ml (normal values < 109 ng/ml) and 24.3 (normal values < 15).

The patient subsequently returned to our attention, reporting a clear improvement of symptoms: clinical and ultrasound examination were at this time negative for any synovitis sign. Therefore the patient was entrusted to the care of the oncologist who prescribed somatostatin analogues (once monthly) and scheduled the follow-up of the other heterotopic lesions.

Three months later the patient returned to the emergency-urgency department complaining of cervical-brachialgia, thus we decided to perform a CT study of the cervical spine that subsequently revealed the presence of lesions suspicious for bone secondarisms at vertebral level.



Fig. 2. Magnetic resonance T2-weighted images showing BME of C1–C5 as well as synovitis of C1–C2.

From the subsequent MRI study, the presence of osteolytic lesions was excluded, however the presence of edema and a slight slipping of the axial tooth was evidenced, consistent with the presence of synovitis and bone marrow edema (BME) at the level of the atlantoaxial joint and extending to the anterior surface of vertebrae from C1 to C5 (Fig. 2).

As soon as the inflammatory arthropathy affecting the atlantoaxial joint was identified, a semi-rigid collar was recommended, and GCs bolus therapy was scheduled. Nevertheless, the patient denied the intravenous bolus therapy with GCs. The subsequent PET/TC DOTATATE did not reveal any new anomalous uptake; therefore, he has been currently attaining to the follow up schedule and sticking to the oncologist therapy.

The latest head-neck MRI after 3 months revealed a spontaneous reduction of BME at atlantoaxial and cervical level. Last serological evaluation of chromogranin documented a significant reduction of chromogranin levels (134 ng/ml).

Discussion

According to results and conclusions of searching of literature which we described above and our case description we may discuss some of define cancers associated with rheumatic diseases.

Gastrointestinal cancers

Stumvoll et al. [5] was the first to report the case report of a patient presenting with symmetrical polyarthritis subsequently diagnosed with a colon adenocarcinoma. Several years later Glinkov et al. [6] described the case of a young women with a two year history erythema nodosum and polyarthritis subsequently diagnosed with an hepatocellular carcinoma.

Mok et Kwan [7] in 2003 described the case of a patients developing a severe symmetrical polyarthritis with RF positivity in a patient soon after diagnosed with peritoneal carcinomatosis; subsequent investigations revealed the presence of a space occupying lesion in the liver, however a clearer diagnosis was not obtained due to the unwillingness of the patients to perdue other invasive maneuvers.

Subsequently, Kumar et al. [8] described the case of a patient presenting with asymmetric polyarthritis and elevated RF with an underlying pancreatic adenocarcinoma. Bahat et al. [9] the same year presented the case of 3 patients (2 with colon adenocarcinoma and 1 with acute immunoblastic leukemia – ALL) with a symmetric polyarthritis and negative RF as well as ACPA, mimicking a seronegative RA.

Ochi et al. [10], Rabah et al. [11], Silvêrio-Antônio et al. [12] described patients affected by gastric cancers developing a symmetrical polyarthritis mimicking a traditional rheumatoid pattern. To date, Levi Sandri et al. [13] are the only group of study who have described an RA-like polyarthritis in a 19 year old boy diagnosed with a follicular dendritic cell sarcoma of the liver.

Finally, Briones-Figueroa et al. [14] in 2019 described the case of a 69 years old man presenting with a symmetrical paraneoplastic arthritis associated with weight loss in a patient subsequently with a squamous cell carcinoma of the tongue.

Pulmonary cancers

From the results of our review (Table II) pulmonary cancers appear as the main kind of tumors linked to the subsequent development of an inflammatory articular disease; histological specimens reveal that the most frequent type of neoplasia is non-small cell lung cancer (NSCLC).

Morel et al. [15] reported 26 patients affected by solid neoplasia presenting with a symmetrical polyarthritis mimicking RA, most of them were affected by lung cancer. Subsequently, the group of study of Cantini et al. [16] in 2007 reported the case of 5 patients affected by NSCLC presenting with a paraneoplastic knee monoarthritis.

The year later Zupancic et al. [17] described the case of an asymmetrical polyarthritis occurring in a patient subsequently diagnosed with small-cell lung cancer (SCLC). Han et al. [18] in 2011, Larson et al. [19], Watson et al. [20] and Bivalacqua et al. [21] in different case reports described the case of patients developing symmetrical polyarthritis mimicking RA subsequently diagnosed with a lung adenocarcinoma.

Kisacik et al. [22] in 2014, presenting the largest case series related to this topic, described 65 paraneoplastic arthritis associated to lung adenocarcinoma as the most frequent type of solid tumor linked to an articular inflammatory involvement.

Sachdev Manjit Singh et al. [23] in 2021 described one of the few cases of paraneoplastic arthritis mimicking an AxSPA in a patient subsequently diagnosed with lung adenocarcinoma and the case of a seronegative symmetrical polyarthritis mimicking RA subsequently diagnosed with ovarian cystoadenocarcinoma.

The only case of a patients diagnosed with pleural mesothelioma developing a paraneoplastic asymmetrical oligoarthritis is described by Eidenschink et al. [24] in 2019.

Urinary tract cancers

The only patients with a prostatic cancer presenting with a paraneoplastic arthritis is described by Kobak

[25]. Other patients presenting with articular inflammatory manifestation relatable to the presence of a neoplasia are reported in the case series of Morel et al. [15] and Kisacik et al. [22].

Hematological malignancies

Hematological malignancies are the most frequent type of tumors associated to an inflammatory articular disease after lung cancer. Main kind of hematological diseases held responsible for this paraneoplastic manifestation are: non-Hodgkin lymphoma (NHL) (8), acute myeloblastic leukemia (AML) (8) and multiple myeloma (MM) (4).

Since Longley et al. [26] who in 1986 were the first authors to diagnose a myeloproliferative disorder in 2 patients with cutaneous angiitis and polyarthritis, many literature evidence has been collected in these years; Cohen et al. 1993 [27] described one of the few cases reported in literature of an AxSPA manifesting as a paraneoplastic syndrome concomitant to a well differentiated lymphocytic lymphoma.

Then, Ardalan et Shoja [28] in 2007 presented the case of a patient with acute kidney injury (AKI) and RA-like symmetrical polyarthritis subsequently diagnosed with MM.

Tedeschi et al. [29] and Iqbal et al. [30] were the only authors to report a case of paraneoplastic arthritis mimicking pseudogout correlated with an underlying myelodysplastic syndrome. Lima et al. [31] described the case of a paraneoplastic polyarthritis, polymyositis and hypercalcemia developing in a patient an acute lymphoblastic leukemia on a chronic myeloid leukemia (CML).

Aruch et Mims [32] in 2013 described the case of a patient presenting with nephrotic syndrome, Hodgkin lymphoma and oligoarthritis. The year later Prashanth et al. [33] described the case of a child presenting with polyarthritis and acute leukemia (not otherwise specified).

In one of the largest case series ever published on the topic, Kisacik et al. [22] in his case series reported 26 hematological malignancies; in this study, the most frequent hematological disease linked to an articular inflammatory involvement was the AML.

Concomitantly, Handy et al. [34] described the only case of a RA-like polyarthritis developing in a patient with a T cell ALL. Furthermore, Erlij et al. [35], reported one of the few cases of Hodgkin lymphoma held responsible for articular paraneoplastic manifestations.

The latest report found in literature lay to Gamage et al. [36] who described the case of paraneoplastic polyarthritis in a patient with diffuse large B cell lymphoma, and Raja et al. [37] who described the only case of a lymphoid granulomatosis responsible of articular paraneoplastic involvement.

Breast cancers

The only two case reports documenting a paraneoplastic polyarthritis in women diagnosed with breast cancer are of Drenth et al. [38] in 1995 and Watson et al. [20] several years later: both documented a paraneoplastic symmetrical polyarthritis mimicking RA in patients with a breast ductal carcinoma.

The largest amount of evidences correlating mammary cancers to inflammatory articular involvement are reported in the case series of Morel et al. [15] and Kisacik et al. [22].

Other neoplastic arthritis

To the best of our knowledge Lambert et Nuki [39] in 1992 was the only to report the case of a patient with a multicentric reticulohistiocytosis presenting concomitantly to a symmetrical polyarthritis affecting hands, feet and wrists.

Wiese et al. [40] in 2004 and subsequently Sachdev Manjit Singh et al. [23] in 2021 are the only authors to have described an articular inflammatory involvement associated to the presence of ovarian tumors: the former identified a patient with an ovarian teratoma, the latter one described a subject with cystadenocarcinoma.

Moreover, in the case series of Kisacik et al. [22] are reported few cases of reproductive apparatus tumors correlated with paraneoplastic articular manifestations.

Paraneoplastic arthritis are an heterogenous group of disorders challenging to rule out in everyday clinical practice since there is a lack of well-structured guidelines to assist the physician during the diagnostic process. Physiopathology of paraneoplastic arthritides remains poorly understood and it seems to involve formation of immunocomplexes and a T cell participation [4].

From the results of our systematic review, we identified that the most frequent phenotype of a paraneoplastic arthritis is represented by a middle-aged man (> 50 years old) with a seronegative symmetrical polyarthritis mimicking RA, often related to the presence of an underlying lung cancer.

That lung tumors are frequently linked to paraneoplastic syndromes is not a novelty in literature and probably this is given to a higher immunogenicity created by the neoplastic epithelial invasion and tumoral microenvironment; however, an interesting observation is that despite the high prevalence of cancers worldwide, very few seems relatable to the presence of paraneoplastic inflammatory joint involvement.

Moreover, rheumatological diseases display a higher prevalence in female gender, conversely in our review we documented that paraneoplastic articular involve-

ment occur more frequently in male subjects. In line with what is already reported, we found that majority of patients affected by a paraneoplastic arthritis did not respond to GCs, NSAIDs and csDMARDs.

Furthermore, in 39.4% of those treated with chemotherapy or surgery, thus reducing or eliminating neoplastic burden, full remission of articular inflammation was achieved. These observations allow to strengthen the hypothesis that paraneoplastic arthritis is immunologically correlated with neoplasm and that tumoral control may mitigate paraneoplastic symptoms.

In our clinical experience, the scarce responsiveness to GCs therapy and the disappearance of inflammatory joint involvement after the gastrointestinal neuroendocrine tumors (GI-NET) removal allowed us to suspect the articular symptoms as being of paraneoplastic nature.

An interesting paper of Hagiwara et al. [41] studied the prevalence of paraneoplastic arthritis in a cohort of patients classifiable as psoriatic according to CASPAR criteria and demonstrated that 19 out of 115 patients had developed arthritis shortly before the tumor was discovered.

Given the challenging clinical assessment in diagnosis of paraneoplastic arthritis, evaluation of neoplastic markers may be of some utility, in fact our patient had abnormally elevated chromogranin levels which dropped near normality after the excision of the primary tumor.

According to the paper of Parperis et al. [42], which proposed an interesting diagnostic flow chart, neoplastic markers are involved during the clinical investigations, nevertheless their research should be done clearly bearing in mind what the physician is trying to rule out.

After several weeks from the GI-NET removal our patient developed a severe cervical-brachialgia which brought to the diagnosis of C1–C2 synovitis with bone marrow edema extending to C5, a condition ever previously described as being of paraneoplastic nature. The involvement of the axial tooth is a typical and rare manifestation of RA which hardly fit the history of our patient.

Study limitations

Main limit of this study lay in the high number of case series and case report included, given the rarity of the disorder.

Conclusions

To date, this is the most up-to-date review regarding paraneoplastic arthritis; bullet points that we found to correctly identify a paraneoplastic disorder are:

- lack of/poor response to NSAIDs, GCs and csDMARD therapy,
- polyarticular symmetrical involvement of limbs,

- absence of RF and ACPA,
- complete remission of paraneoplastic symptoms after treatment of primary tumor,
- recurrence of the symptoms in presence of metastasis,
- age higher than 50 years old,
- major involvement of male subjects.

The authors declare no conflict of interest.

References

- Aletaha D, Neogi T, Silman AJ, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum* 2010; 62: 2569–2581, DOI: 10.1002/art.27584.
- Taylor W, Gladman D, Helliwell P, et al. Classification criteria for psoriatic arthritis: development of new criteria from a large international study. *Arthritis Rheum* 2006; 54: 2665–2673, DOI: 10.1002/art.21972.
- Rudwaleit M, van Der Heijde D, Landewé R, et al the development of Assessment of SpondyloArthritis international Society classification criteria for axial spondyloarthritis (part II): validation and final selection. *Ann Rheum Dis* 2009; 68: 777–783, DOI: 10.1136/ard.2009.108233.
- Manger B, Schett G. Paraneoplastic syndromes in rheumatology. *Nat Rev Rheumatol* 2014; 10: 662–670, DOI: 10.1038/nrrheum.2014.138.
- Stummvoll GH, Aringer M, Machold KP, et al. Cancer polyarthritides resembling rheumatoid arthritis as a first sign of hidden neoplasms. *Scand J Rheumatol* 2001; 30: 40–44, DOI: 10.1080/030097401750065319.
- Glinkov S, Krasnaliev I, Atanassova M, et al. Hepatocellular carcinoma associated with paraneoplastic erythema nodosum and polyarthritides. *J Hepatol* 2003; 39: 656–657, DOI: 10.1016/s0168-8278(03)00248-4.
- Mok CC, Kwan YK. Rheumatoid-like polyarthritides as a presenting feature of metastatic carcinoma: a case presentation and review of the literature. *Clin Rheumatol* 2003; 22: 353–354, DOI: 10.1007/s10067-003-0741-2.
- Kumar S, Sethi S, Irani F, Bode BY. Anticyclic citrullinated peptide antibody-positive paraneoplastic polyarthritides in a patient with metastatic pancreatic cancer. *Am J Med Sci* 2009; 338: 511–512, DOI: 10.1097/MAJ.0b013e3181b0babe.
- Bahat G, Kamali S, Saka B, et al. Paraneoplastic arthritis may mimic rheumatoid arthritis with symmetrical and upper extremity predilecting presentation. *J Clin Rheumatol* 2009; 15: 319–320, DOI: 10.1097/RHU.0b013e3181b18ebd.
- Ochi K, Horiuchi Y, Seki M, et al. Polyarthritides and posterior interosseous nerve palsy associated with gastric carcinoma. *Rheumatol Int* 2012; 32: 2557–2559, DOI: 10.1007/s00296-011-2047-z.
- Rabah S, Bani Hani D, Jilani N. Paraneoplastic arthritis in a patient with gastric cancer: a case report. *Cureus* 2020; 12: 10–13, DOI: 10.7759/cureus.9692.
- Silvério-Antônio M, Parlato F, Martins P, et al. Gastric adenocarcinoma presenting as a rheumatoid factor and anti-cyclic citrullinated protein antibody-positive polyarthritides: a case report and review of literature. *Front Med (Lausanne)* 2021; 8: 627004, DOI: 10.3389/fmed.2021.627004.
- Levi Sandri GB, Colasanti M, Vennarecci G, Ettore GM. Paraneoplastic arthritis as first symptom of a liver inflammatory pseudotumor-like follicular dendritic cell sarcoma. *Liver Int* 2016; 36: 1392, DOI: 10.1111/liv.13148.
- Briones-Figueroa A, Sifuentes-Giraldo WA, Carrillo-Gijón R, Morell-Hita JL. Paraneoplastic polyarthritides as the first manifestation of lingual carcinoma. *Eur J Rheumatol* 2019; 6: 52–53, DOI: 10.5152/eurjrheum.2018.18068.
- Morel J, Deschamps V, Toussiroit E, et al. Characteristics and survival of 26 patients with paraneoplastic arthritis. *Ann Rheum Dis* 2008; 67: 244–247, DOI: 10.1136/ard.2007.070086.
- Cantini F, Niccoli L, Nannini C, et al. Isolated knee monoarthritides heralding resectable non-small-cell lung cancer. a paraneoplastic syndrome not previously described. *Ann Rheum Dis* 2007; 66: 1672–1674, DOI: 10.1136/ard.2007.075333.
- Zupancic M, Annamalai A, Brenneman J, Ranatunga S. Migratory polyarthritides as a paraneoplastic syndrome. *J Gen Intern Med* 2008; 23: 2136–2139, DOI: 10.1007/S11606-008-0794-7.
- Han YM, Fang LZ, Zhang XH, et al. Polyarthritides as a prewarning sign of occult lung cancer. *Kaohsiung J Med Sci* 2012; 28: 54–56, DOI: 10.1016/j.kjms.2011.06.035.
- Larson E, Etwaru D, Siva C, Lawlor K. Report of anti-CCP antibody positive paraneoplastic polyarthritides and review of the literature. *Rheumatol Int* 2011; 31: 1635–1638, DOI: 10.1007/s00296-009-1294-8.
- Watson GA, O'Neill L, Law R, et al. Case report migrating polyarthritides as a feature of occult malignancy: 2 case reports and a review of the literature. *Case Rep Oncol Med* 2015; 2015: 934039, DOI: 10.1155/2015/934039.
- Bivalacqua TJ, Alphs H, Aksentijevich I. Paraneoplastic polyarthritides from non-small-cell lung cancer metastatic to the bladder. *J Clin Oncol* 2007; 26: 2621–2623, DOI: 10.1200/JCO.2007.11.5600.
- Kisacik B, Onat AM, Kasifoglu T, et al. Diagnostic dilemma of paraneoplastic arthritis: case series. *Int J Rheum Dis* 2014; 17: 640–645, DOI: 10.1111/1756-185X.12277.
- Sachdev Manjit Singh B, Wan SA, Cheong YK, et al. Arthritis as an initial presentation of malignancy: two case reports. *J Med Case Rep* 2021; 15: 1–5, DOI: 10.1186/s13256-020-02642-z.
- Eidenschink M, Beard A, Ewart D. Paraneoplastic migratory oligoarthritides in a patient with malignant mesothelioma. *Am J Med* 2019; 132: e801–e802, DOI: 10.1016/j.amjmed.2019.05.020.
- Kobak S. Chronic monoarthritides and foot-drop as a paraneoplastic syndrome in prostate cancer. *Rheumatol Int* 2013; 33: 223–225, DOI: 10.1007/s00296-010-1564-5.
- Longley S, Caldwell JR, Panush RS. Paraneoplastic vasculitides. Unique syndrome of cutaneous angiitides and arthritis associated with myeloproliferative disorders. *Am J Med* 1986; 80: 1027–1030, DOI: 10.1016/0002-9343(86)90660-1.
- Cohen MR, Carrera GE, Lundberg J. Rapidly progressive sacroiliitides in a patient with lymphocytic lymphoma. *Ann Rheum Dis* 1993; 52: 239–240, DOI: 10.1136/ard.52.3.239.

28. Ardalan MR, Shoja MM. Multiple myeloma presented as acute interstitial nephritis and rheumatoid arthritis-like polyarthritis. *Am J Hematol* 2007; 82: 309–313, DOI: 10.1002/ajh.20796.
29. Tedeschi SK, Stone RM, Helfgott SM. Calcium pyrophosphate crystal inflammatory arthritis (pseudogout) with myelodysplastic syndrome: a new paraneoplastic syndrome? *J Rheumatol* 2017; 44: 1101–1102, DOI: 10.3899/jrheum.170106.
30. Iqbal SM, Aslam HM, Faizee F, et al. Pseudogout: an autoimmune paraneoplastic manifestation of myelodysplastic syndrome. *Cureus* 2018; 10: e3372, DOI: 10.7759/cureus.3372.
31. Lima M, Coutinho J, Bernardo L, et al. Philadelphia-positive T cell acute lymphoblastic leukemia with polymyositis, migratory polyarthritis and hypercalcemia following a chronic myeloid leukemia. *Ann Hematol* 2002; 81: 174–177, DOI: 10.1007/s00277-001-0422-7.
32. Aruch DB, Mims MP. Paraneoplastic nephrotic syndrome and inflammatory arthritis at diagnosis in Hodgkin lymphoma. *Clin Lymphoma Myeloma Leuk* 2013; 13: 77–79, DOI: 10.1016/j.clml.2012.09.006.
33. Prashanth GP, Bhandankar M, Patil VD. Migratory polyarthritis as a paraneoplastic syndrome in childhood leukemia. *Rheumatol Int* 2013; 33: 1647–1648, DOI: 10.1007/s00296-011-2345-5.
34. Handy CE, Robles G, Haque U, Houston B. T cell ALL presenting as seropositive rheumatoid arthritis: case report and review of the literature on seropositive paraneoplastic arthritis. *Clin Rheumatol* 2015; 34: 1647–1650, DOI: 10.1007/s10067-014-2697-9.
35. Erlij D, Calderón B, Rivera A, et al. Polyarthritis and membranoproliferative glomerulonephritis as paraneoplastic manifestation of Hodgkin's lymphoma: a case report and literature review. *Reumatol Clin* 2016; 12: 282–284, DOI: 10.1016/j.reuma.2015.10.009 [Article in English, Spanish].
36. Gamage KKK, Rifath MIM, Fernando H. Migratory polyarthritis as a paraneoplastic syndrome in a patient with diffuse large B cell lymphoma: a case report. *J Med Case Rep* 2018; 12: 189, DOI: 10.1186/s13256-018-1700-5.
37. Raja R, Lamont D, Yung A, Solanki K. A can of red herrings. *Int J Rheum Dis* 2010; 13: e46–e50, DOI: 10.1111/j.1756-185X.2010.01535.x.
38. Drenth JP, de Kleijn EH, de Mulder PH, van der Meer JW. Metastatic breast cancer presenting as fever, rash, and arthritis. *Cancer* 1995; 75: 1608–1611, DOI: 10.1002/1097-0142(19950401)75:7<1608::aid-cnrcr2820750709>3.0.co;2-a.
39. Lambert CM, Nuki G. Multicentric reticulohistiocytosis with arthritis and cardiac infiltration: Regression following treatment for underlying malignancy. *Ann Rheum Dis* 1992; 51: 815–817, DOI: 10.1136/ard.51.6.815.
40. Wiese W, Alansari H, Tranchida P, Madrid FF. Paraneoplastic polyarthritis in an ovarian teratoma. *J Rheumatol* 2004; 3: 1854–1857.
41. Hagiwara K, Suyama Y, Fukuda K. Clinical experience in 115 patients with arthritis and/or enthesitis who met the classification criteria for psoriatic arthritis (CASPAR) within the last two years – possible association with malignant disorders. *Mod Rheumatol* 2016; 26: 625–629, DOI: 10.3109/14397595.2015.1097000.
42. Parperis K, Constantinidou A, Panos G. Paraneoplastic arthritides: insights to pathogenesis, diagnostic approach, and treatment. *J Clin Rheumatol* 2016; 27: e505–e509, DOI: 10.1097/RHU.0000000000001202.