

Linking Structural Racism and Discrimination and Breast Cancer Outcomes: A Social Genomics Approach

Ruth C. Carlos, MD, MS¹; Samilia Obeng-Gyasi, MD, MPH²; Steven W. Cole, PhD³; Bradley J. Zebrack, MPH, MSW, PhD¹; Etta D. Pisano, MD⁴; Melissa A. Troester, PhD⁵; Lava Timsina, PhD⁶; Lynne I. Wagner, PhD⁷; Jon A. Steingrimsson, PhD⁸; Ilana Gareen, PhD⁸; Christoph I. Lee, MD, MS, MBA⁹; Alyce S. Adams, PhD, MPP¹⁰; and Consuelo H. Wilkins, MD, MSCI¹¹

Society to Cells to Outcomes

We live in a society where individuals and communities are marginalized because of their race or ethnicity. This structural inequity extracts enormous health and societal costs, decreasing access to cancer care and increasing health disparities, especially among the most vulnerable. In an effort to identify causes of disparities, we have incorporated individual sociodemographic characteristics (eg, income and education) and other social determinants of health (eg, access to care, insurance, and transportation needs), as well as biologic markers (eg, genetic predisposition to disease) that can serve as therapeutic targets into our research.

Although well intentioned, the use of race as a proxy for genetic predisposition is flawed. There are no genes that are common only within a single racial or ethnic group. In a study of the worldwide geographic distribution of 4,000 alleles,¹ 92% were present in two or more regions and almost 50% were present in all seven major geographic regions (Africa, Europe, Middle East, Central/South Asia, East Asia, Oceania, and America). Intergroup allele similarities were more likely than intragroup similarities. As a biologic construct, ancestry has emerged as a preferred term for the genetic variation reflecting one's geographical origins. Thus, without a viable biologic definition and as a social construct subject to bias, race inadequately captures the complex interaction between ancestry, structural racism^{2,3} (eg, higher frequency of hazardous waste producing businesses or food deserts in predominantly Black neighborhoods), and interpersonal and personally mediated racism.

Emerging models of health disparities propose that simply existing in an environment with systematic racial segregation, overt hostility, or heightened vigilance to the threat of hostility induces a level of ambient stress (which has been likened to racial battle fatigue).⁴ The potential effects of this stress may manifest in physiologic dysregulation at the cellular and molecular level, such as initiation of the

inflammatory cascade and increased expression of inflammatory transcriptomes, predisposing to disease and resulting in outcome disparities traditionally ascribed to race. The objective of this commentary is to outline a novel approach that integrates a Society to Cells to Outcomes framework for a more nuanced understanding of the potential biologic mechanism of structural racism and discrimination on health outcome disparities, using breast cancer to illustrate key principles and pathways.

Breast Cancer in Black Women

The significant improvements in screening, diagnosis, and treatment of breast cancer have not translated into better clinical outcomes for Black women,⁵⁻¹² who continue to have up to a 40% higher mortality rate from breast cancer compared with their White counterparts.^{13,14} This racial disparity in mortality has been partially attributed to racial differences in stage of presentation,^{15,16} molecular subtype,¹⁷⁻¹⁹ and disparities in treatment.^{13,20-23} Nevertheless, this mortality disparity persists even in ductal carcinoma in situ,¹⁴ the earliest and curable stage of disease. Black women are more likely to have poor tumor prognostic features such as high-grade, more aggressive breast cancer molecular subtypes (eg, triple-negative breast cancer), and lymph node metastasis at diagnosis.^{10,24} Although hypothesized, the effects of the proportion of African (v European) ancestry markers on tumor biology and clinical outcomes are inconsistent,²⁵ with no differences in the prevalence of germline pathogenic variants in cancer susceptibility genes by race, suggesting that ancestry-related risk insufficiently explains outcome disparities.

Despite breast cancer screening rates that exceed those of White women,²⁶ Black women are diagnosed with more advanced stages of breast cancer. More worrisome, Black women endure higher rates of false-positive screening results, an unfavorable outcome implicated in an increased risk of breast cancer. Additionally, elevated anxiety, stress, financial burden,²⁷⁻³² and paradoxically, reduced breast cancer screening³³

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on January 10, 2022 and published at ascopubs.org/journal/jco on February 2, 2022; DOI <https://doi.org/10.1200/JCO.21.02004>

© 2022 by American Society of Clinical Oncology

from a false-positive diagnosis have been described, but few focused on experiences of Black women.

The National Institute on Minority Health and Health Disparities model of health disparities^{34,35} proposed a schema of the inter-relationship between the larger social environment and individual risk modifiers (Fig 1). Applied to breast cancer, disparities traditionally attributed to race result from a complex interplay between upstream social conditions (such as disproportionate rates of racism) and policies (such as structural racism), midstream social and physical stressors (such as the experience of discrimination), and downstream individual risk and biologic and genetic pathways. Therefore, the contribution of structural racism including residential racial segregation³⁶⁻³⁹ and the experience of discrimination⁴⁰ need to be integrated into any evaluation of disparities in breast cancer outcomes identified by race across the range of clinical outcomes such as molecular subtype, stage at presentation, treatment disparities, and ultimately mortality. However, significant gaps remain in our understanding of this complex relationship between race/racism, genetic and biologic/physiologic factors, false-positive results, and progression from high-risk false-positive lesions to invasive disease.

Linking Structural Racism and Discrimination and Breast Cancer Outcomes: A Social Genomics Approach

We expand on the Society to Cells to Outcomes framework applied to breast cancer using a model where upstream adverse social conditions (eg, structural racism, racial animus, and the experience of discrimination) and the resultant psychologic stress are moderated and/or mediated by genomic/epigenetic changes or allostatic load (physiologic dysregulation) consequently affecting oncogenesis in Black women (Fig 2).

Structural racism can be evaluated using measures of the physical environment, that is, residential racial segregation measures proposed by Massey and Denton,^{41,42} which include (1) dissimilarity measuring distribution evenness by race or ethnicity; (2) isolation, the probability that Black individuals will encounter or be exposed to other Black individuals; (3) concentration, the population density of Black neighborhoods within a geographic area; (4) centralization, the degree to which Black neighborhoods are located in urban centers; and (5) clustering, the extent

to which Black neighborhoods are surrounded by other Black neighborhoods. Recent systematic reviews^{43,44} provide support for the association between higher Black-White segregation and higher breast cancer mortality in Black patients. Landrine et al⁴⁴ summarized the effects of segregation on Black women. Only five studies^{37,39,45-47} used valid segregation measures.^{41,42,48,49} Three of four studies evaluating mortality found that segregation contributed to disparities; one demonstrated associated differences in late-stage breast cancer diagnosis. Segregation contributed to racial disparities in breast cancer screening in some states but not all. Less rigorous studies using the proportion of Black people in the geographic area as a proxy for racial segregation showed that Black women living in neighborhoods with a lower percentage of Black residents are more likely to develop triple-negative breast cancer.⁵⁰ Conversely, Black women residing in neighborhoods with at least 20% Black residents have a lower disease specific and all-cause mortality.^{39,51} Assessment of the effects of segregation on breast cancer incidence yielded mixed results.⁴³

Regarding racial animus as a manifestation of interpersonal racism, Chae et al⁵² evaluated the frequency of Black racial slur searches in media markets as an area racism measure, which exerted the greatest association with Black mortality, exceeding other traditional variables such as poverty and urbanicity. The experience of discrimination as a manifestation of interpersonal racism also contributes to health outcomes. In our nationally representative survey, women’s experiences of discrimination correlated with increased health service utilization and worse health status,⁵³ consistent with recent systematic reviews on discrimination and health.^{40,54-56} We refer to structural racism, racial animus, and the experience of discrimination collectively as structural racism and discrimination.

Health-related social needs (eg, income, employment, education, insurance, and access to care as well as neighborhood socioeconomic status) and resilience factors (eg, social support and psychologic resourcefulness) can intensify or mitigate structural racism and discrimination risk. We showed that Black women with breast cancer^{57,58} were more likely to adhere to endocrine therapy compared with White women controlling for neighborhood socioeconomic status, a composite measure encompassing




FIG 1. Society → cells → outcomes. Conceptual model of environmental-biologic interactions resulting in health disparities. Adapted from the studies by Alvidrez et al³⁵ and Agur-Collins et al.³⁴




FIG 2. Model of SRD, social genomics, and screening outcomes. CRP, C-reactive protein; CTRA, Conserved Transcriptional Response to Adversity; DBP, diastolic blood pressure; ER, estrogen receptor; HER2, human epidermal growth factor receptor 2; HR, heart rate; HRSN, health-related social needs; IL, interleukin; PR, progesterone receptor; PRO, patient-reported outcome; QOL, quality of life; SBP, systolic blood pressure.

area characteristics such as poverty, crowding, and transportation access. Yet, after controlling for neighborhood socioeconomic status, insurance, and endocrine therapy adherence, higher risk for breast cancer recurrence and mortality persisted among Black women.

Despite the significant research on breast cancer and segregation, there are no studies that have evaluated the effects of segregation on unfavorable screening outcomes, nor racial animus effects on breast cancer screening, incidence, or mortality disparities. The separate effects of health-related social needs require further quantification.

Translating Social Constructs into Biologic Effects

Unfortunately, published research on racial disparities in breast cancer continue to conflate race, a social construct⁵⁹ and ancestry, the genomic variation between populations.⁶⁰ Black race as a broad grouping for individuals of African ancestry may not be uniformly endorsed by all who have African ancestry, further adding to the complexity. Emerging research suggests that although race is a social construct, it has biologic implications, which subsequently affects comorbidity and mortality because of SRD. Evidence-based frameworks such as weathering and allostatic load suggest chronic exposure to adverse social determinants of health have implications for physiology and consequently, disease initiation and progression. In the weathering framework, Geronimus⁶¹ postulates that poor health outcomes experienced by Black women such as higher infant and maternal mortality rates, increased comorbidities, and higher overall mortality rates are attributable to the cumulative effects of social and economic marginalization. In this framework, structural racism and discrimination act as environmental stressors and challenges that activate a physiologic stress response.

The weathering framework closely aligns with allostatic load theory, which describes the activation of the hypothalamic-pituitary-adrenal axis and the autonomic nervous system from psychosocial stressors. Allostatic load describes the cumulative physiologic wear and tear because of exposure

to adverse social determinants of health.⁶² Elevated allostatic load has been associated with low educational achievement, low socioeconomic status, and high work-related stress.⁶³ Additionally, higher allostatic load has been noted in historically marginalized groups such as women, Black people, and individuals who identify as lesbian, gay, or bisexual.⁶⁴

Social Genomics: A Potential Biologic Pathway for Structural Racism and Discrimination on Breast Cancer

We have previously proposed the effects of social determinants of health and health-related social needs (eg, poverty, low educational attainment, and social isolation) on health.³ We expand our previous model of health-related social needs including resulting psychologic stress and their potential biologic effects to explicitly account for structural racism and discrimination. The cumulative effects of structural racism and discrimination, health-related social needs, and psychologic stress are hypothesized to result in a biologic stress response with both epigenetic changes and physiologic dysregulation. To this end, a plausible mechanistic pathway for how structural racism and discrimination affects biology can be understood through the lens of social genomics and social epigenomics. Social genomics examines how psychologic (eg, meaning attribution) and social factors (eg, low socioeconomic status and social isolation) influence gene expression. Concomitantly, social epigenomics evaluates how DNA methylation, microRNA expression, and histone modification alter gene expression and consequently affect disease initiation and progression. Social genomics and epigenomics provide an avenue to further evaluate the intersection between race, epigenetics, and health-related social needs.

One proposed potential pathway for linking adverse socio-environmental factors with poor clinical outcomes is the Conserved Transcriptional Response to Adversity (CTRA) RNA profile. Initially identified as the loneliness gene signature for its association with measures of perceived social isolation, the CTRA is a neurobiologically mediated alteration in immune

cell gene regulation involving increased expression of proinflammatory genes (eg, *IL1B*, *IL6*, *IL8/CXCL8*, and *TNF*) and decreased expression of genes involved in innate antiviral responses (eg, *IFI-*, *MX-*, and *OAS*-family genes). The CTRA profile can be measured by genome-wide transcriptional profiling of circulating blood cells (eg, by RNA sequencing), using either prespecified composites of inflammatory and antiviral gene transcripts (eg, a 53-gene composite used in many studies) or by bioinformatic measures of transcription factor activity mediating inflammatory and antiviral gene regulation.⁶⁵ Contributors to poor outcomes among patients with breast cancer have identified African ancestry, increased CTRA expression, higher allostatic load, and elevated interleukin-6 as factors associated with higher morbidity and mortality among Black patients with breast cancer.^{50,65-68} Cole et al showed CTRA expression in breast tumors to be associated with socioenvironmental risk factors, such as social isolation and low socioeconomic status.⁶⁹ For Black patients with breast cancer in particular, allostatic load has been implicated in negative prognostic features such as larger tumor sizes, poorly differentiated tumors, and estrogen-negative subtypes.⁷⁰ Moreover, among noncancer populations, Black women have a higher allostatic load than their White counterparts.⁷¹ Taken together, these studies suggest that breast cancer screening outcomes, tumorigenesis, and disease progression among Black women are a complex interplay between socioenvironmental factors in conjunction with ancestry and their combined effects on physiology through the stress response, DNA modification, and transcription.

We suggest social genomics as a potential biologic pathway including stress-related transcriptional changes (ie, CTRA gene regulation by the sympathetic nervous system) and physiologic dysregulation (ie, allostatic load) through which the social environment mechanistically influences cancer development and progression^{69,72-75} (Fig 2). Several studies have linked socioeconomic disadvantage and experienced racial discrimination to elevated inflammatory gene expression and CTRA.⁶⁹⁻⁷¹ Exploratory studies of mindfulness and cognitive behavioral stress management interventions have found that CTRA downregulation is associated with improved well-being and disease-free survival among breast cancer survivors.⁷²

Relatedly, we hypothesize that structural racism and discrimination may induce psychologic and physiologic stress, DNA modification, and changes to DNA transcription, which has implications for oncogenesis in Black women. Prior

studies suggest the deleterious effects of elevated allostatic load on breast cancer risk, unfavorable breast cancer clinicopathology, and overall survival, mirroring our results in patients with multiple myeloma⁶⁶ and breast cancer. Similar to CTRA, stress reduction interventions have demonstrated allostatic load reduction in conjunction with improved resilience and quality of life in patients with metastatic breast cancer.⁷⁶ Together, these studies implicate CTRA expression and allostatic load as pathways mediating social disparities in breast cancer outcomes.

Summary

Several studies have documented CTRA gene regulation and associated alterations in upstream signal transduction processes in patients with breast cancer.^{51,69,75,77-83} Scant evidence exists on the potential biologic pathway that structural racism and discrimination, that is, structural racism, racial animus, and the experience of discrimination, has on exerting effects that lead to disparities in breast cancer screening outcomes. Superimposed over the effects of structural racism and discrimination, other domains of health-related social needs including poverty, social isolation, and health care access have also been implicated in poor clinical outcomes among Black patients with breast cancer. Although it is known that health outcomes are improved among persons of all races who have adequate support networks and psychologic resourcefulness,⁸⁴⁻⁸⁶ health-related social needs risk and resilience factors have not routinely been evaluated in breast cancer screening.

We call for both mechanistic and intervention trials to better delineate the effects of structural racism and discrimination specifically on breast cancer screening outcomes, including disparities in false-positive diagnoses and aggressive tumor diagnoses. These studies will provide currently unavailable data that will advance our understanding of breast cancer risk and risk-stratified screening strategies on the basis of multilevel factors including SRD. Improved understanding of the biologic underpinnings of SRD could highlight new biologic and molecular targets for interventions. System or individual interventions that target a molecular signaling pathway that mediates deleterious outcomes could be deployed to potentially reduce the negative effects of SRD, social isolation, or other health-related social needs on breast cancer tumor biology to ultimately help enhance clinical outcomes and close persistent disparities gaps.

AFFILIATIONS

¹University of Michigan, Ann Arbor, MI

²The Ohio State University, Columbus, OH

³University of California Los Angeles, Los Angeles, CA

⁴Beth Israel Deaconess Medical Center, Boston, MA

⁵University of North Carolina, Chapel Hill, NC

⁶Indiana University, Bloomington, IN

⁷Wake Forest School of Medicine, Winston-Salem, NC

⁸Brown University, Providence, RI

⁹University of Washington, Seattle, WA

¹⁰Stanford University, Stanford, CA

¹¹Vanderbilt University, Nashville, TN

CORRESPONDING AUTHOR

Ruth C. Carlos, MD, MS, University of Michigan, 500S State St, Ann Arbor, MI 48109; e-mail: rcarlos@umich.edu.

SUPPORT

A.S.A. is supported by the Advancing Geriatrics Infrastructure and Network Growth (AGING) Initiative (R33AG057806) and the Stanford Cancer Institute (P30CA124435). R.C.C., L.I.W., J.A.S., and I.G. are supported by the Eastern Cooperative Oncology Group–American College of Radiology Imaging Network (ECOG-ACRIN) Research Base (UG1CA189828).

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.21.02004>.

AUTHOR CONTRIBUTIONS

Financial support: Lynne I. Wagner

Administrative support: Ruth C. Carlos

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

REFERENCES

- Rosenberg NA, Pritchard JK, Weber JL, et al: Genetic structure of human populations. *Science* 298:2381-2385, 2002
- Flanagin A, Frey T, Christiansen SL, et al: Updated guidance on the reporting of race and ethnicity in medical and science journals. *JAMA* 326:621-627, 2021
- Obeng-Gyasi S, Carlos R: Biology of discrimination. *J Am Coll Radiol* 17:1541-1544, 2020
- Smith WA, Hung M, Franklin JD: Racial battle fatigue and the miseducation of Black men: Microaggressions, societal problems, & environmental stress. *J Negro Educ* 80:63-82, 2011
- Yedjou CG, Sims JN, Miele L, et al: Health and racial disparity in breast cancer. *Adv Exp Med Biol* 1152:31-49, 2019
- Yedjou CG, Tchounwou PB, Payton M, et al: Assessing the racial and ethnic disparities in breast cancer mortality in the United States. *Int J Environ Res Public Health* 14:486, 2017
- Hunt BR, Silva A, Lock D, et al: Predictors of breast cancer mortality among white and black women in large United States cities: An ecologic study. *Cancer Causes Control* 30:149-164, 2019
- Babatunde OA, Eberth JM, Felder T, et al: Social determinants of racial disparities in breast cancer mortality among Black and White women. *J Racial Ethn Health Disparities* 8:147-156, 2021
- Farr DE, Brandt HM, Adams SA, et al: Examining breast cancer screening behavior among Southern Black women after the 2009 US Preventive Services Task Force mammography guideline revisions. *J Community Health* 45:20-29, 2020
- Iqbal J, Ginsburg O, Rochon PA, et al: Differences in breast cancer stage at diagnosis and cancer-specific survival by race and ethnicity in the United States. *JAMA* 313:165-173, 2015
- Williams DR, Mohammed SA, Shields AE: Understanding and effectively addressing breast cancer in African American women: Unpacking the social context. *Cancer* 122:2138-2149, 2016
- Newman LA, Kaljee LM: Health disparities and triple-negative breast cancer in African American women: A review. *JAMA Surg* 152:485-493, 2017
- DeSantis CE, Miller KD, Goding Sauer A, et al: Cancer statistics for African Americans, 2019. *CA Cancer J Clin* 69:211-233, 2019
- Narod SA, Iqbal J, Giannakeas V, et al: Breast cancer mortality after a diagnosis of ductal carcinoma in situ. *JAMA Oncol* 1:888-896, 2015
- Espina C, McKenzie F, Dos-Santos-Silva I: Delayed presentation and diagnosis of breast cancer in African women: A systematic review. *Ann Epidemiol* 27:659-671.e7, 2017
- Howlander N, Altekruse SF, Li CI, et al: US incidence of breast cancer subtypes defined by joint hormone receptor and HER2 status. *J Natl Cancer Inst* 106:dju055, 2014
- Huo D, Hu H, Rhie SK, et al: Comparison of breast cancer molecular features and survival by African and European ancestry in the cancer genome atlas. *JAMA Oncol* 3:1654-1662, 2017
- O'Meara T, Safonov A, Casadevall D, et al: Immune microenvironment of triple-negative breast cancer in African-American and Caucasian women. *Breast Cancer Res Treat* 175:247-259, 2019
- Acheampong T, Kehm RD, Terry MB, et al: Incidence trends of breast cancer molecular subtypes by age and race/ethnicity in the US from 2010 to 2016. *JAMA Netw Open* 3:e2013226, 2020
- Festa K, Hirsch AE, Cassidy MR, et al: Breast cancer treatment delays at an urban safety net hospital among women experiencing homelessness. *J Community Health* 45:452-457, 2020
- Ellis L, Canchola AJ, Spiegel D, et al: Racial and ethnic disparities in cancer survival: The contribution of tumor, sociodemographic, institutional, and neighborhood characteristics. *J Clin Oncol* 36:25-33, 2018
- Patel A, Wang WJ, Warnack E, et al: Surgical treatment of young women with breast cancer: Public vs private hospitals. *Breast J* 25:625-630, 2019
- Coughlin SS, Yoo W, Whitehead MS, et al: Advancing breast cancer survivorship among African-American women. *Breast Cancer Res Treat* 153:253-261, 2015
- DeSantis CE, Ma J, Gaudet MM, et al: Breast cancer statistics, 2019. *CA Cancer J Clin* 69:438-451, 2019
- Lafarga Previdi I, Velez Vega CM: Health disparities research framework adaptation to reflect Puerto Rico's socio-cultural context. *Int J Environ Res Public Health* 17:8544, 2020
- Komen SG: Comparing Breast Cancer Screening Rates Among Different Groups. 2020. <https://ww5.komen.org/BreastCancer/RacialEthnicIssuesinScreening.html?ecid=social:285>
- Bolejko A, Hagell P, Wann-Hansson C, et al: Prevalence, long-term development, and predictors of psychosocial consequences of false-positive mammography among women attending population-based screening. *Cancer Epidemiol Biomarkers Prev* 24:1388-1397, 2015
- Brett J, Bankhead C, Henderson B, et al: The psychological impact of mammographic screening. A systematic review. *Psychooncology* 14:917-938, 2005
- Brewer NT, Salz T, Lillie SE: Systematic review: The long-term effects of false-positive mammograms. *Ann Intern Med* 146:502-510, 2007
- Brodersen J, Siersma VD: Long-term psychosocial consequences of false-positive screening mammography. *Ann Fam Med* 112:106-115, 2013
- Hafslund B, Espehaug B, Nortvedt MW: Effects of false-positive results in a breast screening program on anxiety, depression and health-related quality of life. *Cancer Nurs* 35:E26-E34, 2012
- Tosteson AN, Fryback DG, Hammond CS, et al: Consequences of false-positive screening mammograms. *JAMA Intern Med* 174:954-961, 2014

33. Cullen J, Schwartz MD, Lawrence WF, et al: Short-term impact of cancer prevention and screening activities on quality of life. *J Clin Oncol* 22:943-952, 2004
34. Agurs-Collins T, Persky S, Paskett ED, et al: Designing and assessing multilevel interventions to improve minority health and reduce health disparities. *Am J Public Health* 109:S86-S93, 2019
35. Alvidrez J, Castille D, Laude-Sharp M, et al: The National Institute on Minority Health and Health Disparities research framework. *Am J Public Health* 109: S16-S20, 2019
36. Poulson MR, Beaulieu-Jones BR, Kenzik KM, et al: Residential racial segregation and disparities in breast cancer presentation, treatment, and survival. *Ann Surg* 273:3-9, 2021
37. Haas JS, Earle CC, Orav JE, et al: Racial segregation and disparities in breast cancer care and mortality. *Cancer* 113:2166-2172, 2008
38. Ojinnaka CO, Luo W, Ory MG, et al: Disparities in surgical treatment of early-stage breast cancer among female residents of Texas: The Role of racial residential segregation. *Clin Breast Cancer* 17:e43-e52, 2017
39. Warner ET, Gomez SL: Impact of neighborhood racial composition and metropolitan residential segregation on disparities in breast cancer stage at diagnosis and survival between black and white women in California. *J Community Health* 35:398-408, 2010
40. Williams DR, Lawrence JA, Davis BA, et al: Understanding how discrimination can affect health. *Health Serv Res* 54:1374-1388, 2019 (suppl 2)
41. Massey DS, Denton NA: *American Apartheid: Segregation and the Making of the Underclass*. Cambridge, MA, Harvard University Press, 1993
42. Massey DS, Denton NA: The dimensions of residential segregation. *Social Forces* 67:281-315, 1988
43. Fang CY, Tseng M: Ethnic density and cancer: A review of the evidence. *Cancer* 124:1877-1903, 2018
44. Landrine H, Corral I, Lee JGL, et al: Residential segregation and racial cancer disparities: A systematic review. *J Racial Ethnic Health Disparities* 4:1195-1205, 2017
45. Haas JS, Earle CC, Orav JE, et al: Racial segregation and disparities in cancer stage for seniors. *J Gen Intern Med* 23:699-705, 2008
46. Dai D: Black residential segregation, disparities in spatial access to health care facilities, and late-stage breast cancer diagnosis in metropolitan Detroit. *Health Place* 16:1038-1052, 2010
47. Mobley LR, Kuo TM, Driscoll D, et al: Heterogeneity in mammography use across the nation: Separating evidence of disparities from the disproportionate effects of geography. *Int J Health Geogr* 7:32, 2008
48. Landrine H, Corral I: Advancing research on racial-ethnic health disparities: Improving measurement equivalence in studies with diverse samples. *Front Public Health* 2:282, 2014
49. Wilkes R, Iceland J: Hypersegregation in the twenty-first century. *Demography* 41:23-36, 2004
50. Schneider BP, Shen F, Jiang G, et al: Impact of genetic ancestry on outcomes in ECOG-ACRIN-E5103. *JCO Precis Oncol* [10.1200/PO.17.00059](https://doi.org/10.1200/PO.17.00059) [published online August 21, 2017]
51. Linnenbringer E, Gehlert S, Geronimus AT: Black-White disparities in breast cancer subtype: The intersection of socially patterned stress and genetic expression. *AIMS Public Health* 4:526-556, 2017
52. Chae DH, Clouston S, Hatzenbuehler ML, et al: Association between an internet-based measure of area racism and black mortality. *PLoS One* 10:e0122963, 2015
53. Fazeli Dehkordy S, Hall KS, Dalton VK, et al: The link between everyday discrimination, healthcare utilization, and health status among a national sample of women. *J Womens Health (Larchmt)* 25:1044-1051, 2016
54. de Freitas DF, Fernandes-Jesus M, Ferreira PD, et al: Psychological correlates of perceived ethnic discrimination in Europe: A meta-analysis. *Psychol Violence* 8:712-725, 2018
55. Britt-Spells AM, Slebodnik M, Sands LP, et al: Effects of perceived discrimination on depressive symptoms among black men residing in the United States: A meta-analysis. *Am J Mens Health* 12:52-63, 2018
56. Carter R, Johnson V, Kirkinis K, et al: A meta-analytic review of racial discrimination: Relationships to health and culture. *Race Soc Probl* 11:15-32, 2018
57. Sparano JA, Gray RJ, Makower DF, et al: Adjuvant chemotherapy guided by a 21-gene expression assay in breast cancer. *N Engl J Med* 379:111-121, 2018
58. Sparano JA, Gray RJ, Makower DF, et al: Clinical outcomes in early breast cancer with a high 21-gene recurrence score of 26 to 100 assigned to adjuvant chemotherapy plus endocrine therapy: A secondary analysis of the TAILORx randomized clinical trial. *JAMA Oncol* 6:367-374, 2020
59. Saini A: *Superior: The Return of Race Science*. Boston, MA, Beacon Press, 2019
60. Dries DL: Genetic ancestry, population admixture, and the genetic epidemiology of complex disease. *Circ Cardiovasc Genet* 2:540-543, 2009
61. Geronimus AT: The weathering hypothesis and the health of African-American women and infants: Evidence and speculations. *Ethn Dis* 2:207-221, 1992
62. McEwen BS: Stress, adaptation, and disease. Allostasis and allostatic load. *Ann N Y Acad Sci* 840:33-44, 1998
63. Beckie TM: A systematic review of allostatic load, health, and health disparities. *Biol Res Nurs* 14:311-346, 2012
64. Guidi J, Lucente M, Sonino N, et al: Allostatic load and its impact on health: A systematic review. *Psychother Psychosom* 90:11-27, 2021
65. Bachelot T, Ray-Coquard I, Menetrier-Caux C, et al: Prognostic value of serum levels of interleukin 6 and of serum and plasma levels of vascular endothelial growth factor in hormone-refractory metastatic breast cancer patients. *Br J Cancer* 88:1721-1726, 2003
66. Obeng-Gyasi S, O'Neill AM, Miller K, et al: Social determinants of health, genetic ancestry, and mortality in ECOG-ACRIN E5103. *J Clin Oncol* 39, 2021 (15_suppl; abstr 6527)
67. Sparano JA, O'Neill AM, Graham N, et al: Inflammatory cytokines and distant recurrence in HER2-negative early breast cancer in the ECOG-ACRIN 5103 trial. *J Clin Oncol* 39, 2021 (15_suppl; abstr 520)
68. Akinyemiju T, Wilson LE, Deveaux A, et al: Association of allostatic load with all-cause and cancer mortality by race and body mass index in the REGARDS cohort. *Cancers (Basel)* 12:1695, 2020
69. Bower JE, Shiao SL, Sullivan P, et al: Prometastatic molecular profiles in breast tumors from socially isolated women. *JNCI Cancer Spectr* 2:pk029, 2018
70. Xing CY, Doose M, Qin B, et al: Prediagnostic allostatic load as a predictor of poorly differentiated and larger sized breast cancers among black women in the women's circle of health follow-up study. *Cancer Epidemiol Biomarkers Prev* 29:216-224, 2020
71. Geronimus AT, Hicken M, Keene D, et al: "Weathering" and age patterns of allostatic load scores among blacks and whites in the United States. *Am J Public Health* 96:826-833, 2006
72. Cole SW: Social regulation of human gene expression: Mechanisms and implications for public health. *Am J Public Health* 103:S84-S92, 2013 (suppl 1)
73. Cole SW: The conserved transcriptional response to adversity. *Curr Opin Behav Sci* 28:31-37, 2019
74. Cole SW, Sood AK: Molecular pathways: Beta-adrenergic signaling in cancer. *Clin Cancer Res* 18:1201-1206, 2012
75. Knight JM, Rizzo JD, Wang T, et al: Molecular correlates of socioeconomic status and clinical outcomes following hematopoietic cell transplantation for leukemia. *JNCI Cancer Spectr* 3:pkz073, 2019

76. Ye ZJ, Qiu HZ, Liang MZ, et al: Effect of a mentor-based, supportive-expressive program, be resilient to breast cancer, on survival in metastatic breast cancer: A randomised, controlled intervention trial. *Br J Cancer* 117:1486-1494, 2017
77. Cole SW, Nagaraja AS, Lutgendorf SK, et al: Sympathetic nervous sytem regulation of the tumor microenvironment. *Nat Rev Cancer* 15:563-572, 2015
78. Knight JM, Rizzo JD, Logan BR, et al: Low socioeconomic status, adverse gene expression profiles, and clinical outcomes in hematopoietic stem cell transplant recipients. *Clin Cancer Res* 22:69-78, 2016
79. Bower JE, Kuhlman KR, Ganz PA, et al: Childhood maltreatment and monocyte gene expression among women with breast cancer. *Brain Behav Immun* 88:396-402, 2020
80. Boyle CC, Cole SW, Dutcher JM, et al: Changes in eudaimonic well-being and the conserved transcriptional response to adversity in younger breast cancer survivors. *Psychoneuroendocrinology* 103:173-179, 2019
81. Powell N, Sloan E, Bailey M, et al: Social stress up-regulates inflammatory gene expression in the leukocyte transcriptome via β -adrenergic induction of myelopoiesis. *Proc Natl Acad Sci USA* 110:16574-16579, 2013
82. Lamkin DM, Sloan EK, Patel AJ, et al: Chronic stress enhances progression of acute lymphoblastic leukemia via β -adrenergic signaling. *Brain Behav Immun* 26:635-641, 2012
83. Sloan EK, Priceman SJ, Cox BF, et al: The sympathetic nervous system induces a metastatic switch in primary breast cancer. *Cancer Res* 70:7042-7052, 2010
84. Zebrack BJ, Landier W: The perceived impact of cancer on quality of life for post-treatment survivors of childhood cancer. *Qual Life Res* 20:1595-1608, 2011
85. Taylor EJ: Transformation of tragedy among women surviving breast cancer. *Oncol Nurs Forum* 27:781-788, 2000
86. Taylor SE, Lichtman RR, Wood JV: Attributions, beliefs about control, and adjustment to breast cancer. *J Pers Soc Psychol* 46:489-502, 1984



ASCO[®] AMERICAN SOCIETY OF CLINICAL ONCOLOGY
ASSOCIATION FOR CLINICAL ONCOLOGY

KNOWLEDGE CONQUERS CANCER

We are a global community of nearly 45,000 members from more than 150 countries, serving members from all subspecialties and professional roles in the pursuit of quality cancer care and progress. Membership provides the support, resources, and solutions for your professional needs:

- Stay on the cutting edge of scientific research and advances
- Streamline your pursuit of continuous learning
- Access evidence-based and data-driven quality resources
- Obtain insight into best practices for cancer care teams
- Connect and exchange views with oncology experts

To learn more about the value of membership, visit asco.org/membership. Not a member? Join today at join.asco.org.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Linking Structural Racism and Discrimination and Breast Cancer Outcomes: A Social Genomics Approach

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

Ruth C. Carlos

Travel, Accommodations, Expenses: GE Healthcare

Other Relationship: Journal of the American College of Radiology (Inst)

Uncompensated Relationships: GE Healthcare

Steven W. Cole

Consulting or Advisory Role: BlueNote Therapeutics

Etta D. Pisano

Research Funding: Deep Health, White Rabbit, Therapixel

Patents, Royalties, Other Intellectual Property: I have several patents, none of which have paid me any royalties

Lynne I. Wagner

Stock and Other Ownership Interests: Johnson & Johnson (I), Lilly (I), Gilead Sciences (I)

Consulting or Advisory Role: Celgene, Athenex

Travel, Accommodations, Expenses: Celgene

Christoph I. Lee

Consulting or Advisory Role: GRAIL

Research Funding: GE Healthcare (Inst)

Other Relationship: American College of Radiology, McGraw Hill Inc, Wolters Kluwer, Oxford University Press

Alyce S. Adams

Employment: The Permanente Medical Group NorCal

Research Funding: The Permanente Medical Group NorCal

No other potential conflicts of interest were reported.