

Review

Naringin and Naringenin Polyphenols in Neurological Diseases: Understandings from a Therapeutic Viewpoint

Talha Bin Emran ^{1,2,*}, Fahadul Islam ², Nikhil Nath ³, Hriday Sutradhar ⁴, Rajib Das ⁴, Saikat Mitra ⁴, Mohammed Merae Alshahrani ⁵, Abdulaziz Hassan Alhasaniah ⁵ and Rohit Sharma ⁶

¹ Department of Pharmacy, BGC Trust University Bangladesh, Chittagong 4381, Bangladesh

² Department of Pharmacy, Faculty of Allied Health Sciences, Daffodil International University, Dhaka 1207, Bangladesh

³ Department of Pharmacy, International Islamic University Chittagong, Chittagong 4318, Bangladesh

⁴ Department of Pharmacy, Faculty of Pharmacy, University of Dhaka, Dhaka 1000, Bangladesh

⁵ Department of Clinical Laboratory Sciences, Faculty of Applied Medical Sciences, Najran University, P.O. Box 1988, Najran 61441, Saudi Arabia

⁶ Department of Rasa Shastra and Bhaishajya Kalpana, Faculty of Ayurveda, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India

* Correspondence: talhabmb@bgctub.ac.bd

Abstract: The glycosides of two flavonoids, naringin and naringenin, are found in various citrus fruits, bergamots, tomatoes, and other fruits. These phytochemicals are associated with multiple biological functions, including neuroprotective, antioxidant, anticancer, antiviral, antibacterial, anti-inflammatory, antiadipogenic, and cardioprotective effects. The higher glutathione/oxidized glutathione ratio in 3-NP-induced rats is attributed to the ability of naringin to reduce hydroxyl radical, hydroperoxide, and nitrite. However, although progress has been made in treating these diseases, there are still global concerns about how to obtain a solution. Thus, natural compounds can provide a promising strategy for treating many neurological conditions. Possible therapeutics for neurodegenerative disorders include naringin and naringenin polyphenols. New experimental evidence shows that these polyphenols exert a wide range of pharmacological activity; particular attention was paid to neurodegenerative diseases such as Alzheimer's and Parkinson's diseases, as well as other neurological conditions such as anxiety, depression, schizophrenia, and chronic hyperglycemic peripheral neuropathy. Several preliminary investigations have shown promising evidence of neuroprotection. The main objective of this review was to reflect on developments in understanding the molecular mechanisms underlying the development of naringin and naringenin as potential neuroprotective medications. Furthermore, the configuration relationships between naringin and naringenin are discussed, as well as their plant sources and extraction methods.

Keywords: naringin; naringenin; polyphenols; neurological disease; dietary interventions



Citation: Emran, T.B.; Islam, F.; Nath, N.; Sutradhar, H.; Das, R.; Mitra, S.; Alshahrani, M.M.; Alhasaniah, A.H.; Sharma, R. Naringin and Naringenin Polyphenols in Neurological Diseases: Understandings from a Therapeutic Viewpoint. *Life* **2023**, *13*, 99. <https://doi.org/10.3390/life13010099>

Academic Editor: Jianfeng Xu

Received: 9 November 2022

Revised: 21 December 2022

Accepted: 22 December 2022

Published: 29 December 2022



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The term “neurological disease” is often used to refer to anything that affects the nervous system. In the brain, spinal cord, or other nerves, structural, metabolic, or electrical dysfunctions may cause a broad range of symptoms. These symptoms include altered states of consciousness, convulsions, muscular weakness, poor coordination, a loss of feeling, and a lack of sensation. Other symptoms include disorientation, pain, and discomfort. Extensive study has led to the discovery of several neurological abnormalities, some of which are somewhat common while others are relatively uncommon [1,2]. Some of these problems are inherited while others are not. Neurology and clinical neuropsychology can identify and treat these illnesses using the appropriate diagnostic and therapeutic methods. Although encased in the skull and spinal vertebrae and chemically isolated by the blood–brain barrier, the brain and spinal cord are susceptible organs [3]. Because of

their placement just beneath the epidermis, nerves are nevertheless vulnerable to harm despite the protective layers of tissue around them. Individual neurons and the neural circuits and nerves they form are susceptible to damage to their electrical and structural integrity [4].

Phytochemicals may be broken down into several different classes, such as flavanols, flavan-3-ols, isoflavones, flavanones, anthocyanidins, and flavones [5]. It has been shown that flavonoids promote apoptosis and decrease metastasis, angiogenesis, and proliferation in the setting of carcinogenesis by interfering with various cell signaling pathways. The flavanone-7-O-glycoside naringin is found in many plants, while citrus fruits are the most common place to find it. A flavanone called naringenin is its main component [6,7]. Many different signaling pathways and signaling molecules are affected by this chemical. Its pharmacological properties include but are not limited to an antioxidant, an anti-inflammatory, an anti-apoptotic, an anti-tumor, and an anti-viral, as well as effects on metabolic syndrome, bone regeneration, neurological illnesses, cardiovascular disease, and genetic damage [8]. Naringin may cause drug interactions due to its ability to inhibit cytochrome P450 enzymes, including CYP3A4 and CYP1A2 [9].

Neurological disorders may be alleviated with the help of naringenin (4',5,7-trihydroxy flavanone), a flavonoid abundant in the peels of citrus fruits (particularly grapes and tomatoes) [10]. Among the top dietary sources of this noteworthy flavonoid are grapefruits and oranges. A few of the many medical advantages of this drug include preventing or reversing weight gain, enhancing metabolic health, and restoring typical lipid profiles in patients with dyslipidemia [11]. Among naringenin's many biological properties is an antioxidant effect. It has been shown that they may help with pain, inflammation, kidney health, and even nerve function. Research has revealed that naringenin may help reduce pain and inflammation associated with several medical disorders [12]. Mechanistically, naringenin is a pleiotropic molecule that inhibits leukocyte recruitment, pro-inflammatory cytokine release, and cytokine-induced analgesia by modulating transient receptor potential (TRP) channels and triggering the nitric oxide (NO)/cyclic guanosine monophosphate (cGMP)/protein kinase G(PKG)/adenosine triphosphate (ATP) pathway. The compound's low bioavailability and limited brain access are significant roadblocks to broader use despite naringenin's promise in treating neurological diseases [13,14]. The therapeutic potential of naringin and naringenin for various symptoms related to neurological diseases has been shown by encouraging results from preclinical investigations [15].

The principal objective of this review was to report on the progress made toward understanding the molecular mechanisms behind the development of naringin and naringenin as potential neuroprotective therapeutics. Additionally, the configuration links between naringin and naringenin, as well as their plant sources and extraction methods, are described.

2. Methodology

The keywords naringin and naringenin, neurological diseases, botanical sources, and neuroprotective action were used to search the following databases: PubMed, Scopus, and Web of Science. Up to 2022, English-language research reports, reviews, and original research articles were considered and studied. An algorithm was used that followed the flowchart in Figure 1 and had all of the steps and requirements for choosing the required literature, in accordance with the recommendations of Page et al. [16].




Figure 1. Flowchart depicting the steps to be followed when selecting published data for the current study; n stands for the number of literature reports.

3. Naringin and Naringenin

The flavanone naringenin and the glycosylated derivative naringin are in high concentrations in grapefruit and other citrus fruits. The antioxidant and anti-inflammatory properties of flavonoids have gained widespread recognition [17]. When naringenin is added to SH-SY5Y cells or when 6-OHDA is injected into mice, the cells are protected in an Nrf2-dependent manner, just like the dopaminergic neurons [18]. This is due to the complete elimination of protective effects and expression of Nrf2-dependent cytoprotective genes after treatment with Nrf2 short interfering RNA [19]. Naringenin prevented rotenone-induced structural alterations in muscle and motor impairment in rat models when given after the administration of the medication. The expression of DJ-1 and chaperone-associated E3 ligase was increased in the striatum and SN after treatment with naringenin [20]. Target proteins are ubiquitinated by DJ-1 and a chaperone-associated E3 ligase and then sent on to be degraded by the proteasome. In a rotenone model of Parkinson's disease, naringenin was found to have neuroprotective effects [21,22]. Naringin may aid dopaminergic neuron recovery after injury if given soon after the damage occurs. Dopaminergic neurons were preserved, GDNF levels in the SN were restored, and the number of ionized calcium-binding adaptor molecule 1 (Iba-1) and tumor necrosis factor- α (TNF- α) immunoreactive neurons in the striatum were reduced after pretreatment with naringin in rats with a unilateral MPP+-lesion. Eukaryotic initiation factor 4E-binding protein 1 (4E-BP1) and growth differentiation and neurotrophic factor (GDNF) were upregulated in the SN after a single injection of naringin [23].

On top of that, it is an effective antioxidant. During fasting and stimulated states, rapid glucose uptake is impaired in BC. Insulin stimulates phosphoinositide 3-kinase PIP3/Akt

and mitogen-activated protein kinase (MAPK) activity, and naringenin prevents this MAPK. It decreases TNF- α and COX-2 levels and raises the transcription factor Nrf2 [24,25].

4. Botanical Sources

Flavonoids are phenolic compounds associated with a wide range of biological functions. There are more than 4000 different flavonoids known to science, most of which are found in their natural, unaltered plant-based forms. Flavonoids can be a dietary supplement [26,27]. Grapefruit and other citrus fruits get their distinctive bitter taste from flavonoid naringin. Although the number of flavonoids taken from food may be large and the flavonoids exhibit potential biological action, they have attracted significantly less attention than flavanols and isoflavones [28,29].

In most cases, the researchers focused on flavanols and isoflavones—intracellular cycling of naringenin, hesperidin, and its glycosylated derivatives, naringenin, hesperidin, and rutin. Grapefruit, bergamot, sour orange, tart cherry, tomato, chocolate, Greek oregano, water mint, and beans are all foods and plants that contain norepinephrine or its glycosides [29,30].

5. Neuroprotective Action

5.1. Alzheimer's Disease

Alzheimer's disease (AD) is the most common type of dementia in the industrialized world and is a severe public health problem [31]. Several different molecular processes caused AD, but its exact pathophysiology is still poorly understood. In several pre-clinical studies, naringin and some of its derivatives, such as naringenin, changed these pathways in ways that could be used to treat AD [32,33]. The disease is caused by the death of cholinergic neurons in the frontal lobe and the formation of Amyloid- β (A β) plaques outside the body [34].

Memantine, an antagonist of the NMDA glutamate receptor, breaks down at a much slower rate than acetylcholinesterase (AChE) inhibitors, such as donepezil aricept, which is often used to treat the symptoms of AD [35,36]. Synaptic dysfunction, in which synapses are damaged, cells are killed, and mental impairments occur, results from consuming too much. This dysfunction can be fixed entirely with the proper treatment. It is essential to understand how A β is currently in charge of producing and storing memories and how this affects synaptic plasticity in the brain network. Evidence suggests that calcium/calmodulin-dependent protein kinase II (CaMKII) is a critical synaptic target for A β -induced synaptic depression [37,38]. Several plant species high in flavonoids have been used in traditional medicine for hundreds of years. Epidemiological and dietary studies on both people and animals have shown that these flavonoids protect against and slow down neurodegeneration, especially regarding the cognitive decline that accompanies aging [39]. The flavonoid glycoside naringin, found in citrus fruits in large amounts, is effective against many diseases and conditions, such as cancer, inflammation, ulcers, osteoporosis, and apoptosis. Naringin has been shown to improve behavior and thinking in animal models of epilepsy caused by kainic acid and Huntington's disease caused by 3-nitropropionic acid [40]. The effects of colchicine and D-galactose on learning and memory are also undone by treatment with naringin. Naringenin has been shown to improve insulin signaling and cognitive ability in the brain and reduce the effects of intracerebroventricular-streptozotocin on neurodegeneration caused by AD (Table 1 and Figure 2) [41].




Figure 2. Illustration representing the site of action of naringin and naringenin in Alzheimer's disease pathway.

5.2. Parkinson's Disease

Parkinson's disease (PD) is a degenerative neurological disease that results in impaired motor function due to dopaminergic neuropathy in the substantia nigra [42–44]. Genetic predisposition may play a role in developing mitochondrial damage and oxidative stress; however, additional molecular routes exist. In recent years, oxidative stress has been investigated as a potential mechanism in neurodegeneration, which is only one example of many similarities between AD and PD. The antioxidant properties of flavonoids, alkaloids, and other polyphenols are coming to the fore. Plant chemicals may modulate enzymes and metabolic signaling pathways to reduce ROS production. Bioactive metabolite flavonoids have anti-oxidative actions on the liver's metabolic pathway [45–47].

The antioxidant and neuroprotective properties of naringin have been studied. Asahina and Inubuse found the blueprints for naringin in 1928 [48,49]. Naringenin is linked to naringin through the C-7 hydroxyl group. Metabolites of naringenin may be found in phases I and II of the drug's metabolism. Glucose is the source of naringin's bitter flavor. It reacts with potassium hydroxide or other essential substances to form 1,3-diphenylpropan-1-one, which has a scent reminiscent of menthol [50–52]. It has been shown that naringin may neutralize ROS, scavenge superoxide, suppress xanthine oxidase, decrease lipid peroxidation, and decrease the permeability of oxygen-stimulated K⁺ erythrocytes. Naringin's antioxidant potentials may aid in treating neurology and diabetes. Degeneration of nerve cells in the striatum and substantia nigra pars compacta kills dopamine-producing brain cells, leading to Parkinson's disease (Table 1 and Figure 3) [53]. Neurons in the substantia nigra are affected as microglia are activated, and protein clumps form. Causes of neurodegeneration include oxidative stress, dopamine depletion, and neuroinflammation. In

addition to boosting dopamine, naringenin also reduces inflammation [54]. By activating Nrf2/ARE and its downstream target genes, including HO-1 and glutathione cysteine ligase regulatory subunit, naringenin protects mice against 6-hydroxydopamine-induced dopaminergic neurodegeneration and oxidative damage. By inhibiting and caspase 3, naringenin was able to prevent apoptosis. Grapefruit suppresses CYP3A4 much more so than naringin. With less naringin than grapefruit juice, orange juice also stops CYP3A4. Preclinical tests have shown that naringin is not transformed into naringenin in cultivated cells [55–57].




Figure 3. Illustration representing the site of action of naringin and naringenin in Parkinson's disease.

Table 1. Preclinical findings on the use of naringin and naringenin polyphenols in neurological disorders.

Disease	Compound	Dose/Conc.	Study Model	Findings	References
Alzheimer Disease	Naringin	50, 100 and 200 mg/kg; PO) for twentyone days	ICV-STZ rats	Restoration of cognitive deficits in ICV-STZ rat along with mitigation of mitochondrial dysfunction mediated oxido-nitrosative stress and cytokine release	[58]
	Naringin	50 or 100 mg/kg/day	APP ^{swe} /PSΔE9 transgenic mice	Reduction in plaque burden and an increase in glucose uptake through the inhibition of GSK-3β activity	[59]
	Naringin	40 and 80 mg/kg	Wistar rats	Protection against ICV β-A1–42 and intranasal manganese induced memory dysfunction possibly due to its antioxidant, anti-inflammatory, anti-amyloidogenesis	[60]
	Naringin	100 mg/kg/day	Mice	Neuroprotective effects through a variety of mechanisms, including amyloid β metabolism, Tau protein hyperphosphorylation, acetyl cholinergic system, glutamate receptor system, oxidative stress and cell apoptosis	[61]
	Naringenin	70–210 μg/mL	PC12 cells	Inhibition of AChE activity	[62]
	Naringenin	50 mg/kg	Male albino Wistar rats	Reduced oxidative stress markers: 4-HNE, MDA, TBARS, H ₂ O ₂ , PC, GSH in the hippocampus; Increase antioxidant level: GPx, GR, GST, SOD, CAT and Na ⁺ /K ⁺ -ATPase in the hippocampus	[63]
	Naringenin	25, 50 and 100 mg/kg	Male Sprague-Dawley rats	Increased the mRNA expression of INS and INSR in cerebral cortex and hippocampus. In addition, NAR reversed ICV-STZ induced Tau hyper-phosphorylation in both hippocampus and cerebral cortex through downregulation of GSK-3β activity	[41]
	Naringenin	25, 50 and 100 μM and 1.5, 3.0 and 4.5 mg/kg	PC12 cells and male ICR mice	Decreased ROS level and LDH activity	[64]
	Naringenin	25 and 50 mg/kg	Male Sprague-Dawley rats	Decreased oxidative stress by depleting elevated lipid peroxide and nitric oxide and elevating reduced glutathione levels and exert cholinergic function through the inhibition of elevated ChE activity	[65]
	Naringin dihydrochalcone	100 mg/kg	APP ^{swe} /PS1ΔE9 (APP/PS1) transgenic mice	Reduction in amyloid plaque burden and Aβ levels, suppression of neuroinflammation and promotion of neurogenesis	[66]

Table 1. Cont.

Disease	Compound	Dose/Conc.	Study Model	Findings	References
Alzheimer Disease	Naringin	40 and 80 mg/kg, PO	Male Wistar rats	Improvement in the cognitive performance and attenuated oxidative damage, as evidenced by lowering of malondialdehyde level and nitrite concentration and restoration of superoxide dismutase, catalase, glutathione S-transferase, and reduced glutathione levels, and acetylcholinesterase activity	[67]
	Naringin	30 or 60 mg/kg/day	NMRI male mice	Reduction of A β plaque numbers in CA1, CA3, and DG areas of the hippocampus	[68]
	Naringin	25, 50 and 100 mg/kg PO	Westar rats	Reduced lipid peroxidation, restored reduced superoxide dismutase and catalase) and acetylcholine esterase activity were significantly decreased	[69]
	N,N'-1,10-Bis(Naringin) Triethylenetetraamine	10–200 μ M	PC12 cells	Decreased the level of ROS in Cu ²⁺ -A β ₁₋₄₂ -treated PC12 cells and elevate the SOD activity in Cu ²⁺ -A β ₁₋₄₂ -treated PC12 cells	[70]
	Naringin	2.5, 5 and 10 mg/kg	Swiss mice	Increased the activities of superoxide dismutase and catalase, and glutathione and decreased malondialdehyde and nitrite contents, and reduced brain acetylcholinesterase activity in mice brains	[71]
	Naringin	80 mg/kg	Wistar albino rats	Improvement of the A β -induced cholinergic dysfunction and increase in the activity of AChE in rat hippocampus, prefrontal cortex, and amygdala. Furthermore, naringin attenuated A β -induced decrease in mitochondrial function, integrity, and bioenergetics as well as mitochondrial and cytosolic calcium level in all the brain regions. Moreover, reversal of A β -induced increase in apoptosis and level of mitochondrial calcium uniporter and decrease in the level of hemeoxygenase-1	[72]
	Naringenin	100 mg/kg, orally	male Wistar rats	Lowered hippocampal MDA content	[73]
Parkinson Disease	Naringin	80 mg/kg	rat model	Protection of the nigrostriatal DA projection by increasing glial cell line-derived neurotrophic factor expression and decreasing TNF- α expression in DA neurons and microglia	[74]
	Naringin	8 or 80 mg/kg per day	Female Sprague Dawley (SD) rats	Increased the level of GDNF in DA neurons, contributing to neuroprotection in the MPP+ rat model of PD, with activation of mammalian target of rapamycin complex 1 and pre-treatment with naringin could attenuate the level of TNF- α in the substantia nigra of MPP+-treated brains	[75]

Table 1. Cont.

Disease	Compound	Dose/Conc.	Study Model	Findings	References
	Naringin	80 mg/kg	male Wistar albino rats	Neuroprotective activity against rotenone-induced toxicity in the animals possibly through Nrf2-mediated pathway	[76]
	Naringin	50, 100 and 200 mg/kg	Swiss albino mice	Reduction in haloperidol-induced cataleptic scores in both bar test and block test	[77]
	Naringenin	25, 50, 100 mg/kg/b.w, PO	male C57BL/6j mice	Reversed the toxic effects of MPTP by reducing LPO levels and increasing the activities of glutathione reductase and catalase along with improved behavioral performance	[78]
	Naringenin	50 mg/kg, orally	albino Wister rats	Improved oxidative stress status by decreasing MDA and increasing glutathione content	[79]
	Naringenin	50, 100 mg/kg	male Sprague-Dawley rats BV-2 and MN9D cell lines	Inhibition of microglia-induced neuroinflammation via NLRP3 inflammasome inactivation	[80]
Parkinson Disease	Naringenin	25, 50, and 100 mM	SH-SY5Y Cell Line	Reduction of the ROS production by decreasing oxidative stress markers such as LPO and NO and increasing SOD level. In addition, pretreatment with NGN decreased the inflammatory markers such as TNF- α and NF- κ B in MPP+-treated SH-SY5Y cells. Further, NGN decreased the pro-apoptotic marker—Bax—and increased the anti-apoptotic marker—Bcl-2—in MPP+-induced SH-SY5Y cells	[81]
	Naringenin	40 μ M	Primary rat mesencephalic cultures	Decreased TH-positive neurons and TUNEL positive neurons	[82]
	Naringenin	50 mg/kg	Male Sprague-Dawley rats	Restoration of dopamine concentrations due to neuroprotective effects rather than compensatory effects by remaining TH-positive cells after 6-OHDA lesioning	[83]
	Naringenin	20, 40 and 80 mM (in vitro) 70 mg/kg, orally (in vivo)	Human neuroblastoma SH-SY5Y cells and male C57BL/6 mice	Activated Nrf2/ARE pathway in dopaminergic (in vitro) Up regulated protein levels of Nrf2/ARE genes (in vivo) Reduced striatal oxidative stress and subsequent apoptotic signalling cascades in striatum (in vivo)	[84]
	Naringenin	12.5 μ M and 25 μ M	SH-SY5Y Human Neuroblastoma cell line	Downregulation of the expression of some Parkinsonian genes such as casp9, lrrk2, and polg and upregulate pink1	[85]
	Naringenin	25, 50, and 100 mg/kg/p.o	Male C57BL/6j mice	Reduced NO content and restored SOD activity, also downregulated TNF- α and IL-1 β expression	[86]

Table 1. Cont.

Disease	Compound	Dose/Conc.	Study Model	Findings	References
Parkinson Disease	Naringenin	10 and 50 μ M	Female Wistar rats	Enhanced astroglial neurotrophic effects on DA neurons through the regulation of Nrf2 activation,	[87]
	Naringenin	5, 10 and 20 mg/kg	Male ICR mice	Increased hippocampal 5-HT, NE and GR levels, and reduced serum corticosterone levels	[88]
	Naringenin	5, 10 and 20 mg/kg	Male ICR mice	Up-regulation of BDNF	[89]
Anxiety and depression	Naringenin	10, 25 and 50 mg/kg	Swiss mice	Naringenin (25–50 mg/kg) ameliorated the hypolocomotion, depressive- and anxiety-like behaviors in hypoxic mice. Naringenin (10 mg/kg) increases BDNF expression but did not significantly ($p < 0.05$) alter corticosterone and catalase contents. The increased expressions of iNOS and NF- κ B as well as loss of amygdala neuronal cells were reduced by naringenin (10 mg/kg)	[90]
	Naringin	10 mg/kg	Adult male Swiss mice	Alleviation of the depressive and anxiogenic behaviors evidenced by the increased preference to sucrose and open arm entries and duration in SPT and EPM respectively	[91]
	Naringin	25–100 mg/kg, i.p	Swiss mice	Increased the levels of GAD67, glutathione and decrease AChE activities, pro-inflammatory cytokines (TNF- α , IL-6), malondialdehyde, nitrite concentrations	[92]
	Naringenin	50 mg/kg/day	Adult male Wistar rats	Mitigation of morphological anomalies in the hippocampal CA1 region and cortex and upregulation of BDNF, Shh, GLI1, NKX2.2, and PAX6	[93]
	Naringenin	0.2, 0.4 mM	C3H10T1/2 cells	Suppression of the protein aggregation caused by EGFP-polyQ97 in mammalian cells.	[94]
Huntington's disease	Naringin	50, 100 mg/kg	Male Wistar rats	Protection against 3-nitropropionic acid induced neurotoxicity via nitric oxide mechanism	[95]
	Naringenin	50 mg/kg b.w, PO	Albino Wistar rats	Improvement of the behavioral function and restored the activity of MAO and 5-HT levels and reduction of the activation of astrocytes against 3-NP induced neurotoxicity	[96]
	Naringin	40, and 80 mg/kg	Adult male Sprague-Dawley rats	Modulation of oxido-nitrosative stress, neuroinflammatory, apoptotic markers and mitochondrial complex activity	[97]
	Naringin	(80 mg/kg b.w/day, orally)	Male Wistar rats	Enhancement of phase II and antioxidant gene expressions via Nrf2 activation	[98]
	Naringin	10 μ M	PC12 cells	Modulation in expressions of B-cell lymphoma 2 and Bcl-2-associated X protein and enhancement of the nuclear translocation of Nrf2	[99]

Table 1. Cont.

Disease	Compound	Dose/Conc.	Study Model	Findings	References
Ischemic brain injury	Naringin	50 and 100 mg/kg	Male Wistar rats	Restoration of reduced glutathione and catalase activity and mitochondrial enzyme activities in cortex, striatum, cerebellum	[100]
	Naringin	106 mg/kg/day	male C57BL/6 strain mice	Suppression of neuronal cell death, reversed the reduction in the level of phosphorylated calcium-calmodulin-dependent protein kinase II, had the tendency to reverse the reduction in the level of glutathione, and blockade of excessive activation of microglia and astrocytes	[101]
	Naringin	40, 80 mg/kg	Male Sprague-Dawley rats	Improvement of early brain injury (EBI), including subarachnoid hemorrhage (SAH) severity, neurologic deficits, brain edema and blood-brain barrier integrity by attenuating SAH-induced oxidative stress and apoptosis, and reduction of the oxidative damage and apoptosis by inhibiting the activation of MAPK signaling pathway	[102]
	Naringenin	50 and 100 mg/kg	Male Sprague-Dawley rats	Down-regulation of NOD2, RIP2, NF- κ B, MMP-9 and up-regulation of claudin-5 expression	[103]
	Naringin	80, 120, or 160 mg/kg/day	SH-SY5Y cells	Reduced 3-nitrotyrosine formation, NADPH oxidase, and iNOS expression. Increased nNOS, p47, and p67 expression. Decreased mitophagy	[104]
	Naringin	100 mg/kg/day	Adult Wistar male rats	Continual treatment increased SOD activity, decreased MDA, NO, iNOS, and IL-1 β . It also improved rats' behavioral performance	[105]
Spinal cord injury	Naringin	20, 40 mg/kg	Female Sprague-Dawley rat	Upregulation of the expression of NK χ 2.2 and 2'3'-cyclic nucleotide 3'-phosphodiesterase, and inhibition of β -catenin expression and GSK-3 β phosphorylation	[106]
	Naringenin	5, 10, 15 mM	Male Wistar rats	Suppression of MMP-9 activity and upregulation of GSH, catalase and MMP-2 activation	[107]
	Naringenin	50–100 mg/kg	Female Wistar rats	Repression of miR-223	[108]
	Naringin	25, 50, and 100 mg/kg	Adult Sprague Dawley rats	Reduction of TNF- α , IL-8 as well as MDA content and elevation of IL-10 as well as SOD activity	[109]

Table 1. Cont.

Disease	Compound	Dose/Conc.	Study Model	Findings	References
Chronic hyperglycemic peripheral neuropathy	Naringenin	50, 100 and 200 mg/kg	Male Sprague Dawley rats	Inhibition of upregulated expression of TNF- α , IL-1 β and MCP-1 level; GFAP and Mac-1 mRNA expression	[110]
	Naringenin	25 and 50 mg/kg	Male Sprague Dawley rats	Increase GSH level and decrease MDA and NO level	[111]
	Naringin	50 and 100 mg/kg, b.w	Rat model of OXL-induced peripheral neuropathy	Improved the level of superoxide dismutase, catalase, glutathione peroxidase, nuclear factor erythroid 2-related factor 2, Heme oxygenase-1, nuclear factor- κ B, tumor necrosis factor- α , interleukin-1 β , Bax, Bcl-2, caspase-3, paraoxonase, mitogen-activated protein kinase 14, neuronal nitric oxide synthase (nNOS), acetylcholinesterase, and arginase 2	[112]

4-HNE: 4-hydroxynonenal, TBARS: Thiobarbituric reactive substances, H₂O₂: Hydrogen peroxide, PC: Protein carbonyl, GSH: Reduced glutathione, GPx: Glutathione peroxidase, GR: Glutathione reductase, GST: Glutathione-S-transferase, SOD: Superoxide dismutase, CAT: catalase, GSK-3 β : Glycogen synthase kinase-3 β , ROS: Reactive oxygen species, AchE: Acetylcholine esterase, DA: Dopaminergic, LPO: Lipid peroxidation, MPTP: 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, 5-HT: Serotonin, NE: Norepinephrine, BDNF: brain-derived neurotrophic factor, MDA: Malondialdehyde.

5.3. Cerebral Ischemia

Cerebral ischemia is a disorder that may trigger a cascade of unfavorable biochemical responses in the brain, leading to malfunction of key brain regions and, commonly, neuropathy [113,114]. An inflammatory reaction and the production of ROS following an ischemia event may damage brain tissue and lead to neuronal death [115]. Ischemia-induced damage involves several kinases, including mitogen-activated protein kinases, extracellular signal-regulated kinases, signal transducers and activators of transcription 1, calcium/calmodulin-dependent kinases, etc. ROS initiate the caspase cascade and encourage the synthesis of pro-inflammatory cytokines including interleukin (IL)-1, IL-6, and tumor necrosis factor-alpha, all of which contribute to cell death. Although progress has been made, a full understanding of the molecular mechanisms behind post-ischemic neuronal damage remains elusive. However, naringin and naringenin have been shown to have a neuroprotective effect after ischemia [116–119].

Naringin lowers cholesterol, prevents blood clots, and improves blood circulation and nutrient supply [100]. In addition, naringin protects against central nervous and cardiovascular diseases. To investigate the role of NFKB1 in OGD/R + injured PC12 cells, the previous study measured the components of the HIF-1 α /AKT/mTOR signal path. HIF-1 α , phosphorylated AKT, and mTOR were all higher in the OGD/R + naringin, OGD/R + si-NFKB1, and OGD/R + naringin + si-NFKB1 groups than in the OGD/R group. Significantly higher levels of HIF-1 α , activated kinase AKT, and mTOR signals were expressed in the OGD/R + si-NFKB1 group ($p < 0.01$). In conclusion, naringin targets NFKB1 and modifies PC12 cell proliferation and apoptosis by affecting HIF-1 α , p-AKT, and p-mTOR levels [120–123].

There is evidence that naringenin may help reduce the adverse effects of oxidative stress on the body, making it a potentially beneficial treatment option for various chronic illnesses. In addition to modulating the activity of antioxidant enzymes and regulating the expression of antioxidant genes, the flavonoid may have direct antioxidant effects, such as scavenging reactive species and reducing oxidative damage [124,125]. When Nrf2 is activated, antioxidant defenses are enhanced by maintaining redox equilibrium and decreasing oxidative stress and inflammation. TNF- α and IL-1 β , two key inflammatory activators, are elevated in response to oxidative stress, whereas naringenin suppresses mRNA

expression in the substantia nigra, hippocampus, and BV2 microglial cells [126,127]. For example, studies on rat hippocampus and BV cells have revealed that naringenin reduces NF- κ B activation. Naringenin inhibited NF- κ B, COX-2, iNOS, and their immunoreactivity, preventing the inflammatory cascade that leads to neurological diseases [51,128].

5.4. Anxiety and Depression

Anxiety and depression are two of the most common mental illnesses, both of which have complex origins at the intersection of several biological systems. Li et al. performed the first studies on the antidepressant effects of naringin and naringenin. The chemicals were tested on mice models of depression brought on by chronic unpredictable mild stress (CUMS) [129]. Anxiety may cause a variety of uncomfortable physical and emotional symptoms, including but not limited to: irritation, impatience, weariness, difficulty concentrating, a racing heart, chest pain, and an upset stomach. Anxiety comes in a variety of forms, and each is treated differently [130]. The serotonergic and noradrenergic systems have been connected to mood disorders such as depression and anxiety. The serotonergic system has far-reaching effects on cognitive processes in the brain, in addition to its role in regulating mood and appetite. Memory and focus are only two of the cognitive functions that are controlled by the noradrenergic system. Increases in serotonin (5-HT) and norepinephrine (NE) receptors, activation of brain-derived neurotrophic factor (BDNF), and decreased blood corticosterone are hypothesized to underlie NRG's antidepressant-like effects [131,132]. It inhibits monoamine oxidase, which may also help those who are depressed. Increased rearing activity, decreased immobility, and increased social communication was seen in mice administered NRG intraperitoneally (at doses of 2.5, 5, and 10 mg/kg), which is consistent with anti-depressant-like and anxiolytic-like effects. Reductions were seen in nitrosative stress, lipid peroxidation, and cholinergic transmission. Mental diseases, often known as mental illnesses or psychiatric disorders, are characterized by persistent patterns of thinking or behavior that significantly impair an individual's capacity to function in everyday life. Both the frequency and length of time that these symptoms will persist are unknown at this time. Various diseases and disorders have been identified, and each has its signs. In some instances, seeking the assistance of a clinical psychologist or psychiatrist specializing in evaluating and managing mental health conditions may be beneficial [133–136].

Neuronal inflammatory mediator release exacerbates tissue damage and reactive oxygen and nitrogen species (ROS/RNS) production, perpetuating the neuronal degeneration in stress-induced neuropsychiatric diseases, including depression and cognitive loss. Decreased antioxidant defenses in neurons promote neuroinflammation and neurodegeneration [137]. Several mediators and intracellular signaling molecules have been connected to neuroinflammatory responses to hypoxia damage. When hypoxia occurs, inflammatory transcription factors, including the NF- κ B pathway, are activated, increasing pro-inflammatory cytokine production [137]. Physical changes in the brain's dendritic arborization and synaptic architecture have been linked to psychological and neurological issues, including depression, anxiety, and memory loss. Naringenin, a dietary flavanone, may be abundant in various foods, including citrus fruits, vegetables, and other berries and nuts [138]. Chronic illnesses and ailments may benefit from consuming a diet high in NG-rich fruits and vegetables. Animal pharmacokinetic studies have shown that NG rapidly undergoes intermediate glucuronide metabolism in the liver and readily crosses the blood–brain barrier (BBB). NG's high permeability across the BBB has been attributed to its association with a broad range of CNS effects. However, the oral bioavailability of naringenin is limited by its metabolism in the liver and its degradation by bacterial enzymes in the colon. Lowered levels of inflammatory mediators were seen in rats, including TNF- α , cyclooxygenase-2, and inducible nitric oxide synthase (iNOS). Studies of naringenin's effects on the brain and spinal cord show it may help treat various neurological disorders [139–141].

5.5. Schizophrenia

Schizophrenia is a disabling brain illness characterized by a wide range of neurotic symptoms including but not limited to hallucinations, delusions, cognitive impairment, disorganized speech, and aberrant motor activities [142]. Many different causes contribute to the development of schizophrenia. These include structural brain abnormalities, impaired neurotransmission, and stress-induced signaling cascades. Disruptions in epidermal growth factor (EGF) signaling and abnormalities in the processing or expression of the EGF receptors ErbB1 and ErbB2 are common in all schizophrenias [143]. As the pathophysiology of schizophrenia worsens, oxidative stress has been suspected to be a contributing factor. Several signs of oxidative stress, including ROS, reduced antioxidant enzyme activity (catalase), depleted glutathione, and oxidized lipids, have been associated with schizophrenia [144].

Schizophrenia and apoptosis have been connected via both intrinsic (mitochondrial death) and extrinsic (death receptor) pathways [145]. Cytochrome C interacts with proapoptotic and anti-apoptotic proteins to trigger the release of activated caspase-3 (primarily Bax and Bcl-2). Diabetes was prevented in streptozotocin-treated rats by administration of naringin, which inhibited the production of inflammatory and oxidative stress mediators [146]. Reduced free radical production, decreased release of proinflammatory cytokines (such as interleukin-6 and TNF- α), and down-regulation of inflammatory proteins such as NF- κ B have all been linked to its anti-inflammatory effects in diabetic, chronic bronchitis, and walker carcinosarcoma rats. To evaluate if naringenin protects interendothelial tight junctions, we analyzed the expression and localization of ZO-1, occludin, claudin-1, and claudin-2 across experimental groups. ZO-1 protein expression was significantly reduced in the TNF- α treated group compared to the control group ($p < 0.05$) [147–151].

However, in TNF- α induced RIMVECs ($p < 0.05$), treatment with naringin dramatically increased ZO-1 protein expression. Immunofluorescence's structured cell death also showed ZO-1 distribution. As proposed here, the medication that acts as a positive allosteric modulator of GABA neurotransmission from chandelier neurons is thought to improve the function of dorsolateral prefrontal cortex circuitry in people with schizophrenia by increasing gamma-band synchronization of pyramidal neuron activity [152–155].

6. Concluding Remarks and Future Directions

Although technological advances have substantially sped up research on phytochemicals, we still have a long way to go before we gather more definitive evidence regarding the neurotherapeutic benefits of herbal medicines. Our data and other researchers' data lead us to believe that naringin and naringenin may be beneficial as neurotherapeutic medications because of their ability to alter several signaling pathways. The outcomes thus far are in line with this theory.

Despite the limitations of ongoing clinical studies, naringenin and naringin are promising therapies for various neurological conditions, including AD, PD, cerebral ischemia, anxiety, depression, schizophrenia, and chronic hyperglycemic peripheral neuropathy. Given these obstacles, it is essential that pharmacokinetic research on naringin and naringenin administration be performed, that more accurate dosage designs for different illnesses be developed, and that innovative drug delivery strategies be developed to boost bioavailability in healthcare situations.

Author Contributions: Conceptualization, T.B.E., F.I., R.D. and S.M.; data curation, T.B.E., F.I., R.D. and S.M.; formal analysis, T.B.E., F.I., N.N., H.S., R.D. and S.M.; investigation, T.B.E., F.I., N.N., H.S., R.D. and S.M.; methodology, T.B.E., F.I., N.N., H.S., R.D. and S.M.; project administration, M.M.A., A.H.A. and R.S.; resources, M.M.A., A.H.A., R.S. and T.B.E.; supervision, T.B.E.; validation, M.M.A., A.H.A., R.S. and T.B.E.; visualization, T.B.E., F.I., R.D. and S.M.; writing—original draft, T.B.E., F.I., R.D. and S.M.; writing—review and editing, M.M.A., A.H.A., R.S. and T.B.E. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data used to support the findings of this study are included within the article.

Conflicts of Interest: The authors declare that they have no conflict of interest.

Abbreviations

PD	Parkinson's disease
TRP	Transient receptor potential
NO	Nitric oxide
cGMP	Cyclic guanosine monophosphate
PKG	Protein kinase G
MAPK	Mitogen-activated protein kinase
GDNF	Growth differentiation and neurotrophic factor
AD	Alzheimer's disease
A β	Amyloid- β
AChE	Acetylcholinesterase
CaMKII	Calcium/calmodulin-dependent protein kinase II
GSH	Reduced glutathione
GPx	Glutathione peroxidase
GST	Glutathione-S-transferase
SOD	Superoxide dismutase
ROS	Reactive oxygen species
LPO	Lipid peroxidation
MDA	Malondialdehyde
5-HT	Serotonin
NE	Norepinephrine
BBB	Blood–brain barrier
iNOS	Inducible nitric oxide synthase

References

- Patel, V.; Chisholm, D.; Dua, T.; Laxminarayan, R.; Medina-Mora, M.E. *Mental, Neurological, and Substance Use Disorders: Disease Control Priorities*; The International Bank for Reconstruction and Development/The World Bank: Washington, DC, USA, 2016; Volume 4.
- Zis, P.; Hadjivassiliou, M. Treatment of Neurological Manifestations of Gluten Sensitivity and Coeliac Disease. *Curr. Treat. Options Neurol.* **2019**, *21*, 10. [[CrossRef](#)] [[PubMed](#)]
- Ludvigsson, J.F.; Bai, J.C.; Biagi, F.; Card, T.R.; Ciacci, C.; Ciclitira, P.J.; Green, P.H.R.; Hadjivassiliou, M.; Holdoway, A.; Van Heel, D.A. Diagnosis and Management of Adult Coeliac Disease: Guidelines from the British Society of Gastroenterology. *Gut* **2014**, *63*, 1210–1228. [[CrossRef](#)] [[PubMed](#)]
- Sapone, A.; Bai, J.C.; Ciacci, C.; Dolinsek, J.; Green, P.H.R.; Hadjivassiliou, M.; Kaukinen, K.; Rostami, K.; Sanders, D.S.; Schumann, M. Spectrum of Gluten-Related Disorders: Consensus on New Nomenclature and Classification. *BMC Med.* **2012**, *10*, 13. [[CrossRef](#)] [[PubMed](#)]
- Fuhr, U.; Kummert, A.L. The Fate of Naringin in Humans: A Key to Grapefruit Juice-drug Interactions? *Clin. Pharmacol. Ther.* **1995**, *58*, 365–373. [[CrossRef](#)] [[PubMed](#)]
- Alam, M.A.; Subhan, N.; Rahman, M.M.; Uddin, S.J.; Reza, H.M.; Sarker, S.D. Effect of Citrus Flavonoids, Naringin and Naringenin, on Metabolic Syndrome and Their Mechanisms of Action. *Adv. Nutr.* **2014**, *5*, 404–417. [[CrossRef](#)]
- Zhang, J.; Gao, W.; Liu, Z.; Zhang, Z.; Liu, C. Systematic Analysis of Main Constituents in Rat Biological Samples after Oral Administration of the Methanol Extract of *Fructus aurantii* by HPLC-ESI-MS/MS. *Iran. J. Pharm. Res.* **2014**, *13*, 493–503. [[CrossRef](#)]
- Ramesh, E.; Alshatwi, A.A. Naringin Induces Death Receptor and Mitochondria-Mediated Apoptosis in Human Cervical Cancer (SiHa) Cells. *Food Chem. Toxicol.* **2013**, *51*, 97–105. [[CrossRef](#)]
- Romagnolo, D.F.; Selmin, O.I. Flavonoids and Cancer Prevention: A Review of the Evidence. *J. Nutr. Gerontol. Geriatr.* **2012**, *31*, 206–238. [[CrossRef](#)]
- Bharti, S.; Rani, N.; Krishnamurthy, B.; Arya, D.S. Preclinical Evidence for the Pharmacological Actions of Naringin: A Review. *Planta Med.* **2014**, *80*, 437–451. [[CrossRef](#)]
- Esaki, S.; Nishiyama, K.; Sugiyama, N.; Nakajima, R.; Takao, Y.; Kamiya, S. Preparation and Taste of Certain Glycosides of Flavanones and of Dihydrochalcones. *Biosci. Biotechnol. Biochem.* **1994**, *58*, 1479–1485. [[CrossRef](#)]

12. Felgines, C.; Texier, O.; Morand, C.; Manach, C.; Scalbert, A.; Régerat, F.; Remesy, C.; Felgines, C.; Texier, O.; Morand, C.; et al. Bioavailability of the Flavanone Naringenin and Its Glycosides in Rats. *Am. J. Physiol. Gastrointest. Liver Physiol.* **2000**, *279*, G1148–G1154. [[CrossRef](#)] [[PubMed](#)]
13. Jadeja, R.N.; Devkar, R.V. Chapter 47—Polyphenols and Flavonoids in Controlling Non-Alcoholic Steatohepatitis. In *Polyphenols in Human Health and Disease*; Watson, R.R., Preedy, V.R., Zibadi, S., Eds.; Academic Press: San Diego, CA, USA, 2014; pp. 615–623, ISBN 978-0-12-398456-2.
14. Manchope, M.F.; Ferraz, C.R.; Borghi, S.M.; Rasquel-Oliveira, F.S.; Franciosi, A.; Bagatim-Souza, J.; Dionisio, A.M.; Casagrande, R.; Verri, W.A. Chapter 38—Therapeutic Role of Naringenin to Alleviate Inflammatory Pain. In *Treatments, Mechanisms, and Adverse Reactions of Anesthetics and Analgesics*; Rajendram, R., Patel, V.B., Preedy, V.R., Martin, C.R., Eds.; Academic Press: Cambridge, MA, USA, 2022; pp. 443–455, ISBN 978-0-12-820237-1.
15. Li, Z.-X.; Cao, Y.; Yan, P. Topological Insulators and Semimetals in Classical Magnetic Systems. *Phys. Rep.* **2021**, *915*, 1–64. [[CrossRef](#)]
16. Page, M.J.; McKenzie, J.E.; Bossuyt, P.M.; Boutron, I.; Hoffmann, T.C.; Mulrow, C.D.; Shamseer, L.; Tetzlaff, J.M.; Akl, E.A.; Brennan, S.E.; et al. The PRISMA 2020 Statement: An Updated Guideline for Reporting Systematic Reviews. *Syst. Rev.* **2021**, *10*, 89. [[CrossRef](#)] [[PubMed](#)]
17. Abaza, M.S.I.; Orabi, K.Y.; Al-Quattan, E.; Al-Attiyah, R.J. Growth Inhibitory and Chemo-Sensitization Effects of Naringenin, a Natural Flavanone Purified from *Thymus Vulgaris*, on Human Breast and Colorectal Cancer. *Cancer Cell Int.* **2015**, *15*, 1–19. [[CrossRef](#)] [[PubMed](#)]
18. Ademosun, A.O.; Oboh, G.; Passamonti, S.; Tramer, F.; Ziberna, L.; Boligon, A.A.; Athayde, M.L. Inhibition of Metalloproteinase and Proteasome Activities in Colon Cancer Cells by Citrus Peel Extracts. *J. Basic Clin. Physiol. Pharmacol.* **2015**, *26*, 471–477. [[CrossRef](#)] [[PubMed](#)]
19. Hsiu, S.-L.; Huang, T.-Y.; Hou, Y.-C.; Chin, D.-H.; Chao, P.-D.L. Comparison of Metabolic Pharmacokinetics of Naringin and Naringenin in Rabbits. *Life Sci.* **2002**, *70*, 1481–1489. [[CrossRef](#)]
20. Moghaddam, R.H.; Samimi, Z.; Moradi, S.Z.; Little, P.J.; Xu, S.; Farzaei, M.H. Naringenin and Naringin in Cardiovascular Disease Prevention: A Preclinical Review. *Eur. J. Pharmacol.* **2020**, *887*, 173535. [[CrossRef](#)]
21. Zhang, L.; Song, L.; Zhang, P.; Liu, T.; Zhou, L.; Yang, G.; Lin, R.; Zhang, J. Solubilities of Naringin and Naringenin in Different Solvents and Dissociation Constants of Naringenin. *J. Chem. Eng. Data* **2015**, *60*, 932–940. [[CrossRef](#)]
22. Fujioka, K.; Greenway, F.; Sheard, J.; Ying, Y. The Effects of Grapefruit on Weight and Insulin Resistance: Relationship to the Metabolic Syndrome. *J. Med. Food* **2006**, *9*, 49–54. [[CrossRef](#)]
23. Ishii, K.; Furuta, T.; Kasuya, Y. Determination of Naringin and Naringenin in Human Plasma by High-Performance Liquid Chromatography. *J. Chromatogr. B Biomed. Sci. Appl.* **1996**, *683*, 225–229. [[CrossRef](#)]
24. Lee, C.-H.; Jeong, T.-S.; Choi, Y.-K.; Hyun, B.-H.; Oh, G.-T.; Kim, E.-H.; Kim, J.-R.; Han, J.-I.; Bok, S.-H. Anti-Atherogenic Effect of Citrus Flavonoids, Naringin and Naringenin, Associated with Hepatic ACAT and Aortic VCAM-1 and MCP-1 in High Cholesterol-Fed Rabbits. *Biochem. Biophys. Res. Commun.* **2001**, *284*, 681–688. [[CrossRef](#)] [[PubMed](#)]
25. Renugadevi, J.; Prabu, S.M. Naringenin Protects against Cadmium-Induced Oxidative Renal Dysfunction in Rats. *Toxicology* **2009**, *256*, 128–134. [[CrossRef](#)] [[PubMed](#)]
26. Cook, N.C.; Samman, S. Flavonoids—Chemistry, Metabolism, Cardioprotective Effects, and Dietary Sources. *J. Nutr. Biochem.* **1996**, *7*, 66–76. [[CrossRef](#)]
27. Choudhury, R.; Chowrimootoo, G.; Srail, K.; Debnam, E.; Rice-Evans, C.A. Interactions of the Flavonoid Naringenin in the Gastrointestinal Tract and the Influence of Glycosylation. *Biochem. Biophys. Res. Commun.* **1999**, *265*, 410–415. [[CrossRef](#)] [[PubMed](#)]
28. Croft, K.D. The Chemistry and Biological Effects of Flavonoids and Phenolic Acids A. *Ann. N. Y. Acad. Sci.* **1998**, *854*, 435–442. [[CrossRef](#)]
29. Scalbert, A.; Williamson, G. Dietary Intake and Bioavailability of Polyphenols. *J. Nutr.* **2000**, *130*, 2073S–2085S. [[CrossRef](#)]
30. Pietta, P.; Minoggio, M.; Bramati, L. Plant Polyphenols: Structure, Occurrence and Bioactivity. *Stud. Nat. Prod. Chem.* **2003**, *28*, 257–312. [[CrossRef](#)]
31. Navipour, E.; Neamatshahi, M.; Barabadi, Z.; Neamatshahi, M.; Keykhosravi, A. Epidemiology and Risk Factors of Alzheimer’s Disease in Iran: A Systematic Review. *Iran. J. Public Health* **2019**, *48*, 2133–2139. [[CrossRef](#)]
32. Pedersen, W.A.; McMillan, P.J.; Kulstad, J.J.; Leverenz, J.B.; Craft, S.; Haynatzki, G.R. Rosiglitazone Attenuates Learning and Memory Deficits in Tg2576 Alzheimer Mice. *Exp. Neurol.* **2006**, *199*, 265–273. [[CrossRef](#)]
33. Rajmohan, R.; Reddy, P.H. Amyloid-Beta and Phosphorylated Tau Accumulations Cause Abnormalities at Synapses of Alzheimer’s Disease Neurons. *J. Alzheimer’s Dis.* **2017**, *57*, 975–999. [[CrossRef](#)]
34. Jin, H.; Wang, W.; Zhao, S.; Yang, W.; Qian, Y.; Jia, N.; Feng, G. Aβ-HBc Virus-like Particles Immunization without Additional Adjuvant Ameliorates the Learning and Memory and Reduces Aβ Deposit in PDAPP Mice. *Vaccine* **2014**, *32*, 4450–4456. [[CrossRef](#)] [[PubMed](#)]
35. Gauthier, S.; Scheltens, P.; Cummings, J. *Alzheimer’s Disease and Related Disorders*; CRC Press: Boca Raton, FL, USA, 2005; ISBN 0203931742.
36. Mimura, M.; Yano, M. Memory Impairment and Awareness of Memory Deficits in Early-Stage Alzheimer’s Disease. *Rev. Neurosci.* **2006**, *17*, 253–266. [[CrossRef](#)] [[PubMed](#)]

37. Fakhri, S.; Abbaszadeh, F.; Dargahi, L.; Jorjani, M. Astaxanthin: A Mechanistic Review on Its Biological Activities and Health Benefits. *Pharmacol. Res.* **2018**, *136*, 1–20. [[CrossRef](#)] [[PubMed](#)]
38. Bao, X.-Q.; Li, N.; Wang, T.; Kong, X.-C.; Tai, W.-J.; Sun, H.; Zhang, D. FLZ Alleviates the Memory Deficits in Transgenic Mouse Model of Alzheimer's Disease via Decreasing Beta-Amyloid Production and Tau Hyperphosphorylation. *PLoS ONE* **2013**, *8*, e78033. [[CrossRef](#)]
39. Mimura, M. Memory Impairment and Awareness of Memory Deficits in Early-Stage Alzheimer's Disease. *Tohoku J. Exp. Med.* **2008**, *215*, 133–140. [[CrossRef](#)]
40. Obulesu, M.; Jhansilakshmi, M. Neuroinflammation in Alzheimer's Disease: An Understanding of Physiology and Pathology. *Int. J. Neurosci.* **2014**, *124*, 227–235. [[CrossRef](#)]
41. Yang, W.; Ma, J.; Liu, Z.; Lu, Y.; Hu, B.; Yu, H. Effect of Naringenin on Brain Insulin Signaling and Cognitive Functions in ICV-STZ Induced Dementia Model of Rats. *Neurol. Sci.* **2014**, *35*, 741–751. [[CrossRef](#)]
42. Burke, R.E.; O'Malley, K. Axon Degeneration in Parkinson's Disease. *Exp. Neurol.* **2013**, *246*, 72–83. [[CrossRef](#)]
43. Savitt, J.M.; Dawson, V.L.; Dawson, T.M. Diagnosis and Treatment of Parkinson Disease: Molecules to Medicine. *J. Clin. Investig.* **2006**, *116*, 1744–1754. [[CrossRef](#)]
44. Shulman, J.M.; De Jager, P.L.; Feany, M.B. Parkinson's Disease: Genetics and Pathogenesis. *Annu. Rev. Pathol. Mech. Dis.* **2011**, *6*, 193–222. [[CrossRef](#)]
45. Dexter, D.T.; Jenner, P. Parkinson Disease: From Pathology to Molecular Disease Mechanisms. *Free Radic. Biol. Med.* **2013**, *62*, 132–144. [[CrossRef](#)] [[PubMed](#)]
46. AlDakheel, A.; Kalia, L.V.; Lang, A.E. Pathogenesis-Targeted, Disease-Modifying Therapies in Parkinson Disease. *Neurotherapeutics* **2014**, *11*, 6–23. [[CrossRef](#)] [[PubMed](#)]
47. Olanow, C.W.; Tatton, W.G. Etiology and Pathogenesis of Parkinson's Disease. *Annu. Rev. Neurosci.* **1999**, *22*, 123. [[CrossRef](#)] [[PubMed](#)]
48. Zbarsky, V.; Datla, K.P.; Parkar, S.; Rai, D.K.; Aruoma, O.I.; Dexter, D.T. Neuroprotective Properties of the Natural Phenolic Antioxidants Curcumin and Naringenin but Not Quercetin and Fisetin in a 6-OHDA Model of Parkinson's Disease. *Free Radic. Res.* **2005**, *39*, 1119–1125. [[CrossRef](#)] [[PubMed](#)]
49. Golechha, M.; Chaudhry, U.; Bhatia, J.; Saluja, D.; Arya, D.S. Naringin Protects against Kainic Acid-Induced Status Epilepticus in Rats: Evidence for an Antioxidant, Anti-Inflammatory and Neuroprotective Intervention. *Biol. Pharm. Bull.* **2011**, *34*, 360–365. [[CrossRef](#)]
50. Ahmed, S.; Khan, H.; Aschner, M.; Hasan, M.M.; Hassan, S.T.S. Therapeutic Potential of Naringin in Neurological Disorders. *Food Chem. Toxicol.* **2019**, *132*, 110646. [[CrossRef](#)]
51. Raza, S.S.; Khan, M.M.; Ahmad, A.; Ashafaq, M.; Islam, F.; Wagner, A.P.; Safhi, M.M. Neuroprotective Effect of Naringenin Is Mediated through Suppression of NF-KB Signaling Pathway in Experimental Stroke. *Neuroscience* **2013**, *230*, 157–171. [[CrossRef](#)]
52. Harrison, W.T.A.; Yathirajan, H.S.; Sarojini, B.K.; Narayana, B.; Anilkumar, H.G. Do C—H · · · O and C—H · · · π Interactions Help to Stabilize a Non-Centrosymmetric Structure for Racemic 2, 3-Dibromo-1, 3-Diphenylpropan-1-One? *Acta Crystallogr. Sect. C Cryst. Struct. Commun.* **2005**, *61*, o728–o730. [[CrossRef](#)]
53. Gopinathan, G.; Teravainen, H.; Dambrosia, J.M.; Ward, C.D.; Sanes, J.N.; Stuart, W.K.; Everts, E.V.; Calne, D.B. Lisuride in Parkinsonism. *Neurology* **1981**, *31*, 371. [[CrossRef](#)]
54. Kim, H.D.; Jeong, K.H.; Jung, U.J.; Kim, S.R. Naringin Treatment Induces Neuroprotective Effects in a Mouse Model of Parkinson's Disease in Vivo, but Not Enough to Restore the Lesioned Dopaminergic System. *J. Nutr. Biochem.* **2016**, *28*, 140–146. [[CrossRef](#)]
55. Kensler, T.W.; Wakabayashi, N.; Biswal, S. Cell Survival Responses to Environmental Stresses via the Keap1-Nrf2-ARE Pathway. *Annu. Rev. Pharmacol. Toxicol.* **2007**, *47*, 89–116. [[CrossRef](#)] [[PubMed](#)]
56. Lee, J.-M.; Johnson, J.A. An Important Role of Nrf2-ARE Pathway in the Cellular Defense Mechanism. *BMB Rep.* **2004**, *37*, 139–143. [[CrossRef](#)] [[PubMed](#)]
57. Eagling, V.A.; Profit, L.; Back, D.J. Inhibition of the CYP3A4-Mediated Metabolism and P-Glycoprotein-Mediated Transport of the HIV-1 Protease Inhibitor Saquinavir by Grapefruit Juice Components. *Br. J. Clin. Pharmacol.* **1999**, *48*, 543. [[CrossRef](#)] [[PubMed](#)]
58. Sachdeva, A.K.; Kuhad, A.; Chopra, K. Naringin Ameliorates Memory Deficits in Experimental Paradigm of Alzheimer's Disease by Attenuating Mitochondrial Dysfunction. *Pharmacol. Biochem. Behav.* **2014**, *127*, 101–110. [[CrossRef](#)] [[PubMed](#)]
59. Wang, D.; Gao, K.; Li, X.; Shen, X.; Zhang, X.; Ma, C.; Qin, C.; Zhang, L. Long-Term Naringin Consumption Reverses a Glucose Uptake Defect and Improves Cognitive Deficits in a Mouse Model of Alzheimer's Disease. *Pharmacol. Biochem. Behav.* **2012**, *102*, 13–20. [[CrossRef](#)]
60. Kaur, G.; Prakash, A. Involvement of the Nitric Oxide Signaling in Modulation of Naringin against Intranasal Manganese and Intracerebroventricular β-Amyloid Induced Neurotoxicity in Rats. *J. Nutr. Biochem.* **2020**, *76*, 108255. [[CrossRef](#)]
61. Meng, X.; Fu, M.; Wang, S.; Chen, W.; Wang, J.; Zhang, N. Naringin Ameliorates Memory Deficits and Exerts Neuroprotective Effects in a Mouse Model of Alzheimer's Disease by Regulating Multiple Metabolic Pathways. *Mol. Med. Rep.* **2021**, *23*, 332. [[CrossRef](#)]
62. Heo, H.J.; Kim, M.J.; Lee, J.M.; Choi, S.J.; Cho, H.Y.; Hong, B.; Kim, H.K.; Kim, E.; Shin, D.H. Naringenin from Citrus Junos Has an Inhibitory Effect on Acetylcholinesterase and a Mitigating Effect on Amnesia. *Dement. Geriatr. Cogn. Disord.* **2004**, *17*, 151–157. [[CrossRef](#)]

63. Khan, M.B.; Khan, M.M.; Khan, A.; Ahmed, M.E.; Ishrat, T.; Tabassum, R.; Vaibhav, K.; Ahmad, A.; Islam, F. Naringenin Ameliorates Alzheimer's Disease (AD)-Type Neurodegeneration with Cognitive Impairment (AD-TNDCI) Caused by the Intracerebroventricular- Streptozotocin in Rat Model. *Neurochem. Int.* **2012**, *61*, 1081–1093. [[CrossRef](#)]
64. Heo, H.J.; Kim, D.O.; Shin, S.C.; Kim, M.J.; Kim, B.G.; Shin, D.H. Effect of Antioxidant Flavanone, Naringenin, from Citrus Junos on Neuroprotection. *J. Agric. Food Chem.* **2004**, *52*, 1520–1525. [[CrossRef](#)]
65. Rahigude, A.; Bhutada, P.; Kaulaskar, S.; Aswar, M.; Otari, K. Participation of Antioxidant and Cholinergic System in Protective Effect of Naringenin against Type-2 Diabetes-Induced Memory Dysfunction in Rats. *Neuroscience* **2012**, *226*, 62–72. [[CrossRef](#)] [[PubMed](#)]
66. Yang, W.; Zhou, K.; Zhou, Y.; An, Y.; Hu, T.; Lu, J.; Huang, S.; Pei, G. Naringin Dihydrochalcone Ameliorates Cognitive Deficits and Neuropathology in APP/PS1 Transgenic Mice. *Front. Aging Neurosci.* **2018**, *10*, 169. [[CrossRef](#)] [[PubMed](#)]
67. Kumar, A.; Dogra, S.; Prakash, A. Protective Effect of Naringin, a Citrus Flavonoid, against Colchicine-Induced Cognitive Dysfunction and Oxidative Damage in Rats. *J. Med. Food* **2010**, *13*, 976–984. [[CrossRef](#)] [[PubMed](#)]
68. Jahanshahi, M.; Khalili, M.; Margedari, A. Naringin Chelates Excessive Iron and Prevents the Formation of Amyloid-Beta Plaques in the Hippocampus of Iron-Overloaded Mice. *Front. Pharmacol.* **2021**, *12*, 518. [[CrossRef](#)]
69. Prabhakar, O. Naringin Attenuates Aluminum Induced Cognitive Deficits in Rats. *Indian J. Pharm. Educ. Res.* **2020**, *54*, 674–681. [[CrossRef](#)]
70. Guo, L.; Sun, B. N,N'-1,10-Bis(Naringin) Triethylenetetraamine, Synthesis and as a Cu(II) Chelator for Alzheimer's Disease Therapy. *Biol. Pharm. Bull.* **2021**, *44*, 51–56. [[CrossRef](#)]
71. Ben-Azu, B.; Nwoke, E.; Aderibigbe, A.; Omogbiya, I.; Ajayi, A.; Olonode, E.; Umukoro, S.; Iwalewa, E. Possible Neuroprotective Mechanisms of Action Involved in the Neurobehavioral Property of Naringin in Mice. *Biomed. Pharmacother.* **2019**, *109*, 536–546. [[CrossRef](#)]
72. Varshney, V.; Garabadu, D. Naringin Exhibits Mas Receptor-Mediated Neuroprotection Against Amyloid Beta-Induced Cognitive Deficits and Mitochondrial Toxicity in Rat Brain. *Neurotox. Res.* **2021**, *39*, 1023–1043. [[CrossRef](#)]
73. Ghofrani, S.; Joghataei, M.; Mohseni, S.; Baluchnejadmojarad, T.; Bagheri, M.; Khamse, S.; Roghani, M. Naringenin Improves Learning and Memory in an Alzheimer's Disease Rat Model: Insights into the Underlying Mechanisms. *Eur. J. Pharmacol.* **2015**, *764*, 195–201. [[CrossRef](#)]
74. Jung, U.; Kim, S. Effects of Naringin, a Flavanone Glycoside in Grapefruits and Citrus Fruits, on the Nigrostriatal Dopaminergic Projection in the Adult Brain. *Neural Regen. Res.* **2014**, *9*, 1514.
75. Leem, E.; Nam, J.; Jeon, M.; Shin, W.; Won, S.; Park, S.; Choi, M.; Jin, B.; Jung, U.; Kim, S. Naringin Protects the Nigrostriatal Dopaminergic Projection through Induction of GDNF in a Neurotoxin Model of Parkinson's Disease. *J. Nutr. Biochem.* **2014**, *25*, 801–806. [[CrossRef](#)]
76. Garabadu, D.; Agrawal, N. Naringin Exhibits Neuroprotection Against Rotenone-Induced Neurotoxicity in Experimental Rodents. *NeuroMolecular Med.* **2020**, *22*, 314–330. [[CrossRef](#)] [[PubMed](#)]
77. Kumar, N.; James, R.; Sinha, S.; Kinra, M.; Anuranjana, P.V.; Nandakumar, K. Naringin Exhibited Anti-Parkinsonian like Effect against Haloperidol-Induced Catalepsy in Mice. *Res. J. Pharm. Technol.* **2021**, *14*, 662–666. [[CrossRef](#)]
78. Sugumar, M.; Sevanan, M.; Sekar, S. Neuroprotective Effect of Naringenin against MPTP-Induced Oxidative Stress. *Int. J. Neurosci.* **2018**, *129*, 534–539. [[CrossRef](#)] [[PubMed](#)]
79. Madani, M.; Attia, A.; ElSalalm, R.; El-Shenawy, S.; Arbid, M. Neuropharmacological Effects of Naringenin, Harmine and Adenosine on Parkinsonism Induced in Rats. *Der Pharm. Lett.* **2016**, *8*, 45–57.
80. Chen, C.; Wei, Y.; He, X.; Li, D.; Wang, G.; Li, J. Naringenin Produces Neuroprotection Against LPS-Induced Dopamine Neurotoxicity via the Inhibition of Microglial NLRP3 Inflammasome Activation. *Front. Immunol.* **2019**, *10*, 936. [[CrossRef](#)]
81. Mani, S.; Sekar, S.; Chidambaram, S.B.; Sevanan, M. Naringenin Protects against 1-Methyl-4-Phenylpyridinium-Induced Neuroinflammation and Resulting Reactive Oxygen Species Production in SH - SY5Y Cell Line: An In Vitro Model of Parkinson's Disease. *Pharmacogn. Mag.* **2018**, 458–464. [[CrossRef](#)]
82. Mercer, L.; Kelly, B.; Horne, M.; Beart, P. Dietary Polyphenols Protect Dopamine Neurons from Oxidative Insults and Apoptosis: Investigations in Primary Rat Mesencephalic Cultures. *Biochem. Pharmacol.* **2005**, *69*, 339–345. [[CrossRef](#)]
83. Kabir, M.S.H.; Hossain, M.M.; Kabir, M.I.; Rahman, M.M.; Hasanat, A.; Emran, T.B.; Rahman, M.A. Phytochemical screening, Antioxidant, Thrombolytic, alpha-amylase inhibition and cytotoxic activities of ethanol extract of *Steudnera colocasifolia* K. Koch leaves. *J. Young Pharm.* **2016**, *8*, 391. [[CrossRef](#)]
84. Lou, H.; Jing, X.; Wei, X.; Shi, H.; Ren, D.; Zhang, X. Naringenin Protects against 6-OHDA-Induced Neurotoxicity via Activation of the Nrf2/ARE Signaling Pathway. *Neuropharmacology* **2014**, *79*, 380–388. [[CrossRef](#)]
85. Kesh, S.; Rajesh, R.; Balakrishnan, A. Naringenin Alleviates 6-Hydroxydopamine Induced Parkinsonism in SHSY5Y Cells and Zebrafish Model. *Comp. Biochem. Physiol. Part C Toxicol. Pharmacol.* **2021**, *239*, 108893. [[CrossRef](#)] [[PubMed](#)]
86. Mani, S.; Sekar, S.; Barathidasan, R.; Manivasagam, T.; Thenmozhi, A.J.; Sevanan, M.; Chidambaram, S.B.; Essa, M.M.; Guillemain, G.J.; Sakharkar, M.K. Naringenin Decreases α -Synuclein Expression and Neuroinflammation in MPTP-Induced Parkinson's Disease Model in Mice. *Neurotox. Res.* **2018**, *33*, 656–670. [[CrossRef](#)] [[PubMed](#)]
87. Wang, G.; Zhang, B.; He, X.; Li, D.; Shi, J.; Zhang, F. Naringenin Targets on Astroglial Nrf2 to Support Dopaminergic Neurons. *Pharmacol. Res.* **2018**, *139*, 452–459. [[CrossRef](#)] [[PubMed](#)]

88. Yi, L.; Li, J.; Li, H.; Su, D.; Quan, X.; He, X.; Wang, X. Antidepressant-like Behavioral, Neurochemical and Neuroendocrine Effects of Naringenin in the Mouse Repeated Tail Suspension Test. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **2012**, *39*, 175–181. [[CrossRef](#)]
89. Yi, L.; Liu, B.; Li, J.; Luo, L.; Liu, Q.; Geng, D.; Tang, Y.; Xia, Y.; Wu, D. BDNF Signaling Is Necessary for the Antidepressant-like Effect of Naringenin. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* **2014**, *48*, 135–141. [[CrossRef](#)]
90. Olugbemide, A.S.; Ben-Azu, B.; Bakre, A.G.; Ajayi, A.M.; Femi-Akinlosotu, O.; Umukoro, S. Naringenin Improves Depressive- and Anxiety-like Behaviors in Mice Exposed to Repeated Hypoxic Stress through Modulation of Oxido-Inflammatory Mediators and NF-KB/BDNF Expressions. *Brain Res. Bull.* **2021**, *169*, 214–227. [[CrossRef](#)]
91. Anyanwu, G.E.; Atuadu, V.O.; Esom, E.A.; Nto, J.N.; Katchy, A.U. Putative Role of Monoaminergic Systems in Antidepressant and Anxiolytic Effects of Naringin in Mice: An Interaction Study with Receptor Antagonists. *J. Pharm. Res. Int.* **2021**, *33*, 661–676. [[CrossRef](#)]
92. Oladapo, O.M.; Ben-azu, B.; Ajayi, A.M.; Emokpae, O.; Eneni, A.O. Naringin Confers Protection against Psychosocial Defeat Stress-Induced Neurobehavioral Deficits in Mice: Involvement of Glutamic Acid Decarboxylase Isoform-67, Oxido-Nitregic Stress, and Neuroinflammatory Mechanisms. *J. Mol. Neurosci.* **2020**, *71*, 431–445. [[CrossRef](#)]
93. Tayyab, M.; Farheen, S.; Mariyath, M.; Khanam, P.M.N.; Hossain, M.M. Antidepressant and Neuroprotective Effects of Naringenin via Sonic Hedgehog-GLI1 Cell Signaling Pathway in a Rat Model of Chronic Unpredictable Mild Stress. *Neuromol. Med.* **2019**, *21*, 250–261. [[CrossRef](#)]
94. Yamagishi, N.; Yamamoto, Y.; Noda, C.; Hatayama, T. Naringenin Inhibits the Aggregation of Expanded Polyglutamine Tract-Containing Protein through the Induction of Endoplasmic Reticulum Chaperone GRP78. *Biol. Pharm. Bull.* **2012**, *35*, 1836–1840. [[CrossRef](#)]
95. Kumar, P.; Kumar, A. Protective Effect of Hesperidin and Naringin against 3-Nitropropionic Acid Induced Huntington's like Symptoms in Rats: Possible Role of Nitric Oxide. *Behav. Brain Res.* **2010**, *206*, 38–46. [[CrossRef](#)] [[PubMed](#)]
96. Salman, M.; Sharma, P.; Alam, I.; Tabassum, H. Naringenin Mitigates Behavioral Alterations and Provides Neuroprotection against 3-Nitropropionic Acid-Induced Huntington's Disease like Symptoms in Rats. *Nutr. Neurosci.* **2021**, *25*, 1–11. [[CrossRef](#)]
97. Cui, J.; Wang, G.; Kandhare, A.D.; Mukherjee, A.A.; Bodhankar, S.L. Neuroprotective Effect of Naringin, a Flavone Glycoside in Quinolinic Acid-Induced Neurotoxicity: Possible Role of PPAR- γ , Bax/Bcl-2, and Caspase-3. *Food Chem. Toxicol.* **2018**, *121*, 95–108. [[CrossRef](#)] [[PubMed](#)]
98. Gopinath, K.; Sudhandiran, G. Naringin modulates oxidative stress and inflammation in 3-nitropropionic acid-induced neurodegeneration through the activation of nuclear factor-erythroid 2-related factor-2 signalling pathway. *Neuroscience* **2012**, *227*, 134–143. [[CrossRef](#)]
99. Kulasekaran, G.; Ganapasam, S. Neuroprotective Efficacy of Naringin on 3-Nitropropionic Acid- Induced Mitochondrial Dysfunction through the Modulation of Nrf2 Signaling Pathway in PC12 Cells. *Mol. Cell. Biochem.* **2015**, *409*, 199–211. [[CrossRef](#)] [[PubMed](#)]
100. Gaur, V.; Aggarwal, A.; Kumar, A. Protective Effect of Naringin against Ischemic Reperfusion Cerebral Injury: Possible Neurobehavioral, Biochemical and Cellular Alterations in Rat Brain. *Eur. J. Pharmacol.* **2009**, *616*, 147–154. [[CrossRef](#)] [[PubMed](#)]
101. Okuyama, S.; Yamamoto, K.; Mori, H.; Sawamoto, A.; Amakura, Y. Neuroprotective Effect of Citrus Kawachiensis (Kawachi Bankan) Peels, a Rich Source of Naringin, against Global Cerebral Ischemia/Reperfusion Injury in Mice. *Biosci. Biotechnol. Biochem.* **2018**, *82*, 1216–1224. [[CrossRef](#)]
102. Han, Y.; Su, J.; Liu, X.; Zhao, Y.; Wang, C.; Li, X. Naringin Alleviates Early Brain Injury after Experimental Subarachnoid Hemorrhage by Reducing Oxidative Stress and Inhibiting Apoptosis. *Brain Res. Bull.* **2016**, *133*, 42–50. [[CrossRef](#)]
103. Bai, X.; Zhang, X.; Chen, L. Protective Effect of Naringenin in Experimental Ischemic Stroke: Down-Regulated NOD2, RIP2, NF-KappaB, MMP-9 and Up-Regulated Claudin-5 Expression. *Neurochem. Res.* **2014**, *39*, 1405–1415. [[CrossRef](#)]
104. Feng, J.; Chen, X.; Lu, S.; Li, W.; Yang, D.; Su, W.; Wang, X. Naringin Attenuates Cerebral Ischemia-Reperfusion Injury Through Inhibiting Peroxynitrite-Mediated Mitophagy Activation. *Mol. Neurobiol.* **2018**, *55*, 9029–9042. [[CrossRef](#)]
105. Cui, Q.; Wang, L.; Wei, Z.; Qu, W. Continual Naringin Treatment Benefits the Recovery of Traumatic Brain Injury in Rats Through Reducing Oxidative and Inflammatory Alterations. *Neurochem. Res.* **2014**, *39*, 1254–1262. [[CrossRef](#)] [[PubMed](#)]
106. Rong, W.; Pan, Y.; Cai, X.; Song, F.; Zhao, Z.; Xiao, S.; Zhang, C. The Mechanism of Naringin-Enhanced Remyelination after Spinal Cord Injury. *Neural Regen. Res.* **2017**, *12*, 470. [[PubMed](#)]
107. Fakhri, S.; Sabouri, S.; Kiani, A.; Farzaei, M.H.; Rashidi, K. Intrathecal Administration of Naringenin Improves Motor Dysfunction and Neuropathic Pain Following Compression Spinal Cord Injury in Rats: Relevance to Its Antioxidant and Anti-Inflammatory Activities. *Korean J. Pain* **2022**, *35*, 291–302. [[CrossRef](#)] [[PubMed](#)]
108. Shi, L.; Tang, P.; Zhang, W.; Zhao, Y.; Zhang, L.; Zhang, H. Naringenin Inhibits Spinal Cord Injury-Induced Activation of Neutrophils through MiR-223. *Gene* **2016**, *592*, 128–133. [[CrossRef](#)] [[PubMed](#)]
109. Wang, L.; Zhang, Z.; Wang, H. Naringin Attenuates Cerebral Ischemia-Reperfusion Injury in Rats by Inhibiting Endoplasmic Reticulum Stress. *Transl. Neurosci.* **2021**, *12*, 190–197. [[CrossRef](#)]
110. Hu, C.; Zhao, Y. Analgesic Effects of Naringenin in Rats with Spinal Nerve Ligation-Induced Neuropathic Pain. *Biomed. Rep.* **2014**, *2*, 569–573. [[CrossRef](#)]
111. Kaulaskar, S.; Bhutada, P.; Rahigude, A.; Jain, D.; Harle, U. Effects of Naringenin on Allodynia and Hyperalgesia in Rats with Chronic Constriction Injury-Induced Neuropathic Pain. *J. Chin. Integr. Med.* **2012**, *10*, 1482–1489. [[CrossRef](#)]

112. Semis, H.; Kandemir, F.; Caglayan, C.; Kaynar, O.; Genc, A.; Arıkan, S. Protective Effect of Naringin against Oxaliplatin-induced Peripheral Neuropathy in Rats: A Behavioral and Molecular Study. *J. Biochem. Mol. Toxicol.* **2022**, *36*, e23121. [[CrossRef](#)]
113. Yang, J.; Yuan, L.; Wen, Y.; Zhou, H.; Jiang, W.; Xu, D.; Wang, M. Protective Effects of Naringin in Cerebral Infarction and Its Molecular Mechanism. *Med. Sci. Monit. Int. Med. J. Exp. Clin. Res.* **2020**, *26*, e918772. [[CrossRef](#)]
114. Sveinsson, O.A.; Kjartansson, O.; Valdimarsson, E.M. Cerebral Ischemia/Infarction-Diagnosis and Treatment. *Laeknabladid* **2014**, *100*, 393–401.
115. Olmez, I.; Ozyurt, H. Reactive Oxygen Species and Ischemic Cerebrovascular Disease. *Neurochem. Int.* **2012**, *60*, 208–212. [[CrossRef](#)] [[PubMed](#)]
116. Tsung, A.; Klune, J.R.; Zhang, X.; Jeyabalan, G.; Cao, Z.; Peng, X.; Stolz, D.B.; Geller, D.A.; Rosengart, M.R.; Billiar, T.R. HMGB1 Release Induced by Liver Ischemia Involves Toll-like Receptor 4-Dependent Reactive Oxygen Species Production and Calcium-Mediated Signaling. *J. Exp. Med.* **2007**, *204*, 2913–2923. [[CrossRef](#)] [[PubMed](#)]
117. Ma, L.L.; Song, L.; Yu, X.D.; Yu, T.X.; Liang, H.; Qiu, J.X. The Clinical Study on the Treatment for Acute Cerebral Infarction by Intra-Arterial Thrombolysis Combined with Mild Hypothermia. *Eur. Rev. Med. Pharmacol. Sci.* **2017**, *21*, 1999–2006.
118. Jeong, H.S.; Song, H.-J.; Kim, S.-B.; Lee, J.; Kang, C.W.; Koh, H.-S.; Shin, J.E.; Lee, S.H.; Kwon, H.J.; Kim, J. A Comparison of Stent-Assisted Mechanical Thrombectomy and Conventional Intra-Arterial Thrombolysis for Acute Cerebral Infarction. *J. Clin. Neurol.* **2013**, *9*, 91–96. [[CrossRef](#)] [[PubMed](#)]
119. Bailey, D.G.; Arnold, J.M.O.; Strong, H.A.; Munoz, C.; Spence, J.D. Effect of Grapefruit Juice and Naringin on Nisoldipine Pharmacokinetics. *Clin. Pharmacol. Ther.* **1993**, *54*, 589–594. [[CrossRef](#)] [[PubMed](#)]
120. Cao, W.; Feng, S.-J.; Kan, M.-C. Naringin Targets NFKB1 to Alleviate Oxygen-Glucose Deprivation/Reoxygenation-Induced Injury in PC12 Cells via Modulating HIF-1 α /AKT/MTOR-Signaling Pathway. *J. Mol. Neurosci.* **2021**, *71*, 101–111. [[CrossRef](#)] [[PubMed](#)]
121. Zhang, Y.; Zhang, J.; Wu, C.; Guo, S.; Su, J.; Zhao, W.; Xing, H. Higenamine Protects Neuronal Cells from Oxygen-glucose Deprivation/Reoxygenation-induced Injury. *J. Cell. Biochem.* **2019**, *120*, 3757–3764. [[CrossRef](#)] [[PubMed](#)]
122. Shu, K.; Zhang, Y. Protodioscin Protects PC12 Cells against Oxygen and Glucose Deprivation-Induced Injury through MiR-124/AKT/Nrf2 Pathway. *Cell Stress Chaperones* **2019**, *24*, 1091–1099. [[CrossRef](#)]
123. Wang, C.; Yang, Y.-H.; Zhou, L.; Ding, X.-L.; Meng, Y.-C.; Han, K. Curcumin Alleviates OGD/R-Induced PC12 Cell Damage via Repressing CCL3 and Inactivating TLR4/MyD88/MAPK/NF-KB to Suppress Inflammation and Apoptosis. *J. Pharm. Pharmacol.* **2020**, *72*, 1176–1185. [[CrossRef](#)]
124. Rahman, J.; Tareq, A.M.; Hossain, M.; Sakib, S.A.; Islam, M.N.; Ali, M.H.; Uddin, A.B.M.N.; Hoque, M.; Nasrin, M.S.; Emran, T.B.; et al. Biological evaluation, DFT calculations and molecular docking studies on the antidepressant and cytotoxicity activities of *Cycas pectinata* Buch. - Ham. Compounds. *Pharmaceuticals* **2020**, *13*, 232. [[CrossRef](#)]
125. Tejera, D.; Heneka, M.T. Microglia in Neurodegenerative Disorders. In *Microglia*; Springer: New York, NY, USA, 2019; pp. 57–67.
126. Abdel-Magied, N.; Shedid, S.M. The Effect of Naringenin on the Role of Nuclear Factor (Erythroid-derived 2)-like2 (Nrf2) and Haem Oxygenase 1 (HO-1) in Reducing the Risk of Oxidative Stress-related Radiotoxicity in the Spleen of Rats. *Environ. Toxicol.* **2019**, *34*, 788–795. [[CrossRef](#)] [[PubMed](#)]
127. Drishya, S.; Dhanisha, S.S.; Raghukumar, P.; Guruvayoorappan, C. Amomum Subulatum Mitigates Total Body Irradiation-Induced Oxidative Stress and Associated Inflammatory Responses by Enhancing the Antioxidant Status and Regulating pro-Inflammatory Cytokines. *J. Nutr. Biochem.* **2022**, *107*, 109064. [[CrossRef](#)] [[PubMed](#)]
128. Saha, S.; Buttari, B.; Panieri, E.; Profumo, E.; Saso, L. An Overview of Nrf2 Signaling Pathway and Its Role in Inflammation. *Molecules* **2020**, *25*, 5474. [[CrossRef](#)] [[PubMed](#)]
129. Li, M.; Fu, Q.; Li, Y.; Li, S.; Xue, J.; Ma, S. Emodin Opposes Chronic Unpredictable Mild Stress Induced Depressive-like Behavior in Mice by Upregulating the Levels of Hippocampal Glucocorticoid Receptor and Brain-Derived Neurotrophic Factor. *Fitoterapia* **2014**, *98*, 1–10. [[CrossRef](#)] [[PubMed](#)]
130. Mitra, S.; Anjum, J.; Muni, M.; Das, R.; Rauf, A.; Islam, F.; Emran, T.B.; Semwal, P.; Hemeg, H.A.; Alhumaydhi, F.A. Exploring the Journey of Emodin as a Potential Neuroprotective Agent: Novel Therapeutic Insights with Molecular Mechanism of Action. *Biomed. Pharmacother.* **2022**, *149*, 112877. [[CrossRef](#)]
131. Mitra, S.; Lami, M.S.; Uddin, T.M.; Das, R.; Islam, F.; Anjum, J.; Hossain, M.J.; Emran, T.B. Prospective multifunctional roles and pharmacological potential of dietary flavonoid naringenin. *Biomed. Pharmacother.* **2022**, *150*, 112932. [[CrossRef](#)]
132. Halbreich, U.; Kahn, L.S. Role of Estrogen in the Aetiology and Treatment of Mood Disorders. *CNS Drugs* **2001**, *15*, 797–817. [[CrossRef](#)]
133. Bruce, M.L.; Raue, P.J. Mental Illness as Psychiatric Disorder. In *Handbook of the Sociology of Mental Health*; Springer: Berlin/Heidelberg, Germany, 2013; pp. 41–59.
134. Evans, D.L.; Foa, E.B.; Gur, R.E.; Hendin, H.; O'Brien, C.P.; Seligman, M.E.P.; Walsh, B.T. *Treating and Preventing Adolescent Mental Health Disorders: What We Know and What We Don't Know*; Oxford University Press: Oxford, UK, 2005; ISBN 0198038712.
135. Chtourou, Y.; Gargouri, B.; Kebieche, M.; Fetoui, H. Naringin Abrogates Cisplatin-Induced Cognitive Deficits and Cholinergic Dysfunction through the down-Regulation of AChE Expression and INOS Signaling Pathways in Hippocampus of Aged Rats. *J. Mol. Neurosci.* **2015**, *56*, 349–362. [[CrossRef](#)]

136. Kwatra, M.; Jangra, A.; Mishra, M.; Sharma, Y.; Ahmed, S.; Ghosh, P.; Kumar, V.; Vohora, D.; Khanam, R. Naringin and Sertraline Ameliorate Doxorubicin-Induced Behavioral Deficits through Modulation of Serotonin Level and Mitochondrial Complexes Protection Pathway in Rat Hippocampus. *Neurochem. Res.* **2016**, *41*, 2352–2366. [[CrossRef](#)]
137. Tallei, T.E.; Niode, N.J.; Idroes, R.; Zidan, B.M.R.M.; Mitra, S.; Celik, I.; Nainu, F.; Ağagündüz, D.; Emran, T.B.; Capasso, R.A. Comprehensive Review of the Potential Use of Green Tea Polyphenols in the Management of COVID-19 Evid-based Complement. *Altern. Med.* **2021**, *2021*, 7170736. [[CrossRef](#)]
138. D'Angelo, S.; Mele, E.; Di Filippo, F.; Viggiano, A.; Meccariello, R. Sirt1 Activity in the Brain: Simultaneous Effects on Energy Homeostasis and Reproduction. *Int. J. Environ. Res. Public Health* **2021**, *18*, 1243. [[CrossRef](#)] [[PubMed](#)]
139. Bhattacharya, I.; Domínguez, A.P.; Dräger, K.; Haas, E.; Battagay, E.J. Hypoxia Potentiates Tumor Necrosis Factor- α Induced Expression of Inducible Nitric Oxide Synthase and Cyclooxygenase-2 in White and Brown Adipocytes. *Biochem. Biophys. Res. Commun.* **2015**, *461*, 287–292. [[CrossRef](#)] [[PubMed](#)]
140. Matheus, M.E.; de Almeida Violante, F.; Garden, S.J.; Pinto, A.C.; Fernandes, P.D. Isatins Inhibit Cyclooxygenase-2 and Inducible Nitric Oxide Synthase in a Mouse Macrophage Cell Line. *Eur. J. Pharmacol.* **2007**, *556*, 200–206. [[CrossRef](#)] [[PubMed](#)]
141. Shin, J.-S.; Park, Y.M.; Choi, J.-H.; Park, H.-J.; Shin, M.C.; Lee, Y.S.; Lee, K.-T. Sulfuretin Isolated from Heartwood of Rhus Verniciflua Inhibits LPS-Induced Inducible Nitric Oxide Synthase, Cyclooxygenase-2, and pro-Inflammatory Cytokines Expression via the down-Regulation of NF-KB in RAW 264.7 Murine Macrophage Cells. *Int. Immunopharmacol.* **2010**, *10*, 943–950. [[CrossRef](#)]
142. Feifel, D.; Shilling, P.D.; MacDonald, K. A Review of Oxytocin's Effects on the Positive, Negative, and Cognitive Domains of Schizophrenia. *Biol. Psychiatry* **2016**, *79*, 222–233. [[CrossRef](#)]
143. Lisman, J.E.; Coyle, J.T.; Green, R.W.; Javitt, D.C.; Benes, F.M.; Heckers, S.; Grace, A.A. Circuit-Based Framework for Understanding Neurotransmitter and Risk Gene Interactions in Schizophrenia. *Trends Neurosci.* **2008**, *31*, 234–242. [[CrossRef](#)]
144. Obrosova, I.G. Diabetes and the Peripheral Nerve. *Biochim. Biophys. Acta Mol. Basis Dis.* **2009**, *1792*, 931–940. [[CrossRef](#)]
145. Howes, O.; McCutcheon, R.; Stone, J. Glutamate and Dopamine in Schizophrenia: An Update for the 21st Century. *J. Psychopharmacol.* **2015**, *29*, 97–115. [[CrossRef](#)]
146. Li, P.-F.; Dietz, R.; von Harsdorf, R. P53 Regulates Mitochondrial Membrane Potential through Reactive Oxygen Species and Induces Cytochrome C-Independent Apoptosis Blocked by Bcl-2. *EMBO J.* **1999**, *18*, 6027–6036. [[CrossRef](#)]
147. Zou, W.; Xiao, Z.; Wen, X.; Luo, J.; Chen, S.; Cheng, Z.; Xiang, D.; Hu, J.; He, J. The Anti-Inflammatory Effect of *Andrographis paniculata* (Burm. f.) Nees on Pelvic Inflammatory Disease in Rats through down-Regulation of the NF-KB Pathway. *BMC Complement. Altern. Med.* **2016**, *16*, 483. [[CrossRef](#)]
148. Stenvinkel, P.; Ketteler, M.; Johnson, R.J.; Lindholm, B.; Pecoits-Filho, R.; Riella, M.; Heimbürger, O.; Cederholm, T.; Girndt, M. IL-10, IL-6, and TNF- α : Central Factors in the Altered Cytokine Network of Uremia—The Good, the Bad, and the Ugly. *Kidney Int.* **2005**, *67*, 1216–1233. [[CrossRef](#)] [[PubMed](#)]
149. Yao, Q.; Pecoits-Filho, R.; Lindholm, B.; Stenvinkel, P. Traditional and Non-Traditional Risk Factors as Contributors to Atherosclerotic Cardiovascular Disease in End-Stage Renal Disease. *Scand. J. Urol. Nephrol.* **2004**, *38*, 405–416. [[CrossRef](#)] [[PubMed](#)]
150. Higley, M.J.; Picciotto, M.R. Neuromodulation by Acetylcholine: Examples from Schizophrenia and Depression. *Curr. Opin. Neurobiol.* **2014**, *29*, 88–95. [[CrossRef](#)] [[PubMed](#)]
151. Futamura, T.; Toyooka, K.; Iritani, S.; Niizato, K.; Nakamura, R.; Tsuchiya, K.; Someya, T.; Kakita, A.; Takahashi, H.; Nawa, H. Abnormal Expression of Epidermal Growth Factor and Its Receptor in the Forebrain and Serum of Schizophrenic Patients. *Mol. Psychiatry* **2002**, *7*, 673–682. [[CrossRef](#)] [[PubMed](#)]
152. Zhong, J.; Yu, R.; Zhou, Q.; Liu, P.; Liu, Z.; Bian, Y. Naringenin Prevents TNF- α -Induced Gut-Vascular Barrier Disruption Associated with Inhibiting the NF-KB-Mediated MLCK/p-MLC and NLRP3 Pathways. *Food Funct.* **2021**, *12*, 2715–2725. [[CrossRef](#)]
153. Bitanhirwe, B.K.Y.; Woo, T.-U.W. Oxidative Stress in Schizophrenia: An Integrated Approach. *Neurosci. Biobehav. Rev.* **2011**, *35*, 878–893. [[CrossRef](#)]
154. Jarskog, L.F.; Glantz, L.A.; Gilmore, J.H.; Lieberman, J.A. Apoptotic Mechanisms in the Pathophysiology of Schizophrenia. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* **2005**, *29*, 846–858. [[CrossRef](#)]
155. Missonnier, P.; Hasler, R.; Perroud, N.; Herrmann, F.R.; Millet, P.; Richiardi, J.; Malafosse, A.; Giannakopoulos, P.; Baud, P. EEG Anomalies in Adult ADHD Subjects Performing a Working Memory Task. *Neuroscience* **2013**, *241*, 135–146. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.