

Systematic Review/Meta-Analysis

The Effectiveness of Physical Therapy Interventions for Athletes Post-Concussion: A Systematic Review

Kylie Art¹, Courtney Ridenour¹, Stephanie Durbin¹, Max Bauer¹, Amy Hassen-Miller^{1 a}¹ Physical Therapy, Walsh University

Keywords: athlete, adolescent, concussion, exercise, physical therapy intervention

<https://doi.org/10.26603/001c.68071>

International Journal of Sports Physical Therapy

Vol. 18, Issue 1, 2023

Background

Sports-related concussions are the most common cause of head injury in adolescents and young adults. Typical treatment methods for this injury include cognitive and physical rest. Evidence suggests that physical activity and physical therapy interventions can be beneficial to decrease post-concussion symptoms.

Purpose

The aim of this systematic review was to investigate the effectiveness of physical therapy interventions for adolescent and young adult athletes post-concussion.

Study Design

Systematic Review

Methods

The following databases were utilized for the search: PubMed, CINAHL, Proquest, MEDLINE, SPORTDiscus, and SCOPUS. The search strategy focused on athletes, concussions, and physical therapy interventions. Data extraction from each article included: Authors, subjects, gender, mean age, age range, specific sport, acute or chronic concussion, first or recurrent concussion, treatments for intervention and control group, and outcomes measured.

Results

Eight studies met the inclusion criteria. Six of the eight articles scored seven or higher on the PEDro Scale. Physical therapy intervention(s), such as an aerobic intervention or a multimodal approach, show improvements in time to recovery and reduction of post-concussion symptoms in patients who have had a concussion. Physical activity and physical therapy as early as a couple days following injury, is beneficial at decreasing post-concussion symptoms, allows for earlier return to play, and/or shorter days to recovery, and is considered safe for treating post-concussion symptoms.

Conclusion

This systematic review demonstrates that physical therapy interventions including aerobic exercise and multimodal approaches have been found to be beneficial in treating adolescent and young adult athletes post-concussion. Utilizing aerobic or multimodal interventions for this population allows for a quicker symptom recovery and return to

^a **Corresponding Author:**

Dr. Amy Hassen-Miller
Associate Professor
Walsh University
2020 E. Maple St.
North Canton, OH, 44702
E-Mail: ahassenmiller@walsh.edu

sport than traditional treatment of physical and cognitive rest. Future research should investigate the superior intervention for adolescents and young adults with post-concussion syndrome and determine if a single treatment or a multimodal approach is more beneficial.

INTRODUCTION

Concussion is a mild traumatic brain injury. Concussion occurs as a result of direct trauma, whiplash, or blast injury that causes the brain to move forward and backward. Symptoms associated with concussion include headache, dizziness, vertigo, nausea, fatigue, light and noise sensitivity, as well as altered mental status.^{1,2} The leading cause of head injury in adults occurs from car accidents, whereas in the younger population (15-24 years old), injury occurs from a sport related accident. Of the forty-four million children involved in various sports-related activities, around 3.8 million experience a concussion related to sport each year.² Emergency department visits due to sport-related concussions increased by more than double between 2001-2012.^{1,3} The most common cause of sport-related concussion is player to player contact and happens frequently in football and ice hockey. Female athletes have a higher incidence of concussion compared to male athletes. Athletes who have had a prior concussion also have a higher rate of experiencing additional concussions.^{2,4} Adolescent athletes are at higher risk of suffering a concussion than adults and may also require a longer period of time to recover.⁵ If an athlete is diagnosed with a concussion, they must be cleared by a health care professional before returning to play.

Traditional treatment for concussion includes physical and cognitive rest.^{6,7} Unfortunately, individuals prescribed with rest can demonstrate withdrawal from life activities, deconditioning, and mental health issues.⁸ Research is emerging that shows the benefit of exercise and early physical activity after a concussion.² Specifically, early physical activity is advantageous and lower rates of post-concussive symptoms are associated with higher levels of physical activity and this research states that patients suffering from concussion should be encouraged to exercise as tolerated 1-2 days after concussion and initial rest.² Individuals with a concussion present with varying symptoms and presentations requiring patient-specific intervention strategies. According to a clinical practice guideline, recommended frequency, intensity, timing, and type(s) of intervention is dependent on the severity of a patient's concussion, their level of irritability, and how they respond to treatment. Patients should also be educated on symptom reporting so that interventions can be adjusted resulting in earlier return to sport.⁹

Existing literature suggests that cervical and vestibular interventions as well as sub-symptom threshold exercise may be beneficial for individuals 24-48, hours after concussion. Aerobic exercise has been shown to improve autonomic nervous system function suggesting that sub-threshold exercise treatment could be a beneficial treatment for concussion. Systematic reviews have shown that exercise post-concussion can improve symptom scores after concussion and that sub-symptomatic aerobic exer-

cise programs can be more effective in reducing post-concussion syndrome symptoms than traditional standard of care.¹⁰

There is a gap in the current literature regarding sports-related concussions in adolescent and young adult athletes and the appropriate interventions for treatment. The aim of this systematic review was to investigate the effectiveness of physical therapy interventions for adolescent and young adult athletes post-concussion.

METHODS

This systematic review was registered with Prospero and completed using guidelines from Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA). PRISMA is a 27-item checklist based on evidence that focuses on reporting the impact of interventions in systematic reviews and meta-analysis.¹¹ Prospero provides the opportunity to determine if a similar review already exists or is in the process of being published to prevent unwanted duplication.¹² The registration ID for Prospero for this systematic review is CRD42021256465.

ELIGIBILITY CRITERIA

In order to be included in this systematic review, articles had to meet the following inclusion criteria: be original research, a randomized controlled-trial, include participants that were athletes under the age of thirty, have an acute or chronic sport-related concussion, and include physical therapy intervention(s). Randomized controlled-trial articles were chosen due to high placement on the hierarchy of evidence and to strengthen the results of this systematic review. Articles were excluded if they were not of a randomized controlled-trial study design, not reported in English, or did not fit the inclusion criteria stated.

SEARCH STRATEGIES

The following databases were utilized for the search: PubMed, CINAHL, Proquest, MEDLINE, SPORTDiscus, and SCOPUS. The search strategy focused on three key concepts: athletes, concussions, and physical therapy interventions. After these concepts were identified, similar terms were noted, defined, and used in the searches. These terms were then utilized in each database and any database-specific terms such as Subject Heading (SH) or Medical Subject Heading (MESH) when relevant. Appendix 1 provides the specific search strategies utilized in this search. A hand search was completed by looking through the references of articles included in this systematic review. The initial search was started in March 2021 and the final search concluded in January 2022.

STUDY SELECTION

The total number of abstracts, titles, and full texts from the search strategy were divided in half and allocated to two teams of two authors. Each title, abstract, and full text for each article was assessed independently and screened by two separate authors to be reviewed for inclusion in the systematic review. If a tie breaker was needed due to a disagreement between team members, then a third author would serve as a tie breaker. Any duplicates found during the search were removed. Cohen's unweighted Kappa was calculated for agreement during the title, abstract, and full text review. "Kappa is a measurement that is used to determine agreement among raters when scores are repeated."¹³ Recommended values for Kappa include: <0.00 is poor agreement, (0.01-0.20) is slight agreement among raters, (0.21-0.40) is considered fair agreement, (0.41-0.60) is moderate, (0.61-0.80) is considered substantial agreement among raters, and (0.81-1.0) is almost perfect agreement."¹³

RISK OF BIAS IN INCLUDED STUDIES

Risk of bias assessment for included studies was performed by two authors independently and was reviewed for each of the articles. If a disagreement was present, a third author would serve as a tiebreaker to reach consensus. Risk of bias assessment in the included studies was assessed using the PEDro scale. The PEDro scale is an appropriate tool to assess the quality of randomized controlled trials. A higher score on the PEDro scale indicates a lower risk of bias in an article, with the highest score possible being a 10 and lowest score of a 0. The PEDro scale has 11 criterion but is only scored using 10, thus has a score out of 10. One item, the item on eligibility, is not counted towards the score because it is related to external validity. The PEDro scale does not assess external validity or effect size of treatment.¹⁴

DATA EXTRACTION

Two authors performed data extraction independently. Consensus was reached after independent extraction. Data extraction included: Authors, subjects, gender, mean age, acute or chronic concussion, treatment for intervention group versus control group, and outcomes measured.

RESULTS

STUDY SELECTION

The initial search results produced 7,034 titles. There were 3,253 titles found after all duplicates were removed. Two authors screened titles and abstracts and 3,193 articles were excluded. A hand search was completed, and two additional randomized controlled trials were included. After title and abstract review, 60 articles were chosen for full-text review with an observed kappa value of one and confidence interval of one to one demonstrating perfect agreement between reviewers. Non-randomized control trials were excluded as well as one ongoing study, which resulted in eight

articles that fit the inclusion criteria to be included in the systematic review. Details of the study selection can be seen in [Figure 1](#).

STUDY CHARACTERISTICS

Aerobic intervention for the treatment groups was used for four of the eight articles, and multimodal intervention for the treatment groups was used for the remaining four articles. The control groups for each of the articles either participated in education, rest, sub-therapeutic exercise, placebo, or sham treatment. The outcome measures varied throughout the eight studies and included: days from injury to recovery, PCSS, PCSI, Health Related Quality of Life, Beck Depression Inventory for Youth, Pediatric Quality of Life Multidimensional Fatigue Scale, the BESS, ImpACT, time to return to play, health and demographic questionnaires, Borg CR10, Rate of Perceived Exertion (RPE), and Post Ride Symptom Change Rating. See [Table 1](#) for a description of the outcome measures that were used in the articles included in this systematic review. In total, there were 375 subjects that participated in the eight studies with 163 females and 212 males. The age range of participants included was 11.2-21.2 years old. Four articles included subjects who had acute post-concussion symptoms, one article included participants with chronic post-concussion symptoms, two articles included participants with both acute and chronic post-concussion symptoms, and one article did not indicate severity of concussion. Six of the eight articles included participants who have had one concussion or a recurrence of concussions while the other two articles did not indicate occurrence of concussion. Study characteristics can be found in [Table 2](#).

QUALITY ASSESSMENT

Two reviewers assessed the risk of bias of the eight articles included in this systematic review using the PEDro Scale. Each of the articles was given a PEDro Scale score, shown in [Table 3](#). To evaluate the level of agreement between the two reviewers, an unweighted kappa was calculated totaling one, (CI = 1:1), which demonstrates perfect agreement. Five of the eight articles were of a higher quality relating to risk of bias compared to the other three included articles, with scores ranging from a 5 to a 10 out of 10 items. Three of the five higher quality articles found significant improvements in the treatment groups compared to the control groups.^{20,21,24} Although only three articles demonstrated statistically significant results for the treatment group compared to the control group, all articles demonstrated improved symptoms with time and there were no adverse effects reported with early physical activity and therapy. Three of the eight articles had a lower quality relating to risk of bias scoring a 3 to 4 out of 10 items.^{19,25,26} Criterion 1, 2, 4 and 10 were met by all articles included in this study.




Figure 1. Study Selection.

RESULTS OF INDIVIDUAL STUDIES (INTERVENTIONS)

AEROBIC INTERVENTION

Four of the eight studies included in the systematic review used aerobic interventions for the treatment group.^{19-21,26} Leddy et al.²⁰ included aerobic exercise on a stationary bike or treadmill if available, or walking/jogging outdoors if they did not have access to exercise equipment. The aerobic exercise would last 20 minutes or until their symptoms increased by two or more points from their pre-exercise symptom level (based on a 10-point Visual Analogue Scale). Participants in the aerobic exercise group recovered in a median of 13 days compared to the control group who recovered in 17 days ($p = 0.009$). Leddy et al.²¹ included aerobic exercise such as walking, jogging, or stationary biking. The aerobic group would exercise for 20 minutes and up to 90% of their maximum heart rate achieved on the Buffalo Concussion Treadmill Test. They were instructed to stop when reaching 90% of their maximum heart rate or when they experienced concussion-like symptoms. Median days to recovery for the aerobic exercise group was 14 days compared to 19 days for the stretching group. Four out of 43 participants in the aerobic exercise group (9%) had persistent post-concussive symptoms, compared to 12 out of 39 participants in the stretching group (31%), ($p=0.014$ with

Fisher's exact test). Maerlender et al.¹⁹ had participants ride on a stationary bike with an RPE of 0 - 6 for 20 minutes or until they felt uncomfortable. There was no significant difference between groups for days of recovery, with a median of 13 days to recovery for the standard group, and 15 days to recovery for the exertion group. Micay et al.²⁶ had participants complete eight sessions that progressed with duration and intensity using a stationary bike that was connected to a heart rate monitor. Each session started with a five-minute warm up and ended with a five-minute cool down. If symptoms worsened, intervention was terminated. The aerobic exercise group showed greater symptom resolution compared to the control group over time, however there was not a significant difference between groups. Independent t-tests were performed for or between-group difference in mean time to medical clearance which was 36.1 ± 18.5 days for the aerobic exercise group and 29.6 ± 15.8 days for the control group ($t(13)=0.03$; $p=0.87$).

MULTIMODAL INTERVENTION

Four of the eight studies included in the systematic review used multimodal intervention for the intervention provided to the treatment group.²²⁻²⁵ Chan et al.²² utilized an active rehabilitation program including submaximal aerobic training, light coordination, sport-specific exercises, visu-

Table 2. Study Characteristic

Author	Intervention	Control	Outcome Measure	Acute vs chronic and definition	Intervention group		Control Group	
					N	Mean Age (SD) or [Range]	N	Mean Age (SD)
Leddy ²⁰	Aerobic Exercise on bike or treadmill	Placebo-like stretching regimen	Days from injury to recovery; symptom severity score, proportion of participants with delayed recovery	Acute, starting within 10 days	52 (24 females, 28 male)	15.3 (1.6)	51 (24 female, 27 male)	15.4 (1.7)
Leddy, Master ²¹	Aerobic exercise: walking, jogging, or stationary bike	Placebo-like stretching regimen	PSCI, days to recovery, development of post-concussive symptoms	Acute, 24 hrs-10 days post-concussion diagnosis)	61 (23 females, 38 males)	15.5 (1.4)	57 (21 females, 36 males)	15.9 (1.4)
Chan ²²	Aerobic Training: coordination exercises and visualization/imagery techniques	Usual Treatment (education about symptom management, school consultation with a hospital affiliated teacher who facilitated return to school, physician consult for medication and referral for community therapy)	PCSS, health-related quality of life, beck depression inventory for youth, pediatric quality of life, multidimensional fatigue scale, balance error scoring system, immediate post-concussion assessment	post-concussion symptoms \geq 1 month; \geq 2 persistent post-concussion symptoms	10 (8 females, 2 males)	15.1 (1.47)	9 (6 females, 3 males)	15.9 (1.66)
Reneker ²³	Pragmatically designed individualized and progressive treatment plan	sham, subtherapeutic, and non-progressive therapeutic techniques to minimally progressive therapy	PCSS, time to return to play	acute (up to 14 days post concussion to start treatment)	22 (14 males, 8 females)	16.5 (2.9)	19 (11 males, 8 females)	15.9 (2.9)
Schneider ²⁴	Postural education, range of motion exercises and cognitive and physical rest, until asymptomatic followed by a protocol of graded exertion. Along with cervical spine and vestibular rehab	Same as the intervention group except the cervical spine and vestibular rehab.	Return to play (days)	Both, but more chronic than acute	15 (11 males, 4 females)	15 [12-27]	16 (7 males, 9 females)	16 [7 males, 9 females]

Author	Intervention	Control	Outcome Measure	Acute vs chronic and definition	Intervention group		Control Group	
					N	Mean Age (SD) or [Range]	N	Mean Age (SD)
Dobney ²⁵	4 components: (1) aerobic activity, (2) coordination/skill practice, (3) visualization, and (4) education beginning 2 weeks post injury	4 components: (1) aerobic activity, (2) coordination/skill practice, (3) visualization, and (4) education beginning at the standard 4 week post injury	PCSI	Acute. Starting either 2 or 4 weeks after concussion	10 (4 females, 6 males)	15.6 (1.8)	10 (4 females, 6 males)	14.3 (2.7)
Maerlender ¹⁹	Exertion: Stationary bike ride for 20 min unless uncomfortable for 0-6 on RPE scale (mild to moderate exertion)	No physical exertion other than walking to school or daily activities.	Recovery length-symptom free from bike ride, IMPACT, neurocognitive test battery, BORG CR10 RPE.	n/a	15 (12 females, 3 males)	N/a, Collegiate athletes	13 (8 females, 5 males)	N/a, Collegiate athletes
Micay ²⁶	Aerobic exercise-progressive intensity/duration on cycle ergometer;	usual care-rest followed by physician advised progressions of activity	Intervention feasibility: symptom status pre-post exercise sessions and completion of intervention; Clinical recovery: symptom status at weeks 1,2,3, and 4 post injury and medical clearance date using self reported symptom severity scores on PCSS and days to clearance	acute starting at day 6 post injury	7 (all males)	15.8 (1.2)	8 (all Males)	15.6 (1.0)

N=Number of participants; SD=Standard Deviation; PCSS=Post-concussion symptom scale; PCSI=Post-concussion symptom inventory

Table 3. Quality assessment using PEDro scale.

	Leddy ²⁰	Schneider ²⁴	Leddy, Master ²¹	Chan ²²	Reneker ²³	Dobney ²⁵	Maerlender ¹⁹	Micay ²⁶
Criterion 1	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Criterion 2	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Criterion 3	No	Yes	No	Yes	No	No	No	No
Criterion 4	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Criterion 5	No	Yes	No	Yes	Yes	No	No	No
Criterion 6	Yes	Yes	No	Yes	No	No	No	No
Criterion 7	Yes	Yes	Yes	Yes	Yes	No	No	No
Criterion 8	Yes	Yes	Yes	Yes	Yes	No	No	Yes
Criterion 9	Yes	Yes	Yes	Yes	Yes	No	No	No
Criterion 10	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Criterion 11	No	No	Yes	Yes	No	No	No	No
Score	7	9	7	10	7	3	3	4

1. Eligibility criteria were specified. 2. Subjects were randomly allocated to groups. 3. Allocation was concealed. 4. The groups were similar at baseline regarding the most important prognostic indicators. 5. There was blinding of all subjects. 6. There was blinding of all therapists who administered the therapy. 7. There was blinding of all assessors who measured at least one key outcome. 8. Measures of at least one key outcome were obtained from more than 85% of the subjects initially allocated to groups. 9. All subjects for whom outcome measures were available received the treatment or control condition as allocated or, where this was not the case, data for at least one key outcome was analyzed by "intention to treat." 10. The results of between group statistical comparisons are reported for at least one key outcome. 11. The study provides both point measures and measures of variability for at least one key outcome.

alization and imagery techniques, and a home exercise program. The active rehabilitation group had a mean \pm SD change of -24.7 ± 19.1 on the PCSS from baseline to follow-up. The control group had a mean \pm SD change of -15.8 ± 12.5 on the PCSS from baseline to follow-up. These results were associated with a significant difference between groups with a Cohen's *d* treatment medium effect size of 0.55. Reneker et al.²³ study's treatment group received manual techniques including soft tissue release, mobilizations, and thrust manipulations, and vestibular rehabilitation techniques including habituation, adaptation, oculomotor control, neuromotor control, and balance exercises. The interventions provided were tailored to each specific patient's needs and progressed over time as the patient's symptoms would allow. The median number of days to post-concussion syndrome (PCS) recovery for the experimental group was 13.5 days and 17 days for the control group, with a log-rank of 0.13 representing no statistical significance between groups. Schneider et al.²⁴ had participants in the treatment group receive cervical spine physical therapy interventions and vestibular rehabilitation accompanied by non-provocative range of motion exercises, stretching, and postural education. The cervical spine physical therapy interventions included manual therapy of the thoracic and cervical spine and therapeutic exercises including cervical neuromotor retraining exercises. The vestibular rehabilitation was individualized to each person and consisted of habituation, gaze stabilization, adaptation exercises, standing and dynamic balance exercises, as well as canalith repositioning maneuvers. There was a greater number of participants that were medically cleared to return to sport within eight weeks of treatment in the treatment group (11 of the 15 individuals), compared to the control group (one out of the 14 individuals). Individuals in the treatment group were 10.27 (95% CI 1.51 to 69.56) times more likely to be medically cleared to return to sport within eight weeks than the participants in the control group ($p < 0.001$). Dobney et al.²⁵ had participants in both the treatment group and control group complete 15 minutes on a stationary bike or treadmill, 10 minutes of coordination and skill practice related to sport preference, 5 to 10 minutes of visualization related to sport preference, and education about recovery, coping with post-concussion symptoms, and the process of returning to school or sport. The only difference between groups was that the treatment group started two weeks post-concussion and the control started treatment four weeks post-concussion. Participants were instructed to discontinue activity if symptoms increased. Post-concussion symptoms improved over time for both groups, with no significant difference between the two groups.

OVERALL

When looking at physical therapy interventions, whether it be aerobic interventions or a multimodal approach, the included studies show improvements in outcomes of time to recovery and reduction of post-concussion symptoms in patients who have had a concussion. Four of the studies had significant between-group differences. The other four

did not have significant between group differences but did show improvements overall with no significant adverse effects.

DISCUSSION

The aim of this systematic review was to investigate the effectiveness of physical therapy interventions for athletes post-concussion. There is a gap in the current literature regarding sports-related concussions in athletes and appropriate interventions for treatment. This review aims to reveal physical therapy interventions to address post-concussive symptoms. Overall, this review demonstrates decreased time to recovery and reduced post-concussion symptoms when patients receive active physical therapy interventions post-concussion. None of the studies demonstrated any significant adverse effects from early physical activity and physical therapy intervention, indicating that this treatment approach is safe for adolescent and young adult athletes post-concussion. Physical therapists should incorporate physical activity and physical therapy interventions earlier to optimize post-concussion recovery.

One limitation of this systematic review is that three articles were at high risk of bias according to PEDro, including Maerlender,¹⁹ Micay,²⁶ and Dobney.²⁵ Another limitation is that all studies included a small sample size that ranged from 15 to 118 total participants. This could result in larger effect sizes, decrease the power of the data, and be unrepresentative of the population. One article included only male participants decreasing the generalizability of the results.²⁶ Another limitation of this systematic review is that the articles reviewed incorporated both acute and chronic concussions. By including both, the variation in time to resolve symptoms and return to play following the concussion may impact the overall validity and generalizability of the results. Because some of the articles included a combination of treatment interventions for athletes post-concussion, this made determining which treatment was most effective and beneficial for the participant difficult to determine.

One strength of this systematic review is that only randomized control trials were included. Kappa statistics were calculated among authors and resulted in perfect or near perfect agreement and this review was conducted following PRISMA guidelines. Lastly, available articles for this systematic review are from the year 2014 to present, and attempt to provide the most up to date research on athletes with post-concussion syndrome.

A weakness of this review is the results from the search strategy captured physical therapy interventions that were similar in nature being either aerobic or multimodal, as opposed to a broader spectrum of potential interventions. Also, limited evidence is available regarding athletes post-concussion resulting in only eight randomized control trials available for this review.

There are multiple published systematic reviews that present similar findings. One included 14 articles comparing rest and exercise and concluded that exercise was beneficial at improving PCSS score, decreasing symptoms, and

Table 4. Outcomes.

Measured Outcome	Author	Intervention Baseline Outcome Measure Results	Intervention Final Outcome Measure Results	Control Baseline Outcome Measure Results	Control Final Outcome Measure Results	95% Confidence Interval	P Values and/or Effect Size
PCSS							
	Chan ²²	51.50 ±27.80 (SD) (mean)	25.00 ±19.40 (SD) (mean)	56.90 ±31.00 (SD) (mean)	40.30 ±29.40 (SD) (mean)	-	Effect size of .55
	Micay ²⁶	-	18.80 +4.90 (SD) (mean)	-	10.00 ±6.10 (SD) (mean)	-	Effect size of 1.6039
PCSI							
PCSI-SR13 for Youth	Dobney ²⁵	-	3 (median) (range 0-9)	-	2.25 (median) (range 0-24)	-	-
Recovery Time (Days)							
	Leddy ²⁰	-	13 days (median)	-	17 days (median)	-	p = 0.009
	Reneker ²³	-	13.5 days (median)	-	17 days (median)	(0.95-4.15)	-
	Leddy ²¹	-	14 days (median) (IQR 10-25)	-	19 days (median) (IQR 13-31)	-	-
	Maerlender ¹⁹	-	15 days (median) (range 5-61)	-	13 days (median) (range 6-56)	-	p=0.705
Return to Sport (Weeks)							
	Micay ²⁶	-	36.10±18.50 (SD) (mean)	-	29.60 ±15.80 (SD) (mean)	-	p = 0.87; Effect size of .3756
Amount medically cleared for return to sport within 8 weeks (number of participants)							
	Schneider ²⁴	-	11	-	1	(40-92.3)	p = < 0.001
Development of Persistent Post-Concussive Symptoms (number of participants)							
	Leddy ²¹	-	13	-	18	-	p=0.039

PCSS=Post-concussion symptom scale; PCSI=Post-concussion symptom inventory; SD=Standard deviation.

decreasing the duration of lingering post-concussion symptoms.²⁷ A systematic review that included five articles assessing athletes post-concussion with aerobic training demonstrated improvements in post-concussion symptoms.³ Along with these, a systematic review by Makdissi et al²⁸ also found evidence supporting the importance of including physical therapy and aerobic exercise in the treatment of a sports related concussion. This review specifically assessed athletes with persistent symptoms, but they found evidence that treating the patient’s specific persistent symptoms that was beneficial and safe. Lastly, a systematic review that assessed mode and intensity of treatment post sport-related concussion found that light, moderate, and high-intensity levels of exercises can be safe and beneficial for patients during treatment.

The results of this systematic review suggest that physical therapists should incorporate early physical activity

and physical therapy interventions into their treatment for adolescent and young adult athletes with post-concussion syndrome. Practicing therapists should determine the individualized impairment(s) present, treat the impairment(s), and use symptom reporting as a guideline to return the athlete to sport to ensure proper timing of interventions.^{8,9} Due to limitations of this current study, future research should investigate the impact of physical therapy interventions for adolescent and young adult athletes status post chronic versus acute concussion rather than mixing the two and research should be conducted with a larger sample size to allow for a more accurate effect size to be calculated. Future research should also investigate the superior intervention for adolescents and young adults suffering from post-concussion syndrome and determine if single treatment is more beneficial or if multimodal approach is more beneficial to the participants’ recovery. Lastly, future research

should include more randomized controlled trials to improve validity and enhance professional decision making.

CONCLUSION

The results of this systematic review indicate that there is evidence supporting the use of physical therapy interventions including aerobic or multi-modal treatments in the treatment of adolescent and young adult athletes post concussion. The evidence included in this systematic review suggests that aerobic exercise or multi-modal interventions may lead to quicker recovery and return to sport in adolescent and young adult athletes with post-concussive symptoms when compared to traditional treatments such

as physical and cognitive rest. Physical therapists should consider incorporation of these interventions into their plan of care for this population to allow for quicker recovery and return to sport.

.....

CONFLICTS OF INTEREST

The authors indicate no conflicts of interest in the reporting of this systematic review.

Submitted: April 20, 2022 CST, Accepted: November 30, 2022 CST



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

Table 1. Description of Outcome Measures Utilized.

Outcome Measures	Description
Balance Error Scoring System (BESS)	Measures postural stability and vestibular dysfunction related to return to play. ¹⁵ Used primarily in youth, athletes, and patients with a traumatic brain injury. ¹⁵
Beck Depression Inventory for Youth	Self-reported outcome measure with 21 items ranked on a four point scale to assess depression. ¹⁶ Score >10 is positive for depression which is common in athletes. ¹⁶ Used commonly for Parkinson's Disease, stroke, spinal cord, and non-specific populations.
Post-Concussion Symptom Inventory (PCSI)	26 item self-reported outcome measure for individuals aged 5-18. ¹⁷ Assess cognition, sleep, emotions, and physical domains.
Post-Concussion Symptom Scale (PCSS)	21 item self-reported outcome measure using a 7 point likert scale. Used to determine whether athletes have a concussion or not. ¹⁷
Immediate Post-Concussion Assessment and Cognitive Testing (ImPACT)	Online computerized neuropsychological series including patient demographics, determine if any of the 22 listed concussion symptoms are present and at what severity, and neuropsychological modules to assess attention, memory, processing speed, reaction time, sequencing, and learning. Used primarily with the athlete population. ¹⁷
BORG CR 10 RPE	Measures exercise intensity perceived by the patient during exercise and used primarily with patients with breathlessness, angina, and musculoskeletal pain. ¹⁸
Post Ride Symptom Change Rating	Experiential likert scale that ranks perceived exertion after biking. ¹⁹

REFERENCES

- Baker B, Koch E, Vicari K, Walenta K. Mode and intensity of physical activity during the postacute phase of sport-related concussion: a systematic review. *J Sport Rehabil.* 2020;30(3):492-500. doi:10.1123/jsr.2019-0323
- Mullally WJ. Concussion. *Am J Med.* 2017;130(8):885-892. doi:10.1016/j.amjmed.2017.04.016
- Powell C, McCaulley B, Scott Brosky Z, Stephenson T, Hassen-Miller A. The effect of aerobic exercise on adolescent athletes post-concussion: a systematic review and meta-analysis. *Int J Sports Phys Ther.* 2020;15(5):650-658. doi:10.26603/ijsp20200650
- Reneker JC, Babl R, Flowers MM. History of concussion and risk of subsequent injury in athletes and service members: a systematic review and meta-analysis. *Musculoskelet Sci Pract.* 2019;42:173-185. doi:10.1016/j.msksp.2019.04.004
- Kimble DE, Murphy M, Dhandapani KM. Concussion and the adolescent athlete. *J Neurosci Nurs.* 2011;43(6):286-290. doi:10.1097/jnn.0b013e31823858a6
- Leddy J, Hinds A, Sirica D, Willer B. The role of controlled exercise in concussion management. *PM R.* 2016;8(3 Suppl):S91-S100. doi:10.1016/j.pmrj.2015.10.017
- Silverberg ND, Iverson GL. Is rest after concussion "the best medicine?": recommendations for activity resumption following concussion in athletes, civilians, and military service members. *J Head Trauma Rehabil.* 2013;28(4):250-259. doi:10.1097/ht.r.0b013e31825ad658
- Sawyer Q, Vesci B, McLeod TC. Physical activity and intermittent postconcussion symptoms after a period of symptom-limited physical and cognitive rest. *J Athl Train.* 2016;51(9):739-742. doi:10.4085/1062-6050-51.12.01
- Quatman-Yates CC, Hunter-Giordano A, Shimamura KK, et al. Physical therapy evaluation and treatment after concussion/mild traumatic brain injury. *J Orthop Sports Phys Ther.* 2020;50(4):CPG1-CPG73. doi:10.2519/jospt.2020.0301
- Leddy JJ, Wilber CG, Willer BS. Active recovery from concussion. *Curr Opin Neurol.* 2018;31(6):681-686. doi:10.1097/wco.0000000000000611
- Prisma. PRISMA. Accessed July 28, 2021. <http://prisma-statement.org/>
- Register a review. PROSPERO. Accessed July 28, 2021. <https://www.crd.york.ac.uk/PROSPERO/>
- Jewell DV. *Guide to Evidence-Based Physical Therapist Practice.* Jones & Bartlett Learning; 2018.

14. Pedro scale. PEDro. Published September 9, 2020. Accessed August 29, 2021. <https://pedro.org.au/english/resources/pedro-scale/>
15. Balance error scoring system. Shirley Ryan AbilityLab. Accessed July 1, 2021. <https://www.srala.b.org/rehabilitation-measures/balance-error-scoring-system>
16. Beck depression inventory. Shirley Ryan AbilityLab. Accessed July 1, 2021. <https://www.srala.b.org/rehabilitation-measures/beck-depression-inventory>
17. Graham R, Rivara FP, Ford MA, Committee on Sports-Related Concussions in Youth; Board on Children, Youth, and Families; Institute of Medicine. *Sports-Related Concussions in Youth: Improving the Science, Changing the Culture*. National Academies Press (US); 2014.
18. Borg rating of perceived exertion. Physiopedia. Accessed July 15, 2021. https://www.physio-pedia.com/Borg_Rating_Of_Perceived_Exertion
19. Maerlender A, Rieman W, Lichtenstein J, Condiracci C. Programmed physical exertion in recovery from sports-related concussion: a randomized pilot study. *Dev Neuropsychol*. 2015;40(5):273-278. doi:10.1080/87565641.2015.1067706
20. Leddy JJ, Haider MN, Ellis MJ, et al. Early subthreshold aerobic exercise for sport-related concussion: a randomized clinical trial. *JAMA Pediatr*. 2019;173(4):319-325. doi:10.1001/jamapediatrics.2018.4397
21. Leddy JJ, Master CL, Mannix R, et al. Early targeted heart rate aerobic exercise versus placebo stretching for sport-related concussion in adolescents: a randomised controlled trial. *Lancet Child Adolesc Health*. 2021;5(11):792-799. doi:10.1016/s2352-4642(21)00267-4
22. Chan C, Iverson GL, Purtzki J, et al. Safety of active rehabilitation for persistent symptoms after pediatric sport-related concussion: a randomized controlled trial. *Arch Phys Med Rehabil*. 2018;99(2):242-249. doi:10.1016/j.apmr.2017.09.108
23. Reneker JC, Hassen A, Phillips RS, Moughiman MC, Donaldson M, Moughiman J. Feasibility of early physical therapy for dizziness after a sports-related concussion: a randomized clinical trial. *Scand J Med Sci Sports*. 2017;27(12):2009-2018. doi:10.1111/sms.12827
24. Schneider KJ, Meeuwisse WH, Nettel-Aguirre A, et al. Cervicovestibular rehabilitation in sport-related concussion: a randomised controlled trial. *Br J Sports Med*. 2014;48(17):1294-1298. doi:10.1136/bjsports-2013-093267
25. Dobney DM, Grilli L, Beaulieu C, et al. Feasibility of early active rehabilitation for concussion recovery in youth: a randomized trial. *Clin J Sport Med*. 2020;30(6):519-525. doi:10.1097/jsm.0000000000000671
26. Micay R, Richards D, Hutchison MG. Feasibility of a postacute structured aerobic exercise intervention following sport concussion in symptomatic adolescents: a randomised controlled study. *BMJ Open Sport Exerc Med*. 2018;4(1):e000404. doi:10.1136/bmjsem-2018-000404
27. Lal A, Kolakowsky-Hayner SA, Ghajar J, Balamane M. The effect of physical exercise after a concussion: a systematic review and meta-analysis. *Am J Sports Med*. 2018;46(3):743-752. doi:10.1177/0363546517706137
28. Makdissi M, Schneider KJ, Feddermann-Demont N, et al. Approach to investigation and treatment of persistent symptoms following sport-related concussion: a systematic review. *Br J Sports Med*. 2017;51(12):958-968. doi:10.1136/bjsports-2016-097470

SUPPLEMENTARY MATERIALS

Appendix

Download: <https://ijspt.scholasticahq.com/article/68071-the-effectiveness-of-physical-therapy-interventions-for-athletes-post-concussion-a-systematic-review/attachment/135447.docx>
