



# The importance of perioperative team familiarity and its contribution to surgical efficiency and outcomes in microsurgical breast reconstruction

Jevan Cevik<sup>1,2</sup>, David J. Hunter-Smith<sup>1,2</sup>, Warren M. Rozen<sup>1,2</sup><sup>^</sup>

<sup>1</sup>Department Surgery, Peninsula Clinical School, Central Clinical School, Faculty of Medicine, Monash University, Frankston, VIC, Australia;

<sup>2</sup>Monash University Plastic and Reconstructive Surgery Group (Peninsula Clinical School), Peninsula Health, Frankston, VIC, Australia

*Correspondence to:* Professor Warren M. Rozen, MBBS, BMedSc, MD, PhD, FRACS. Department Surgery, Peninsula Clinical School, Central Clinical School, Faculty of Medicine, Monash University, 2 Hastings Road, Frankston, VIC 3199, Australia; Monash University Plastic and Reconstructive Surgery Group (Peninsula Clinical School), Peninsula Health, 2 Hastings Road, Frankston, VIC 3199, Australia. Email: warrenrozen@hotmail.com.

*Comment on:* Speck NE, Dreier K, Fluetsch A, *et al.* Comparing complications and perioperative teams in microsurgical breast reconstruction: retrospective cohort study. *Gland Surg* 2022;11:1754-63.

**Keywords:** Microsurgery; breast reconstruction; anatomy

Submitted Nov 15, 2022. Accepted for publication Dec 16, 2022. Published online Dec 26, 2022.

doi: 10.21037/gs-22-672

**View this article at:** <https://dx.doi.org/10.21037/gs-22-672>

A surgical operation involves the effective interaction and collaboration of many different specialized personnel. A typical operating room may contain nurses, technicians, anaesthetists, surgeons, and surgical assistants. These individuals together form the surgical team that must function cohesively to ensure not only optimal efficiency of the surgical procedure but also to achieve ideal outcomes for patients and avoid complications. This is particularly important in microsurgical breast reconstruction. Given the often-greater complexity and specialization of procedures performed during microsurgical breast reconstruction, greater familiarity is required from all members of the surgical team to achieve optimal results.

In the modern day, the demand of medicine is increasing. With an aging and increasing population, the operative process must progress and become more efficient whilst, simultaneously, complications must be minimised. Surgeons have thus been required to refine their surgical technique and develop adjuvant technological adjuncts to assist them. Additionally, in recent times, surgeons have broadened their attention to all aspects of the operation including optimisation of the perioperative team.

It is a common experience among surgeons that a greater familiarity and expertise of each perioperative team member facilitates a smoother and less complicated operation. In support of this, it has been previously shown that surgical teams whose members have a greater familiarity with each other display shorter operating times and fewer surgical complications (1-7). Hence, optimisation of the operative environment by increasing familiarity amongst team members is warranted given the potential improvement in patient outcomes and cost-benefits to the public healthcare system.

On this note, a recent study by Speck *et al.* outlines a novel study designed to assess the impact that a perioperative team may have on operative outcomes in microsurgical breast reconstruction (8). A retrospective review of cases conducted at two different institutions was performed. One institution was a smaller, private clinic, the other was a larger conventional hospital. These two institutions differed most significantly in the size of the pool of perioperative staff that could be involved in each procedure, the conventional hospital having a pool of staff almost four times larger than the small clinic.

<sup>^</sup> ORCID: 0000-0002-4092-182X.

The author's found that there were lower rates of flap loss and flap revision among the reconstructive procedures performed at the smaller clinic when compared to the larger hospital. The author's hypothesized that this outcome may be because the pool of staff at the smaller clinic was smaller, meaning each individual member of the surgical team would work together more often and be exposed to the surgical procedures more frequently increasing the familiarity of each individual and thereby improving the performance of the team as a whole.

This study offers novel quantitative evidence to support the importance of experienced perioperative teams in microsurgical breast reconstruction (8). Yet, while this study assessed the variation in outcomes among various perioperative teams, a potentially interesting finding that was unfortunately omitted from the study by Speck *et al.* is the impact that the perioperative team may have on the operative duration in microsurgical breast reconstruction. However, it was discussed in the article that greater familiarity and experience amongst surgical team members would be likely to increase efficiency and reduce operation durations. Given the known association between operation duration and complication rates in breast reconstructive surgery, improved operative efficiency would also likely play a role in improving surgical outcomes (9).

On a broader level, the study written by Speck *et al.* contributes to the growing development of surgical efficiency and optimisation. Along with improvements in understanding of the importance of cohesive perioperative teamwork, including the use of two microsurgeons in microsurgical cases, throughout history, surgeons have made significant advancements in numerous other areas of the operative process to improve surgical efficiency and outcomes in microsurgical breast reconstruction (10). Many improvements in efficiency have stemmed from implementation of a technique known as process mapping.

Process mapping is a technique that involves deconstructing a process and identifying its key individual components. When considering a microsurgical breast reconstruction procedure, improving the rate of an operation involves more than simply operating faster but rather it involves elimination of wasted effort and time (11-13). In the context of breast reconstruction, the process includes the entirety of the pre-operative, intra-operative and post-operative events (11-13). Once the key components in each of these events are identified, one can then implement strategies to optimise each step such that the entire process becomes more efficient. Through the

use of process mapping, the rate of microsurgical breast reconstruction has increased drastically, such that now up to three microsurgical operations can be performed in a day in a single theatre (14,15).

Despite the importance of optimising perioperative teams and process mapping to identify areas of inefficiency, it is undeniable that technological advancements have similarly had a major impact on the improved efficiency seen in microsurgical breast reconstruction. Two technological advancements that have undoubtedly had such an impact are the use of computed tomography angiographic (CTA) imaging pre-operatively and the use of venous coupler devices intraoperatively.

The use of CTA pre-operatively, particularly for deep inferior epigastric perforators (DIEP) flaps, has had marked implications for breast reconstructive surgery. Pre-operative imaging with CTA allows surgeons to survey the vascular anatomy, identify perforators and plan flap design prior to the operation. Given more effective pre-operative planning, operative efficiency has improved, and operation durations decreased (16-20). More specifically, perforator identification and selection time intraoperatively is significantly shorter if CTA is used pre-operatively to assess the vasculature (21).

During microsurgical breast reconstruction, effective and careful anastomosis of the vessels of the flap and the recipient site is crucial for flap success. Inadequate anastomosis of these vessels can predispose to thrombosis and subsequent flap failure. Traditionally, anastomosis is performed using very fine 8-0 or 9-0 sutures to meticulously connect the two ends of the vessels together. Yet, after introduction of intraoperative venous anastomotic coupler devices, the speed of anastomosis of veins has significantly increased while maintaining adequate patency and perfusion (22-25).

Ultimately, the process of microsurgical breast reconstruction has developed significantly since its conception. Throughout its development, surgeons have continually refined the art of breast reconstruction to reduce unnecessary steps, optimise efficiency and avoid complications. On one hand, these advancements have stemmed from extraordinary progress in technology and surgical expertise, yet, additionally, surgeons have recently realised the importance of an effective perioperative team and the role this plays in operative success. As a consequence of these advances, countless more patients have received treatment than they would have otherwise. It is without doubt that as surgeons strive for excellence in the operating room, the art of microsurgical breast reconstruction will continue to evolve and improve the outcomes experienced

by patients.

## Acknowledgments

*Funding:* None.

## Footnote

*Provenance and Peer Review:* This article was commissioned by the editorial office, *Gland Surgery*. The article did not undergo external peer review.

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <https://gs.amegroups.com/article/view/10.21037/gS-22-672/coif>). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Kurmann A, Keller S, Tschan-Semmer F, et al. Impact of team familiarity in the operating room on surgical complications. *World J Surg* 2014;38:3047-52.
2. Xu R, Carty MJ, Orgill DP, et al. The teaming curve: a longitudinal study of the influence of surgical team familiarity on operative time. *Ann Surg* 2013;258:953-7.
3. Mazzocco K, Petitti DB, Fong KT, et al. Surgical team behaviors and patient outcomes. *Am J Surg* 2009;197:678-85.
4. Witmer HDD, Dhiman A, Jones AD, et al. A Systematic Review of Operative Team Familiarity on Metrics of Efficiency, Patient Outcomes, Cost, and Team Satisfaction. *Ann Surg* 2022;276:e674-81.
5. Henaux PL, Michinov E, Rochat J, et al. Relationships Between Expertise, Crew Familiarity and Surgical Workflow Disruptions: An Observational Study. *World J Surg* 2019;43:431-8.
6. Catchpole K, Mishra A, Handa A, et al. Teamwork and error in the operating room: analysis of skills and roles. *Ann Surg* 2008;247:699-706.
7. Gillespie BM, Chaboyer W, Fairweather N. Factors that influence the expected length of operation: results of a prospective study. *BMJ Qual Saf* 2012;21:3-12.
8. Speck NE, Dreier K, Fluetsch A, et al. Comparing complications and perioperative teams in microsurgical breast reconstruction: retrospective cohort study. *Gland Surg* 2022;11:1754-63.
9. Allan J, Goltsman D, Moradi P, et al. The effect of operative time on complication profile and length of hospital stay in autologous and implant-based breast reconstruction patients: An analysis of the 2007-2012 ACS-NSQIP database. *J Plast Reconstr Aesthet Surg* 2020;73:1292-8.
10. Cevik J, Hunter-Smith DJ, Rozen WM. Current Advances in Breast Reconstruction. *J Clin Med* 2022;11:3328.
11. Chung RD, Hunter-Smith DJ, Spychal RT, et al. A systematic review of intraoperative process mapping in surgery. *Gland Surg* 2017;6:715-25.
12. Sharma HR, Rozen WM, Mathur B, et al. 100 Steps of a DIEP Flap-A Prospective Comparative Cohort Series Demonstrating the Successful Implementation of Process Mapping in Microsurgery. *Plast Reconstr Surg Glob Open* 2019;7:e2016.
13. Haddock NT, Teotia SS. Deconstructing the Reconstruction: Evaluation of Process and Efficiency in Deep Inferior Epigastric Perforator Flaps. *Plast Reconstr Surg* 2020;145:717e-24e.
14. Marsh D, Patel NG, Rozen WM, et al. Three routine free flaps per day in a single operating theatre: principles of a process mapping approach to improving surgical efficiency. *Gland Surg* 2016;5:107-14.
15. Acosta R, Enajat M, Rozen WM, et al. Performing two DIEP flaps in a working day: an achievable and reproducible practice. *J Plast Reconstr Aesthet Surg* 2010;63:648-54.
16. Colakoglu S, Tebockhorst S, Freedman J, et al. CT angiography prior to DIEP flap breast reconstruction: a randomized controlled trial. *J Plast Reconstr Aesthet Surg* 2022;75:45-51.
17. Malhotra A, Chhaya N, Nsiah-Sarbeng P, et al. CT-guided deep inferior epigastric perforator (DIEP) flap loc-ization -- better for the patient, the surgeon, and the hospital.

- Clin Radiol 2013;68:131-8.
18. Rozen WM, Ashton MW, Grinsell D, et al. Establishing the case for CT angiography in the preoperative imaging of abdominal wall perforators. *Microsurgery* 2008;28:306-13.
  19. Fitzgerald O'Connor E, Rozen WM, Chowdhry M, et al. Preoperative computed tomography angiography for planning DIEP flap breast reconstruction reduces operative time and overall complications. *Gland Surg* 2016;5:93-8.
  20. Ohkuma R, Mohan R, Baltodano PA, et al. Abdominally based free flap planning in breast reconstruction with computed tomographic angiography: systematic review and meta-analysis. *Plast Reconstr Surg* 2014;133:483-94.
  21. Haddock NT, Dumestre DO, Teotia SS. Efficiency in DIEP Flap Breast Reconstruction: The Real Benefit of Computed Tomographic Angiography Imaging. *Plast Reconstr Surg* 2020;146:719-23.
  22. Pratt GF, Rozen WM, Westwood A, et al. Technology-assisted and sutureless microvascular anastomoses: evidence for current techniques. *Microsurgery* 2012;32:68-76.
  23. Fitzgerald O'Connor E, Rozen WM, Chowdhry M, et al. The microvascular anastomotic coupler for venous anastomoses in free flap breast reconstruction improves outcomes. *Gland Surg* 2016;5:88-92.
  24. Rozen WM, Whitaker IS, Acosta R. Venous coupler for free-flap anastomosis: outcomes of 1,000 cases. *Anticancer Res* 2010;30:1293-4.
  25. Jandali S, Wu LC, Vega SJ, et al. 1000 consecutive venous anastomoses using the microvascular anastomotic coupler in breast reconstruction. *Plast Reconstr Surg* 2010;125:792-8.

**Cite this article as:** Cevik J, Hunter-Smith DJ, Rozen WM. The importance of perioperative team familiarity and its contribution to surgical efficiency and outcomes in microsurgical breast reconstruction. *Gland Surg* 2023;12(1):1-4. doi: 10.21037/gs-22-672