




# Development and applications of a CRISPR activation system for facile genetic overexpression in *Candida albicans*

Nicholas C. Gervais <sup>1</sup>, Alyssa A. La Bella,<sup>2</sup> Lauren F. Wensing,<sup>1</sup> Jehoshua Sharma,<sup>1</sup> Victoria Acquaviva,<sup>1</sup> Madison Best,<sup>1</sup> Ricardo Omar Cadena López,<sup>1</sup> Meea Fogal <sup>1</sup>, Deeva Uthayakumar,<sup>1,3</sup> Alejandro Chavez,<sup>4</sup> Felipe Santiago-Tirado,<sup>2</sup> Ana L. Flores-Mireles,<sup>2</sup> Rebecca S. Shapiro <sup>1,\*</sup>

<sup>1</sup>Department of Molecular and Cellular Biology, University of Guelph, Guelph, ON N1H 5N4, Canada

<sup>2</sup>Department of Biological Sciences, University of Notre Dame, Notre Dame, IN 46556, USA

<sup>3</sup>Present address: Department of Immunology, University of Toronto, Toronto, ON, Canada

<sup>4</sup>Department of Pathology and Cell Biology, Columbia University College of Physicians and Surgeons, New York, NY 10032, USA

\*Corresponding author. Department of Molecular and Cellular Biology, University of Guelph, 50 Stone Road East, Guelph, ON, Canada, N1H 5N4. Email: shapiror@uoguelph.ca

## Abstract

For the fungal pathogen *Candida albicans*, genetic overexpression readily occurs via a diversity of genomic alterations, such as aneuploidy and gain-of-function mutations, with important consequences for host adaptation, virulence, and evolution of antifungal drug resistance. Given the important role of overexpression on *C. albicans* biology, it is critical to develop and harness tools that enable the analysis of genes expressed at high levels in the fungal cell. Here, we describe the development, optimization, and application of a novel, single-plasmid-based CRISPR activation (CRISPRa) platform for targeted genetic overexpression in *C. albicans*, which employs a guide RNA to target an activator complex to the promoter region of a gene of interest, thus driving transcriptional expression of that gene. Using this system, we demonstrate the ability of CRISPRa to drive high levels of gene expression in *C. albicans*, and we assess optimal guide RNA targeting for robust and constitutive overexpression. We further demonstrate the specificity of the system via RNA sequencing. We highlight the application of CRISPR activation to overexpress genes involved in pathogenesis and drug susceptibility, and contribute toward the identification of novel phenotypes. Consequently, this tool will facilitate a broad range of applications for the study of *C. albicans* genetic overexpression.

**Keywords:** fungal genetics, CRISPR activation, *Candida albicans*, gene regulation, fungal pathogens

## Introduction

Fungal pathogens are an increasingly important cause of human illness, with significant impacts on morbidity and mortality (Fisher et al. 2020). *Candida albicans* is an opportunistic fungal pathogen that inhabits the microbiome of most healthy adults as a commensal organism (Gow et al. 2011; Pérez 2019). However, it is also responsible for invasive infections, particularly among immunocompromised individuals, which are often fatal (Bongomin et al. 2017). Instances of *C. albicans* infections are becoming more prevalent globally, in part due to an increasing population of susceptible elderly and immunocompromised individuals (Bongomin et al. 2017). Illustratively, systemic *Candida* infections have been identified in up to 15% of patients hospitalized for COVID-19, where patients with invasive candidiasis spent longer in the hospital intensive care unit on average and had more severe symptoms than those COVID-19 patients without a *Candida* infection (Segrelles-Calvo et al. 2021). The economic cost of *Candida* infections is estimated to be ~1.4 billion dollars yearly in the United States of America alone, hospitalizing >26,000

people (Benedict et al. 2018). With limited development of novel antifungal drugs (Perfect 2017) and increasing prevalence of antifungal drug-resistant strains and species (Sanguinetti et al. 2015; Arendrup and Patterson 2017; Geddes-McAlister and Shapiro 2018), *Candida* pathogens remain a serious threat to human health.

Given the importance of *C. albicans* and other fungal pathogens, it is imperative that tools are developed and applied to manipulate these organisms and dissect genetic functions. For *C. albicans*, along with countless other microbial species, CRISPR-based genetic manipulation tools have revolutionized the ability of scientists to modify and study gene function (Shapiro, Chavez, Collins 2018; Uthayakumar et al. 2021). Such CRISPR platforms typically rely on a CRISPR-associated endonuclease protein, often Cas9, targeted to a genomic region of interest via a single-guide RNA (sgRNA), resulting in targeted DNA double-strand breaks, and subsequent repair via nonhomologous end-joining or homology-directed repair via a donor DNA repair template (Adli 2018; Bock et al. 2022). Since the initial development of CRISPR tools for

Received: September 12, 2022. Accepted: November 04, 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of the Genetics Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

genetic editing in *C. albicans* (Vyas et al. 2015), numerous systems have been developed to modify this fungal genome with efficiency and specificity (Min et al. 2016, 2018; Huang and Mitchell 2017; Ng and Dean 2017; Nguyen et al. 2017; Shapiro, Chavez, Porter, et al. 2018; Vyas et al. 2018; Román, Prieto, et al. 2019; Morio et al. 2020; Uthayakumar et al. 2021), and to build and screen genetic mutant libraries (Shapiro, Chavez, Porter, et al. 2018; Rosiana et al. 2021). Together, these gene editing systems have played an important role in providing novel biological insight into this fungal pathogen.

In addition to these canonical CRISPR-based genetic modification systems that introduce mutations, insertions, or deletions into the genome, alternative CRISPR technologies have been developed that modulate the expression of *C. albicans* genes (Román, Coman, et al. 2019; Wensing et al. 2019). These CRISPR systems rely on the inactivation of the Cas9 endonuclease, to generate a nuclease-dead (dCas9) protein, which can be fused to constructs from a transcriptional repressor or activator that generate CRISPR interference (CRISPRi) or CRISPR activation (CRISPRa) systems, respectively (Dominguez et al. 2016; Wang et al. 2016). These dCas9 fusions, when targeted to the promoter region of a gene via the sgRNA, can repress or activate gene transcription from a gene of interest, and have been widely applied in diverse organisms (Peters et al. 2015; Kampmann 2018; Xu and Qi 2019; Schultenkämper et al. 2020; Todor et al. 2021; Bock et al. 2022). In *C. albicans*, these modified CRISPR systems have enabled the modulation and study of important fungal genes, including essential genes (Román, Coman, et al. 2019; Wensing et al. 2019; Wensing and Shapiro 2022).

While a majority of CRISPR systems and other genetic manipulation platforms focus on gene mutation, deletion, or downregulation, it is also critical to develop techniques that enable the upregulation or overexpression of genes. Indeed, genetic overexpression in *C. albicans* occurs readily via gene copy number amplification and aneuploidy, and these copy number amplification events are frequently observed in genetically diverse clinical isolates (Coste et al. 2006; Forche et al. 2018; Todd et al. 2019). The upregulation of gene expression plays a critical role in diverse aspects of fungal biology (Rai et al. 2021), including the evolution of antifungal drug resistance (Selmecki et al. 2008; Sanglard et al. 2009; Schubert et al. 2011; Flowers et al. 2012; Lohberger et al. 2014; Todd and Selmecki 2020), host colonization (Prieto et al. 2017; Znaidi et al. 2018), and virulence (Fu et al. 2008; Lohberger et al. 2014). Additionally, in other microbial systems, including bacteria, protozoa, and the model yeast *Saccharomyces cerevisiae*, genetic overexpression libraries have proved to be critical tools for identifying targets of drugs with uncharacterized mechanisms of action, as mutants overexpressing the drug target are typically more resistant to the drug in question (Li et al. 2004; Luesch et al. 2005; Smith et al. 2010; Begolo et al. 2014). Overexpression screens can also help dissect molecular pathways (Sopko et al. 2006; Sahni et al. 2010; Polvi et al. 2019). In *C. albicans*, several techniques exist for genetic overexpression (Rai et al. 2021), including promoter replacement strategies (Delgado et al. 2003; Park and Morschhäuser 2005; Eckert and Mühlischlegel 2009; Sahni et al. 2010), the generation of ORFeome collection strains (Chauvel et al. 2012; Legrand et al. 2018), and CRISPR-based methods (Román, Coman, et al. 2019). While these tools have made important contributions to understanding overexpression in *C. albicans*, there remains a need for simple and rapid methods to efficiently target fungal genes of interest for overexpression, and to do so in diverse strain backgrounds.

Here, we introduce a CRISPR-based tool for genetic overexpression in *C. albicans*, to bolster the existing functional genetic toolbox

available to manipulate genes in this critical fungal pathogen. Our CRISPRa tool exploits a *C. albicans*-optimized tripartite activator complex fused to dCas9, which was previously developed for highly efficient transcriptional regulation in *S. cerevisiae* (Chavez et al. 2015). This CRISPRa system is distinct from other CRISPR-based gene activation systems for *C. albicans*, as it is an efficient single-plasmid system designed for rapid Golden Gate cloning and facile fungal strain generation, which can be readily applied to diverse strain backgrounds including clinical isolates. We test CRISPRa guide targeting principles based on predicted transcriptional start sites, confirm on-target guide efficiency, and demonstrate the ability of this CRISPRa system to drive high levels of expression of *C. albicans* genes. Using this optimized CRISPRa technique, we validate its ability to drive the overexpression of genes with established roles in antifungal drug susceptibility and biofilm formation. Together, this work introduces a novel CRISPRa system for genetic overexpression in *C. albicans*, with a wide range of future applications.

## Materials and methods

### Plasmid design and cloning

The plasmid backbone used in this study was the *C. albicans*-optimized CRISPR-dCas9 plasmid (pRS143, Addgene #122377) used in our previous study (Wensing et al. 2019), containing the *NEUT5L* integration site, sgRNA cloning site (SNR52 promoter, SapI cloning locus, and sgRNA tail), and dCas9. The dCas9-VP64-p65-Rta (VPR) fusion construct was generated via Gibson assembly. The VPR tripartite complex was codon-optimized for *C. albicans* expression and synthesized as gBlocks gene fragments from Integrated DNA Technologies (IDT). This gene fragment was cloned with Gibson assembly into the dCas9 plasmid backbone. We have made the CRISPRa (dCas9-VPR) plasmid available via Addgene (reference #182707). All plasmids are listed in [Supplementary Table 1](#).

### sgRNA design and cloning

CRISPRa strain construction was broadly performed as described in detail (Wensing and Shapiro 2022), with some minor modifications. The sgRNA CRISPR RNAs (crRNA; 20 nucleotide sequence complementary to the target DNA genomic DNA) were designed based on efficiency and predicted specificity via the sgRNA design tool Eukaryotic Pathogen CRISPR gRNA Design Tool (EuPaGDT; <http://grna.ctegd.uga.edu>; Peng and Tarleton 2015). Generally, designing sgRNAs around -100 to -350 bp upstream of both the start codon (ATG) and transcriptional start site of the target gene led to most successful overexpression. In all cases, employing 4-6 sgRNAs in total (2-3 sgRNAs per site) led to significant overexpression in at least one of the corresponding strains. These sgRNA crRNA N20 sequences were cloned into the dCas9-VPR CRISPRa plasmid at the sgRNA cloning locus (SapI cloning site, flanked by SNR52 promoter, and sgRNA tail) using Golden Gate cloning (Engler et al. 2008), as previously described (Halder et al. 2019; Wensing and Shapiro 2022). crRNA N20 sequences were obtained as two oligonucleotides from IDT in forward and reverse complement orientation, each containing a SapI cloning site, and were reconstituted to 100  $\mu$ M in a nuclease-free duplex buffer from IDT. Equal volumes of the two complementary oligonucleotides were combined after being heated separately at 94°C, and then duplexed by heating to 94°C for 2 min and cooling to room temperature. The duplexed fragment was then cloned into the CRISPRa plasmid with the following Golden Gate cloning reaction: 10  $\mu$ l miniprep CRISPRa plasmid, 1  $\mu$ l duplexed oligonucleotide,

2  $\mu$ l 10X CutSmart buffer, 2  $\mu$ l ATP, 1  $\mu$ l SapI, 1  $\mu$ l T4 DNA ligase, and 3  $\mu$ l nuclease-free water. This mixture was incubated in a thermocycler under the following cycling conditions: (37°C, 2 min; 16°C, 5 min) for 99 cycles; 65°C, 15 min; 80°C, 15 min. After cycling, 1  $\mu$ l of additional SapI was added to each reaction mixture, and the reaction was incubated at 37°C for 1 h. All sgRNAs are listed in [Supplementary Table 1](#).

## Media and growth conditions

*Escherichia coli* DH5 $\alpha$  cells were grown at 37°C in Lysogeny Broth (LB) and LB plates supplemented with 100 mg/ml ampicillin (AMP) and 250 mg/ml nourseothricin (NAT) for plasmid selection. *Candida albicans* cells were grown at 30°C or 37°C in yeast peptone dextrose (YPD) broth and YPD plates supplemented with 250 mg/ml NAT for plasmid selection.

## Bacterial transformation

High-efficiency 5-alpha competent *E. coli* (NEB) cells were thawed on ice. 1  $\mu$ l of Golden Gate cloning reagents was added to 50  $\mu$ l of competent cells and incubated on ice for 30 min, heat shocked for 30 s at 42°C, and incubated on ice for an additional 5 min. About 950  $\mu$ l of SOC outgrowth media was added to each cell culture and incubated at 30°C for 1.5 h at 250 RPM. Transformed cells were then plated on LB media containing AMP and NAT and grown at 30°C for 1 day.

## *Candida albicans* transformation

Plasmids were transformed into *C. albicans* via a chemical transformation strategy, as previously described ([Halder et al. 2019](#); [Wensing and Shapiro 2022](#)). Briefly, miniprep CRISPRa plasmids were linearized via a 1.5X restriction digest mix using the PacI enzyme. A transformation master mix was prepared as follows: 800  $\mu$ l of 50% polyethylene glycol, 100  $\mu$ l of 10X Tris-EDTA buffer solution, 100  $\mu$ l of 1 M lithium acetate, 40  $\mu$ l of 10 mg/ml salmon sperm DNA, and 20  $\mu$ l of 1 M dithiothreitol. The transformation mix was added to *C. albicans* cells and linearized CRISPRa plasmid and left to incubate at 30°C for 1 h, then heat shocked at 42°C for 45 min. Cells were washed with fresh YPD, then grown in YPD for 4 h to allow for expression of the NAT resistance construct. Transformed cells were then plated on YPD media containing NAT, and grown at 30°C for 2 days. All *C. albicans* strains are listed in [Supplementary Table 1](#).

## Growth curves

Overnight cultures of *C. albicans* grown in YPD were diluted to an OD<sub>600</sub> of 0.05 in fresh YPD and added to a 96-well flat-bottomed plate with a total volume of 200  $\mu$ l per well. Growth was measured at 30°C via optical density at 600 nm at 15 min intervals over the course of 24 h using an Infinite 200 PRO microplate reader (Tecan), and plates were shaken orbitally for 900 s at a 4-mm amplitude in between growth measurements.

## Minimum inhibitory concentration assays

Fluconazole minimum inhibitory concentration (MIC) assays were performed in 96-well flat-bottomed plates. A suspension of 40  $\mu$ g/ml fluconazole was prepared in water, of which 100  $\mu$ l was added to the first column of each plate containing 100  $\mu$ l of culture, to obtain a total volume of 200  $\mu$ l per well. The first column was serially diluted 2-fold across the plate in water. The gradient of fluconazole, therefore, ranged from 20 to 0  $\mu$ g/ml. Overnight cultures of *C. albicans* grown in YPD were diluted to an OD<sub>600</sub> of 0.1 in 2X RPMI-1640 with 40 g/l of added D-glucose and mixed into the plates in an equal volume such that the starting OD<sub>600</sub> values of

each strain were 0.05 and the growth media was diluted to RPMI-1640 with 20 g/l of added D-glucose. Plates were incubated at 37°C at 900 RPM, and absorbance values at 600 nm were read after 24 h using an Infinite 200 PRO microplate reader (Tecan). The amphotericin B MIC assays were also performed in 96-well flat-bottomed plates similar to the protocol described above with a few modifications. A starting concentration of 5  $\mu$ g/ml of amphotericin B was used to create a gradient ranging from 2.5 to 0  $\mu$ g/ml. Strains were instead grown in YPD media using a starting OD<sub>600</sub> of 0.001. Plates were incubated at 37°C statically and absorbance values at 600 nm were read after 72 h using an Infinite 200 PRO microplate reader (Tecan).

## Biofilm assays

RPMI-based biofilm assays were performed as previously described ([Razzaq et al. 2021](#)), with minor modifications. Overnight cultures of *C. albicans* grown in YPD were diluted to an OD<sub>600</sub> of 0.001 in 5 ml of RPMI-1640 with 20 g/l of supplemented D-glucose. 100  $\mu$ l of RPMI-1640 was mixed with 100  $\mu$ l of each diluted overnight culture into a 96-well flat-bottomed plate (12 wells per strain). Plates were grown statically at 37°C for 24 h to allow biofilms to form. About 120  $\mu$ l of planktonic cell supernatant was removed from each well and moved to a new 96-well flat-bottomed plate. Density of the planktonic cells was measured at 600 nm using an Infinite 200 PRO microplate reader (Tecan). Original plates from which planktonic cells were removed were washed by adding 200  $\mu$ l 1X PBS to each well with an electronic multichannel pipette at a moderate dispense speed. 1X PBS was then discarded, and this wash step was repeated. Plates were then left to dry for 1–2 h. Once biofilms were dry, 90  $\mu$ l of 1 mg/ml tetrazolium salt (XTT) (prepared in 1X PBS and centrifuged to remove sediment prior to use) and 10  $\mu$ l of 0.32 mg/ml PMS (prepared in water) were added to each well. Plates were incubated statically at 30°C for 2 h to allow biofilms to reduce XTT, measured at 490 nm using an Infinite 200 PRO microplate reader (Tecan), and normalized to the growth of planktonic cells harvested previously.

For urine-based biofilm assays, biofilm formations were performed in 96-well flat-bottomed plates. Plates were coated with 150  $\mu$ g/ml of fibrinogen and incubated overnight at 4°C. The *C. albicans* strains were cultured at 37°C with aeration in 5 ml of YPD broth. The inoculum was normalized to  $\sim 1 \times 10^6$  CFU/ml and then diluted (1:10) in human urine (female, pH  $\sim 6.5$ ). Urine was supplemented with 20 mg/ml of bovine serum albumin (BSA) for carbon and nitrogen sources mimicking the catheterized bladder environment. 200  $\mu$ l of the urine with inoculum was incubated in each well of the 96-well plate at 37°C for 48 h while static. Following the 48 h incubation, the cultures were removed and the wells were washed 3 $\times$  with 200  $\mu$ l 1X PBS to remove unbound fungi. Plates were incubated with 200  $\mu$ l of 0.5% crystal violet for 15 min. Crystal violet was removed, and plates were washed with water and dried. 200  $\mu$ l of 33% acetic acid was added to the wells and 100  $\mu$ l was transferred to a new 96-well plate. Absorbance was measured at 595 nm using a SpectraMax ABS Plus microplate reader (Molecular Devices).

## RNA extraction and real-time quantitative PCR

To detect differences in gene expression, overnight cultures of *C. albicans* grown in YPD were diluted to an OD<sub>600</sub> of 0.05 in fresh YPD and grown to an OD<sub>600</sub> of  $>0.2$  at 30°C. ALS3 CRISPRa strains were diluted to an OD<sub>600</sub> of 0.05 in fresh RPMI and grown for 5 h at 37°C as well to measure overexpression in conditions that favor

high levels of basal expression. Cultures were pelleted and frozen at  $-80^{\circ}\text{C}$  before RNA was extracted. RNA extractions were performed using either a Presto Mini RNA Yeast kit from FroggaBio (cat. RBYD050) or an RNeasy Mini kit from Qiagen (cat. 74104). To assess RNA integrity, an RNA ScreenTape assay was performed on all samples using the TapeStation 4150 system following manufacturer's instructions (Agilent Technologies). Only samples with RIN values of  $>5.0$  were used for reverse-transcription quantitative PCR (RT-qPCR). Synthesis of cDNA from 1,000 ng (10  $\mu\text{l}$ ) RNA was performed using a High Capacity cDNA Reverse-Transcription kit from Applied Biosystems (cat. 4368814). Briefly, 20  $\mu\text{l}$  reaction mixtures were prepared with 2  $\mu\text{l}$  10X reverse-transcription buffer, 0.8  $\mu\text{l}$  of 25X dNTPs at a concentration of 100 mM, 2  $\mu\text{l}$  of random primers, 1  $\mu\text{l}$  of MultiScribe Reverse Transcriptase at 50 IU/ $\mu\text{l}$ , 4.2  $\mu\text{l}$  of nuclease-free water, and 10  $\mu\text{l}$  of RNA (1  $\mu\text{g}$ ). Reverse-transcription run conditions were as follows: 10 min at  $25^{\circ}\text{C}$ , 120 min at  $37^{\circ}\text{C}$ , 5 min at  $85^{\circ}\text{C}$ . Real-time PCR assays were conducted using a QuantStudio 7 Pro Real-Time PCR system from Thermo Fisher Scientific Inc. 20  $\mu\text{l}$  reaction mixtures containing 10  $\mu\text{l}$  2X SsoAdvanced Universal Inhibitor-Tolerant SYBR supermix (Bio-Rad, cat: 172-5017), 0.8  $\mu\text{l}$  of PCR forward and reverse primer mix at 5  $\mu\text{M}$  (final concentration of primer at 200 nM), 4.2  $\mu\text{l}$  of water, and 5  $\mu\text{l}$  of diluted cDNA. The run conditions were as follows: 3 min at  $98^{\circ}\text{C}$  polymerase activation step, followed by 40 cycles of a two-step qPCR (10 s of  $98^{\circ}\text{C}$  denaturation, 30 s of  $60^{\circ}\text{C}$  combined annealing/extension). Primers are listed in [Supplementary Table 1](#). Expression profiling calculations were performed according to the comparative  $C_T$  method ([Schmittgen and Livak 2008](#)). Briefly, expression values for the experimental gene of interest were compared with the housekeeping gene ACT1 to obtain a  $\Delta C_T$  value within each strain. The  $\Delta C_T$  values in the experimental CRISPRa strains were then compared with the nontargeting CRISPRa control strain to obtain a  $\Delta\Delta C_T$  value and finally a fold difference in expression of the experimental gene, and are listed in [Supplementary Table 3](#).

## RNA sequencing

RNA preparation and sequencing (RNA-seq) were performed as previously described, with minor modifications ([Razzaq et al. 2021](#)). Strains were grown in YPD at  $30^{\circ}\text{C}$  overnight preceding RNA extraction. RNA extraction, sample QC, library preparations, and sequencing reactions were conducted at GENEWIZ, LLC./Azenta US, Inc. as follows: Total RNA was extracted using Qiagen RNeasy Plus Universal kit following manufacturer's instructions (Qiagen). RNA samples were quantified using Qubit 2.0 Fluorometer (Thermo Fisher Scientific) and RNA integrity was checked with 4200 TapeStation (Agilent Technologies). ERCC RNA Spike-In Mix kit (cat. 4456740) from Thermo Fisher Scientific was added to normalized total RNA prior to library preparation following the manufacturer's protocol. The RNA-seq library was prepared using the NEBNext Ultra II RNA Library Prep kit for Illumina using the manufacturer's instructions (New England Biolabs). Briefly, mRNAs were initially enriched with Oligod(T) beads. Enriched mRNAs were fragmented for 15 min at  $94^{\circ}\text{C}$ . First-strand and second-strand cDNAs were subsequently synthesized. cDNA fragments were end-repaired and adenylated at 3' ends, and universal adapters were ligated to cDNA fragments, followed by index addition and library enrichment by PCR with limited cycles. The sequencing library was validated on the Agilent TapeStation (Agilent Technologies), and quantified by using Qubit 2.0 Fluorometer (Thermo Fisher Scientific) as well as by quantitative PCR (KAPA Biosystems). The sequencing libraries were multiplexed and clustered onto a flowcell. After

clustering, the flowcell was loaded onto the Illumina HiSeq 4000 or equivalent instrument according to the manufacturer's instructions. The samples were sequenced using a  $2 \times 150$  bp paired end configuration. Image analysis and base calling were conducted by the Illumina Control Software.


Raw sequence data (.bcl files) generated from the Illumina instrument was converted into fastq files and de-multiplexed using Illumina bcl2fastq 2.20 software. One mis-match was allowed for index sequence identification. Sequence reads were trimmed to remove possible adapter sequences and nucleotides with poor quality using Trimmomatic v.0.36, and trimmed reads were mapped to the *Candida albicans* reference genome available on ENSEMBL using the STAR aligner v.2.5.2b. The STAR aligner is a splice aligner that detects splice junctions and incorporates them to help align the entire read sequences. BAM files were generated as a result of this step. Unique gene hit counts were then calculated by using featureCounts from the Subread package v.1.5.2. The hit counts were summarized and reported using the gene\_id feature in the annotation file. Only unique reads that fell within exon regions were counted. After extraction of gene hit counts, the gene hit counts table was used for downstream differential expression analysis. Using DESeq2, a comparison of gene expression between the control and experimental groups of samples was performed. The Wald test was then used to generate P-values and log<sub>2</sub>-fold changes. Genes with an adjusted P-value  $<0.05$  and absolute log<sub>2</sub>-fold change  $>1$  were called as differentially expressed genes for each comparison. The data have been deposited in NCBI's Gene Expression Omnibus (GEO) ([Edgar et al. 2002](#)) and are accessible through GEO Series accession number GSE201904 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE201904>).

## Results

### Development of a single-plasmid CRISPRa tool in *C. albicans*

First, we designed a single-plasmid CRISPRa system for targeted gene activation, optimized for *C. albicans*. We exploited an integrating plasmid backbone, which we previously used for CRISPRi in *C. albicans*, containing a codon-optimized dCas9 and a Golden Gate assembly compatible sgRNA cloning site ([Wensing et al. 2019](#)). This plasmid can be linearized to integrate at the *C. albicans* NEUT5L locus (a large intergenic region whose disruption does not affect fungal fitness; [Gerami-Nejad et al. 2013](#)), and contains a dual SapI cloning site for facile and highly efficient Golden Gate cloning of the sgRNA targeting sequence. This plasmid also contains a dominant NAT resistance cassette, to enable strain generation in a diversity of fungal strains, including clinical isolates.

To generate a version of dCas9 that could lead to strong constitutive activation of genes of interest, we generated a plasmid with dCas9 fused to the tripartite activator complex, VPR ([Fig. 1, a and b](#)). VPR consists of 3 previously described transcriptional activators: (1) VP64, a viral transcriptional activator domain composed of 4 tandem copies of VP16 (Herpes simplex viral protein 16), which has been used extensively in CRISPRa systems ([La Russa and Qi 2015](#)); (2) p65, the NF- $\kappa$ B subunit responsible for the strong transcription activation in mammalian cells ([Schmitz and Baeuerle 1991](#)); and (3) Rta, the Epstein-Barr virus R transactivator, with strong transcriptional activation properties ([Hardwick et al. 1992](#); [Fig. 1, a and b](#)). Together the VPR activator complex fused to dCas9 and targeted to the promoter region of genes has previously been demonstrated to generate robust gene activation in *S. cerevisiae* ([Chavez et al. 2015](#)). Exploiting the ability of dCas9 to




**Fig. 1.** Design of a CRISPRa system for *Candida albicans*. a) The single-plasmid system contains a catalytically inactive (“dead”) dCas9 protein, paired with the VPR complex of activator domains to cause upregulation at a target locus. The sgRNA region can be subjected to Golden Gate cloning with a custom N20 sequence in order to target the dCas9-VPR complex to a desired region of DNA through complementary base-pairing. The PaclI enzyme is used to linearize the plasmid at the NEUT5L locus, which allows the plasmid to be integrated into the cell’s chromosome and remain in the cell following cell division. Panel created with BioRender.com. b) Once the sgRNA recognizes and binds to a target sequence of DNA, the dCas9-VPR complex will cause the recruitment of transcription initiation complex factors to the adjacent genetic material, thus increasing gene expression from that region. Panel created with BioRender.com. c) Wild-type *C. albicans* cells and cells containing a nontargeting CRISPRa plasmid were grown in YPD at 30°C to compare fitness under standard laboratory conditions. Growth was recorded via optical density at 600 nm at 15 min intervals over the course of 24 h. Data points show the mean and standard deviation at each time point ( $n = 32$ ).

bind and be targeted to genomic loci via sgRNAs enables us to design sgRNAs that will target dCas9 along with these transcriptional activators to any gene promoter of interest.

This plasmid was generated by homology-based cloning and validated via Sanger sequencing, and has been deposited to Addgene (Supplementary Table 1, Addgene plasmid #182707). To confirm if the expression of the dCas9-VPR construct had an impact on fungal fitness, we compared the growth of a wild-type *C. albicans* strain with a strain transformed with the dCas9-VPR CRISPRa plasmid (expressing a nontargeting sgRNA: a random sequence of nucleotides that have no predicted complementarity to any region in the *C. albicans* genome). We found no difference in growth between these strains (Fig. 1c), suggesting no significant impact of the CRISPRa system on *C. albicans* fitness.

### Validation and sgRNA design principles for CRISPRa overexpression in *C. albicans*

Next, we aimed to validate that the CRISPRa system could successfully overexpress genes in *C. albicans*, and sought to determine the optimal sgRNA design strategy to maximize genetic regulation. Our previous *C. albicans* CRISPRi system exploited sgRNAs ~50–150 bp upstream of the start codon to functionally repress genes of interest (Wensing et al. 2019). However, CRISPRa systems are known to have different optimal sgRNA targeting regions (Jensen 2018; Sanson et al. 2018), and our previous design did not take into account the transcription start site (TSS), which is known to be an important predictor of regulatory efficiency for CRISPRi/a systems (Smith et al. 2016; Jensen 2018), but which had not been robustly characterized across all *C. albicans* genes at the time.



**Fig. 2.** Determining optimal guide-design principles for CRISPRa overexpression in *Candida albicans*. CRISPRa strains were created using 6 different sgRNA molecules designed to target different positions in the promoter region of each of a) *KSP1*, b) *ALS3*, and c) *FGR28* upstream of either the start codon or the predicted dominant TSS. Fold change in the expression of the target gene relative to the housekeeping gene *ACT1* in the experimental strains and the nontargeting CRISPRa control strain was measured with RT-qPCR and compared. Data points represent the mean fold difference and standard deviation in expression of the respective target gene in the CRISPRa strains compared with the nontargeting CRISPRa control ( $n=3$ ). d) Fold change in the basal expression level of *ALS3* relative to the housekeeping gene *ACT1* in the nontargeting CRISPRa control strain grown in either standard laboratory conditions (YPD at 30°C) or conditions known to favor a high level of *ALS3* expression (RPMI at 37°C) was measured with RT-qPCR. Data represent the mean fold difference and standard deviation in expression of *ALS3* in the nontargeting CRISPRa control strain in both conditions normalized to growth in YPD at 30°C ( $n=3$ ). e) Fold change in the expression of *ALS3* relative to the housekeeping gene *ACT1* in two CRISPRa strains targeting *ALS3* for overexpression as well as the nontargeting CRISPRa control strain grown in either YPD at 30°C or RPMI at 37°C was measured with RT-qPCR. Data represent the mean fold difference and standard deviation in expression of *ALS3* in all strains in both conditions normalized to the nontargeting CRISPRa control strain (WT;  $n=3$ ).

More recently, the TSS for many *C. albicans* genes has been predicted with single-nucleotide resolution using nAnt-iCAGE technology, which specifically identifies the 5' end of RNA fragments (Lu and Lin 2021). Thus, we sought to determine the optimal design for CRISPRa sgRNAs based on the start codon and the TSS. We identified 3 *C. albicans* genes with a predicted TSS that was particularly far upstream of the start codon compared with all other genes in the genome (801–952 bp upstream): *KSP1*, *ALS3*, and *FGR28*. We designed 6 sgRNAs for each gene, targeting upstream of the start codon (as previously designed for CRISPRi; Wensing et al. 2019), as well as upstream of the TSS (based on predicted TSS; Lu and Lin 2021).

We generated CRISPRa plasmids based on these sgRNAs, used these to create *C. albicans* strains, and monitored overexpression of each gene via RT-qPCR. This analysis validated our novel

CRISPRa system and its ability to overexpress *C. albicans* genes ~2- to 12-fold, based on different sgRNA designs and target genes (Fig. 2). We found that in some instances, such as for *KSP1*, the TSS was in fact a better predictor of CRISPRa overexpression, as the most robust overexpression of this gene was observed with sgRNAs targeted upstream of the TSS, while no overexpression was observed based on targeting upstream of the start codon (Fig. 2a). In other instances, such as for *ALS3*, sgRNA targeting upstream of the start codon led to very high levels of overexpression, while sgRNAs targeted to the TSS region led to moderate overexpression (Fig. 2b). For *FGR28*, modest overexpression (~2-fold) was achieved with sgRNAs targeted to either the start codon or the TSS upstream regions (Fig. 2c), together suggesting that optimal sgRNA design can be predicted based on targeting the TSS and/or start codon, but varies by gene. This data also suggest

that our CRISPRa tool may have additional applications in the functional assessment of key transcriptional regulatory regions in *C. albicans*. These differences in sgRNA targeting may be based on differences in overall sgRNA efficiency or may be attributed to inaccurate TSS predictions, or complex promoter dynamics and the preferential use of different TSS regions in different conditions (McMillan et al. 2019).

Given that our CRISPRa system has the benefit of exploiting a gene's endogenous promoter for overexpression, we next wanted to determine if our CRISPRa system could be used to promote enhanced expression of genes under conditions that already favored high gene expression. We focused on *ALS3*, where expression levels are known to be significantly enhanced upon growth in conditions that favor filamentous or biofilm growth, such as growth in RPMI media at 37°C (Bahn et al. 2007; Ruben et al. 2020; Deng et al. 2021). We compared basal *ALS3* transcript levels of the *C. albicans* control strain grown in YPD at 30°C compared with RPMI at 37°C via RT-qPCR, and confirmed that *ALS3* exhibits ~100-fold higher expression in RPMI at 37°C (Fig. 2d). To determine whether our CRISPRa system could promote enhanced overexpression of *ALS3* in these conditions, we monitored *ALS3* expression in two CRISPRa *ALS3* strains (from Fig. 2b) in RPMI at 37°C. We found *ALS3* transcript levels in the CRISPRa strain were further overexpressed ~4-fold in RPMI at 37°C, despite the existing high level of *ALS3* under these conditions (Fig. 2e). However, while *ALS3* is overexpressed in RPMI via CRISPRa, the degree of overexpression is relatively lower than what occurs in YPD conditions (Fig. 2e). Together this suggests that our CRISPRa system can effectively be used to increase transcription under conditions that already favor high levels of expression of a target gene of interest, but that there may be a maximal level of overexpression possible under these conditions.

### Use of CRISPRa to study antifungal drug susceptibility phenotypes

To validate that the CRISPRa system could recapitulate phenotypes associated with *C. albicans* gene overexpression, we targeted genes with known phenotypes associated with high levels of expression. Initially, we focused on *CDR1*, a well-characterized *C. albicans* gene, which encodes a drug efflux pump (Prasad et al. 2015), and whose overexpression is known to play a role in resistance to azole antifungals in laboratory and clinical isolates (Prasad et al. 1995; Sanglard et al. 1995; White 1997; Marr et al. 1998, 2001; Niimi et al. 2004; Manoharlal et al. 2008). To determine an optimal range in which to target our sgRNA in the *CDR1* promoter, we designed 3 sgRNAs targeting loci ~90, 200, and 290 bp upstream of the *CDR1* TSS. These sgRNAs were each cloned into our CRISPRa dCas9-VPR backbone and transformed into *C. albicans*. Next, we quantified the expression of *CDR1* in the 3 different fungal strains via RT-qPCR, compared with a control strain containing the CRISPRa plasmid with a nontargeting sgRNA. We found that targeting the dCas9-VPR construct to the *CDR1* promoter enhanced transcription by 2.9- to 5.6-fold (Fig. 3a), indicating that this CRISPRa system is able to drive enhanced levels of gene expression from this target gene of interest. For *CDR1*, we found the highest level of overexpression was obtained by targeting the dCas9-VPR construct ~200 bp upstream of the TSS (Fig. 3a), which helped further inform the design of sgRNAs for subsequent target genes.


In order to determine if *CDR1*-overexpressing strains had measurable changes in antifungal drug susceptibility phenotypes, we performed MIC assays with these strains in the presence of the antifungal drug fluconazole. We found that strains

overexpressing *CDR1* exhibited decreased sensitivity to fluconazole compared with the control strain and that higher levels of overexpression corresponded with lower levels of susceptibility (Fig. 3b). Together, this confirms that our CRISPRa system can be exploited to functionally overexpress target genes of interest, and recapitulate phenotypes associated with high levels of gene expression. Further, targeting dCas9-VPR to different promoter loci leads to variable levels of overexpression, which could be used to study correlations between expression levels and associated phenotypes.

We next sought to determine the ability of our CRISPRa system to be employed to characterize a gene whose overexpression has never been profiled with regard to antifungal drug susceptibility, to our knowledge. We selected the gene *YAK1* as a target for overexpression with CRISPRa. *Candida albicans* *YAK1* is a predicted serine-threonine protein kinase, inactivation of which renders cells more sensitive to the antifungal amphotericin B (Chen et al. 2018). While *YAK1* overexpression has been shown to promote filamentation in *C. albicans* (MacAlpine et al. 2021), and there is emerging evidence for a direct relationship between the 2 phenotypes (Sharma et al. 2019), the role of overexpressing this factor in antifungal drug susceptibility has not previously been established. We generated a CRISPRa strain overexpressing *YAK1* by ~2-fold (Fig. 3c) and profiled the ability of this strain to grow in the presence of amphotericin B. We found that the *YAK1* CRISPRa strain was less sensitive to amphotericin B based on MIC testing on this strain compared with the control strain (Fig. 3d). This highlights our capacity to exploit this CRISPRa system to investigate novel phenotypes in *C. albicans*.

### Validation of CRISPRa overexpression targeting fungal pathogenesis traits

Next, we aimed to validate the CRISPRa system as a means to assess *C. albicans* genes involved in fungal pathogenesis traits. We focused on *ALS1*, a well-characterized *C. albicans* adhesin gene, with important roles in adherence, biofilm formation, and virulence (Fu et al. 2002; Kamai et al. 2002; Alberti-Segui et al. 2004; Zhao et al. 2004; Nobile et al. 2008; Rosiana et al. 2021). Previous work has demonstrated that overexpression of *ALS1* via promoter replacement increases fungal adherence (Fu et al. 2002); therefore, we chose to overexpress this gene using our CRISPRa system. We designed and cloned 2 sgRNAs targeting ~130 and 260 bp upstream of the *ALS1* TSS, and generated *C. albicans* *ALS1* CRISPRa strains with each of these constructs. We performed RT-qPCR to confirm the overexpression of these strains, which enhanced the expression of *ALS1* by ~25- and ~6-fold, respectively (Fig. 4a). To confirm phenotypes associated with overexpression of *ALS1*, we performed two distinct biofilm growth assays that compared the *ALS1* CRISPRa strain to the CRISPRa control strain. In the first assay, we allowed biofilms to establish over 24 h in flat-bottom 96-well polystyrene plates, removed planktonic cells, and quantified the metabolic activity of the biofilm via XTT reduction, relative to the planktonic cell population. We found that the *ALS1* CRISPRa strain formed significantly more robust biofilms, relative to the control strain ( $P < 0.0005$ ; Fig. 4b). Interestingly, biofilm growth was not correlated with *ALS1* overexpression levels, as the strain with ~25-fold *ALS1* overexpression formed less robust biofilms compared with the strain expressing ~6-fold more *ALS1*. This suggests that very high levels of *ALS1* may ultimately impair biofilm formation, or that the level of *ALS1* expression varies in the conditions used for RT-qPCR analysis compared with biofilm growth, perhaps due to condition-specific repositioning in the TSS.




**Fig. 3.** Using CRISPRa to validate antifungal drug susceptibility phenotypes. a) Three *C. albicans* CRISPRa strains were created targeting the drug transporter gene *CDR1* for overexpression. Fold change in the expression of *CDR1* relative to the housekeeping gene *ACT1* in the 3 experimental strains and the nontargeting CRISPRa control strain was measured with RT-qPCR. All 3 experimental strains are represented by the position of their sgRNA in the promoter region upstream of the proposed dominant *CDR1* transcriptional start site. Data represent the mean fold difference and standard deviation in expression of *CDR1* in the CRISPRa strains compared with the nontargeting CRISPRa control (WT;  $n = 3$ ). b) The fitness of all 3 CRISPRa strains targeting *CDR1* for overexpression was measured in the presence of the antifungal drug fluconazole with a minimum inhibitory concentration assay. The  $OD_{600}$  value of each strain in each concentration of fluconazole was measured to represent growth in differing conditions, where  $OD_{600}$  values approaching 1.0 reflect the highest fitness. c) Fold change in the expression of *YAK1* relative to the housekeeping gene *ACT1* in the experimental strain and the nontargeting CRISPRa control strain was measured with RT-qPCR. Data represent the mean fold difference and standard deviation in expression of *YAK1* in the CRISPRa strains compared with the nontargeting CRISPRa control (WT;  $n = 3$ ). d) The fitness of the CRISPRa strain targeting *YAK1* for overexpression was measured in the presence of the antifungal drug amphotericin B with a minimum inhibitory concentration assay. The  $OD_{600}$  value of each strain in each concentration of amphotericin B was measured to represent growth in differing conditions, where  $OD_{600}$  values approaching 1.0 reflect the highest fitness. ND: no drug.

Next, to mimic conditions associated with fungal biofilm formation in catheter-associated urinary tract infections (CAUTIs), we used a urine-based assay and analyzed biofilm formation of a *C. albicans* strain overexpressing *ALS1*, relative to its control strain, using crystal violet staining. For this assay, we followed up on the *ALS1* CRISPRa strain with ~6-fold overexpression of *ALS1*, which had more robust biofilm growth than the control strain (Fig. 4, a and b). The *C. albicans* strains were incubated for 48 h in human urine supplemented with BSA in fibrinogen-coated 96-well plates. Fibrinogen has been shown to be a critical protein for pathogen binding in CAUTIs, by providing a scaffold for biofilm formation and a nutritional source for pathogens (Flores-Mireles et al. 2016; La Bella et al. 2021; Andersen et al. 2022). The *ALS1* CRISPRa overexpression strain showed significantly greater biofilm formation compared with the corresponding wild-type strain ( $P < 0.0001$ ; Fig. 4c). This difference in biofilm formation suggests that *ALS1* promotes binding to fibrinogen and plays a vital role in biofilm formation in urine.

To assess the specificity of this CRISPRa system and ensure that there were no off-target effects from the dCas9-sgRNA complex, we monitored global changes in transcription in the *ALS1* CRISPRa strain via RNA-seq. Since *ALS1* is a member of the large agglutinin-like sequence (ALS) gene family (Hoyer et al. 2008; Hoyer and Cota 2016), it may be a likely candidate for any

potential off-target effects. We also considered *ALS1* as a strong candidate for this RNA-seq off-target profiling since *ALS1* is not known to be a transcriptional regulator, and therefore its overexpression is unlikely to lead to direct downstream transcriptional alterations that would obscure our analysis of CRISPRa off-target effects. Therefore, we performed RNA-seq on the validated CRISPRa strain overexpressing *ALS1* by ~25-fold as well as on a control strain containing the dCas9 cassette and a nontargeting sgRNA when grown in standard laboratory conditions. This analysis confirmed the specific overexpression of *ALS1* (log<sub>2</sub>-fold change ~3.3 via RNA-seq analysis, adjusted  $P$ -value  $3.07E-142$ ; Fig. 4d, Supplementary Table 2). Five additional transcripts were identified as significantly modulated (adjusted  $P$ -value  $< 0.05$ , absolute log<sub>2</sub>-fold change  $> 1$ ; Supplementary Table 2). However, their altered expression was much less significant than *ALS1* (Fig. 4d), and none of them had an adjusted  $P$ -value of  $< 0.03$ , differing from the adjusted  $P$ -value of  $3E-142$  for *ALS1*. We suspect these additional transcripts' altered expression is likely unrelated to the CRISPRa system based on the fact that 4/5 genes were down-regulated instead of up-regulated, and none of the genes are in proximity to the *ALS1* locus, nor do they appear to share any sgRNA target similarities in their promoter regions. Together this suggests that this CRISPRa system is likely specific with regards to gene targeting.





**Fig. 4.** Validation of CRISPRa overexpression and target specificity through targeting fungal pathogenesis traits. a) Two *C. albicans* CRISPRa strains targeting different positions in the promoter region of the adhesin protein gene *ALS1* were created. Fold change in the expression of *ALS1* relative to the housekeeping gene *ACT1* in both experimental strains and the nontargeting CRISPRa control strain was measured with RT-qPCR. Experimental strains are represented by the position of their sgRNA in the promoter region upstream of the proposed dominant *ALS1* transcriptional start site. Data points represent the mean fold difference and standard deviation in expression of *ALS1* in the CRISPRa strains compared with the nontargeting CRISPRa control (WT;  $n = 3$ ). b) Two CRISPRa strains targeting *ALS1* and a nontargeting CRISPRa control strain were grown in the biofilm-inducing media RPMI and left to grow statically at 37°C for 24 h to allow biofilms to form. The robustness of biofilms formed by all strains was elucidated through an XTT reduction assay and measuring of the resulting  $OD_{490}$  values before normalizing final values to nonbiofilm-forming planktonic cells. Data points represent the mean and the standard error of the mean ( $n = 72$ ). Differences between groups were tested for significance using Dunnett's multiple comparison test. \*\*\*\* $P < 0.0001$ . \*\*\* $P < 0.001$ . c) A *C. albicans* CRISPRa strain overexpressing *ALS1* ~6-fold and a nontargeting CRISPRa strain were grown in human urine supplemented with fibrinogen and BSA at 37°C for 48 h to simulate a catheterized bladder environment. The robustness of biofilms formed by both strains was measured by reading the  $OD_{595}$  value following a crystal violet assay. Data points are combined from 3 independent experiments where  $n = 32$  for the nontargeting CRISPRa control and  $n = 48$  for the CRISPRa *ALS1* overexpression strain. Differences between groups were tested for significance using the Mann-Whitney *U* test. \*\*\*\* $P < 0.0001$ . d) Global gene expression of a CRISPRa strain overexpressing *ALS1* ~25-fold, as well as a nontargeting CRISPRa control strain, was measured using RNA-seq. Differential gene expression analysis showed that *ALS1* had a  $\log_2$ -fold change of ~3.3 in the *ALS1* overexpression strain compared with the CRISPRa control strain, with an adjusted *P*-value of  $3.07E-142$ , and showed no additional transcripts that had differential expression with an adjusted *P*-value of  $< 0.03$ .

## Discussion

Here, we present the development, validation, and application of a single-plasmid-based CRISPRa platform for efficient genetic overexpression in *C. albicans*. We demonstrate the ability of this system to overexpress genes ~2- to 20-fold, and recapitulate important biological phenotypes associated with genetic overexpression, including biofilm formation and antifungal drug susceptibility. In addition, we assess sgRNA design principles based on predicted TSS, and validate the specificity of this system by monitoring off-target effects via RNA-seq. Together, this work demonstrates an effective, efficient, and specific tool for genetic overexpression in *C. albicans* to enhance the available techniques for genetic modulation in this important fungal pathogen. We believe this tool will have practical application for many researchers, and can be easily applied in other laboratories by following plasmid and strain construction protocols for *C. albicans*-based dCas9 strategies, which we previously described in detail (Wensing and Shapiro 2022). Based on our results, we advise designing and testing 2-3 sgRNAs in regions -100 to -350 bp upstream of both the start codon (ATG) and TSS of the target gene of interest to achieve high levels of overexpression.

Studying genetic overexpression is an important strategy for functional genetic analysis, as overexpressing genes can mimic natural biological states where genes are expressed at high levels. In *C. albicans*, there are numerous scenarios in which the overexpression of genes occurs, including mutations that modify the function of transcriptional regulators and promote the expression of downstream genes (Coste et al. 2004, 2006), as well as aneuploidies such as trisomies (Selmecki et al. 2006; Forche et al. 2009, 2018; Ford et al. 2015; Li et al. 2015; Anderson et al. 2017; Ene et al. 2018), isochromosome formation (Selmecki et al. 2006, 2008), and copy number variations (Ford et al. 2015; Todd and Selmecki 2020). These overexpression events are frequently detected among *C. albicans* clinical isolates (Selmecki et al. 2006; Ford et al. 2015; Anderson et al. 2017) and can play an important role in drug resistance (Selmecki et al. 2006, 2008; Li et al. 2015; Todd and Selmecki 2020), host adaptation (Forche et al. 2018, 2019), and other stress tolerance phenotypes (Yang et al. 2019, 2021). For example, amplification of part or all of chromosome 5, via trisomy or isochromosome formation has been tightly linked to the development of resistance to azole antifungals in *C. albicans* via the increased copy number of genes encoding the azole target, Erg11, and the transcriptional regulator of drug efflux, Tac1 (Selmecki et al. 2008). The prevalence and importance of these phenomena associated with increased gene expression highlight the relevance of developing new tools to targetedly assess phenotypes associated with genetic overexpression in *C. albicans*.

Harnessing the cell's natural transcriptional machinery to increase gene expression with CRISPRa also leads to a unique set of obstacles. While our RNA-seq results suggest that our CRISPRa system is specific when paired with one sgRNA targeted to ALS1, there remains the possibility of off-target effects occurring in different CRISPRa constructs (Zhang et al. 2015). Using complementary genetic tools to confirm phenotypes observed from CRISPRa can help mitigate this. One phenomenon shared among many different CRISPRa activation domains, including VPR, is a reduced capacity to induce overexpression in genes that are already expressed at a high levels (Chavez et al. 2016). This may explain why we achieved only modest overexpression of YAK1, the gene with the highest basal expression level of all genes tested, but also why a ~2-fold change in transcript abundance still resulted in a clear decrease in susceptibility to amphotericin B

(Fig. 3d). Incidentally, the gene that we achieved the lowest overexpression with across all of our testing, *FGR28*, had the lowest basal expression (Supplementary Table 3). However, efficiency of the sgRNA molecule and its exact position in the target region of the gene of interest, as well as a host of promoter dynamics, can also influence CRISPR-Cas9 efficacy (Radziszewska et al. 2016; Cámara et al. 2020; Lu and Lin 2021). This illustrates why multiple sgRNAs need to be designed when targeting a gene with CRISPRa to maximize successful overexpression of the target gene. While the rapid and inexpensive nature of CRISPRa strain construction justifies the need to design several sgRNA molecules for a given gene, limitations resulting from the specific "NGG" PAM recognition site by *Streptococcus pyogenes* Cas9 proteins leads to the restraint that, on rare occasions, it may not be possible to produce any efficient sgRNA molecules in a suitable region of the promoter (Uthayakumar et al. 2021). It may therefore be prudent to further expand the CRISPR toolbox in *C. albicans* by introducing novel Cas9 variants with differing PAM recognition sites, or by constructing novel CRISPR systems exploiting alternative Cas proteins with different and broader PAM recognition capacities, offering the ability to target regions in the genome that cannot be manipulated with current CRISPR-Cas9 tools (Hu et al. 2018; Ciurkot et al. 2021).

Alternative overexpression strategies in *C. albicans* that involve introducing an extra copy of a gene of interest or introducing an alternative promoter have been used to construct and screen libraries composed of hundreds and even thousands of protein-coding genes (Rai et al. 2021). These studies have revealed novel insights into gene function related to morphogenesis (Chauvel et al. 2012), cell cycle progression (Jaitly et al. 2022), drug tolerance (Delarze et al. 2020), and biofilm formation (Cabral et al. 2014). Previous comparisons between CRISPRa and ORF-based overexpression libraries in other species have highlighted the potential for these complementary strategies to both confirm findings and unveil distinct genetic hits (Sansom et al. 2018). CRISPRa offers the advantage of being high throughput and easily scalable into a large-scale pooled format, and mutant libraries in different strain backgrounds can therefore be quickly generated (Sanjana 2017; Momen-Roknabadi et al. 2020). In contrast, ORF-based overexpression strategies may be preferable if attempting to overexpress only a single transcript variant or isoform (La Russa and Qi 2015), and promoter replacement strategies may be more valuable to overexpress diverse genes to similar levels. Both of these strategies would be advantageous as well in rare instances when a target sequence lacks a PAM site. Non-CRISPR overexpression libraries previously assembled in *C. albicans* have the potential weakness of missing phenotypes if the level of expression achieved in mutant strains is too low (Znaidi et al. 2018). The ability of our system to cause upwards of 25-fold overexpression of gene targets is thus an important feature, and future iterations of this CRISPRa system could also exploit an inducible dCas9-VPR construct for tunability of gene expression. Furthermore, phenotypes in mutant strains may only be observable if the level of overexpression closely mimics the level of upregulation that naturally occurs in the cell in a given condition (Znaidi et al. 2018). That our CRISPRa system exploits the endogenous promoter to drive upregulation may thus be critical in understanding natural overexpression mechanisms and regulatory circuitry that other systems are not capable of studying.

While large-scale CRISPRa libraries have not been generated in *C. albicans*, their application in other organisms, namely mammalian cell lines, has enabled important discoveries in cancer

biology, stem cell reprogramming, cellular fitness, and drug resistance (Gilbert et al. 2014; Konermann et al. 2015; Horlbeck et al. 2016; Jung et al. 2017; Jost et al. 2017; Kampmann 2018; Sanson et al. 2018; Yang et al. 2019). The development of CRISPRa systems in *C. albicans* could enable the development of larger overexpression libraries for functional genomic screens to complement existing fungal gene deletion and depletion libraries (Roemer et al. 2003; Homann et al. 2009; Blankenship et al. 2010; Noble et al. 2010), that have been subject to robust screening and analysis under diverse conditions (Gervais et al. 2021) to reveal critical new insight into fungal biology (Lohse et al. 2016; Hossain et al. 2020), pathogenesis (Noble et al. 2010; Nobile et al. 2012; Ryan et al. 2012; O'Meara et al. 2015; Lee et al. 2016; Hossain et al. 2020), and mechanisms of drug and stress tolerance (Motaung et al. 2017; Mount et al. 2018; Caplan et al. 2018; Fu et al. 2021). The application of similar CRISPRa systems to other fungal organisms would lend further insight into fungal biology across this kingdom. Many CRISPR tools have been developed in a diversity of fungal organisms (Muñoz et al. 2019; Cai et al. 2019; Román, Prieto, et al. 2019; Schuster and Kahmann 2019; Morio et al. 2020; Ouedraogo and Tsang 2020; Shan et al. 2021), including numerous non-*albicans* *Candida* species (Enkler et al. 2016; Lombardi et al. 2017, 2019; Zoppo et al. 2019; Maroc and Fairhead 2019; Uthayakumar et al. 2021; Ennis et al. 2021; Santana and O'Meara 2021), and other prominent pathogens such as *Cryptococcus neoformans* (Fan and Lin 2018; Wang 2018; Huang et al. 2021) and *Aspergillus fumigatus* (Fuller et al. 2015; Zhang et al. 2016; Al Abdallah et al. 2018; van Rhijn et al. 2020). These CRISPR platforms could be adapted to similar CRISPRa tools in these organisms, with many possible applications for the analysis of pathogen biology or drug target identification (Li et al. 2004; Luesch et al. 2005; Smith et al. 2010; Arnoldo et al. 2014; Begolo et al. 2014).

Genetic overexpression systems also have many useful applications for industrially important fungi, as targeted genetic overexpression can be exploited for metabolic engineering to enhance the production of commercially important metabolites (Donohoue et al. 2018; Zhao et al. 2018; Ouedraogo and Tsang 2020; Jiang et al. 2021). CRISPR techniques have already been adapted to numerous industrially relevant fungi (Nødvig et al. 2015; Kwon et al. 2019; Jiang et al. 2021; Wilson and Harrison 2021), and CRISPRa has recently been optimized for use in the filamentous fungi *Aspergillus nidulans* and *Penicillium rubens* (Roux et al. 2020; Mózsik et al. 2021). These CRISPRa systems were applied to activate transcriptionally silent biosynthetic gene clusters in these filamentous fungi, with important applications for the discovery of novel bioactive molecules (Roux et al. 2020; Mózsik et al. 2021). As many *Candida* species play important industrial roles in manufacturing metabolites for food and pharmaceutical industries, as well as remediating wastewater via hydrocarbon degradation (Joo et al. 2008; Gargouri et al. 2015; Kieliszek et al. 2017; Payen and Thompson 2019; García-Béjar et al. 2020), species-specific adaptations to the CRISPRa tool presented here could produce optimized *Candida* strains for valuable biomanufacturing and bioremediation applications.

## Data availability

The *C. albicans* CRISPRa plasmid backbone was deposited to Addgene as plasmid number #182707. RNA-seq data were deposited in NCBI GEO repository (accession number GSE201904; <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE201904>).

Supplemental material available at G3 online.

## Acknowledgments

The authors thank Shapiro lab members, past and present, for helpful discussions and support of this project, as well as Dr Anna Selmecki and Dr Petra Vande Zande (University of Minnesota) for helpful feedback. The authors thank Dr Zhenguo Lin (Saint Louis University) for helpful discussions on TSS prediction, and Kieran Shah for programming support on this project. They thank Jing Zhang from the University of Guelph Advanced Analysis Centre for RT-qPCR support.

## Funding

This work was supported by an Natural Sciences and Engineering Research Council of Canada (NSERC Discovery) Grant (RGPIN-2018-4914), an Ontario Early Research Award, and a CIFAR Azrieli Global Scholar award to R.S.S. N.C.G., L.F.W., and M.F. are EvoFunPath Fellows (NSERC CREATE), L.F.W. is supported by an Ontario Graduate Scholarship, M.F. is supported by an NSERC-CGS-M scholarship, and R.O.C.L. was supported by a MITACS Globalink Research Internship. A.C. is supported by a Career Award for Medical Scientists from the Burroughs Wellcome Fund. Work by A.L.F.M. and F.S.T. was supported by institutional funds from the University of Notre Dame as well as National Institute of Health grant R01-DK128805 (to A.L.F.M. and A.A.L.B.), and the Arthur J. Schmitt Presidential Leadership Foundation (to A.A.L.B.).

## Conflicts of interest

The authors declare no conflict of interest.

## Literature cited

- Adli M. The CRISPR tool kit for genome editing and beyond. *Nat Commun.* 2018;9(1):1911. doi:10.1038/s41467-018-04252-2
- Al Abdallah Q, Souza ACO, Martin-Vicente A, Ge W, Fortwendel JR. Whole-genome sequencing reveals highly specific gene targeting by *in vitro* assembled Cas9-ribonucleoprotein complexes in *Aspergillus fumigatus*. *Fungal Biol Biotechnol.* 2018;5(1):11. doi:10.1186/s40694-018-0057-2
- Alberti-Segui C, Morales AJ, Xing H, Kessler MM, Willins DA, et al. Identification of potential cell-surface proteins in *Candida albicans* and investigation of the role of a putative cell-surface glycosidase in adhesion and virulence. *Yeast.* 2004;21(4):285–302. doi:10.1002/yea.1061
- Andersen MJ, Fong C, La Bella AA, Molina JJ, Molesan A, et al. Inhibiting host-protein deposition on urinary catheters reduces associated urinary tract infections. *eLife.* 2022;11:e75798. doi:10.7554/eLife.75798
- Anderson MZ, Saha A, Haseeb A, Bennett RJ. A chromosome 4 trisomy contributes to increased fluconazole resistance in a clinical isolate of *Candida albicans*. *Microbiology.* 2017;163(6):856–865. doi:10.1099/mic.0.000478
- Arendrup MC, Patterson TF. Multidrug-resistant *Candida*: epidemiology, molecular mechanisms, and treatment. *J Infect Dis.* 2017; 216(suppl 3):S445–S451. doi:10.1093/infdis/jix131
- Arnoldo A, Kittanakom S, Heisler LE, Mak AB, Shukalyuk AI, et al. A genome scale overexpression screen to reveal drug activity in human cells. *Genome Med.* 2014;6(4):32. doi:10.1186/gm549
- Bahn Y-S, Molenda M, Staab JF, Lyman CA, Gordon LJ, et al. Genome-wide transcriptional profiling of the cyclic AMP-dependent signaling pathway during morphogenic

- transitions of *Candida albicans*. Eukaryot Cell. 2007;6(12):2376–2390. doi:10.1128/EC.00318-07
- Begolo D, Erben E, Clayton C. Drug target identification using a trypanosome overexpression library. Antimicrob Agents Chemother. 2014;58(10):6260–6264. doi:10.1128/AAC.03338-14
- Benedict K, Jackson BR, Chiller T, Beer KD. Estimation of direct healthcare costs of fungal diseases in the United States. Clin Infect Dis. 2018;68(11):1791–1797. doi:10.1093/cid/ciy776
- Blankenship JR, Fanning S, Hamaker JJ, Mitchell AP. An extensive circuitry for cell wall regulation in *Candida albicans*. PLoS Pathogens. 2010;6:e1000752. doi:10.1371/journal.ppat.1000752
- Bock C, Datlinger P, Chardon F, Coelho MA, Dong MB, et al. High-content CRISPR screening. Nat Rev Methods Primers. 2022;2(1):1–1. doi:10.1038/s43586-021-00093-4
- Bongomin F, Gago S, Oladele R, Denning D. Global and multi-national prevalence of fungal diseases—estimate precision. J Fungi (Basel). 2017;3(4):57. doi:10.3390/jof3040057
- Cabral V, Znaidi S, Walker LA, Martin-Yken H, Dague E, et al. Targeted changes of the cell wall proteome influence *Candida albicans* ability to form single- and multi-strain biofilms. PLoS Pathog. 2014;10(12):e1004542. doi:10.1371/journal.ppat.1004542
- Cai P, Gao J, Zhou Y. CRISPR-mediated genome editing in non-conventional yeasts for biotechnological applications. Microb Cell Fact. 2019;18(1):63. doi:10.1186/s12934-019-1112-2
- Cámara E, Lenitz I, Nygård Y. A CRISPR activation and interference toolkit for industrial *Saccharomyces cerevisiae* strain KE6–12. Sci Rep. 2020;10(1):14605. doi:10.1038/s41598-020-71648-w
- Caplan T, Polvi EJ, Xie JL, Buckhalter S, Leach MD, et al. Functional genomic screening reveals core modulators of echinocandin stress responses in *Candida albicans*. Cell Rep. 2018;23:2292–2298. doi:10.1016/j.celrep.2018.04.084
- Chauvel M, Neseir A, Cabral V, Znaidi S, Goyard S, et al. A versatile overexpression strategy in the pathogenic yeast *Candida albicans*: identification of regulators of morphogenesis and fitness. PLoS One. 2012;7(9):e45912. doi:10.1371/journal.pone.0045912
- Chavez A, Scheiman J, Vora S, Pruitt BW, Tuttle M, et al. Highly efficient Cas9-mediated transcriptional programming. Nat Methods. 2015;12(4):326–328. doi:10.1038/nmeth.3312
- Chavez A, Tuttle M, Pruitt BW, Ewen-Campen B, Chari R, et al. Comparison of Cas9 activators in multiple species. Nat Methods. 2016;13(7):563–567. doi:10.1038/nmeth.3871
- Chen Y, Mallick J, Maqnas A, Sun Y, Choudhury BI, et al. Chemogenomic profiling of the fungal pathogen *Candida albicans*. Antimicrob Agents Chemother. 2018;62(2):e02365-17. doi:10.1128/AAC.02365-17
- Ciurkot K, Gorochofski TE, Roubos JA, Verwaal R. Efficient multiplexed gene regulation in *Saccharomyces cerevisiae* using dCas12a. Nucleic Acids Res. 2021;49(13):7775–7790. doi:10.1093/nar/gkab529
- Coste AT, Karababa M, Ischer F, Bille J, Sanglard D. TAC1, Transcriptional activator of CDR genes, is a new transcription factor involved in the regulation of *Candida albicans* ABC transporters CDR1 and CDR2. Eukaryot Cell. 2004;3(6):1639–1652. doi:10.1128/EC.3.6.1639-1652.2004
- Coste A, Turner V, Ischer F, Morschhäuser J, Forche A, et al. A mutation in Tac1p, a transcription factor regulating CDR1 and CDR2, is coupled with loss of heterozygosity at chromosome 5 to mediate antifungal resistance in *Candida albicans*. Genetics. 2006;172(4):2139–2156. doi:10.1534/genetics.105.054767
- Delarze E, Brandt L, Trachsel E, Patxot M, Pralong C, et al. Identification and characterization of mediators of fluconazole tolerance in *Candida albicans*. Front Microbiol. 2020;11:591140. doi:10.3389/fmicb.2020.591140
- Delgado ML, Luisa Delgado M, Luisa Gil M, Gozalbo D. *Candida albicans* TDH3 gene promotes secretion of internal invertase when expressed in *Saccharomyces cerevisiae* as a glyceraldehyde-3-phosphate dehydrogenase-invertase fusion protein. Yeast. 2003;20(8):713–722. doi:10.1002/yea.993
- Deng K, Jiang W, Jiang Y, Deng Q, Cao J, et al. ALS3 Expression as an indicator for *Candida albicans* biofilm formation and drug resistance. Front Microbiol. 2021;12:655242. doi:10.3389/fmicb.2021.655242
- Dominguez AA, Lim WA, Qi LS. Beyond editing: repurposing CRISPR-Cas9 for precision genome regulation and interrogation. Nat Rev Mol Cell Biol. 2016;17(1):5–15. doi:10.1038/nrm.2015.2
- Donohoue PD, Barrangou R, May AP. Advances in industrial biotechnology using CRISPR-Cas systems. Trends Biotechnol. 2018;36(2):134–146. doi:10.1016/j.tibtech.2017.07.007
- Eckert SE, Mühlshlegel FA. Promoter regulation in *Candida albicans* and related species. FEMS Yeast Res. 2009;9(1):2–15. doi:10.1111/j.1567-1364.2008.00455.x
- Edgar R, Domrachev M, Lash AE. Gene expression omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res. 2002;30:207–210. doi:10.1093/nar/30.1.207
- Ene IV, Farrer RA, Hirakawa MP, Agwamba K, Cuomo CA, et al. Global analysis of mutations driving microevolution of a heterozygous diploid fungal pathogen. Proc Natl Acad Sci U S A. 2018;115(35):8688–8697. doi:10.1073/pnas.1805234115
- Engler C, Kandzia R, Marillonnet S. A one pot, one step, precision cloning method with high throughput capability. PLoS One. 2008;3(11):e3647. doi:10.1371/journal.pone.0003647
- Enkler L, Richer D, Marchand AL, Ferrandon D, Jossinet F. Genome engineering in the yeast pathogen *Candida glabrata* using the CRISPR-Cas9 system. Sci Rep. 2016;6(1):35766. doi:10.1038/srep35766
- Ennis CL, Hernday AD, Nobile CJ. A markerless CRISPR-mediated system for genome editing in *Candida auris* reveals a conserved role for Cas5 in the caspofungin response. Microbiol Spectr. 2021;9(3):e0182021. doi:10.1128/Spectrum.01820-21
- Fan Y, Lin X. Multiple applications of a transient CRISPR-Cas9 coupled with electroporation (TRACE) system in the *Cryptococcus neoformans* species complex. Genetics. 2018;208(4):1357–1372. doi:10.1534/genetics.117.300656
- Fisher MC, Gurr SJ, Cuomo CA, Blehert DS, Jin H, et al. Threats posed by the fungal kingdom to humans, wildlife, and agriculture. mBio. 2020;11:e00449-20. doi:10.1128/mBio.00449-20
- Flores-Mireles AL, Walker JN, Bauman TM, Potretzke AM, Schreiber HL 4th, et al. Fibrinogen release and deposition on urinary catheters placed during urological procedures. J Urol. 2016;196(2):416–421. doi:10.1016/j.juro.2016.01.100
- Flowers SA, Barker KS, Berkow EL, Toner G, Chadwick SG, et al. Gain-of-function mutations in UPC2 are a frequent cause of ERG11 upregulation in azole-resistant clinical isolates of *Candida albicans*. Eukaryot Cell. 2012;11(10):1289–1299. doi:10.1128/EC.00215-12
- Forche A, Cromie G, Gerstein AC, Solis NV, Pisithkul T, et al. Rapid phenotypic and genotypic diversification after exposure to the oral host niche in *Candida albicans*. Genetics. 2018;209(3):725–741. doi:10.1534/genetics.118.301019
- Forche A, Magee PT, Selmecki A, Berman J, May G. Evolution in *Candida albicans* populations during a single passage through a mouse host. Genetics. 2009;182(3):799–811. doi:10.1534/genetics.109.103325
- Forche A, Solis NV, Swidergall M, Thomas R, Guyer A, et al. Selection of *Candida albicans* trisomy during oropharyngeal infection results

- in a commensal-like phenotype. *PLoS Genet.* 2019;15(5): e1008137. doi:10.1371/journal.pgen.1008137
- Ford CB, Funt JM, Abbey D, Issi L, Guiducci C, et al. The evolution of drug resistance in clinical isolates of *Candida albicans*. *eLife.* 2015;4:e00662. doi:10.7554/eLife.00662
- Fu Y, Ibrahim AS, Sheppard DC, Chen Y-C, French SW, et al. *Candida albicans* Als1p: an adhesin that is a downstream effector of the EFG1 filamentation pathway. *Mol Microbiol.* 2002;44(1):61–72. doi:10.1046/j.1365-2958.2002.02873.x
- Fu Y, Luo G, Spellberg BJ, Edwards JE Jr, Ibrahim AS. Gene overexpression/suppression analysis of candidate virulence factors of *Candida albicans*. *Eukaryot Cell.* 2008;7(3):483–492. doi:10.1128/EC.00445-07
- Fuller KK, Chen S, Loros JJ, Dunlap JC. Development of the CRISPR/Cas9 system for targeted gene disruption in *Aspergillus fumigatus*. *Eukaryot Cell.* 2015;14(11):1073–1080. doi:10.1128/EC.00107-15
- Fu C, Zhang X, Veri AO, Iyer KR, Lash E, et al. Leveraging machine learning essentiality predictions and chemogenomic interactions to identify antifungal targets. *Nat Commun.* 2021;12:6497. doi:10.1038/s41467-021-26850-3
- García-Béjar B, Arévalo-Villena M, Guisantes-Batan E, Rodríguez-Flores J, Briones A. Study of the bioremediatory capacity of wild yeasts. *Sci Rep.* 2020;10(1):11265. doi:10.1038/s41598-020-68154-4
- Gargouri B, Mhiri N, Karray F, Aloui F, Sayadi S. Isolation and characterization of hydrocarbon-degrading yeast strains from petroleum contaminated industrial wastewater. *Biomed Res Int.* 2015; 2015:929424. doi:10.1155/2015/929424
- Geddes-McAlister J, Shapiro RS. New pathogens, new tricks: emerging, drug-resistant fungal pathogens and future prospects for antifungal therapeutics. *Ann N Y Acad Sci.* 2018;1435(1):57–78. doi:10.1111/nyas.13739
- Gerami-Nejad M, Zacchi LF, McClellan M, Matter K, Berman J. Shuttle vectors for facile gap repair cloning and integration into a neutral locus in *Candida albicans*. *Microbiology.* 2013;159(Pt 3):565–579. doi:10.1099/mic.0.064097-0
- Gervais NC, Halder V, Shapiro RS. A data library of *Candida albicans* functional genomic screens. *FEMS Yeast Res.* 2021;21:foab060. doi:10.1093/femsyr/foab060
- Gilbert LA, Horlbeck MA, Adamson B, Villalta JE, Chen Y, et al. Genome-scale CRISPR-mediated control of gene repression and activation. *Cell.* 2014;159:647–661. doi:10.1016/j.cell.2014.09.029
- Gow NAR, van de Veerdonk FL, Brown AJP, Netea MG. *Candida albicans* morphogenesis and host defence: discriminating invasion from colonization. *Nat Rev Microbiol.* 2011;10:112–122. doi:10.1038/nrmicro2711
- Halder V, Porter CBM, Chavez A, Shapiro RS. Design, execution, and analysis of CRISPR–Cas9-based deletions and genetic interaction networks in the fungal pathogen *Candida albicans*. *Nat Protoc.* 2019;14(3):955–975. doi:10.1038/s41596-018-0122-6
- Hardwick JM, Tse L, Applegren N, Nicholas J, Veluona MA. The Epstein-Barr virus R transactivator (Rta) contains a complex, potent activation domain with properties different from those of VP16. *J Virol.* 1992;66(9):5500–5508. doi:10.1128/jvi.66.9.5500-5508.1992
- Homann OR, Dea J, Noble SM, Johnson D. A phenotypic profile of the *Candida albicans* regulatory network. *PLoS Genet.* 2009;5: e1000783. doi:10.1371/journal.pgen.1000783
- Horlbeck MA, Gilbert LA, Villalta JE, Adamson B, Pak RA, et al. Compact and highly active next-generation libraries for CRISPR-mediated gene repression and activation. *eLife.* 2016;5: e19760. doi:10.7554/eLife.19760
- Hossain S, Veri AO, Cowen LE. The proteasome governs fungal morphogenesis via functional connections with Hsp90 and cAMP-protein kinase a signaling. *mBio.* 2020;11:e00290-20. doi:10.1128/mBio.00290-20
- Hoyer LL, Cota E. *Candida albicans* agglutinin-like sequence (als) family vignettes: a review of als protein structure and function. *Front Microbiol.* 2016;7:280. doi:10.3389/fmicb.2016.00280
- Hoyer LL, Green CB, Oh S-H, Zhao X. Discovering the secrets of the *Candida albicans* agglutinin-like sequence (ALS) gene family—a sticky pursuit. *Med Mycol.* 2008;46(1):1–15. doi:10.1080/13693780701435317
- Hu JH, Miller SM, Geurts MH, Tang W, Chen L, et al. Evolved Cas9 variants with broad PAM compatibility and high DNA specificity. *Nature.* 2018;556(7699):57–63. doi:10.1038/nature26155
- Huang MY, Joshi MB, Boucher MJ, Lee S, Loza LC, et al. Short homology-directed repair using optimized Cas9 in the pathogen *Cryptococcus neoformans* enables rapid gene deletion and tagging. *Genetics.* 2021;220(1):iyab180. doi:10.1093/genetics/iyab180
- Huang MY, Mitchell AP. Marker recycling in *Candida albicans* through CRISPR-Cas9-induced marker excision. *mSphere.* 2017;2: e00050-17. doi:10.1128/mSphere.00050-17
- Jaitly P, Legrand M, Das A, Patel T, Chauvel M, et al. A phylogenetically-restricted essential cell cycle progression factor in the human pathogen *Candida albicans*. *Nat Commun.* 2022; 13(1):4256. doi:10.1038/s41467-022-31980-3
- Jensen MK. Design principles for nuclease-deficient CRISPR-based transcriptional regulators. *FEMS Yeast Res.* 2018;18(4):foy039. doi:10.1093/femsyr/foy039
- Jiang C, Lv G, Tu Y, Cheng X, Duan Y, et al. Applications of CRISPR/Cas9 in the synthesis of secondary metabolites in filamentous fungi. *Front Microbiol.* 2021;12:638096. doi:10.3389/fmicb.2021.638096
- Joo H-S, Ndegwa PM, Shoda M, Phae C-G. Bioremediation of oil-contaminated soil using *Candida catenulata* and food waste. *Environ Pollut.* 2008;156(3):891–896. doi:10.1016/j.envpol.2008.05.026
- Jost M, Chen Y, Gilbert LA, Horlbeck MA, Krenning L, et al. Combined CRISPRi/a-based chemical genetic screens reveal that rigosertib is a microtubule-destabilizing agent. *Mol Cell.* 2017;68:210–223.e6. doi:10.1016/j.molcel.2017.09.012
- Joung J, Engreitz JM, Konermann S, Abudayyeh OO, Verdine V, et al. Genome-scale activation screen identifies a lncRNA locus regulating a gene neighbourhood. *Nature.* 2017;548:343–346. doi:10.1038/nature23451
- Kamai Y, Kubota M, Kamai Y, Hosokawa T, Fukuoka T, et al. Contribution of *Candida albicans* ALS1 to the pathogenesis of experimental oropharyngeal candidiasis. *Infect Immun.* 2002; 70(9):5256–5258. doi:10.1128/IAI.70.9.5256-5258.2002
- Kampmann M. CRISPRi and CRISPRa screens in mammalian cells for precision biology and medicine. *ACS Chem Biol.* 2018;13(2): 406–416. doi:10.1021/acscchembio.7b00657
- Kieliszek M, Kot AM, Bzducha-Wróbel A, Błazejak S, Gientka I, et al. Biotechnological use of *Candida* yeasts in the food industry: a review. *Fungal Biol Rev.* 2017;31(4):185–198. doi:10.1016/j.fbr.2017.06.001
- Konermann S, Brigham MD, Trevino AE, Joung J, Abudayyeh OO, et al. Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature.* 2015;517:583–588. doi:10.1038/nature14136
- Kwon MJ, Schütze T, Spohner S, Haefner S, Meyer V. Practical guidance for the implementation of the CRISPR genome editing tool in filamentous fungi. *Fungal Biol Biotechnol.* 2019;6(1):15. doi:10.1186/s40694-019-0079-4
- La Bella AA, Andersen MJ, Gervais NC, Molina JJ, Molesan A, et al. Catheterized bladder environment induces *Candida albicans*

- hyphal formation and promotes colonization and persistence through Als1. *bioRxiv*. 2021. doi:10.1101/2021.06.01.446547, 9 May 2022, preprint: not peer reviewed.
- La Russa MF, Qi LS. The new state of the art: Cas9 for gene activation and repression. *Mol Cell Biol*. 2015;35:3800–3809. doi:10.1128/MCB.00512-15
- Lee JA, Robbins N, Xie JL, Ketela T, Cowen LE. Functional genomic analysis of *Candida albicans* adherence reveals a key role for the Arp2/3 complex in cell wall remodelling and biofilm formation. *PLoS Genet*. 2016;12:e1006452. doi:10.1371/journal.pgen.1006452
- Legrand M, Bachellier-Bassi S, Lee KK, Chaudhari Y, Tournu H, et al. Generating genomic platforms to study *Candida albicans* pathogenesis. *Nucleic Acids Res*. 2018;46(14):6935–6949. doi:10.1093/nar/gky594
- Li X, Yang F, Li D, Zhou M, Wang X, et al. Trisomy of chromosome R confers resistance to triazoles in *Candida albicans*. *Med Mycol*. 2015;53(3):302–309. doi:10.1093/mmy/myv002
- Li X, Zolli-Juran M, Cechetto JD, Daigle DM, Wright GD, et al. Multicopy suppressors for novel antibacterial compounds reveal targets and drug efflux susceptibility. *Chem Biol*. 2004;11(10):1423–1430. doi:10.1016/j.chembiol.2004.08.014
- Lohberger A, Coste AT, Sanglard D. Distinct roles of *Candida albicans* drug resistance transcription factors *TAC1*, *MRR1*, and *UPC2* in virulence. *Eukaryot Cell*. 2014;13(1):127–142. doi:10.1128/EC.00245-13
- Lohse MB, Ene IV, Craik VB, Hernday AD, Mancera E, et al. Systematic genetic screen for transcriptional regulators of the *Candida albicans* white-opaque switch. *Genetics*. 2016;203:1679–1692. doi:10.1534/genetics.116.190645
- Lombardi L, Oliveira-Pacheco J, Butler G. Plasmid-based CRISPR-Cas9 gene editing in multiple *Candida* species. *mSphere*. 2019;4(2):e00125–19. doi:10.1128/mSphere.00125-19
- Lombardi L, Turner SA, Zhao F, Butler G. Gene editing in clinical isolates of *Candida parapsilosis* using CRISPR/Cas9. *Sci Rep*. 2017;7(1):8051. doi:10.1038/s41598-017-08500-1
- Lu Z, Lin Z. The origin and evolution of a distinct mechanism of transcription initiation in yeasts. *Genome Res*. 2021;31(1):51–63. doi:10.1101/gr.264325.120
- Luesch H, Wu TYH, Ren P, Gray NS, Schultz PG, et al. A genome-wide overexpression screen in yeast for small-molecule target identification. *Chem Biol*. 2005;12(1):55–63. doi:10.1016/j.chembiol.2004.10.015
- MacAlpine J, Daniel-Ivadi M, Liu Z, Yano J, Revie NM, et al. A small molecule produced by *Lactobacillus* species blocks *Candida albicans* filamentation by inhibiting a DYRK1-family kinase. *Nat Commun*. 2021;12(1):6151. doi:10.1038/s41467-021-26390-w
- Manoharlal R, Gaur NA, Panwar SL, Morschhäuser J, Prasad R. Transcriptional activation and increased mRNA stability contribute to overexpression of *CDR1* in azole-resistant *Candida albicans*. *Antimicrob Agents Chemother*. 2008;52(4):1481–1492. doi:10.1128/AAC.01106-07
- Maroc L, Fairhead C. A new inducible CRISPR-Cas9 system useful for genome editing and study of double-strand break repair in *Candida glabrata*. *Yeast*. 2019;36(12):723–731. doi:10.1002/yea.3440
- Marr KA, Lyons CN, Ha K, Rustad TR, White TC. Inducible azole resistance associated with a heterogeneous phenotype in *Candida albicans*. *Antimicrob Agents Chemother*. 2001;45(1):52–59. doi:10.1128/AAC.45.1.52-59.2001
- Marr KA, Lyons CN, Rustad TR, Bowden RA, White TC. Rapid, transient fluconazole resistance in *Candida albicans* is associated with increased mRNA levels of *CDR*. *Antimicrob Agents Chemother*. 1998;42(10):2584–2589. doi:10.1128/AAC.42.10.2584
- McMillan J, Lu Z, Rodriguez JS, Ahn T-H, Lin Z. YeasTSS: an integrative web database of yeast transcription start sites. *Database*. 2019;2019:baz048. doi:10.1093/database/baz048
- Min K, Biermann A, Hogan DA, Konopka JB. Genetic analysis of NDT80 family transcription factors in *Candida albicans* using new CRISPR-Cas9 approaches. *mSphere*. 2018;3:e00545-18. doi:10.1128/mSphere.00545-18
- Min K, Ichikawa Y, Woolford CA, Mitchell AP. *Candida albicans* gene deletion with a transient CRISPR-Cas9 system. *mSphere*. 2016;1:e00130-16. doi:10.1128/mSphere.00130-16
- Momen-Roknabadi A, Oikonomou P, Zegans M, Tavazoie S. An inducible CRISPR interference library for genetic interrogation of *Saccharomyces cerevisiae* biology. *Commun Biol*. 2020;3(1):723. doi:10.1038/s42003-020-01452-9
- Morio F, Lombardi L, Butler G. The CRISPR toolbox in medical mycology: state of the art and perspectives. *PLoS Pathog*. 2020;16(1):e1008201. doi:10.1371/journal.ppat.1008201
- Motaung TE, Ells R, Pohl CH, Albertyn J, Tsilo TJ. Genome-wide functional analysis in *Candida albicans*. *Virulence*. 2017;8:1563–1579. doi:10.1080/21505594.2017.1292198
- Mount HO, Revie NM, Todd RT, Anstett K, Collins C, et al. Global analysis of genetic circuitry and adaptive mechanisms enabling resistance to the azole antifungal drugs. *PLoS Genet*. 2018;14:e1007319. doi:10.1371/journal.pgen.1007319
- Mózsik L, Hoekzema M, de Kok NAW, Bovenberg RAL, Nygård Y, et al. CRISPR-based transcriptional activation tool for silent genes in filamentous fungi. *Sci Rep*. 2021;11(1):1118. doi:10.1038/s41598-020-80864-3
- Muñoz IV, Sarrocco S, Malfatti L, Baroncelli R, Vannacci G. CRISPR-Cas for fungal genome editing: a new tool for the management of plant diseases. *Front Plant Sci*. 2019;10:135. doi:10.3389/fpls.2019.00135
- Ng H, Dean N. Dramatic improvement of CRISPR/Cas9 editing in *Candida albicans* by increased single guide RNA expression. *mSphere*. 2017;2:e00385-16. doi:10.1128/mSphere.00385-16
- Nguyen N, Quail MMF, Hernday AD. An efficient, rapid, and recyclable system for CRISPR-mediated genome editing in *Candida albicans*. *mSphere*. 2017;2:e00149-17. doi:10.1128/mSphereDirect.00149-17
- Niimi M, Niimi K, Takano Y, Holmes AR, Fischer FJ, et al. Regulated overexpression of *CDR1* in *Candida albicans* confers multidrug resistance. *J Antimicrob Chemother*. 2004;54(6):999–1006. doi:10.1093/jac/dkh456
- Noble SM, French S, Kohn LA, Chen V, Johnson AD. Systematic screens of a *Candida albicans* homozygous deletion library decouple morphogenetic switching and pathogenicity. *Nat Genet*. 2010;42:590–598. doi:10.1038/ng.605
- Nobile CJ, Fox EP, Nett JE, Sorrells TR, Mitrovich QM, et al. A recently evolved transcriptional network controls biofilm development in *Candida albicans*. *Cell*. 2012;148:126–138. doi:10.1016/j.cell.2011.10.048
- Nobile CJ, Schneider HA, Nett JE, Sheppard DC, Filler SG, et al. Complementary adhesin function in *C. albicans* biofilm formation. *Curr Biol*. 2008;18(14):1017–1024. doi:10.1016/j.cub.2008.06.034
- Nødvig CS, Nielsen JB, Kogle ME, Mortensen UH. A CRISPR-Cas9 system for genetic engineering of filamentous fungi. *PLoS One*. 2015;10(7):e0133085. doi:10.1371/journal.pone.0133085
- O'Meara TR, Veri AO, Ketela T, Jiang B, Roemer T, et al. Global analysis of fungal morphology exposes mechanisms of host cell escape. *Nat Commun*. 2015;6:6741.
- Quedraogo J-P, Tsang A. CRISPR Cas systems for fungal research. *Fungal Biol Rev*. 2020;34(4):189–201. doi:10.1016/j.fbr.2020.10.002

- Park Y-N, Morschhäuser J. Tetracycline-inducible gene expression and gene deletion in *Candida albicans*. *Eukaryot Cell*. 2005;4(8):1328–1342. doi:10.1128/EC.4.8.1328-1342.2005
- Payen C, Thompson D. The renaissance of yeasts as microbial factories in the modern age of biomanufacturing. *Yeast*. 2019;36(12):685–700. doi:10.1002/yea.3439
- Peng D, Tarleton R. EuPaGDT: a web tool tailored to design CRISPR guide RNAs for eukaryotic pathogens. *Microb Genom*. 2015;1:e000033. doi:10.1099/mgen.0.000033
- Pérez JC. *Candida albicans* dwelling in the mammalian gut. *Curr Opin Microbiol*. 2019;52:41–46. doi:10.1016/j.mib.2019.04.007
- Perfect JR. The antifungal pipeline: a reality check. *Nat Rev Drug Discov*. 2017;16(9):603–616. doi:10.1038/nrd.2017.46
- Peters JM, Silvis MR, Zhao D, Hawkins JS, Gross CA, et al. Bacterial CRISPR: accomplishments and prospects. *Curr Opin Microbiol*. 2015;27:121–126. doi:10.1016/j.mib.2015.08.007
- Polvi EJ, Veri AO, Liu Z, Hossain S, Hyde S, et al. Functional divergence of a global regulatory complex governing fungal filamentation. *PLoS Genet*. 2019;15(1):e1007901. doi:10.1371/journal.pgen.1007901
- Prasad R, Banerjee A, Khandelwal NK, Dhamgaye S. The ABCs of *Candida albicans* multidrug transporter Cdr1. *Eukaryot Cell*. 2015;14(12):1154–1164. doi:10.1128/EC.00137-15
- Prasad R, De Wergifosse P, Goffeau A, Balzi E. Molecular cloning and characterization of a novel gene of *Candida albicans*, CDR1, conferring multiple resistance to drugs and antifungals. *Curr Genet*. 1995;27(4):320–329. doi:10.1007/BF00352101
- Prieto D, Román E, Alonso-Monge R, Pla J. Overexpression of the transcriptional regulator *WOR1* increases susceptibility to bile salts and adhesion to the mouse gut mucosa in *Candida albicans*. *Front Cell Infect Microbiol*. 2017;7:389. doi:10.3389/fcimb.2017.00389
- Radzishuskaya A, Shlyueva D, Müller I, Helin K. Optimizing sgRNA position markedly improves the efficiency of CRISPR/dCas9-mediated transcriptional repression. *Nucleic Acids Res*. 2016;44(18):e141. doi:10.1093/nar/gkw583
- Rai LS, van Wijlick L, Chauvel M, d'Enfert C, Legrand M, et al. Overexpression approaches to advance understanding of *Candida albicans*. *Mol Microbiol*. 2021;117(3):589–599. doi:10.1111/mmi.14818
- Razzaq I, Berg MD, Jiang Y, Genereaux J, Uthayakumar D, et al. The SAGA and NuA4 component Tra1 regulates *Candida albicans* drug resistance and pathogenesis. *Genetics*. 2021;219(2):iyab131. doi:10.1093/genetics/iyab131
- Roemer T, Jiang B, Davison J, Ketela T, Veillette K, et al. Large-scale essential gene identification in *Candida albicans* and applications to antifungal drug discovery. *Mol Microbiol*. 2003;50:167–181. doi:10.1046/j.1365-2958.2003.03697.x
- Román E, Coman I, Prieto D, Alonso-Monge R, Pla J. Implementation of a CRISPR-based system for gene regulation in *Candida albicans*. *mSphere*. 2019;4:e00001-19. doi:10.1128/mSphere.00001-19
- Román E, Prieto D, Alonso-Monge R, Pla J. New insights of CRISPR technology in human pathogenic fungi. *Future Microbiol*. 2019;14:1243–1255. doi:10.2217/fmb-2019-0183
- Rosiana S, Zhang L, Kim GH, Revtovich AV, Uthayakumar D, et al. Comprehensive genetic analysis of adhesin proteins and their role in virulence of *Candida albicans*. *Genetics*. 2021;217(2):iyab003. doi:10.1093/genetics/iyab003
- Roux I, Woodcraft C, Hu J, Wolters R, Gilchrist CLM, et al. CRISPR-mediated activation of biosynthetic gene clusters for bioactive molecule discovery in filamentous fungi. *ACS Synth Biol*. 2020;9(7):1843–1854. doi:10.1021/acssynbio.0c00197
- Ruben S, Garbe E, Mogavero S, Albrecht-Eckardt D, Hellwig D, et al. *Ahr1* and *Tup1* contribute to the transcriptional control of virulence-associated genes in *Candida albicans*. *mBio*. 2020;11(2):e00206–20. doi:10.1128/mBio.00206-20
- Ryan O, Shapiro RS, Kurat CF, Mayhew D, Baryshnikova A, et al. Global gene deletion analysis exploring yeast filamentous growth. *Science*. 2012;337:1353–1356. doi:10.1126/science.1224339
- Sahni N, Yi S, Daniels KJ, Huang G, Srikantha T, et al. *Tec1* mediates the pheromone response of the white phenotype of *Candida albicans*: insights into the evolution of new signal transduction pathways. *PLoS Biol*. 2010;8(5):e1000363. doi:10.1371/journal.pbio.1000363
- Sanglard D, Coste A, Ferrari S. Antifungal drug resistance mechanisms in fungal pathogens from the perspective of transcriptional gene regulation. *FEMS Yeast Res*. 2009;9(7):1029–1050. doi:10.1111/j.1567-1364.2009.00578.x
- Sanglard D, Kuchler K, Ischer F, Pagani JL, Monod M, et al. Mechanisms of resistance to azole antifungal agents in *Candida albicans* isolates from AIDS patients involve specific multidrug transporters. *Antimicrob Agents Chemother*. 1995;39(11):2378–2386. doi:10.1128/AAC.39.11.2378
- Sanguinetti M, Posteraro B, Lass-Flörl C. Antifungal drug resistance among *Candida* species: mechanisms and clinical impact. *Mycoses*. 2015;58(2):2–13. doi:10.1111/myc.12330
- Sanjana NE. Genome-scale CRISPR pooled screens. *Anal Biochem*. 2017;532:95–99. doi:10.1016/j.ab.2016.05.014
- Sanson KR, Hanna RE, Hegde M, Donovan KF, Strand C, et al. Optimized libraries for CRISPR-Cas9 genetic screens with multiple modalities. *Nat Commun*. 2018;9(1):5416. doi:10.1038/s41467-018-07901-8
- Santana DJ, O'Meara TR. Forward and reverse genetic dissection of morphogenesis identifies filament-competent *Candida auris* strains. *Nat Commun*. 2021;12(1):7197. doi:10.1038/s41467-021-27545-5
- Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc*. 2008;3(6):1101–1108. doi:10.1038/nprot.2008.73
- Schmitz ML, Baeuerle PA. The p53 subunit is responsible for the strong transcription activating potential of NF-kappa B. *EMBO J*. 1991;10(12):3805–3817. doi:10.1002/j.1460-2075.1991.tb04950.x
- Schubert S, Barker KS, Znaidi S, Schneider S, Dierolf F, et al. Regulation of efflux pump expression and drug resistance by the transcription factors *Mrr1*, *Upc2*, and *Cap1* in *Candida albicans*. *Antimicrob Agents Chemother*. 2011;55(5):2212–2223. doi:10.1128/AAC.01343-10
- Schultenkämper K, Brito LF, Wendisch VF. Impact of CRISPR interference on strain development in biotechnology. *Biotechnol Appl Biochem*. 2020;67(1):7–21. doi:10.1002/bab.1901
- Schuster M, Kahmann R. CRISPR-Cas9 genome editing approaches in filamentous fungi and oomycetes. *Fungal Genet Biol*. 2019;130:43–53. doi:10.1016/j.fgb.2019.04.016
- Segrelles-Calvo G, de S Araújo GR, Llopis-Pastor E, Carrillo J, Hernández-Hernández M, et al. *Candida* spp. Co-infection in COVID-19 patients with severe pneumonia: prevalence study and associated risk factors. *Respir Med*. 2021;188:106619. doi:10.1016/j.rmed.2021.106619
- Selmecki A, Forche A, Berman J. Aneuploidy and isochromosome formation in drug-resistant *Candida albicans*. *Science*. 2006;313(5785):367–370. doi:10.1126/science.1128242
- Selmecki A, Gerami-Nejad M, Paulson C, Forche A, Berman J. An isochromosome confers drug resistance in vivo by amplification of two genes, *ERG11* and *TAC1*. *Mol Microbiol*. 2008;68(3):624–641. doi:10.1111/j.1365-2958.2008.06176.x

- Shan L, Dai Z, Wang Q. Advances and opportunities of CRISPR/Cas technology in bioengineering non-conventional yeasts. *Front Bioeng Biotechnol.* 2021;9:765396. doi:10.3389/fbioe.2021.765396
- Shapiro RS, Chavez A, Collins JJ. CRISPR-based genomic tools for the manipulation of genetically intractable microorganisms. *Nat Rev Microbiol.* 2018;16(6):333–339. doi:10.1038/s41579-018-0002-7
- Shapiro RS, Chavez A, Porter CBM, Hamblin M, Kaas CS, et al. A CRISPR-Cas9-based gene drive platform for genetic interaction analysis in *Candida albicans*. *Nat Microbiol.* 2018;3(1):73–82. doi:10.1038/s41564-017-0043-0
- Sharma J, Rosiana S, Razzaq I, Shapiro RS. Linking cellular morphogenesis with antifungal treatment and susceptibility in *Candida* pathogens. *J Fungi.* 2019;5(1):17. doi:10.3390/jof5010017
- Smith AM, Ammar R, Nislow C, Giaever G. A survey of yeast genomic assays for drug and target discovery. *Pharmacol Ther.* 2010;127(2):156–164. doi:10.1016/j.pharmthera.2010.04.012
- Smith JD, Suresh S, Schlecht U, Wu M, Wagih O, et al. Quantitative CRISPR interference screens in yeast identify chemical-genetic interactions and new rules for guide RNA design. *Genome Biol.* 2016;17(1):45. doi:10.1186/s13059-016-0900-9
- Sopko R, Huang D, Preston N, Chua G, Papp B, et al. Mapping pathways and phenotypes by systematic gene overexpression. *Mol Cell.* 2006;21(3):319–330. doi:10.1016/j.molcel.2005.12.011
- Todd RT, Selmecki A. Expandable and reversible copy number amplification drives rapid adaptation to antifungal drugs. *eLife.* 2020;9:e58349. doi:10.7554/eLife.58349
- Todd RT, Wikoff TD, Forche A, Selmecki A. Genome plasticity in *Candida albicans* is driven by long repeat sequences. *eLife.* 2019;8:e45954. doi:10.7554/eLife.45954
- Todor H, Silvis MR, Osadnik H, Gross CA. Bacterial CRISPR screens for gene function. *Curr Opin Microbiol.* 2021;59:102–109. doi:10.1016/j.mib.2020.11.005
- Uthayakumar D, Sharma J, Wensing L, Shapiro RS. CRISPR-based genetic manipulation of *Candida* species: historical perspectives and current approaches. *Front Genome Ed.* 2021;2:606281. doi:10.3389/fgeed.2020.606281
- van Rhijn N, Furukawa T, Zhao C, McCann BL, Bignell E, et al. Development of a marker-free mutagenesis system using CRISPR-Cas9 in the pathogenic mould *Aspergillus fumigatus*. *Fungal Genet Biol.* 2020;145:103479. doi:10.1016/j.fgb.2020.103479
- Vyas VK, Barrasa MI, Fink GR. A *Candida albicans* CRISPR system permits genetic engineering of essential genes and gene families. *Sci Adv.* 2015;1(3):e1500248. doi:10.1126/sciadv.1500248
- Vyas VK, Guy Bushkin G, Bernstein DA, Getz MA, Sewastianik M, et al. New CRISPR mutagenesis strategies reveal variation in repair mechanisms among fungi. *mSphere.* 2018;3:e00154-18.
- Wang P. Two distinct approaches for CRISPR-Cas9-mediated gene editing in *Cryptococcus neoformans* and related species. *mSphere.* 2018;3:e00208-18. doi:10.1128/mSphereDirect.00208-18
- Wang H, La Russa M, Qi LS. CRISPR/Cas9 in genome editing and beyond. *Annu Rev Biochem.* 2016;85(1):227–264. doi:10.1146/annurev-biochem-060815-014607
- Wensing L, Shapiro RS. Design and generation of a CRISPR interference system for genetic repression and essential gene analysis in the fungal pathogen *Candida albicans*. *Methods Mol Biol.* 2022;2377:69–88. doi:10.1007/978-1-0716-1720-5\_4
- Wensing L, Sharma J, Uthayakumar D, Proteau Y, Chavez A, et al. A CRISPR interference platform for efficient genetic repression in *Candida albicans*. *mSphere.* 2019;4(1):e00002-19. doi:10.1128/mSphere.00002-19
- White TC. Increased mRNA levels of ERG16, CDR, and MDR1 correlate with increases in azole resistance in *Candida albicans* isolates from a patient infected with human immunodeficiency virus. *Antimicrob Agents Chemother.* 1997;41(7):1482–1487. doi:10.1128/AAC.41.7.1482
- Wilson FM, Harrison RJ. CRISPR/Cas9 mediated editing of the Quorn fungus *Fusarium venenatum* A3/5 by transient expression of Cas9 and sgRNAs targeting endogenous marker gene PKS12. *Fungal Biol Biotechnol.* 2021;8(1):15. doi:10.1186/s40694-021-00121-8
- Xu X, Qi LS. A CRISPR-dCas toolbox for genetic engineering and synthetic biology. *J Mol Biol.* 2019;431(1):34–47. doi:10.1016/j.jmb.2018.06.037
- Yang F, Teoh F, Tan ASM, Cao Y, Pavelka N, et al. Aneuploidy enables cross-adaptation to unrelated drugs. *Mol Biol Evol.* 2019;36(8):1768–1782. doi:10.1093/molbev/msz104
- Yang F, Todd RT, Selmecki A, Jiang Y-Y, Cao Y-B, et al. The fitness costs and benefits of trisomy of each *Candida albicans* chromosome. *Genetics.* 2021;218(2):iyab056. doi:10.1093/genetics/iyab056
- Zhang C, Meng X, Wei X, Lu L. Highly efficient CRISPR mutagenesis by microhomology-mediated end joining in *Aspergillus fumigatus*. *Fungal Genet Biol.* 2016;86:47–57. doi:10.1016/j.fgb.2015.12.007
- Zhang X-H, Tee LY, Wang X-G, Huang Q-S, Yang S-H. Off-target effects in CRISPR/Cas9-mediated genome engineering. *Mol Ther Nucleic Acids.* 2015;4:e264. doi:10.1038/mtna.2015.37
- Zhao X, Oh S-H, Cheng G, Green CB, Nuessen JA, et al. ALS3 And ALS8 represent a single locus that encodes a *Candida albicans* adhesin; functional comparisons between Als3p and Als1p. *Microbiology.* 2004;150(7):2415–2428. doi:10.1099/mic.0.26943-0
- Zhao X-Q, Zhang X-Y, Zhang F, Zhang R, Jiang B-J, et al. 2018. Metabolic engineering of fungal strains for efficient production of cellulolytic enzymes. In: Fang X, Qu Y, editors. *Fungal Cellulolytic Enzymes: Microbial Production and Application*. Singapore: Springer. p. 27–41.
- Znaidi S, van Wijlick L, Hernández-Cervantes A, Sertour N, Desseyn J-L, et al. Systematic gene overexpression in *Candida albicans* identifies a regulator of early adaptation to the mammalian gut. *Cell Microbiol.* 2018;20(11):e12890. doi:10.1111/cmi.12890
- Zoppo M, Luca MD, Villarreal SN, Poma N, Barrasa MI, et al. A CRISPR/Cas9-based strategy to simultaneously inactivate the entire ALS gene family in *Candida orthopsilosis*. *Future Microbiol.* 2019;14(16):1383–1396. doi:10.2217/fmb-2019-0168