



# HHS Public Access

Author manuscript

*Ann Hum Biol.* Author manuscript; available in PMC 2023 February 10.

Published in final edited form as:

*Ann Hum Biol.* 2021 June ; 48(4): 346–349. doi:10.1080/03014460.2021.1951351.

## Body size and composition of Samoan toddlers aged 18–25 months in 2019

Sakurako Oyama<sup>1,2</sup>, Rachel L. Duckham<sup>3,4,5</sup>, Kendall J. Arslanian<sup>6</sup>, Erin E. Kershaw<sup>7</sup>, Joshua A. Strayer<sup>7</sup>, Ulai T. Fidow<sup>8</sup>, Take Naseri<sup>9</sup>, Nicola L. Hawley<sup>6</sup>

<sup>1</sup>Department of Anthropology, Yale University, New Haven, CT

<sup>2</sup>Yale School of Medicine, Yale University, New Haven, CT

<sup>3</sup>Institute of Physical Activity and Nutrition, Deakin University, Geelong, Australia

<sup>4</sup>Australian Institute for Musculoskeletal Science (AIMSS), The University of Melbourne and Western Health, St Albans, Victoria

<sup>5</sup>Department of General Practice, Monash University, Melbourne, AUS

<sup>6</sup>Department of Chronic Disease Epidemiology, Yale School of Public Health, New Haven, CT

<sup>7</sup>Division of Endocrinology, Department of Medicine, University of Pittsburgh, Pittsburgh, PA

<sup>8</sup>Department of Obstetrics & Gynecology, Tupua Tamasese Meaole Hospital, Samoa National Health Services, Apia, Samoa

<sup>9</sup>Ministry of Health, Apia, Samoa

### Abstract

**Background:** The “*Foafoaga O le Ola* (Beginning of Life)” study is a prospective birth cohort of n=160 Samoan mother-infant dyads established in 2017–2018. A primary study aim is to explore how a missense variant at *CREBRF*<sub>rs373863828</sub> impacts growth in early life, given its association with increased body size but lower risk of diabetes in adult Samoans. Here, we examine body size and composition by genotype among toddlers aged 18.7–24.5 months.

**Methods:** Height, weight, head circumference, mid-upper-arm circumference, and abdominal circumference, as well as subscapular, triceps, iliac crest and thigh skinfold thickness were measured among 107 toddlers with known *rs373863828* genotype; 42 of these toddlers completed dual-energy X-ray absorptiometry (DXA) scans from which body composition (total body less head fat mass, lean mass, bone mass, % fat mass and % fat-free mass) was estimated.

**Results:** After controlling for sex and age, toddlers with at least one copy of the *CREBRF* minor allele (AA/AG) were 1.31cm taller (SE=0.64, p=0.045) than toddlers with the GG genotype.

---

**Corresponding Author:** Dr. Nicola Hawley, Yale School of Public Health, 60 College Street, New Haven, CT 06520, Tel: +1 203 737 7176; nicola.hawley@yale.edu.

**Conflict of Interest Disclosure:** The authors have no conflicts of interest to disclose.

Access to the dataset

Requests to access the dataset for the purpose of comparative studies are encouraged and should be addressed to the corresponding author.

**Conclusion:** Whether greater linear growth in early childhood could contribute to the metabolically protective effects associated with the *CREBRF* variant in adulthood should be explored in future studies.

### Keywords

dual-energy X-ray absorptiometry; body composition; toddlers; Pacific Islands

---

## Background

Across the Pacific Islands, obesity prevalence has increased dramatically over the past several decades (Hawley & McGarvey 2015). A genome-wide association study in Samoan adults identified a missense variant in CREB3 regulatory factor (*CREBRF*) rs373863828, associated with increased BMI but decreased odds of type 2 diabetes mellitus (T2DM). Little is known about how the variant influences body composition and growth during critical periods of early development. Recently, Arslanian et al. (2020) reported no association between *CREBRF* genotype and birthweight among Samoan infants. However, the variant was associated with greater lean and bone mass at age 4 months. Here, we examine body size and composition by rs373863828 genotype among Samoan toddlers at age 18.7–24.5 months.

## Sample

Participants belong to a prospective birth cohort of n=160 Samoan mother-infant dyads (Arslanian et al. 2020). Mothers were recruited at 35–40 weeks gestation, and dyads participated in assessments at four time points: immediately after birth (0–41 days) and at approximately 2 (1.8–3.4), 4 (3.6–6.5) and 21 (18.7–24.5) months. *CREBRF* genotyped toddlers (n=107) who participated in the 21-month assessment are included in this analysis. Mothers gave written informed consent for participation. Protocols were approved by the Yale University Institutional Review Board (HIC #2000021076) and the Health Research Committee of the Samoa Ministry of Health.

## Data Collection

Toddler height was measured to the nearest 0.1cm using a portable GPM anthropometer (Pfister Imports, USA) and weight to the nearest 0.1kg using a digital weight scale (HD351, Tanita Corporation of America, USA). Left subscapular, triceps, iliac crest, and thigh skinfold thickness were measured using a Harpenden caliper (FG1056, UK). The sum of skinfolds was calculated as a proxy for subcutaneous fat. Head, left mid-upper-arm (MUAC), and abdominal circumference were measured using a standard tape measure (201, SECA, Germany). Abdominal circumference-to-height ratio was calculated as a proxy for abdominal visceral fat. All measurements were collected in duplicate and averaged for analyses.

Dual x-ray absorptiometry (DXA) scans were performed on toddlers wearing clean diapers and standardized clothing (Lunar iDXA, version enCORE 17, GE Healthcare Medicine, USA). Scans were aborted if movement artefacts interfered with scan readability. While

obtaining DXA images of neonates is relatively straightforward, as they develop motor skills, avoiding movement artefacts becomes difficult until old enough to reliably follow directions. Thus, only 42 of 107 toddlers completed successful DXA scans. DXA-derived body composition outcomes were absolute total body less head (TBLH) fat, lean and bone mass, as well as percent (%) fat and fat-free (lean + bone) mass.

## Data Management and Statistical Analysis

Age- and sex- standardized z-scores were calculated using World Health Organization Child Growth Standards (WHO Multicentre Growth Reference Study Group, 2006). Depending on normality, independent two-sample *t*-tests or Mann-Whitney *U* tests were used to examine differences in anthropometric characteristics by DXA completion status. Multivariable linear regression was used to test whether *CREBRF* genotype was associated with body size and composition, controlling for age and sex. Non-parametric measures were normalized using box-cox transformations. Given the expected small sample size of toddlers with AA genotype, and because both AG and AA genotypes are associated with increased body size in Samoan adults (Minster et al. 2016; Carlson et al. 2020), AG and AA genotypes were pooled for all analyses, providing a comparison of toddlers with no copies of the variant (GG) versus those with at least one copy (AA or AG) (Arslanian *et al.*, 2020). All analyses were conducted in RStudio version 1.2.1335 ([www.rstudio.com](http://www.rstudio.com)). Reported probabilities are two-sided and the threshold for statistical significance was set at  $\alpha=0.05$ .

## Results

On average, Samoan toddlers were 0.82 standard deviations shorter and 0.56 standard deviations heavier than WHO standards (table 1). After adjustment for sex and age, multivariable linear regression revealed the *CREBRF* variant was associated with 1.31cm greater height (SE=0.64,  $p=0.045$ ) and 1.23mm lower thigh skinfold thickness (SE=0.49,  $p=0.014$ ). Other anthropometric measures did not vary significantly by genotype (table 2).

DXA-scanned toddlers were 0.83 months younger (95% CI[0.26–1.45],  $p=0.006$ ), and were consequently 1.85cm shorter (95% CI[0.62–3.09],  $p=0.004$ ), and 0.60kg lighter (95% CI[0.05–1.20],  $p=0.04$ ) than unscanned toddlers. Scanned toddlers also had 0.90mm thinner triceps skinfolds (95% CI[0.20–1.55],  $p=0.007$ ), 0.95mm thinner iliac crest skinfolds (95% CI[0.25–1.70],  $p=0.005$ ), 1.20mm thinner thigh skinfolds (95% CI[0.19–2.21],  $p=0.02$ ), and thereby exhibited 3.65mm less subcutaneous fat (95% CI[1.35–5.70],  $p=0.002$ ) compared to unscanned toddlers. Differences in skinfold thickness between scanned and unscanned toddlers remained significant after controlling for age and sex. DXA-estimated body composition measures did not vary significantly by genotype (table 2).

## Comments

Among Samoan toddlers aged 18.7–24.5 months, having at least one copy of the *CREBRF* rs373863828 variant was associated with greater height. Similar findings have been documented in both Samoan (Carlson et al. 2020) and other Polynesian adults (Metcalf et al. 2020), and Samoan children aged 5–18 years (Carlson et al. 2020). Among children in

New Zealand, the *CREBRF* variant was associated with increased height at 4 years, but not at 2 years (Berry et al. 2018). However, Major et al. (2018) previously called in to question the validity of their findings, given the unexpectedly high rs37386328 A allele frequencies reported among the European (1.5%) and Asian (1.1%) samples. Further, the European rs37386328 genotype frequencies reported by Berry et al. deviate from Hardy-Weinberg equilibrium, suggesting genotyping errors or population groupings which do not accurately reflect genetic ancestry (Major et al. 2018). The pooling of all ethnicities prior to testing for association with body size further compounds these issues. If their study results are indeed valid, our results suggest that height differences by rs37386328 genotype may emerge earlier among Samoans living in Samoa, potentially due to gene-environment interactions. The association between rs37386328 variant and taller stature may appear at different ages according to the nutritional context encountered during development. The *CREBRF* variant was also associated with decreased thigh skinfold thickness in our sample. However, the biological significance of genotype differences in skinfold thickness at one site alone is unclear.

While toddlers with the minor allele were taller, the *CREBRF* variant was not significantly associated with greater bone mass. In contrast, the minor allele was associated with greater lean and bone mass in this cohort at age 4 months (Arslanian et al. 2020). It is possible that genotype differences in body composition are transient in infancy. However, this discrepancy is most likely due to the small sample size of DXA-scanned toddlers (n=42 versus n=110 at 4 months), the primary limitation of this study. Further, successful DXA scan completion was associated with lower age, weight, height and skinfold thicknesses. This is likely because younger and smaller toddlers were more likely to remain still during the DXA scan. Elucidating the relationship between rs37386328 genotype and body composition in toddlers and young children will require studies with larger sample sizes. Further, longitudinal study designs are required to determine whether the *CREBRF* variant promotes accelerated growth throughout childhood or only during specific developmental windows, and whether greater linear growth helps mediate the metabolically protective effects associated with the variant in adulthood. Future studies should also aim to refine our understanding of how variation in nutritional conditions and infectious disease loads interact with the rs37386328 genotype.

## Acknowledgements

The authors are incredibly grateful to the families who participated in this study. In addition, they thank the Samoan Ministry of Health for their support of this study. Finally, the authors thank the Samoan field team for their critical assistance in implementing the study protocol.

## References

- Arslanian KJ, Fidow UT, Atanoa T, Unasa-Apelu F, Naseri T, Wetzel AI, Pomer A, Duckham RL, McGarvey ST, Strayer JA, et al. 2021. A missense variant in *CREBRF*, rs37386328, is associated with fat-free mass, not fat mass in Samoan infants. *Int J Obes.* 45:45–55.
- Berry SD, Walker CG, Ly K, Snell RG, Atatoa Carr PE, Bandara D, Mohal J, Castro TG, Marks EJ, Morton SMB, Grant CC. 2018. Widespread prevalence of a *CREBRF* variant amongst M ori and Pacific children is associated with weight and height in early childhood. *Int J Obes.* 42(4):603–607.

- Carlson JC, Rosenthal SL, Russell EM, Hawley NL, Sun G, Cheng H, Naseri T, Reupena MS, Tuitele J, Deka R, et al. 2020. A missense variant in CREBRF is associated with taller stature in Samoans. *Am J Hum Biol.* 32(6):e23414. [PubMed: 32190945]
- Hawley NL, McGarvey ST. 2015. Obesity and Diabetes in Pacific Islanders: the Current Burden and the Need for Urgent Action. *Curr Diab Rep.* 15(5):1–10.
- Major TJ, Krishnan M, Topless RK, Dewes O, Thompson J, de Zoysa J, Stamp LK, Dalbeth N, Deka R, Weeks DE, et al. 2018. Re: “Widespread prevalence of a CREBRF variant among M ori and Pacific children is associated with weight and height in early childhood.” *Int J Obes.* 42(7):1389–1391.
- Metcalfe LK, Krishnan M, Turner N, Yaghootkar H, Merry TL, Dewes O, Hindmarsh JH, De Zoysa J, Dalbeth N, Stamp LK, et al. 2020. The M ori and Pacific specific CREBRF variant and adult height. *Int J Obes.* 44(3):748–752.
- Minster RL, Hawley NL, Su C, Sun G, Kershaw EE, Cheng H, Buhule OD, Lin J, Tuitele J, Naseri T, et al. 2016. A thrifty variant in CREBRF strongly influences body mass index in Samoans. *Nat Genet.* 48(9):1049–1054. [PubMed: 27455349]
- WHO Multicentre Growth Reference Study Group. WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development. 2006. Geneva. <http://www.who.int/childgrowth/en/>.

**Table 1:**

Toddler age and z-scores by sex at 18–25 months

	Girls		Boys		All	
	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n
<b>Age (months)</b>	21.35 (1.43)	49	21.27 (1.58)	58	21.31 (1.51)	107
<b>Height-for-age z-score</b>	−0.64 (1.08)	48	−0.97 (1.11)	58	−0.82 (1.11)	106
<b>Weight-for-age z-score</b>	0.70 (1.02)	48	0.44 (1.00)	58	0.56 (1.02)	106
<b>Weight-for-height z-score</b>	1.34 (1.02)	48	1.23 (0.92)	58	1.28 (0.96)	106
<b>BMI-for-age z-score</b>	1.51 (1.04)	48	1.45 (0.92)	58	1.48 (0.97)	106
<b>Head circumference-for-age z-score</b>	0.80 (1.18)	49	0.38 (1.29)	58	0.57 (1.25)	107
<b>MUAC-for-age z-score</b>	1.14 (1.00)	49	1.20 (1.20)	58	1.17 (1.11)	107
<b>Subscapular skinfold-for-age z-score</b>	1.29 (0.93)	47	1.09 (0.98)	58	1.18 (0.96)	105
<b>Triceps skinfold-for-age z-score</b>	0.67 (1.05)	48	0.38 (0.95)	58	0.51 (1.00)	106

**Table 2:**

Toddler body size and composition by sex and genotype at 18–25 months

	Girls						Boys						All GG		All AG/AA	
	All girls		GG		AG/AA		All boys		GG		AG/AA		mean (SD)	n	mean (SD)	n
	mean (SD)	n	mean (SD)	n	mean (SD)	n	mean (SD)	n	mean (SD)	n	mean (SD)	n				
<b>Height (cm)</b>	82.0 (3.4)	48	81.8 (3.3)	31	82.4 (3.7)	17	82.5 (3.5)	58	81.9 (3.6)	33	83.3 (3.2)	25	81.9 (3.4)	64	82.9 (3.4)	42
<b>Weight (kg)</b>	12.0 (1.6)	48	11.9 (1.2)	31	12.3 (2.2)	17	12.3 (1.4)	58	12.2 (1.2)	33	12.4 (1.6)	25	12.1 (1.2)	64	12.3 (1.8)	42
<b>BMI (kg/m<sup>2</sup>)</b>	17.9 (1.6)	48	17.8 (1.4)	31	18.0 (2.0)	17	18.0 (1.4)	58	18.1 (1.1)	33	17.8 (1.7)	25	18.0 (1.3)	64	17.9 (1.8)	42
<b>Head circumference (cm)</b>	47.9 (1.7)	49	47.9 (1.7)	31	48.0 (1.6)	18	48.4 (1.8)	58	48.5 (1.8)	33	48.3 (1.8)	25	48.2 (1.8)	64	48.2 (1.7)	43
<b>MUAC (cm)</b>	16.2 (1.4)	49	16.0 (1.3)	31	16.5 (1.5)	18	16.5 (1.6)	58	16.7 (1.6)	33	16.3 (1.2)	25	16.4 (1.5)	64	16.4 (1.6)	43
<b>Abdominal circumference (cm)</b>	49.9 (3.8)	49	49.4 (3.3)	31	50.8 (4.5)	18	49.1 (3.3)	58	49.0 (2.9)	33	49.2 (3.9)	25	49.2 (3.0)	64	49.9 (4.2)	43
<b>Abdominal circumference-to-height ratio</b>	.61 (.05)	48	.61 (.05)	31	.61 (.05)	17	.60 (.04)	58	.60 (.04)	33	.59 (.05)	25	.60 (.04)	64	.60 (.05)	42
<b>Subscapular skinfold thickness (mm)</b>	8.3 (2.0)	47	8.0 (1.6)	29	8.7 (2.6)	18	7.6 (1.5)	58	7.7 (1.4)	33	7.5 (1.6)	25	7.8 (1.5)	62	8.0 (2.1)	43
<b>Triceps skinfold thickness (mm)</b>	9.2 (2.1)	48	8.9 (1.7)	30	9.6 (2.7)	18	8.8 (6.0)	58	8.6 (1.5)	33	8.2 (1.6)	25	8.8 (1.6)	63	8.8 (2.2)	43
<b>Iliac crest skinfold thickness (mm)</b>	6.8 (2.5)	47	6.5 (2.1)	29	7.5 (3.0)	18	6.2 (1.9)	58	6.4 (2.1)	33	6.0 (1.5)	25	6.4 (2.1)	62	6.6 (2.3)	43
<b>Thigh skinfold thickness (mm)</b>	13.8 (2.8)	47	14.4 (2.8)	29	12.8 (2.7)	18	13.4 (2.3)	58	13.8 (2.0)	33	12.8 (2.6)	25	14.0 (2.4)	62	12.8 (2.6)	43
<b>Subcutaneous fat (mm)</b>	38.0 (7.1)	47	37.7 (6.4)	29	38.5 (8.3)	18	35.7 (4.9)	58	36.5 (4.8)	33	34.6 (4.9)	25	37.1 (5.6)	62	36.2 (6.7)	43
<b>TBLH fat mass (g)</b>	2788.16 (529.09)	19	2793.42 (484.53)	12	2779.14 (639.72)	7	2717.13 (455.59)	23	2757.07 (354.99)	15	2645.12 (625.27)	8	2773.22 (409.29)	27	2707.67 (612.91)	15
<b>TBLH lean mass (g)</b>	6342.16 (734.96)	19	6341.92 (596.49)	12	6342.57 (1004.06)	7	6800.17 (622.59)	23	6904.13 (558.27)	15	6605.25 (726.85)	8	6654.26 (631.98)	27	6482.67 (845.35)	15
<b>TBLH bone mass (g)</b>	170.32 (24.35)	19	170.58 (14.49)	12	169.86 (37.33)	7	176.57 (24.75)	23	182 (24.05)	15	166.38 (22.81)	8	176.93 (20.83)	27	168.00 (29.89)	15
<b>Percent fat mass</b>	29.80 (3.13)	19	29.84 (2.96)	12	29.73 (3.64)	7	27.92 (2.96)	23	27.97 (2.58)	15	27.81 (3.76)	8	28.80 (2.86)	27	28.71 (3.70)	15
<b>Percent fat-free mass</b>	70.20 (3.13)	19	70.16 (2.96)	12	70.27 (3.64)	7	72.08 (2.96)	23	72.03 (2.58)	15	72.19 (3.76)	8	71.20 (2.86)	27	71.29 (3.70)	15