




Brief Communication

Enhancing gene editing and gene targeting efficiencies in *Arabidopsis thaliana* by using an intron-containing version of *ttLbCas12a*

Patrick Schindele¹ , Laura Merker¹ , Tom Schreiber², Anja Prange², Alain Tissier² and Holger Puchta^{1,*} ¹Botanical Institute, Karlsruhe Institute of Technology, Karlsruhe, Germany²Department of Cell and Metabolic Biology, Leibniz Institute of Plant Biochemistry, Halle (Saale), Germany

Received 20 September 2022;

revised 4 November 2022;

accepted 11 November 2022.

*Correspondence (Tel +49 721 608 48894; fax +49 721 608 44874; email holger.puchta@kit.edu)

Keywords: CRISPR/Cas12a, *ttLbCas12a*, gene editing, gene targeting, intron.

The CRISPR/Cas12a system can be used in a wide range of organisms. However, in plants, efficient editing by Cas12a was hampered due to the lower temperatures required for plant cultivation (Malzahn *et al.*, 2019). Recently, we engineered an improved, temperature-tolerant variant of Cas12a from *Lachnospiraceae bacterium*, *ttLbCas12a*, which has shown dramatically increased editing efficiencies compared to *LbCas12a* in *Arabidopsis* at the standard cultivation temperature of 22 °C (Schindele and Puchta, 2020). Furthermore, we could show that, by using *ttLbCas12a*, gene targeting (GT) efficiencies can be elevated considerably, both in *Arabidopsis* and tobacco (Huang *et al.*, 2021; Merker *et al.*, 2020). Nevertheless, even when using *ttLbCas12a*, some target sites remain highly resistant to editing. Evidence is accumulating that especially heterochromatic sites are more recalcitrant to editing (Weiss *et al.*, 2022). Recently, it was demonstrated that the incorporation of 13 introns into the Cas9 coding sequence dramatically improves the editing efficiency in plants (Grützner *et al.*, 2021). Based on these findings, an intron-containing *ttLbCas12a* construct (*ttLbCas12a-i*) was established to investigate whether editing in plants could be further improved.

The *ttLbCas12a-i* variant was generated similarly to the respective Cas9-intron variant (Grützner *et al.*, 2021). Short introns from *Arabidopsis* genes with canonical donor and acceptor sites (AGGT) were equally placed into the coding sequence with an average distance of 200–400 nt (exon size). Due to the smaller size of *ttLbCas12a* compared to SpCas9, only 10 introns were introduced into the coding sequence (Appendix S1). The resulting *ttLbCas12a-i* ORF was optimized for improved splicing efficiency using the NetGene2 intron splice site prediction software (<http://www.cbs.dtu.dk/services/NetGene2/>). The intron-containing *ttLbCas12a* coding sequence was cloned into a T-DNA expression vector (Figure 1a, top) as previously described for *ttLbCas12a* (Schindele and Puchta, 2020). First, we selected seven different target sites to compare the editing efficiency of *ttLbCas12a-i* with *ttLbCas12a*. Here, we

mainly focused on target sites with low to medium editing efficiencies of *ttLbCas12a*. Two sites were located in the *ECA3* gene and five in heterochromatic, pericentromeric DNA adjacent to centromere I, III and IV (Figure 1a, bottom). Using the floral dip method, *Arabidopsis thaliana* were transformed with T-DNAs containing the respective crRNAs as well as the two variants. To determine the editing efficiency, 17–20 transgenic plants each were grown for 2 weeks at 22 °C. Subsequently, the gDNA of each individual whole plant was extracted and then assessed by TIDE analysis. As shown in Figure 1b,c, three target sites showed hardly any editing in case of *ttLbCas12a* (T1–T3), another three showed low-to-medium editing efficiencies (T4–T6) and one showed medium-to-high editing efficiencies (T7). The new construct *ttLbCas12a-i* surpassed *ttLbCas12a* significantly in terms of editing efficiency at every single target. On average, *ttLbCas12a-i* was between 1.3- and 6.8-fold more efficient than *ttLbCas12a*. This improvement is also highlighted at the population level, where, for the majority of the respective targets and plants, higher and more consistent editing efficiencies could be observed in case of *ttLbCas12a-i*. However, the most tremendous improvements were obtained in case of the three targets that had been barely edited by *ttLbCas12a*. By using *ttLbCas12a-i*, all of these targets became accessible for editing. Beyond that, almost every single plant showed much higher editing efficiencies in case of *ttLbCas12a-i*. Based on these results, we aimed to investigate whether also GT could be further improved by using *ttLbCas12a-i*. Therefore, we performed GT experiments in *Arabidopsis*, using the ALS selection system and same setup as previously described for *ttLbCas12a* (Merker *et al.*, 2020). Using the floral dip method, *A. thaliana* were transformed with T-DNAs containing the respective GT constructs. For both variants, 100 transgenic plants each were grown at 22 °C, propagated and, in the following generation, analysed for GT events. We determined the percentage of lines revealing GT events and assessed the total GT efficiency in terms of percentage of GT events per line (Figure 1d). The percentage of lines with GT events was only slightly higher in case of *ttLbCas12a-i* compared to *ttLbCas12a*. However, concerning the total GT efficiency, a dramatic increase could be observed in case of *ttLbCas12a-i* compared with *ttLbCas12a*, which corresponded to a 2.7-fold increase. If the GT efficiencies of the individual lines are taken into account, the difference between *ttLbCas12a* and *ttLbCas12a-i* becomes even more apparent (Figure 1e): While in case of *ttLbCas12a*, less than a third of the positive lines showed GT efficiencies above 1%, in case of *ttLbCas12a-i*, more than three-quarter of the positive lines did. To analyse the nature of GT events, that is to assess whether GT events constitute perfect or ectopic GT




Figure 1 Improvement of gene editing and GT using *ttLbCas12a-i*. (a) Cas12a expression cassettes for gene editing. The D156R mutation is highlighted, introns are represented by grey bars. The target sequences and regions are given. (b) Table showing the editing efficiencies determined by TIDE analysis as mean values for the respective target sites and variants. Editing efficiencies of the variants were determined for each of the tested target sites and individual transgenic plants grown at 22 °C. (c) Distribution of editing efficiencies of individual plants for both *ttLbCas12a* variants. Illustrated are the editing efficiencies of each of the analysed transgenic plants at the tested target sites (T1-T7) for both variants at 22 °C. Data are presented as boxplots, each box represents the 25th and 75th percentile, the median is indicated by a black line, whiskers encompass the 1.5× interquartile range and data beyond, that is shown as outliers. Fold increases for *ittLbCas12a-i* compared with *ttLbCas12a* at each target site are shown. T1: *n* = 20, T2: *n* = 20, T3: *n* = 17, T4: *n* = 17, T5: *n* = 18, T6: *n* = 20, T7: *n* = 20. *P* values were calculated using the one-way ANOVA test: **P* < 0.05, ***P* < 0.01, *****P* < 0.0001. (d) Table showing the GT efficiencies and molecular analysis of GT events. The number of analysed T1 lines, the number and percentage of positive T1 lines and the total GT efficiency are shown on the left side of the table. The total GT efficiency is defined as the average of GT efficiencies of each individual line including the lines without GT events. The number of analysed positive T2 plants and the number and percentage of T2 plants with perfect GT events are shown on the right side of the table. (e) Distribution of GT efficiencies of the individual lines for both variants at 22 °C. The GT efficiencies of the individual positive lines are arranged in descending order for both variants. GT events were detected for 83 out of 100 and 87 out of 100 lines in case of *ttLbCas12a* and *ttLbCas12a-i*, respectively.

events, molecular analysis of selected positive plants was performed (Figure 1d). It turned out that perfect GT events were obtained at similar frequencies by using either of the variants.

Taken together, our results demonstrate that the introduction of introns into the coding sequence of *ttLbCas12a* dramatically improves both gene editing and gene targeting in *Arabidopsis*, so that even heterochromatic and other hardly accessible targets become editable. In an independent study, it has very recently been shown that the addition of 8 introns into the *ttLbCas12a* coding sequence can enhance editing efficiencies in barley (Lawrenson *et al.*, 2022). Thus, in general, the inclusion of introns into ORFs of Cas nucleases seems to be a very promising way to enhance the editing efficiency in mono- as well as dicotyledonous plant species.

It has already been shown before that the inclusion of introns leads to higher mutation frequencies by Cas9 in *Arabidopsis* due to a related increase in protein levels (Grützner *et al.*, 2021). We assume that the enhanced accessibility of heterochromatic sites to editing is also due to the presence of higher amounts of nuclease in the cells. This expression boost is promoter-independent, making *ttLbCas12a-i* a valuable novel tool for the plant community, enabling a wide variety of applications. It is especially promising in cases where the induction of several or multiple breaks with high efficiency is required, such as multiplexing. Moreover, the use of *ttLbCas12a-i* might also improve gene editing efficiencies in other organisms that require cultivation temperatures similar to plants.

Acknowledgements

We thank Aleya Abinik and Carolin Brechtel for excellent technical assistance and Michelle Rönspies for critical reading of the manuscript. The *ttLbCas12a-i* constructs (MoClo L0 Golden Gate module pAGT8163; AATG_ *ttLbCas12a-i*_TTCG) will be available via Addgene (ID 192453). Open Access funding enabled and organized by Projekt DEAL.

Conflict of interest

The authors declare no conflict of interest.

Author contribution

H.P. and A.T. conceived the research. P.S. and L.M. designed the analysis. T.S. designed the *ttLbCas12a* with introns. A.P. generated the original *ttLbCas12a-i* construct. P.S. and L.M. executed the experiments. All authors wrote the article.

References

- Grützner, R., Martin, P., Horn, C., Mortensen, S., Cram, E.J., Lee-Parsons, C.W.T., Stuttmann, J. *et al.* (2021) High-efficiency genome editing in plants mediated by a Cas9 gene containing multiple introns. *Plant Commun.* **2**, 135.
- Huang, T.-K., Armstrong, B., Schindele, P. and Puchta, H. (2021) Efficient gene targeting in *Nicotiana tabacum* using CRISPR/SaCas9 and temperature tolerant LbCas12a. *Plant Biotechnol. J.* **19**, 1314–1324.
- Lawrenson, T., Hinchliffe, A., Forner, M. and Harwood, W. (2022) *Highly efficient genome editing in barley using novel Lb Cas12a variants and impact of sgRNA architecture.* *bioRxiv*, <https://doi.org/10.1101/2022.04.28.489853v1>
- Malzahn, A.A., Tang, X., Lee, K., Ren, Q., Sretenovic, S., Zhang, Y., Chen, H. *et al.* (2019) Application of CRISPR-Cas12a temperature sensitivity for improved genome editing in rice, maize, and Arabidopsis. *BMC Biol.* **17**, 9.
- Merker, L., Schindele, P., Huang, T.-K., Wolter, F. and Puchta, H. (2020) Enhancing in planta gene targeting efficiencies in *Arabidopsis* using temperature-tolerant CRISPR/LbCas12a. *Plant Biotechnol. J.* **18**, 2382–2384.
- Schindele, P. and Puchta, H. (2020) Engineering CRISPR/LbCas12a for highly efficient, temperature-tolerant plant gene editing. *Plant Biotechnol. J.* **18**, 1118–1120.
- Weiss, T., Crisp, P.A., Rai, K.M., Song, M., Springer, N.M. and Zhang, F. (2022) Epigenetic features drastically impact CRISPR-Cas9 efficacy in plants. *Plant Physiol.* **190**, 1153–1164.

Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Appendix S1 Nucleotide sequence of the open reading frame of the intron-containing *ttLbCas12a-i*; all introns are highlighted in green.