

# Myomatrix arrays for high-definition muscle recording

\*Bryce Chung<sup>1</sup>, \*Muneeb Zia<sup>2</sup>, Kyle A. Thomas<sup>3</sup>, Jonathan A. Michaels<sup>4</sup>, Amanda Jacob<sup>1</sup>, Andrea Pack<sup>5</sup>, Matthew J. Williams<sup>3</sup>, Kailash Nagapudi<sup>1</sup>, Lay Heng Teng<sup>1</sup>, Eduardo Arrambide<sup>1</sup>, Logan Ouellette<sup>1</sup>, Nicole Oey<sup>1</sup>, Rhuna Gibbs<sup>1</sup>, Philip Anschutz<sup>6</sup>, Jiaao Lu<sup>7</sup>, Yu Wu<sup>2</sup>, Mehrdad Kashefi<sup>4</sup>, Tomomichi Oya<sup>4</sup>, Rhonda Kersten<sup>4</sup>, Alice C. Mosberger<sup>8</sup>, Sean O'Connell<sup>3</sup>, Runming Wang<sup>9</sup>, Hugo Marques<sup>10</sup>, Ana Rita Mendes<sup>10</sup>, Constanze Lenschow<sup>10,11</sup>, Gayathri Kondakath<sup>12</sup>, Jeong Jun Kim<sup>13</sup>, William Olson<sup>13</sup>, Kiara N. Quinn<sup>14</sup>, Pierce Perkins<sup>14</sup>, Graziana Gatto<sup>15,16</sup>, Ayesha Thanawalla<sup>15</sup>, Susan Coltman<sup>17</sup>, Taegyo Kim<sup>18</sup>, Trevor Smith<sup>18</sup>, Ben Binder-Markey<sup>19</sup>, Martin Zaback<sup>20</sup>, Christopher K. Thompson<sup>20</sup>, Simon Giszter<sup>18</sup>, Abigail Person<sup>17</sup>, Martyn Goulding<sup>15</sup>, Eiman Azim<sup>15</sup>, Nitish Thakor<sup>14</sup>, Daniel O'Connor<sup>13</sup>, Barry Trimmer<sup>12</sup>, Susana Q. Lima<sup>10</sup>, Megan R. Carey<sup>10</sup>, Chethan Pandarinath<sup>9</sup>, Rui M. Costa<sup>8,21</sup>, J. Andrew Pruszyński<sup>4</sup>, Muhannad Bakir<sup>2</sup>, Samuel J. Sober<sup>1</sup>

\*Equal contributions

<sup>1</sup> Department of Biology, Emory University (Atlanta, GA, USA).

<sup>2</sup> School of Electrical and Computer Engineering, Georgia Institute of Technology (Atlanta, GA, USA).

<sup>3</sup> Graduate Program in Biomedical Engineering at Emory University and Georgia Tech (Atlanta, GA, USA).

<sup>4</sup> Department of Physiology and Pharmacology, Western University (London, ON, Canada).

<sup>5</sup> Neuroscience Graduate Program, Emory University (Atlanta, GA, USA).

<sup>6</sup> Graduate Program in BioEngineering, Georgia Tech (Atlanta, GA, USA).

<sup>7</sup> Graduate Program in Electrical and Computer Engineering, Georgia Tech (Atlanta, GA, USA).

<sup>8</sup> Zuckerman Mind Brain Behavior Institute at Columbia University (New York, NY, USA).

<sup>9</sup> Department of Biomedical Engineering at Emory University and Georgia Tech (Atlanta, GA, USA).

<sup>10</sup> Champalimaud Neuroscience Programme, Champalimaud Foundation (Lisbon, Portugal).

<sup>11</sup> current address: Institute of Biology, Otto-von-Guericke University, (Magdeburg, Germany).

<sup>12</sup> Department of Biology, Tufts University (Medford, MA, USA).

<sup>13</sup> Solomon H. Snyder Department of Neuroscience, Johns Hopkins School of Medicine (Baltimore, MD, USA).

<sup>14</sup> Departments of Biomedical Engineering and Neurology, Johns Hopkins School of Medicine (Baltimore, MD, USA).

<sup>15</sup> Salk Institute for Biological Studies (La Jolla, CA, USA).

<sup>16</sup> current address: Department of Neurology, University Hospital of Cologne (Cologne, Germany).

<sup>17</sup> Department of Physiology and Biophysics, University of Colorado Anschutz Medical Campus (Aurora, CO, USA).

<sup>18</sup> Department of Neurobiology & Anatomy, Drexel University, College of Medicine (Philadelphia, PA, USA).

<sup>19</sup> Department of Physical Therapy and Rehabilitation Sciences, Drexel University College of Nursing and Health Professions (Philadelphia, PA).

<sup>20</sup> Department of Health and Rehabilitation Sciences, Temple University (Philadelphia, PA, USA).

<sup>21</sup> Allen Institute (Seattle, WA, USA).

# Myomatrix arrays for high-definition muscle recording


## Abstract

Neurons coordinate their activity to produce an astonishing variety of motor behaviors. Our present understanding of motor control has grown rapidly thanks to new methods for recording and analyzing populations of many individual neurons over time. In contrast, current methods for recording the nervous system's actual motor output – the activation of muscle fibers by motor neurons – typically cannot detect the individual electrical events produced by muscle fibers during natural behaviors and scale poorly across species and muscle groups. Here we present a novel class of electrode devices (“Myomatrix arrays”) that record muscle activity at unprecedented resolution across muscles and behaviors. High-density, flexible electrode arrays allow for stable recordings from the muscle fibers activated by a single motor neuron, called a “motor unit”, during natural behaviors in many species, including mice, rats, primates, songbirds, frogs, and insects. This technology therefore allows the nervous system's motor output to be monitored in unprecedented detail during complex behaviors across species and muscle morphologies. We anticipate that this technology will allow rapid advances in understanding the neural control of behavior and in identifying pathologies of the motor system.

## Introduction

1 Recent decades have seen tremendous advances in our understanding of the physiological mechanisms by which the  
2 brain controls complex motor behaviors. Critical to these advances have been tools to record neural activity at  
3 scale<sup>4,5</sup>, which, when combined with novel algorithms for behavioral tracking<sup>6-10</sup>, can reveal how neural activity  
4 shapes behavior<sup>11,12</sup>. In contrast, current methods for observing the nervous system's motor output lag far behind  
5 neural recording technologies. The nervous system's control of skeletal motor output is ultimately mediated by “motor  
6 units”, each of which consists of a single motor neuron and the muscle fibers it activates, producing motor unit action  
7 potentials (**Fig. 1a**) that generate muscle force to produce movement<sup>13</sup>. Because each action potential in a motor  
8 neuron reliably evokes a single spike in its target muscle fibers, action potentials recorded from muscle provide a  
9 high-resolution readout of motor neuron activity in the spinal cord and brainstem. However, our understanding of  
10 motor unit activity during natural behaviors is rudimentary due to the difficulty of recording spike trains from motor unit  
11 populations.

12  
13 Traditional methods for recording muscle fiber activity via electromyography (EMG) include fine wires inserted into  
14 muscles and electrode arrays placed on the surface of the skin<sup>14</sup>. These methods can resolve the activity of individual  
15 motor units in only a limited range of settings. First, to prevent measurement artifacts, traditional EMG methods  
16 require that a subject's movements be highly restricted, typically in “isometric” force tasks where subjects contract  
17 their muscles without moving their bodies<sup>15-18</sup>. Moreover, fine wire electrodes typically cannot detect single motor unit  
18 activity in small muscles, including the muscles of widely used model systems such as mice or songbirds<sup>19-21</sup>, and  
19 surface electrode arrays are poorly tolerated by freely behaving animal subjects. These limitations have impeded our



**Figure 1: Myomatrix arrays record muscle activity at motor unit resolution.** (a) The nervous system controls behavior via motor units, each consisting of a single motor neuron and the muscle fibers it innervates. Each motor neuron's spiking evokes motor unit action potentials in the corresponding muscle fibers. Myomatrix arrays (right) bearing 32 electrode contacts on a flexible substrate (**Supplemental Fig. 1**) can be targeted to one or more muscles and yield high-resolution recordings of motor activity during free behavior. Motor neurons, muscle fibers, and electrode arrays are not shown to scale. (b,c) Example recordings from the right triceps muscle of a freely behaving mouse. (b) Top, bipolar Myomatrix recording from the mouse triceps during locomotion. Blue dots indicate the spike times of one motor unit isolated from the data using a spike sorting method based on principal components analysis (**Supplemental Fig. 2a-d**). Bottom, example data (from <sup>1</sup>, used with permission) from traditional fine-wire EMG recording of triceps activity during locomotion. Applying the PCA-based spike sorting method to the fine-wire data did not isolate any individual motor units. (c) Unipolar Myomatrix recording during quiet stance. Colored boxes illustrate motor unit action potentials from four identified units. Spike waveforms from some units, including those highlighted with gray and orange boxes, appear on multiple electrode channels, requiring the use of a multi-channel spike sorting algorithm (Kilosort 2.5, see **Supplemental Fig. 2e-h**). (d) Spiking pattern (tick marks) of six individual motor units recorded simultaneously during locomotion on a treadmill. The three bursts of motor unit action potentials correspond to triceps activity during three stride cycles. Motor unit 4 (cyan) is the same motor unit represented by cyan dots in (b). The other motor units in this recording, including the smaller amplitude units at top in (b), were isolated using Kilosort but could not be isolated with the PCA-based method applied to data from only the single recording channel shown (b).

20 understanding of fundamental questions in motor control, including how the nervous system coordinates populations  
 21 of motor units to produce skilled movements, how this coordination degrades in pathological states, and how motor  
 22 unit activity is remapped when animals learn new tasks or adapt to changes in the environment.

24 Here, we present a novel approach (**Fig. 1**) to recording populations of individual motor units from many different  
25 muscle groups and species during natural behaviors. Flexible multielectrode (“Myomatrix”) arrays were developed to  
26 achieve the following goals:

- (a) Record muscle activity at motor unit resolution
- (b) Record motor units during active movements
- (c) Record from a wide range of muscle groups, species, and behaviors
- (d) Record stably over time and with minimal movement artifact

27 To achieve these goals, we developed a variety of array configurations for use across species and muscle groups.  
28 Voltage waveforms from individual motor units (**Fig. 1b,c**) can be readily extracted from the resulting data using a  
29 range of spike-sorting algorithms, including methods developed to identify the waveforms of individual neurons in  
30 high-density arrays<sup>2,22</sup>. Below, we show how Myomatrix arrays provide high-resolution measures of motor unit  
31 activation in a variety of species and muscle groups including forelimb, hindlimb, orofacial, pelvic, vocal, and  
32 respiratory muscles.

33

## 34 Results

35 We developed methods to fabricate flexible, high-density EMG (“Myomatrix”) arrays, as detailed in the Methods and  
36 schematized in **Supplemental Figure 1**. We selected polyimide as a substrate material due to its strength and  
37 flexibility and the ease with which we could define electrode contacts, suture holes, and other sub-millimeter features  
38 that facilitate ease of implantation and recording stability (**Supplemental Fig. 1a-e**). Moreover, simple modifications to  
39 the fabrication pipeline allowed us to rapidly design, test, and refine different array morphologies targeting a range of  
40 muscle sizes, shapes, and anatomical locations (**Supplemental Fig. 1c, f, g**).

41

### 42 Myomatrix arrays record muscle activity at motor unit resolution


43 Myomatrix arrays robustly record the activity of individual motor units in freely behaving mice. Arrays specialized for  
44 mouse forelimb muscles include four thin “threads” (8 electrodes per thread, 32 electrodes total) equipped with suture  
45 holes, flexible “barbs,” and other features to secure the device within or onto a muscle (**Fig. 1a, Supplemental Fig.**  
46 **1c, d, e, h**). These devices yielded well-isolated waveforms from individual motor units (**Fig. 1b, top**), which were  
47 identified using open-source spike sorting tools<sup>2,22</sup>. As detailed in **Supplemental Figure 2a-d**, in some cases the  
48 spike times of individual motor units (cyan dots, **Fig. 1b**) can be isolated from an individual electrode channel with  
49 simple spike sorting approaches including single-channel waveform clustering<sup>22</sup>. In other cases, waveforms from  
50 individual motor units appeared on multiple electrode channels (**Fig. 1c**), allowing – and in many cases necessitating  
51 – more advanced spike-sorting approaches that leverage information from multiple channels to identify larger

52 numbers of units and resolve overlapping spike waveforms <sup>2</sup>, as detailed in **Supplemental Figure 2e-h**. These  
53 methods allow the user to record simultaneously from ensembles of single motor units (**Fig. 1c,d**) in freely behaving  
54 animals, even from small muscles including the lateral head of the triceps muscle in mice (approximately 9 mm in  
55 length with a mass of 0.02 g <sup>23</sup>). Myomatrix recordings isolated single motor units for extended periods (greater than  
56 two months, **Supplemental Fig. 3e**), although highest unit yield was typically observed in the first 1-2 weeks after  
57 chronic implantation. Because recording sessions from individual animals were often separated by several days  
58 during which animals were disconnected from data collection equipment, we are unable to assess based on the  
59 present data whether the same motor units can be recorded over multiple days.

60

### 61 **Myomatrix arrays record motor units during active movements**

62 Myomatrix arrays outperform traditional fine-wire electrodes in mice by reliably recording isolated single units in  
63 behaving animals. First, Myomatrix arrays isolate the activity of multiple individual motor units during freely moving  
64 behavior (**Fig. 1c-d**). In contrast, wire electrodes typically cannot resolve individual motor units during muscle  
65 lengthening and shorting, as occurs in naturalistic movements such as locomotion <sup>1,24</sup>. **Figure 1b** illustrates a  
66 comparison between Myomatrix (top) and fine-wire (bottom) data recorded during locomotion in the mouse triceps.  
67 Spike-sorting identified well-isolated motor unit spikes in the Myomatrix data (cyan dots in **Fig. 1b, top**) but failed to  
68 extract any isolated motor units in the fine wire data (**Supplemental Fig. 2a,b**). Similarly, while Myomatrix recordings  
69 robustly isolated motor units from a songbird vocal muscle, fine wire EMG electrodes applied to the same muscle did  
70 not yield isolatable units (**Supplemental Fig. 2c,d**). This lack of resolution, which is typical of fine wire EMG, severely  
71 limits access to motor unit activity during active behavior, although wire electrodes injected through the skin can  
72 provide excellent motor unit isolation during quiet stance in mice <sup>25</sup>. Second, because wire-based EMG requires  
73 inserting an additional wire for each additional electrode contact, only a single pair of wires (providing a single bipolar  
74 recording channel, **Fig. 1b, bottom**) can be inserted into an individual mouse muscle in most cases <sup>1,24,26</sup>. In contrast,  
75 at least four Myomatrix “threads” (**Fig 1a**), bearing a total of 32 electrodes, can be inserted into one muscle (**Fig. 1c**  
76 shows five of 32 channels recorded simultaneously from mouse triceps), greatly expanding the number of recording  
77 channels within a single muscle. Single motor units were routinely isolated during mouse locomotion in our Myomatrix  
78 recordings (**Fig. 1**), but never in the fine-wire datasets from <sup>1</sup> we re-analyzed or, to our knowledge, in any prior study.  
79 Moreover, in multiunit recordings, Myomatrix arrays have significantly higher signal-to-noise ratios than fine-wire EMG  
80 arrays (**Supplemental Fig. 3**). Myomatrix arrays therefore far exceed the performance of wire electrodes in mice in  
81 terms of both the quality of recordings and the number of channels that can be recorded simultaneously from one  
82 muscle.



**Figure 2: Myomatrix recordings across muscles and species.** (a) Example recording from **mouse triceps** during a head-fixed pellet reaching task. Arrows at top indicate the approximate time that the animal's paw leaves a rest position and first contacts the target. Bottom, colored boxes highlight motor unit action potentials identified using Kilosort<sup>2</sup>. Different box colors on the same voltage trace indicate distinct motor units. (b) Recordings from the **mouse superficial masseter** muscle were obtained in anesthetized, head-fixed mice when passive mandible displacement evoked reflexive muscle contractions. Top trace shows the lateral component of jaw displacement, with arrows indicating the direction and approximate time of displacement onset. (c) In a recording from **rat triceps** during head-free locomotion, the arrowhead indicates the time that the mouse's paw touched the treadmill surface, marking the beginning of the stance phase. (d) Recording from the **rat flexor digitorum profundus** muscle during a pellet reaching task, arrow indicates the time of grasp initiation. (e) Myomatrix recording of motor unit activity in the **mouse bulbospongiosus** muscle evoked by optical stimulation of spinal motor neurons, producing motor unit spikes at latencies between 10-15 msec, consistent with results obtained from traditional fine-wire electrodes in mice<sup>3</sup>. (f-j) Recordings from the **cat soleus** (f) during sensory nerve stimulation, **songbird vocal (ventral syringeal) muscle** (g) and **expiratory muscle** (h) during quiet respiration, **hawkmoth larva dorsal internal medial (DIM) muscle** (i) during fictive locomotion, and **bull frog semimembranosus (SM) muscle** (j) in response to cutaneous (foot) stimulation. Spike times from individual motor units are indicated by colored tick marks under each voltage trace in f-j. Recordings shown in panels (a, c, g, h, i, and j) were collected using bipolar amplification, data in panels (b, d, e, and f) were collected using unipolar recording. See Methods for details of each experimental preparation.

83 **Myomatrix arrays record from a wide range of muscle groups, species, and behaviors**

84 Myomatrix arrays provide high-resolution EMG recordings across muscle targets and experimental preparations (Fig.


85 2). Beyond the locomotor and postural signals shown in Figure 1, Myomatrix recordings from mouse triceps also

86 provided single-unit EMG data during a head-fixed reaching task (**Fig. 2a**). In addition to recording single motor units  
87 during these voluntary behaviors, Myomatrix arrays also allow high-resolution recordings from other muscle groups  
88 during reflex-evoked muscle activity. **Figure 2b** shows single motor unit EMG data recorded from the superficial  
89 masseter (jaw) muscle when reflexive muscle contraction was obtained by passively displacing the jaw of an  
90 anesthetized mouse. To extend these methods across species, we collected Myomatrix recordings from muscles of  
91 the rat forelimb, obtaining isolated motor units from the triceps during locomotion (**Fig. 2c**) and a digit-flexing muscle  
92 in the lower forearm during head-free reaching (**Fig. 2d**). Myomatrix arrays can furthermore isolate motor unit  
93 waveforms evoked by direct optogenetic stimulation of spinal motor neurons. **Figure 2e** shows recordings of light  
94 evoked spikes in the mouse bulbospongiosus muscle (a pelvic muscle that wraps around the base of the penis in  
95 male mice), demonstrating that optogenetic stimulation of the spinal cord evokes spiking in single motor units with  
96 millisecond-scale timing jitter (**Fig. 2e**, center) and with latencies (**Fig. 2e**, right) consistent with the latencies of  
97 recordings obtained with fine-wire electrodes<sup>3</sup>. Beyond rodents, simple modifications of the basic electrode array  
98 design (**Supplemental Fig. 1f**) allowed us to obtain high-resolution recordings from hindlimb muscles in cats (**Fig. 2f**),  
99 vocal and respiratory muscles in songbirds (**Fig. 2g,h**, see also<sup>27</sup>), body wall muscles in moth larvae (**Fig. 2i**), and leg  
100 muscles in frogs (**Fig. 2j**).

101  
102 In addition to isolating individual motor units, Myomatrix arrays provide stable multi-unit recordings of comparable or  
103 superior quality to conventional fine wire EMG. Although single-unit recordings are essential to identify individual  
104 motor neurons' contributions to muscle activity<sup>28,29</sup>, for other lines of inquiry a multi-unit signal is preferred as it  
105 reflects the combined activity of many motor units within a single muscle. Although individual Myomatrix channels are  
106 often dominated by spike waveforms from one or a small number of motor units (**Fig 1b**), other channels reflect the  
107 combined activity of multiple motor units as typically observed in fine-wire EMG recordings<sup>30</sup>. As shown in  
108 **Supplemental Figure 3a** and **b**, these multi-unit Myomatrix signals are stable over multiple weeks of recordings,  
109 similar to the maximum recording longevity reported for wire-based systems in mice and exceeding the 1-2 weeks of  
110 recording more typically obtained with wire electrodes in mice<sup>1,24,26</sup>, and with significantly greater recording quality  
111 than that obtained from wire electrodes at comparable post-implantation timepoints (**Supplemental Fig. 3d**).

112

113 To record from larger muscles than those described above, we also created designs targeting the forelimb and  
 114 shoulder muscles of rhesus macaques (**Fig. 3**). Although fine wire electrodes have been used to isolate individual  
 115 motor units in both humans and monkeys <sup>14,31</sup>, and skin-surface electrode arrays robustly record motor unit  
 116 populations in human subjects <sup>15,32</sup>,  
 117 this resolution is limited to isometric  
 118 tasks – that is, muscle contraction  
 119 without movement – due to the  
 120 sensitivity of both fine-wire and surface  
 121 array electrodes to electrical artifacts  
 122 caused by body movement. For ease  
 123 of insertion into larger muscles, we  
 124 modified the “thread” design used in  
 125 our mouse arrays so that each  
 126 Myomatrix array could be loaded into a  
 127 standard hypodermic syringe and  
 128 injected into the muscle  
 129 (**Supplemental Fig. 1g,i**), inspired by  
 130 earlier work highlighting the  
 131 performance of injectable arrays in  
 132 primates <sup>14,33,34</sup>. As shown in **Figure**  
 133 **3a-d**, injectable Myomatrix arrays  
 134 yielded motor unit recordings during  
 135 arm movements. Tick marks in **Figure**  
 136 **3d** show the activity of 13 motor units  
 137 recorded simultaneously during a



**Figure 3: Motor unit recordings during active movement in primates. (a)** An injectable version of the Myomatrix array (**Supplemental Fig. 1g**) was inserted percutaneously (**Supplemental Fig. 1i**) into the right biceps of a rhesus macaque performing a cued reaching task. Green and red dots: reach start and endpoints, respectively; grey regions: start and target zones. **(b)** Recording from five of 32 unipolar channels showing spikes from three individual motor units isolated from the multichannel recording using Kilosort (**Supplemental Fig. 2**). **(c)** At trial onset (dotted line), a sudden force perturbation extends the elbow, signaling the animal to reach to the target. **(d)** Spike times (tick marks) from 13 simultaneously recorded motor units. **(e)** Example voltage data from a Myomatrix array (top) and traditional fine-wire EMG (middle, bottom) collected from the same biceps muscle in the same animal performing the same task, but in a separate recording session. Gray traces (bottom) show smoothed EMG data from the fine-wire electrodes in all trials, orange trace shows trial-averaged smoothed fine-wire EMG, dark gray trace represents the fine-wire trial shown at middle. **(f)** Spike times of four motor units (of the 13 shown in **d**) recorded simultaneously over 144 trials.



138 single trial in which a monkey was cued (by a force perturbation which causes an extension of the elbow joint) to  
139 reach to a target. In contrast to the ensemble of spike times obtained with a Myomatrix probe, conventional fine-wire  
140 EMGs inserted into the same muscle (in a separate recording session) yielded only a single trace reflecting the  
141 activity of an unknown number of motor units (**Fig 3e**, middle). Moreover, although fine-wire EMG signals varied  
142 across trials (**Fig. 3e**, bottom), the lack of motor unit resolution makes it impossible to assess how individual motor  
143 units vary (and co-vary) across trials. In contrast, Myomatrix recordings provide spiking resolution of multiple  
144 individual motor units across many trials, as illustrated in **Figure 3f**.

145

#### 146 **Myomatrix arrays record stably over time and with minimal movement artifact**

147 Myomatrix arrays provide EMG recordings that are resistant to movement artifacts and stable over time. In recordings  
148 from rodents during locomotion (**Fig. 1b**, **Fig. 2c**), we did not observe voltage artifacts at any point in the stride cycle  
149 (e.g. when the paw of the recorded limb first touches the ground in each stride cycle; **Fig. 2c**, black arrowhead).  
150 Movement artifacts were similarly absent during active arm movements in monkeys (peak fingertip speeds ~15  
151 cm/sec; peak elbow angle velocity ~40 deg/sec, **Fig 3b,e**), during passive jaw displacement in anesthetized mice  
152 (**Fig. 2b**), or in the other anesthetized preparations shown in **Figure 2**. Individual motor units could typically be  
153 isolated for the entire duration of an acute or chronic recording session. In triceps recordings during locomotion in  
154 rodents (**Fig. 1**, **Fig. 2c**), isolated motor units were recorded for up to 4,791 stride cycles (mice) or 491 stride cycles  
155 (rats) during continuous recording sessions lasting 10-60 minutes. Myomatrix recordings in behaving nonhuman  
156 primates were similarly long-lived, as in the dataset shown in **Figure 3**, where single-unit isolation was maintained  
157 across 1,292 reaching trials collected over 97 minutes. In each of these datasets, the duration of single-unit EMG  
158 recording was limited by the willingness of the animal to continue performing the behavior, rather than a loss of signal  
159 isolation. Recordings in acute preparations were similarly stable. For example, the songbird dataset shown in **Figure**  
160 **2g** includes single-unit data from 8,101 respiratory cycles collected over 74 minutes, and, like the other acute  
161 recordings shown in **Figure 2**, recordings were ended by the experimenter rather than because of a loss of signal  
162 from individual motor units.

163

164 The diversity of applications presented here demonstrates that Myomatrix arrays can obtain high-resolution EMG  
165 recordings across muscle groups, species, and experimental conditions including spontaneous behavior, reflexive  
166 movements, and stimulation-evoked muscle contractions. Although this resolution has previously been achieved in  
167 moving subjects by directly recording from motor neuron cell bodies in vertebrates<sup>35-37</sup> and by using fine-wire  
168 electrodes in moving insects<sup>38,39</sup>, both methods are extremely challenging and can only target a small subset of

169 species and motor unit populations. Exploring additional muscle groups and model systems with Myomatrix arrays will  
170 allow new lines of investigation into how the nervous system executes skilled behaviors and coordinates the  
171 populations of motor units both within and across individual muscles. These approaches will be particularly valuable in  
172 muscles in which each motor neuron controls a very small number of muscle fibers, allowing fine control of  
173 oculomotor muscles in mammals as well as vocal muscles in songbirds (**Fig. 2g**), in which most individual motor  
174 neurons innervate only 1-3 muscle fibers<sup>40</sup>. Of further interest will be combining high-resolution EMG with precise  
175 measurement of muscle length and force output to untangle the complex relationship between neural control, body  
176 kinematics, and muscle force that characterizes dynamic motor behavior. Similarly, combining Myomatrix recordings  
177 with high-density brain recordings or targeted manipulations of neural activity can reveal how central circuits shape  
178 and reshape motor activity and – in contrast to the multi-unit signals typically obtained from traditional EMG in animals  
179 – reveal how neural dynamics in cortical, subcortical, and spinal circuits shape the spiking patterns of individual motor  
180 neurons.

181

182 Applying Myomatrix technology to human motor unit recordings, particularly by using the minimally invasive injectable  
183 designs shown in **Figure 3** and **Supplemental Figure 1g,i**, will create novel opportunities to diagnose motor  
184 pathologies and quantify the effects of therapeutic interventions in restoring motor function. Moreover, because  
185 Myomatrix arrays are far more flexible than the rigid needles commonly used to record clinical EMG, our technology  
186 might significantly reduce the risk and discomfort of such procedures while also greatly increasing the accuracy with  
187 which human motor function can be quantified. This expansion of access to high-resolution EMG signals – across  
188 muscle groups, species, and behaviors – is the chief impact of the Myomatrix project.

## 189 Methods

190

### 191 Myomatrix array fabrication

192 The microfabrication process (schematized in **Supplemental Fig. 1a**) consists of depositing and patterning a series of  
193 polymer (polyimide) and metal (gold) layers, using a combination of spin coating, photolithography, etching, and  
194 evaporation processes, as described previously<sup>27,41,42</sup>. These methods allow very fine pitch escape routing (<10  $\mu\text{m}$   
195 spacing between the thin “escape” traces connecting electrode contacts to the connector), spatial alignment between  
196 the multiple layers of polyimide and gold that constitute each device, and precise definition of “via” pathways that  
197 connect different layers of the device. Once all the metal and polyimide layers have been deposited and patterned on  
198 carrier wafers, the gold EMG recording electrode sites are formed by removing the top polyimide layer over each  
199 electrode site using reactive ion etching process ( $\text{O}_2$  and  $\text{SF}_6$  plasma, 5:1 ratio). Electrode sites are then coated with a  
200 conductive polymer, PEDOT:PSS (Poly(3,4-ethylenedioxythiophene)-poly(styrene-sulfonate)),<sup>43-45</sup> to reduce the  
201 electrode impedance<sup>46</sup>. PEDOT:PSS was deposited on the electrode contacts to a thickness of 100 nm using spin  
202 coating, resulting in final electrode impedances of 5 kOhms or less (100 x 200  $\mu\text{m}$  electrode sites). Once all layers  
203 have been deposited on the carrier wafer, the wafer is transferred to an Optec Femtosecond laser system, which is  
204 used to cut the electrode arrays into the shape/pattern needed based on the target muscle group and animal species.  
205 The final device thickness was  $\sim 40 \mu\text{m}$  for the injectable (primate forelimb) design and  $\sim 20 \mu\text{m}$  for all other design  
206 variants. The final fabrication step is bonding a high-density connector (Omnetics, Inc.) to the surface of the electrode  
207 array using a Lambda flip-chip bonder (Finetech, Inc.). This fabrication pipeline allows the rapid development and  
208 refinement of multiple array designs (**Supplemental Fig. 1c-g**).

209

### 210 Myomatrix array implantation

211 For chronic EMG recording in mice and rats (**Fig. 1, Fig. 2a, c, d**), arrays such as those shown in **Supplemental**  
212 **Figure 1c-f** were implanted by first making a midline incision (approximately 10 mm length) in the scalp of an  
213 anesthetized animal. The array’s connector was then secured to the skull using dental cement (in some cases along  
214 with a headplate for later head-fixed chronic recordings), and the electrode array threads were routed subcutaneously  
215 to a location near the target muscle or muscles (**Supplemental Fig. 1h**). In some electrode array designs,  
216 subcutaneous routing is facilitated with “pull-through tabs” that can be grasped with a forceps to pull multiple threads  
217 into position simultaneously. For some anatomical targets a small additional incision was made to allow surgical  
218 access to individual muscles (e.g. a 2-5 mm incision near the elbow to facilitate implantation into the biceps and/or  
219 triceps muscles). Once each thread has been routed subcutaneously and positioned near its target, any pull-through  
220 tabs are cut off with surgical scissors and discarded. Each thread can then either be sutured to the surface of the thin  
221 sheet of elastic tissue that surrounds muscles (“epimysial attachment”) or inserted into the muscle using a suture  
222 needle (“intramuscular implantation”). For epimysial attachment, each electrode thread is simply sutured to the  
223 surface of each muscle (suture sizes ranging from 6-0 to 11-0) in one of the proximal suture holes (located on the  
224 depth-restrictor tabs) and one of the distal suture holes. For intramuscular implantation (**Supplemental Fig. 1h**), a  
225 suture (size 6-0 to 11-0 depending on anatomical target) is tied to the distal-most suture hole. The needle is then  
226 passed through the target muscle and used to pull the attached array thread into the muscle. In some designs, a  
227 “depth-restrictor tab” (**Supplemental Fig. 1d**) prevents the thread from being pulled any further into the muscle,  
228 thereby limiting the depth at which the electrodes are positioned within the target muscle. The array is then secured  
229 within the muscle by the passive action of the flexible polyimide “barbs” lining each thread and/or by adding additional  
230 sutures to the proximal and distal suture holes.

231

232 Acute recording in small animals (including rodents, songbirds, cats, frogs, and caterpillars; **Fig. 2b,e-j**) used the  
233 same arrays as chronic recordings. However, for both epimysial and intramuscular acute recordings, the Myomatrix  
234 array traces were simply placed on or within the target muscle after the muscle was exposed via an incision in the  
235 overlying skin of the anesthetized animal (rather than routed subcutaneously from the skull as in chronic applications).

236

237 For acute recordings in nonhuman primates, prior to recording, the “tail” of the injectable array (**Supplemental Fig.**  
238 **1g**) was loaded into a sterile 23-gauge cannula (1” long) until fully seated. The upper half of the cannula bevel, where  
239 contact is made with the electrode, was laser-blunted to prevent breakage of the tail<sup>34</sup>. During insertion  
240 (**Supplemental Fig. 1i**), the tail was bent over the top of the cannula and held tightly, and the electrode was inserted  
241 parallel to bicep brachii long head muscle fibers at an angle of  $\sim 45$  degrees to the skin. Once the cannula was fully  
242 inserted, the tail was released, and the cannula slowly removed. After recording, the electrode and tail were slowly  
243 pulled out of the muscle together. Insertion and removal of injectable Myomatrix devices appeared to be comparable  
244 or superior to traditional fine-wire EMG electrodes (in which a “hook” is formed by bending back the uninsulated tip of  
245 the recording wire) in terms of both ease of injection, ease of removal of both the cannula and the array itself, and  
246 animal comfort. Moreover, in over 100 Myomatrix injections performed in rhesus macaques, there were zero cases in  
247 which Myomatrix arrays broke such that electrode material was left behind in the recorded muscle, representing a  
248 substantial improvement over traditional fine-wire approaches, in which breakage of the bent wire tip regularly occurs

249 14.

250

251 For all Myomatrix array designs, a digitizing, multiplexing headstage (Intan, Inc.) was plugged into the connector,  
252 which was cemented onto the skull for chronic applications and attached to data collection devices via a flexible  
253 tether, allowing EMG signals to be collected during behavior. By switching out different headstages, data from the  
254 same 32 electrode channels on each Myomatrix array could be recorded either as 32 unipolar channels or as 16  
255 bipolar channels, where each bipolar signal is computed by subtracting the signals from physically adjacent electrode  
256 contacts.

257

#### 258 Data analysis: spike sorting

259 Motor unit action potential waveforms from individual motor units were identified with analysis methods previously  
260 used to sort spikes from neural data. In all cases, Myomatrix signals (sampling rate 30 or 40 kHz) were first band-  
261 passed between 350-7,000 Hz. When the voltage trace from a single Myomatrix channel is dominated by a single  
262 high-amplitude action potential waveform (as in **Fig. 1b**), single units can be isolated using principal components  
263 analysis (PCA) to detect clusters of similar waveforms, as described previously<sup>22</sup>. As detailed in **Supplemental**  
264 **Figure 2a-d**, this method provides a simple quantitative measure of motor unit isolation by quantifying the overlap  
265 between clusters of spike waveforms in the space of the first two principal components.

266

267 In other cases (as in **Fig. 1c**), the spikes of individual motor units appear on multiple channels and/or overlap with  
268 each other in time, requiring a more sophisticated spike sorting approach to identifying the firing times of individual  
269 motor units. We therefore adapted Kilosort version 2.5<sup>2,47</sup> and wrote custom MATLAB and Python code to sort  
270 waveforms into clusters arising from individual motor units (**Supplemental Fig. 2e-h**). Our modifications to Kilosort  
271 reflect the different challenges inherent in sorting signals from neurons recorded with Neuropixels probes and motor  
272 units recorded with Myomatrix arrays<sup>14</sup>. These modifications include the following:

273

274 Modification of spatial masking: Individual motor units contain multiple muscle fibers (each of which is  
275 typically larger than a neuron's soma), and motor unit waveforms can often be recorded across  
276 spatially distant electrode contacts as the waveforms propagate along muscle fibers. In contrast,  
277 Kilosort - optimized for the much more local signals recorded from neurons - uses spatial masking to  
278 penalize templates that are spread widely across the electrode array. Our modifications to Kilosort  
279 therefore include ensuring that Kilosort search for motor unit templates across all (and only) the  
280 electrode channels inserted into a given muscle. In the Github repository linked above, this is  
281 accomplished by setting parameter `nops.sigmaMask` to infinity, which effectively eliminates spatial  
282 masking in the analysis of the 32 unipolar channels recorded from the injectable Myomatrix array  
283 schematized in **Supplemental Figure 1g**. In cases including chronic recording from mice where only  
284 a single 8-contact thread is inserted into each muscle, a similar modification can be achieved with a  
285 finite value of `nops.sigmaMask` by setting parameter `NchanNear`, which represents the number of  
286 nearby EMG channels to be included in each cluster, to equal the number of unipolar or bipolar data  
287 channels recorded from each thread. Finally, note that in all cases Kilosort parameter `NchanNearUp`  
288 (which defines the maximum number of channels across which spike templates can appear) must be  
289 reset to be equal to or less than the total number of Myomatrix data channels.

290

291 Allowing more complex spike waveforms: We also modified Kilosort to account for the greater  
292 duration and complexity (relative to neural spikes) of many motor unit waveforms. In the code  
293 repository linked above, Kilosort 2.5 was modified to allow longer spike templates (151 samples  
294 instead of 61), more spatiotemporal PCs for spikes (12 instead of 6), and more left/right eigenvector  
295 pairs for spike template construction (6 pairs instead of 3) to account for the greater complexity and  
296 longer duration of motor unit action potentials<sup>14</sup> compared to the neural action potentials for which  
297 Kilosort was initially created. These modifications were crucial for improving sorting performance in  
298 the nonhuman primate dataset shown in **Figure 3**, and in a subset of the rodent datasets (although  
299 they were not used in the analysis of mouse data shown in **Fig. 1** and **Supplemental Fig. 2a-f**).

300

301

302 We therefore used Kilosort version 2.5<sup>2,47</sup> and custom MATLAB and Python code to sort waveforms into clusters  
303 arising from individual motor units (**Supplemental Fig. 2e-h**). Kilosort 2.5 was modified to allow longer spike  
304 templates (151 samples instead of 61), more spatiotemporal PCs for spikes (12 instead of 6), and more left/right  
305 eigenvector pairs for spike template construction (6 pairs instead of 3) to account for the greater complexity and  
306 longer duration of motor unit action potentials<sup>14</sup> compared to the neural action potentials for which Kilosort was  
307 initially created.

308

309 Individual motor units were identified from "candidate" units by assessing motor unit waveform consistency, SNR, and  
310 spike count, by inspecting auto-correlograms to ensure that each identified units displayed an absolute refractory

311 period of less than 1 msec, and by examining cross-correlograms with other sorted units to ensure that each motor  
312 unit's waveforms were being captured by only one candidate unit. Candidate units with inconsistent waveforms or  
313 >1% of inter-spike intervals above 1 msec were discarded. Candidate units with highly similar waveform shapes and  
314 cross-correlation peaks at lag zero were merged, resulting in sorted units with well-differentiated waveform shapes  
315 and firing patterns (**Supplemental Fig. 2e,f**). Our spike sorting code, which includes the above-mentioned  
316 modifications to Kilosort, is available at <https://github.com/JonathanAMichaels/PixelProcessingPipeline>.

317  
318 Our approach to spike sorting shares the same ultimate goal as prior work using skin-surface electrode arrays to  
319 isolate signals from individual motor units but pursues this goal using different hardware and analysis approaches. A  
320 number of groups have developed algorithms for reconstructing the spatial location and spike times of active motor  
321 units<sup>18,48</sup> based on skin-surface recordings, in many cases drawing inspiration from earlier efforts to localize cortical  
322 activity using EEG recordings from the scalp<sup>49</sup>. Our approach differs substantially. In Myomatrix arrays, the close  
323 electrode spacing and very close proximity of the contacts to muscle fibers ensure that each Myomatrix channel  
324 records from a much smaller volume of tissue than skin-surface arrays. This difference in recording volume in turn  
325 creates different challenges for motor unit isolation: compared to skin-surface recordings, Myomatrix recordings  
326 include a smaller number of motor units represented on each recording channel, with individual motor units appearing  
327 on a smaller fraction of the sensors than typical in a skin-surface recording. Because of this sensor-dependent  
328 difference in motor unit source mixing, different analysis approaches are required for each type of dataset.  
329 Specifically, skin-surface EMG analysis methods typically use source-separation approaches that assume that each  
330 sensor receives input from most or all of the individual sources within the muscle as is presumably the case in the  
331 data. In contrast, the much sparser recordings from Myomatrix are better decomposed using methods like Kilosort,  
332 which are designed to extract waveforms that appear only on a small, spatially restricted subset of recording  
333 channels.

#### 334 335 336 Additional recording methods – mouse forelimb muscle

337 All procedures described below were approved by the Institutional Animal Care and Use Committee at Emory  
338 University (data in **Fig. 1c, Supplemental Fig 3**) or were approved by European Committee Council Directive, the  
339 Animal Care and Users Committee of the Champalimaud Neuroscience Program, and the Portuguese National  
340 Authority for Animal Health (data in **Fig. 1b, Supplemental Fig. 2e**). Individual Myomatrix threads were implanted in  
341 the triceps muscle using the “intramuscular” method described above under isoflurane anesthesia (1-4% at flow rate 1  
342 L/min). EMG data were then recorded either during home cage exploration or while animals walked on a custom-built  
343 linear treadmill<sup>50</sup> at speeds ranging from 15-25 cm/sec. A 45° angled mirror below the treadmill allowed simultaneous  
344 side and bottom views of the mouse<sup>6</sup> using a single monochrome usb3 camera (Grasshopper3, Teledyne FLIR) to  
345 collect images 330 frames per second. We used DeepLabCut<sup>51</sup> to track paw, limb, and body positions. These tracked  
346 points were used to identify the stride cycles of each limb, defining stance onset as the time at which each paw  
347 contacts the ground and swing onset as the time when each paw leaves the ground.

#### 348 349 Additional recording methods – mouse orofacial muscle

350 All procedures described below were approved by The Institutional Animal Care and Use Committee at Johns  
351 Hopkins University. Individual Myomatrix threads were implanted on the masseter muscle using the “epimysial”  
352 method described above. A ground pin was placed over the right visual cortex. As described previously<sup>52</sup>, EMG  
353 signals and high-speed video of the orofacial area were recorded simultaneously in head-fixed animals under  
354 isoflurane anesthesia (0.9-1.5% at flow rate 1L/min). During data collection, the experimenter used a thin wooden  
355 dowel to gently displace the mandible to measure both jaw displacement and muscle activity from the jaw jerk reflex.  
356 Jaw kinematics were quantified using a high-speed camera (PhotonFocus DR1-D1312-200-G2-8) at 400 frames per  
357 second using an angled mirror to collect side and bottom views simultaneously. Jaw displacement was quantified by  
358 tracking eleven keypoints along the jaw using DeepLabCut<sup>51</sup>.

#### 359 360 Additional recording methods – rat forelimb muscle

361 All procedures described below were approved by The Institutional Animal Care and Use Committee at Emory  
362 University. Anesthesia was induced with an initial dose of 4% isoflurane in oxygen provided in an induction chamber  
363 with 2 L/minute rate and maintained with 3% isoflurane at 1 L/minute. Following this, rats received a subcutaneous  
364 injection of 1mg/kg Meloxicam, a subcutaneous injection of 1% Lidocaine and topical application of lidocaine ointment  
365 (5%) at each incision site. Myomatrix threads were implanted in the triceps muscle using the “intramuscular” method.  
366 EMG data were then recorded while animals walked on a treadmill at speeds ranging from 8-25 cm/sec. Kinematics  
367 were quantified using a circular arrangement of four high-speed FLIR Black Fly S USB3 cameras (BFS-U3-16S2M-  
368 CS, Mono), each running at 125 FPS. We used DeepLabCut to label pixel locations of each of ten anatomical  
369 landmarks on the limbs and body, which we then transformed into 3D cartesian coordinates using Anipose<sup>51,53</sup>. We  
370 then defined the onset of each swing/stance cycle by using local minima in the rat's forelimb endpoint position along  
371 the direction of locomotion.

372

#### 373 Additional recording methods – rat forelimb muscle

374 All procedures described below were approved by The Institutional Animal Care and Use Committee at Johns  
375 Hopkins University. Prior to electrode implantation, rodents were trained for 4-6 weeks to perform a single pellet reach  
376 task<sup>54</sup>. Rodents were food restricted for 17-18 hours prior to training. During the task, rats used the right arm to reach  
377 for sucrose pellets through a vertical slit (width = 1 cm) in a custom-built acrylic chamber. Individual MyoMatrix  
378 threads were implanted on the right flexor digitorum profundus (FDP) muscle using the “epimysial” method described  
379 above under 2.0-3.0% isoflurane anesthesia in oxygen gas. The arrays were then connected to data collection  
380 hardware via a flexible tether and EMG data were recorded while unrestrained animals performed the reaching task.  
381 Kinematics were quantified using a smartphone camera running at 60 fps. Two raters then manually labelled the  
382 frames of grasp onset. Grasp initiation was defined when a frame of full digit extension was immediately followed by a  
383 frame of digit flexion.

384

#### 385 Additional recording methods – mouse pelvic muscle

386 All experimental procedures were carried out in accordance with the guidelines of the European Committee Council  
387 Directive and were approved by the Animal Care and Users Committee of the Champalimaud Neuroscience Program,  
388 the Portuguese National Authority for Animal Health. As described in detail elsewhere<sup>3</sup>, spinal optogenetic stimulation  
389 of the motor neurons innervating the bulbospongiosus muscle (BSM) was performed on 2-3 month old male BL6  
390 mice that had received an injection of a rAAVCAG-ChR2 into the BSM on postnatal day 3-6. Individual Myomatrix  
391 threads were implanted in the BSM using the “intramuscular” method described above. During EMG recording an  
392 optrode was moved on top of the spinal cord along the rostral-caudal axis while applying optical stimulation pulses (10  
393 msec duration, power 1-15 mW).

394

#### 395 Additional recording methods – songbird vocal and respiratory muscles

396 All procedures described below were approved by The Institutional Animal Care and Use Committee at Emory  
397 University. As described previously<sup>20,27,42</sup>, adult male Bengalese finches (>90 d old) were anesthetized using  
398 intramuscular injections of 40 mg/kg ketamine and 3 mg/kg midazolam injected and anesthesia was maintained using  
399 1-5% isoflurane in oxygen gas. To record from the expiratory (respiratory) muscles, an incision was made dorsal to  
400 the leg attachment and rostral to the pubic bone and the electrode array was placed on the muscle surface using the  
401 “epimysial” approach described above. To record from syringeal (vocal) muscles, the vocal organ was accessed for  
402 electrode implantation via a midline incision into the intraclavicular air sac as described previously<sup>20</sup> to provide access  
403 to the ventral syringeal (VS) muscle located on the ventral portion of the syrinx near the midline.

404

#### 405 Additional recording methods – cat soleus muscle

406 All procedures described below were approved by The Institutional Animal Care and Use Committee at Temple  
407 University. As described previously<sup>55</sup>, an adult female cat was provided atropine (0.05 mg/kg IM) and anesthetized  
408 with isoflurane (1.5–3.5% in oxygen), during which a series of surgical procedures were performed including L3  
409 laminectomy, implantation of nerve cuffs on the tibial and sural nerve, and isolation of hindlimb muscles. Individual  
410 Myomatrix threads were implanted in hindlimb muscles using the “intramuscular” method described above. Following  
411 these procedures, a precollicular decerebration was performed and isoflurane was discontinued. Following a recovery  
412 period, the activity of hindlimb motor units were recorded in response to electrical stimulation of either the  
413 contralateral tibial nerve or the ipsilateral sural nerve.

414

#### 415 Additional recording methods – hawkmoth larva (caterpillar) body wall muscle

416 EMG recordings were obtained from 5th instar larvae of the tobacco hornworm *Manduca sexta* using a semi-intact  
417 preparation called the “flaterpillar” as described previously (Metallo, White, and Trimmer 2011). Briefly, after chilling  
418 animals on ice for 30 minutes, an incision was made along the cuticle, allowing the nerve cord and musculature to be  
419 exposed and pinned down in a Sylgard dish under cold saline solution. This preparation yields spontaneous muscle  
420 activity (fictive locomotion) which was recorded from the dorsal intermediate medial (DIM) muscle using the  
421 “epimysial” method described above, with the modification that sutures were not used to hold the muscle in place.

422

#### 423 Additional recording methods – frog hindlimb muscles

424 Spinal bullfrogs were prepared under anesthesia in accordance with USDA and PHS guidelines and regulations  
425 following approval from the Institutional Animal Care and Use Committee at Drexel University as described previously  
426<sup>56</sup>. Bull frogs were anesthetized with 5% tricaine (MS-222, Sigma), spinalized, and decerebrated. The frog was placed  
427 on a support and Myomatrix arrays were implanted into the semimembranosus (SM) hindlimb muscle using the  
428 “intramuscular” method described above. Epidermal electrical stimulation at the heel dorsum (500 msec train of 1  
429 msec, 5V biphasic pulses delivered at 40 Hz) or foot pinch was used to evoke reflexive motor activity.

430

#### 431 Additional recording methods – rhesus macaque forelimb muscle

432 All procedures described below were approved by The Institutional Animal Care and Use Committee at Western  
433 University. One male rhesus monkey (Monkey M, *Macaca mulatta*, 10 kg) was trained to perform a range of reaching  
434 tasks while seated in a robotic exoskeleton (NHP KINARM, Kingston, Ontario). As described previously<sup>57,58</sup>, this  
435 robotic device allows movements of the shoulder and elbow joints in the horizontal plane and can independently apply  
436 torque at both joints. Visual cues and hand feedback were projected from an LCD monitor onto a semi-silvered mirror  
437 in the horizontal plane of the task and direct vision of the arm was blocked with a physical barrier.  
438

439 An injectable Myomatrix array (**Supplemental Fig. 1g**) was inserted percutaneously as shown in **Supplemental**  
440 **Figure 1i**. Then, using his right arm, Monkey M performed a reaching task similar to previous work<sup>57</sup>. On each trial  
441 the monkey waited in a central target (located under the fingertip when the shoulder and elbow angles were 32° and  
442 72°, respectively; size = 0.6 cm diameter) while countering a constant elbow load (-0.05 Nm). The monkey was  
443 presented with one of two peripheral goal targets (30/84° and 34/60° shoulder/elbow, 8cm diameter), and after a  
444 variable delay (1.2-2s) received one of two unpredictable elbow perturbations ( $\pm 0.15$ Nm step-torque) which served as  
445 a go cue to reach to the goal target. At the time of perturbation onset, all visual feedback was frozen until the hand  
446 remained in the goal target for 800ms, after which a juice reward was given. On 10% of trials no perturbation was  
447 applied, and the monkey had to maintain the hand in the central target. In addition to Myomatrix injectables, we  
448 acquired bipolar electromyographic activity from nonhuman primates using intramuscular fine-wire electrodes in the  
449 biceps brachii long head as described previously<sup>59</sup>, recording in this instance from the same biceps muscle in the  
450 same animal from which we also collected Myomatrix data, although in a separate recording session. Fine-wire  
451 electrodes were spaced ~8 mm apart and aligned to the muscle fibers, and a reference electrode was inserted  
452 subcutaneously in the animal's back. Muscle activity was recorded at 2,000 Hz, zero-phase bandpass filtered (25–500  
453 Hz, fourth order Butterworth) and full-wave rectified.  
454

#### 455 **Data and code availability:**

456 A data archive including two EMG datasets recorded with Myomatrix arrays from behaving animals is available at  
457 [doi.org/10.5061/dryad.66t1g1k70](https://doi.org/10.5061/dryad.66t1g1k70). This archive includes the mouse triceps data shown in **Figure 1**  
458 **(b,d)** and **Supplemental Figure 2(a,b,e,f)**, the rhesus macaque biceps data shown in **Figure 3** and **Supplemental**  
459 **Figure 2g**, and associated metadata.  
460


## REFERENCES

- 461 1 Miri, A. *et al.* Behaviorally Selective Engagement of Short-Latency Effector Pathways by Motor Cortex. *Neuron* **95**, 683-696 e611 (2017). <https://doi.org/10.1016/j.neuron.2017.06.042>
- 462
- 463 2 Pachitariu, M., Sridhar, S. & Stringer, C. Solving the spike sorting problem with Kilosort. *bioRxiv*,  
464 2023.2001.2007.523036 (2023). <https://doi.org/10.1101/2023.01.07.523036>
- 465 3 Lenschow, C. *et al.* A galanin-positive population of lumbar spinal cord neurons modulates sexual behavior  
466 and arousal. *bioRxiv*, 2022.2010.2004.510783 (2022). <https://doi.org/10.1101/2022.10.04.510783>
- 467 4 Steinmetz, N. A., Koch, C., Harris, K. D. & Carandini, M. Challenges and opportunities for large-scale  
468 electrophysiology with Neuropixels probes. *Curr Opin Neurobiol* **50**, 92-100 (2018).  
469 <https://doi.org/10.1016/j.conb.2018.01.009>
- 470 5 Urai, A. E., Doiron, B., Leifer, A. M. & Churchland, A. K. Large-scale neural recordings call for new insights to  
471 link brain and behavior. *Nat Neurosci* **25**, 11-19 (2022). <https://doi.org/10.1038/s41593-021-00980-9>
- 472 6 Machado, A. S., Darmohray, D. M., Fayad, J., Marques, H. G. & Carey, M. R. A quantitative framework for  
473 whole-body coordination reveals specific deficits in freely walking ataxic mice. *Elife* **4** (2015).  
474 <https://doi.org/10.7554/eLife.07892>
- 475 7 Berman, G. J., Choi, D. M., Bialek, W. & Shaevitz, J. W. Mapping the stereotyped behaviour of freely moving  
476 fruit flies. *J R Soc Interface* **11** (2014). <https://doi.org/10.1098/rsif.2014.0672>
- 477 8 Pereira, T. D. *et al.* Fast animal pose estimation using deep neural networks. *Nat Methods* **16**, 117-125  
478 (2019). <https://doi.org/10.1038/s41592-018-0234-5>
- 479 9 Mathis, A., Schneider, S., Lauer, J. & Mathis, M. W. A Primer on Motion Capture with Deep Learning:  
480 Principles, Pitfalls, and Perspectives. *Neuron* **108**, 44-65 (2020). <https://doi.org/10.1016/j.neuron.2020.09.017>
- 481 10 Wiltschko, A. B. *et al.* Revealing the structure of pharmacobehavioral space through motion sequencing. *Nat*  
482 *Neurosci* **23**, 1433-1443 (2020). <https://doi.org/10.1038/s41593-020-00706-3>
- 483 11 Hernandez, D. G., Sober, S. J. & Nemenman, I. Unsupervised Bayesian Ising Approximation for decoding  
484 neural activity and other biological dictionaries. *Elife* **11** (2022). <https://doi.org/10.7554/eLife.68192>
- 485 12 Vyas, S., Golub, M. D., Sussillo, D. & Shenoy, K. V. Computation Through Neural Population Dynamics. *Annu*  
486 *Rev Neurosci* **43**, 249-275 (2020). <https://doi.org/10.1146/annurev-neuro-092619-094115>
- 487 13 Manuel, M., Chardon, M., Tysseling, V. & Heckman, C. J. Scaling of Motor Output, From Mouse to Humans.  
488 *Physiology (Bethesda)* **34**, 5-13 (2019). <https://doi.org/10.1152/physiol.00021.2018>
- 489 14 Loeb, G. E. & Gans, C. *Electromyography for Experimentalists, First ed.* (The University of Chicago Press,  
490 1986).
- 491 15 Bracklein, M. *et al.* The control and training of single motor units in isometric tasks are constrained by a  
492 common input signal. *Elife* **11** (2022). <https://doi.org/10.7554/eLife.72871>
- 493 16 Farina, D. & Holobar, A. Characterization of Human Motor Units From Surface EMG Decomposition. *IEEE*  
494 *Proceedings* **104**, 353-373 (2016).
- 495 17 Marshall, N. J. *et al.* Flexible neural control of motor units. *Nat Neurosci* **25**, 1492-1504 (2022).  
496 <https://doi.org/10.1038/s41593-022-01165-8>
- 497 18 Negro, F., Muceli, S., Castronovo, A. M., Holobar, A. & Farina, D. Multi-channel intramuscular and surface  
498 EMG decomposition by convolutive blind source separation. *J Neural Eng* **13**, 026027 (2016).  
499 <https://doi.org/10.1088/1741-2560/13/2/026027>
- 500 19 Pearson, K. G., Acharya, H. & Fouad, K. A new electrode configuration for recording electromyographic  
501 activity in behaving mice. *J Neurosci Methods* **148**, 36-42 (2005).  
502 <https://doi.org/10.1016/j.jneumeth.2005.04.006>
- 503 20 Srivastava, K. H., Elemans, C. P. & Sober, S. J. Multifunctional and Context-Dependent Control of Vocal  
504 Acoustics by Individual Muscles. *J Neurosci* **35**, 14183-14194 (2015).  
505 <https://doi.org/10.1523/JNEUROSCI.3610-14.2015>
- 506 21 Pack, A. R., Yan, J. S., Pasquali, M., Sober, S. J. & Elemans, C. P. H. A flexible carbon nanotube electrode  
507 array for acute in vivo EMG recordings. *J Neurophysiol* (2023). <https://doi.org/10.1152/jn.00262.2022>
- 508 22 Sober, S. J., Wohlgemuth, M. J. & Brainard, M. S. Central contributions to acoustic variation in birdsong. *J*  
509 *Neurosci* **28**, 10370-10379 (2008). <https://doi.org/10.1523/JNEUROSCI.2448-08.2008>
- 510 23 Mathewson, M. A., Chapman, M. A., Hentzen, E. R., Friden, J. & Lieber, R. L. Anatomical, architectural, and  
511 biochemical diversity of the murine forelimb muscles. *J Anat* **221**, 443-451 (2012).  
512 <https://doi.org/10.1111/j.1469-7580.2012.01559.x>
- 513 24 Tysseling, V. M. *et al.* Design and evaluation of a chronic EMG multichannel detection system for long-term  
514 recordings of hindlimb muscles in behaving mice. *J Electromyogr Kinesiol* **23**, 531-539 (2013).  
515 <https://doi.org/10.1016/j.jelekin.2012.11.014>
- 516 25 Ritter, L. K., Tresch, M. C., Heckman, C. J., Manuel, M. & Tysseling, V. M. Characterization of motor units in  
517 behaving adult mice shows a wide primary range. *J Neurophysiol* **112**, 543-551 (2014).  
518 <https://doi.org/10.1152/jn.00108.2014>



- 519 26 Tysseling, V. M. *et al.* Constitutive activity of 5-HT<sub>2C</sub> receptors is present after incomplete spinal cord injury but is not modified after chronic SSRI or baclofen treatment. *J Neurophysiol* **118**, 2944-2952 (2017).  
520 <https://doi.org/10.1152/jn.00190.2017>
- 521 27 Zia, M., Chung, B., Sober, S. & Bakir, M. S. Flexible Multielectrode Arrays With 2-D and 3-D Contacts for In  
522 Vivo Electromyography Recording. *IEEE Trans Compon Packaging Manuf Technol* **10**, 197-202 (2020).  
523 <https://doi.org/10.1109/tcpmt.2019.2963556>
- 524 28 Sober, S. J., Sponberg, S., Nemenman, I. & Ting, L. H. Millisecond Spike Timing Codes for Motor Control.  
525 *Trends Neurosci* **41**, 644-648 (2018). <https://doi.org/10.1016/j.tins.2018.08.010>
- 526 29 Srivastava, K. H. *et al.* Motor control by precisely timed spike patterns. *Proc Natl Acad Sci U S A* **114**, 1171-  
527 1176 (2017). <https://doi.org/10.1073/pnas.1611734114>
- 528 30 Quinlan, K. A. *et al.* Chronic electromyograms in treadmill running SOD1 mice reveal early changes in muscle  
529 activation. *J Physiol* **595**, 5387-5400 (2017). <https://doi.org/10.1113/JP274170>
- 530 31 Marshall, N. J. *et al.* Flexible neural control of motor units. *bioRxiv*, 2021.2005.2005.442653 (2021).  
531 <https://doi.org/10.1101/2021.05.05.442653>
- 532 32 Farina, D. & Merletti, R. Comparison of algorithms for estimation of EMG variables during voluntary isometric  
533 contractions. *J Electromyogr Kinesiol* **10**, 337-349 (2000). [https://doi.org/10.1016/s1050-6411\(00\)00025-0](https://doi.org/10.1016/s1050-6411(00)00025-0)
- 534 33 Farina, D., Yoshida, K., Stieglitz, T. & Koch, K. P. Multichannel thin-film electrode for intramuscular  
535 electromyographic recordings. *J Appl Physiol* (1985) **104**, 821-827 (2008).  
536 <https://doi.org/10.1152/jappphysiol.00788.2007>
- 537 34 Muceli, S. *et al.* Accurate and representative decoding of the neural drive to muscles in humans with multi-  
538 channel intramuscular thin-film electrodes. *J Physiol* **593**, 3789-3804 (2015).  
539 <https://doi.org/10.1113/JP270902>
- 540 35 Hoffer, J. A., O'Donovan, M. J., Pratt, C. A. & Loeb, G. E. Discharge patterns of hindlimb motoneurons during  
541 normal cat locomotion. *Science* **213**, 466-467 (1981). <https://doi.org/10.1126/science.7244644>
- 542 36 Robinson, D. A. Oculomotor unit behavior in the monkey. *J Neurophysiol* **33**, 393-403 (1970).  
543 <https://doi.org/10.1152/jn.1970.33.3.393>
- 544 37 Hyngstrom, A. S., Johnson, M. D., Miller, J. F. & Heckman, C. J. Intrinsic electrical properties of spinal  
545 motoneurons vary with joint angle. *Nat Neurosci* **10**, 363-369 (2007). <https://doi.org/10.1038/nn1852>
- 546 38 Pflugger, H. J. B., M. Locusts use the same basic motor pattern in swimming as in jumping and kicking. *Journal*  
547 *of experimental biology* **75**, 81-93 (1978).
- 548 39 Putney, J., Niebur, T., Wood, L., Conn, R. & Sponberg, S. An information theoretic method to resolve  
549 millisecond-scale spike timing precision in a comprehensive motor program. *PLOS Computational Biology* **19**,  
550 e1011170 (2023).
- 551 40 Adam, I. *et al.* One-to-one innervation of vocal muscles allows precise control of birdsong. *Curr Biol* **31**, 3115-  
552 3124 e3115 (2021). <https://doi.org/10.1016/j.cub.2021.05.008>
- 553 41 Lu, J. *et al.* High-performance Flexible Microelectrode Array with PEDOT:PSS Coated 3D Micro-cones for  
554 Electromyographic Recording. *Annu Int Conf IEEE Eng Med Biol Soc* **2022**, 5111-5114 (2022).  
555 <https://doi.org/10.1109/EMBC48229.2022.9871052>
- 556 42 Zia, M., Chung, B., Sober, S. J. & Bakir, M. S. Fabrication and Characterization of 3D Multi-Electrode Array  
557 on Flexible Substrate for In Vivo EMG Recording from Expiratory Muscle of Songbird. *Tech Dig Int Electron*  
558 *Devices Meet* **2018**, 29 24 21-29 24 24 (2018). <https://doi.org/10.1109/IEDM.2018.8614503>
- 559 43 Cui, X. Y. & Martin, D. C. Electrochemical deposition and characterization of poly(3,4-  
560 ethylenedioxythiophene) on neural microelectrode arrays. *Sensor Actuat B-Chem* **89**, 92-102 (2003).  
561 [https://doi.org/10.1016/S0925-4005\(02\)00448-3](https://doi.org/10.1016/S0925-4005(02)00448-3)
- 562 44 Dijk, G., Rutz, A. L. & Malliaras, G. G. Stability of PEDOT:PSS-Coated Gold Electrodes in Cell Culture  
563 Conditions. *Adv Mater Technol-Us* **5** (2020). <https://doi.org/10.1002/admt.201900662>
- 564 45 Rossetti, N. *et al.* Poly(3,4-ethylenedioxythiophene) (PEDOT) Coatings for High-Quality Electromyography  
565 Recording. *Acs Appl Bio Mater* **2**, 5154-5163 (2019). <https://doi.org/10.1021/acsabm.9b00809>
- 566 46 Ludwig, K. A. *et al.* Poly(3,4-ethylenedioxythiophene) (PEDOT) polymer coatings facilitate smaller neural  
567 recording electrodes. *J Neural Eng* **8**, 014001 (2011). <https://doi.org/10.1088/1741-2560/8/1/014001>
- 568 47 Steinmetz, N. A. *et al.* Neuropixels 2.0: A miniaturized high-density probe for stable, long-term brain  
569 recordings. *Science* **372** (2021). <https://doi.org/10.1126/science.abf4588>
- 570 48 van den Doel, K., Ascher, U. M. & Pai, D. K. Computed myography: three-dimensional reconstruction of motor  
571 functions from surface EMG data. *Inverse Problems* **24**, 065010 (2008).
- 572 49 Michel, C. M. *et al.* EEG source imaging. *Clin Neurophysiol* **115**, 2195-2222 (2004).  
573 <https://doi.org/10.1016/j.clinph.2004.06.001>
- 574 50 Darmohray, D. M., Jacobs, J. R., Marques, H. G. & Carey, M. R. Spatial and Temporal Locomotor Learning in  
575 Mouse Cerebellum. *Neuron* **102**, 217-231 e214 (2019). <https://doi.org/10.1016/j.neuron.2019.01.038>
- 576 51 Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat*  
577 *Neurosci* **21**, 1281-1289 (2018). <https://doi.org/10.1038/s41593-018-0209-y>
- 578

- 579 52 Severson, K. S. *et al.* Active Touch and Self-Motion Encoding by Merkel Cell-Associated Afferents. *Neuron*  
580 **94**, 666-676 e669 (2017). <https://doi.org/10.1016/j.neuron.2017.03.045>
- 581 53 Karashchuk, P. *et al.* Anipose: A toolkit for robust markerless 3D pose estimation. *Cell Rep* **36**, 109730  
582 (2021). <https://doi.org/10.1016/j.celrep.2021.109730>
- 583 54 Whishaw, I. Q., Pellis, S. M., Gorny, B., Kolb, B. & Tetzlaff, W. Proximal and distal impairments in rat forelimb  
584 use in reaching follow unilateral pyramidal tract lesions. *Behav Brain Res* **56**, 59-76 (1993).  
585 [https://doi.org/10.1016/0166-4328\(93\)90022-i](https://doi.org/10.1016/0166-4328(93)90022-i)
- 586 55 Zaback, M. *et al.* Toward Assessing the Functional Connectivity of Spinal Neurons. *Front Neural Circuits* **16**,  
587 839521 (2022). <https://doi.org/10.3389/fncir.2022.839521>
- 588 56 Kim, T. *et al.* Highly Flexible Precisely Braided Multielectrode Probes and Combinatorics for Future  
589 Neuroprostheses. *Front Neurosci* **13**, 613 (2019). <https://doi.org/10.3389/fnins.2019.00613>
- 590 57 Pruszynski, J. A., Omrani, M. & Scott, S. H. Goal-dependent modulation of fast feedback responses in  
591 primary motor cortex. *J Neurosci* **34**, 4608-4617 (2014). <https://doi.org/10.1523/JNEUROSCI.4520-13.2014>
- 592 58 Scott, S. H. Apparatus for measuring and perturbing shoulder and elbow joint positions and torques during  
593 reaching. *J Neurosci Methods* **89**, 119-127 (1999). [https://doi.org/10.1016/s0165-0270\(99\)00053-9](https://doi.org/10.1016/s0165-0270(99)00053-9)
- 594 59 Maeda, R. S., Kersten, R. & Pruszynski, J. A. Shared internal models for feedforward and feedback control of  
595 arm dynamics in non-human primates. *Eur J Neurosci* **53**, 1605-1620 (2021).  
596 <https://doi.org/10.1111/ejn.15056>




597

598 **Supplemental Figure 1: Myomatrix fabrication, design variations, and implantation.**

599 (a) Workflow for electrode array fabrication. Layers of insulating polymer (polyimide) and conductive metal (gold) are successively  
600 deposited on a carrier wafer to form a flexible, 20 or 40 μm thick electrode array of gold electrode contacts, which receive a surface  
601 treatment of PEDOT to improve recording properties (see Methods). Electrodes are connected via thin gold traces to a receiving  
602 pad for a high-density connector (Omnetics Inc.) which is then bonded to the array. The completed array is then peeled off the

603 carrier wafer. **(b)** Photo showing two different Myomatrix designs (left) as well as “blank” arrays comprised only of the flexible  
604 polyimide substrate for surgical practice and design optimization. **(c-e)** Expanded views of the electrode array also shown in **Figure**  
605 **1** of the main manuscript text, which has four “threads” each bearing eight electrode contacts. This array design can be used for  
606 either acute or chronic recordings. For chronic implantation, the surgeon grasps the “pull-through tabs” when tunneling the threads  
607 subcutaneously. For intramuscular implantation in either acute or chronic settings, a needle is used to pull each thread through the  
608 target muscle. In this use case, the “depth-restrictor tabs” prevent the thread from being pulled any further into the muscle, thereby  
609 determining the depth of the electrode contacts within the muscle. **(d,e)** Detail views highlighting sub-millimeter features used to  
610 increase electrode stability within the muscle (barbs, suture holes) and labels to indicate which channel/thread labels have been  
611 implanted in which muscles. **(f,g)** Design variations. The fabrication process shown in **(a)** can easily be modified to alter size and  
612 shape of the electrode array. Each Myomatrix design in **f** and **g** has 32 electrode contacts. **(f)** Two array designs customized for  
613 chronic recording applications in different muscle groups in rodents. **(g)** Injectable array for recording forelimb muscles in  
614 nonhuman primates. **(h)** For chronic implantation in mice, the connector end of the array is attached to the skull using dental acrylic  
615 **(1)** and the flexible array threads are then routed subcutaneously to a small distal incision located near the targeted muscle or  
616 muscles **(2)**. For intramuscular implantation, the surgeon secures each thread to a suture and needle, which are then inserted through  
617 the target area of muscle tissue **(3)**. The surgeon then pulls the suture further through the muscle, eventually drawing the array  
618 thread into the muscles such that the depth-restrictor tabs prevent further insertion and ensure that the electrode contacts are  
619 positioned at the correct depth within the muscle **(4)**. In contrast, for epimysial (as opposed to intramuscular) implantation, the array  
620 threads are sutured to the surface of the muscle fascia rather than being inserted with a suture needle. After all threads are secured  
621 to the muscle, the distal incision site is sutured closed **(5)**. **(i)** For percutaneous insertion of injectable arrays, the array’s thin “tail”  
622 is loaded into a modified hypodermic syringe <sup>14,34</sup>. During insertion, the tail is secured by bending it back over the plastic needle  
623 holder and securing it with either the surgeon’s fingers or an additional syringe inserted into the cannula. After electrode array  
624 insertion the needle is gently pulled out of the muscle, leaving the electrode-bearing part of the array thread within the target muscle  
625 for the duration of the recording session. After recording, the electrode and tail are gently pulled out of the muscle together as with  
626 injectable fine-wire EMG <sup>14</sup>.




627

## 628 Supplemental Figure 2: Spike sorting

629 Action potential (“spike”) waveforms from individual motor units can be identified (“sorted”) using analysis methods that are  
 630 commonly used to sort spikes from neural data. **(a-d)** Single channel spike sorting. In some cases, a single motor unit’s spike will

631 dominate the recording on an individual Myomatrix channel, as shown in an example bipolar recording from mouse triceps during  
632 locomotion (**a**, top). In such cases, a simple voltage threshold (dashed line) can be used to isolate spike times of the largest  
633 recorded unit (blue dots) from a single channel. In contrast (**a**, bottom), fine-wire EMG typically does not yield isolated single  
634 units during active behaviors. (**b**) Single-channel spike sorting using principal components analysis (PCA) of the data shown in  
635 (**a**). Each data point in (**b**) represents a single voltage waveform represented in the dimensions defined by the first two principal  
636 components (PC1 and PC2) of the set of all spike waveforms. As described previously<sup>22</sup>, k-means clustering can discriminate the  
637 waveforms from individual motor units (cyan dots in **a** and **b**) and waveforms from other motor units and/or background noise  
638 (black dots in **b**). If one of the clusters has less than 1% overlap with any other cluster (based on fitting each cluster with a 2D  
639 Gaussian as described previously) and displays an absolute refractory period (less than 1% of inter-spike intervals less than 1  
640 msec), it is classified as a single unit<sup>22</sup>. When applied to the Myomatrix data in (**a**), PCA-based sorting method produced identical  
641 spike times as the thresholding method (cyan dots in **a**). In contrast, the same analysis applied to the fine-wire data shown in **a** did  
642 not produce any well-isolated clusters in PCA space (**b**, right), indicating that this method could not extract any single motor  
643 units. Myomatrix and fine-wire data shown in (**a**,**b**) are from the same datasets as the examples shown in main text Figure 1a,b.  
644 (**c,d**) Single-channel spike sorting applied to bipolar Myomatrix recordings from the ventral syringeal (VS) muscle, a songbird  
645 vocal muscle<sup>20</sup>. Here again, PCA-based sorting of Myomatrix data method produced identical spike times as the thresholding  
646 method (orange dots in **c** and **d**). In contrast, the same analysis applied to fine-wire data recorded from VS shown in **c** did not  
647 produce any well-isolated clusters in PCA space (**d**, right), Other plotting conventions for (**c**,**d**) are the same as for the mouse data  
648 in (**a**,**b**).  
649  
650 (**e-h**) Multichannel spike sorting using Kilosort. We used Kilosort version 2.5<sup>2,47</sup> and custom MATLAB and Python code to sort  
651 waveforms into clusters arising from individual motor units. (**e**) Spike times (top) and mean waveforms (bottom) of six motor  
652 units recorded simultaneously from mouse triceps during locomotion (same dataset as **Fig. 1** in the main text). Mean waveforms  
653 for the six motor units (columns at bottom) are shown from six different EMG channels (rows) and illustrate the distinct pattern of  
654 spike waveforms across channel associated with the discharge of each identified motor unit. (**f**) Left, feature space projection of  
655 individual waveforms (colored dots), projected onto the space of singular values (“factors”) that describe the space of all recorded  
656 waveforms. The clustering of waveforms from kilosort-identified units (colors) further illustrates the distinctness of voltage  
657 waveforms assigned to each of the identified motor units. Right, autocorrelograms (colors) and cross correlograms (gray) of the  
658 six motor units shown in (**e**). In addition to examining the consistency of each candidate motor unit’s spike waveforms we also  
659 inspected autocorrelations to ensure that each identified unit showed an absolute refractory period (zero or near-zero  
660 autocorrelations at lag zero) and that cross-correlograms did not have strong peaks at zero lag (which might indicate the same  
661 motor unit being detected by multiple Kilosort clusters). (**g,h**) Myomatrix recordings from nonhuman primate and rat (unipolar  
662 and bipolar recordings respectively, same datasets as in main text **Fig. 3** and **Fig. 2c**), respectively. These examples (along with  
663 the mouse data in main text **Fig. 1c**) highlight the finding that Myomatrix arrays typically record the same motor unit on multiple  
664 channels simultaneously. This redundancy is critical for Kilosort and related methods to isolate single motor unit waveforms,  
665 particularly when waveforms from multiple units overlap in time.

666



667

### 668 Supplemental Fig 3: Longevity of Myomatrix recordings

669 In addition to isolating individual motor units, Myomatrix arrays also provide stable multi-unit recordings of comparable or  
 670 superior quality to conventional fine wire EMG. **(a,b)** Bipolar Myomatrix recordings from the triceps muscle of a mouse recorded  
 671 during treadmill locomotion over a period 61 days. Colored regions in **(b)** highlight the “stance” phase (when the paw from the  
 672 recorded forelimb is in contact with the treadmill surface) and “swing” phase (when the paw is lifted off the treadmill surface). To  
 673 quantify changes in recording quality over time, we computed a “signal-to-noise ratio (SNR)” for each of each stride cycle as  
 674 described previously<sup>21</sup>. Here, the “locomotor SNR” for each swing-stance-cycle is defined as the root mean square (RMS)  
 675 amplitude of the multi-unit EMG signal during each single stance cycle divided by the RMS of the EMG signal during the  
 676 immediately subsequent swing phase. **(c)** Fine-wire EMG data recorded from the triceps muscle during locomotion (reproduced  
 677 with permission from<sup>1</sup>). Note that all horizontal gray bars in **(b,c)** represent 100 msec. **(d)** Mean +/- standard error of locomotor  
 678 SNR across five mouse subjects implanted with Myomatrix arrays. Filled symbols indicate EMG implantation in the right triceps  
 679 muscle, unfilled symbols indicate EMG implantation in the left triceps. The black trace with unfilled symbols represents the  
 680 animal whose data are also shown in panel **(b)**. In some cases, error bars are hidden behind plotting symbols. Blue symbols  
 681 indicate the locomotor SNR from the fine-wire data from<sup>1</sup>, with each symbol representing a single day’s recording from one of  
 682 four individual mice. SNR values from Myomatrix arrays are significantly greater than those from fine-wire EMG, both when all  
 683 data shown in **(d)** are pooled and when only data from day 14 are included (2-sample KS-test,  $p=0.002$  and  $p=0.038$ ,  
 684 respectively). **(e)** Although individual motor units were most frequently recorded in the first two weeks of chronic recordings  
 685 (see main text), Myomatrix arrays also isolate individual motor units after much longer periods of chronic implantation, as shown  
 686 here where spikes from two individual motor units (colored boxes in bottom trace) were isolated during locomotion 65 days after  
 687 implantation. This bipolar recording was collected from the subject plotted with unfilled black symbols in panel **(d)**.

688

## **Acknowledgements:**

The participants of the Emory-SKAN Remote Workshop for Advanced EMG Methods, which brought together over 100 researchers from around the world, for their critical feedback on how to improve and refine the electrode technology described here.

Dr. Andrew Miri for helpful discussions and for sharing locomotor EMG data from Miri et al. (2017).

Dr. Cinzia Metallo for the initial *Manduca* studies using flexible electrode arrays.

Drs. Gabriela J. Martins and Mariana Correia for project and colony coordination.

Dr. Ana Gonçalves for technical assistance in mouse locomotion experiments.

Mattia Rigotti, Margo Shen, Nevin Aresh, and Manikandan Venkatesh for assistance in collecting rat forelimb data.

Components of all figures were created using BioRender.com.

## **Funding sources:**

Azim	National Institutes of Health (“NIH”) DP2NS105555, R01NS111479, RF1NS128898, U19NS112959 Searle Scholars Program Pew Charitable Trusts McKnight Foundation
Bakir	NIH R01NS109237, U24NS126936 McKnight Foundation
Carey	The Simons Foundation (as part of the Simons-Emory International Consortium on Motor Control) Portuguese Fundação para a Ciência e a Tecnologia (SFRH/BPD/119404/2016) to Hugo G. Marques. European Research Council Consolidator Grant #866237
Costa	The Simons Foundation (as part of the Simons-Emory International Consortium on Motor Control) NIH U19NS104649
Gatto	DFG SFB1451 iBehave Netzwerke 2021 Halle Institute for Global Research
Giszter	NIH R01NS104194, F31NS124347 PA Dept Health CURE award Coulter-Drexel Translational Research Partnership award
Goulding	NIH R01NS112959, R01NS111643
Lenschow	Human Frontier Science Program Postdoctoral Fellowship (LT000353/2018-L4) 2020 Marie Skłodowska-Curie Actions Individual Fellowship (799973)
Lima	Research infrastructure Congento, co-financed by the Lisboa Regional Operational Programme (Lisboa2020), under the PORTUGAL 2020 Partnership Agreement, through the European Regional Development Fund (ERDF) FCT under the project LISBOA-01-0145-FEDER-022170 InterEmerging Actions 2020 (301137) European Research Council (ERC) Consolidator Grant (772827)
Mendes	Fundação para a Ciência e a Tecnologia (FCT) PhD Fellowship (PD/BD141576/2018)
Michaels	Banting Postdoctoral Fellowship BrainsCAN (Canada First Research Excellence Fund) Postdoctoral Fellowship Vector Institute Postgraduate Affiliate Program
Mosberger	Swiss National Science Foundation P400PM_183904, P2EZIP3_172128, NIH K99NS126307
O’Connor	NIH R01NS104834, F32MH120873



Pandarinath	The Simons Foundation (as part of the Simons-Emory International Consortium on Motor Control) Emory Neuromodulation and Technology Innovation Center (ENTICe) National Science Foundation ("NSF") NCS 1835364 DARPA PA-18-02-04-INI-FP-021 NIH Eunice Kennedy Shriver NICHD K12HD073945 NIH-NINDS/OD DP2NS127291 Alfred P. Sloan Foundation
Person	The Simons Foundation (as part of the Simons-Emory International Consortium on Motor Control)
Pruszynski	The Simons Foundation (as part of the Simons-Emory International Consortium on Motor Control) Canadian Institutes of Health Research, Project Grant: PJT-175010 Canada Research Chair Program BrainsCAN (Canada First Research Excellence Fund) Natural Sciences and Engineering Research Council of Canada (RGPIN-2022-04421)
Sober	The Simons Foundation (as part of the Simons-Emory International Consortium on Motor Control) NIH R01NS084844, R01NS109237, R01NS099375, R01EB022872, U24NS126936 NSF 1822677 NSF DGE-1937971 (to Kyle Thomas) NSF DGE-1444932 (to Andrea Pack) McKnight Foundation Kavli Foundation Azrieli Foundation Novo Nordisk Foundation Halle Institute for Global Research
Thakor	NSF DGE-2139757 (to Kiara Quinn) Johns Hopkins Discovery Award
Thanawalla	Salk Alumni Fellowship Award
Thompson	NIH R01NS124820
Trimmer	NIH R34NS118412 NSF 1050908