



Results of the COVID-19 mental health international for the health professionals (COMET-HP) study: depression, suicidal tendencies and conspiracism

Konstantinos N. Fountoulakis¹ · Grigorios N. Karakatsoulis¹ · Seri Abraham^{2,3,4} · Kristina Adorjan⁵ · Helal Uddin Ahmed⁶ · Renato D. Alarcón^{7,8} · Kiyomi Arai⁹ · Sani Salihu Auwal^{10,11} · Julio Bobes^{15,16} · Teresa Bobes-Bascaran^{17,18} · Julie Bourgin-Duchesnay¹⁹ · Cristina Ana Bredicean²⁰ · Laurynas Bukelskis²¹ · Akaki Burkadze^{22,23} · Indira Indiana Cabrera Abud²⁴ · Ruby Castilla-Puentes²⁵ · Marcelo Cetkovich^{26,27} · Hector Colon-Rivera²⁸ · Ricardo Corral^{29,30} · Carla Cortez-Vergara³¹ · Piirika Crepin³² · Domenico de Berardis^{33,34,35} · Sergio Zamora Delgado³⁶ · David de Lucena³⁷ · Avinash de Sousa^{38,39} · Ramona di Stefano⁴⁰ · Seetal Dodd^{12,13,41} · Livia Priyanka Elek⁴² · Anna Elissa⁴³ · Berta Erdelyi-Hamza⁴² · Gamze Erzin⁴⁴ · Martin J. Etchevers⁴⁵ · Peter Falkai⁵ · Adriana Farcas⁴⁶ · Ilya Fedotov⁴⁷ · Viktoriia Filatova⁴⁸ · Nikolaos K. Fountoulakis⁴⁹ · Iryna Frankova⁵⁰ · Francesco Franza^{51,52} · Pedro Frias⁵³ · Tatiana Galako⁵⁴ · Cristian J. Garay⁴⁵ · Leticia Garcia-Álvarez¹⁸ · Paz García-Portilla^{15,55} · Xenia Gonda⁴² · Tomasz M. Gondek⁵⁶ · Daniela Morera González⁵⁷ · Hilary Gould⁵⁸ · Paolo Grandinetti³³ · Arturo Grau^{36,59} · Violeta Groudeva⁶⁰ · Michal Hagin⁶¹ · Takayuki Harada⁶² · Tasdik M. Hasan^{63,64} · Nurul Azreen Hashim⁶⁵ · Jan Hilbig²¹ · Sahadat Hossain⁶⁶ · Rossitza Iakimova⁶⁷ · Mona Ibrahim⁶⁸ · Felicia Iftene⁶⁹ · Yulia Ignatenko⁷⁰ · Matias Irrarazaval⁷¹ · Zaliha Ismail⁷² · Jamila Ismayilova⁷³ · Asaf Jacobs^{74,75} · Miro Jakovljević⁷⁶ · Nenad Jakšić¹⁴ · Afzal Javed^{77,78,79} · Helin Yilmaz Kafali⁸⁰ · Sagar Karia³⁸ · Olga Kazakova⁸¹ · Doaa Khalifa⁶⁸ · Olena Khaustova⁵⁰ · Steve Koh⁵⁸ · Svetlana Kopishinskaia^{82,83} · Korneliia Kosenko⁸⁴ · Sotirios A. Koupidis⁸⁵ · Illes Kovacs⁴² · Barbara Kulig⁴² · Alisha Lalljee³⁹ · Justine Liewig¹⁹ · Abdul Majid⁸⁶ · Evgeniia Malashonkova¹⁹ · Khamelia Malik⁴³ · Najma Iqbal Malik⁸⁷ · Gulay Mammadzada⁸⁸ · Bilvesh Mandalia³⁹ · Donatella Marazziti^{89,90,91} · Darko Marčinko^{14,76} · Stephanie Martinez⁵⁸ · Eimantas Matiekus²¹ · Gabriela Mejia⁵⁸ · Roha Saeed Memon⁹² · Xarah Elenne Meza Martínez⁹³ · Dalia Mickevičiūtė⁹⁴ · Roumen Milev⁶⁹ · Muftau Mohammed⁹⁵ · Alejandro Molina-López⁹⁶ · Petr Morozov⁹⁷ · Nuru Suleiman Muhammad⁹⁸ · Filip Mustač¹⁴ · Mika S. Naor⁹⁹ · Amira Nassieb⁶⁸ · Alvydas Navickas²¹ · Tarek Okasha⁶⁸ · Milena Pandova⁶⁷ · Anca-Livia Panfil¹⁰⁰ · Liliya Panteleeva¹⁰¹ · Ion Papava²⁰ · Mikaela E. Patsali^{102,103} · Alexey Pavlichenko⁷¹ · Bojana Pejuskovic^{104,105} · Mariana Pinto da Costa¹⁰⁶ · Mikhail Popkov¹⁰⁷ · Dina Popovic¹⁰⁸ · Nor Jannah Nasution Raduan⁶⁵ · Francisca Vargas Ramírez^{36,59} · Elmars Rancans^{109,110} · Salmi Razali⁶⁵ · Federico Rebok^{111,112} · Anna Rewekant¹¹³ · Elena Ninoska Reyes Flores¹¹⁴ · María Teresa Rivera-Encinas¹¹⁵ · Pilar A. Saiz¹⁵ · Manuel Sánchez de Carmona¹¹⁶ · David Saucedo Martínez¹¹⁷ · Jo Anne Saw⁶⁵ · Görkem Saygili¹¹⁸ · Patricia Schneiderit¹¹⁹ · Bhumika Shah¹²⁰ · Tomohiro Shirasaka¹²¹ · Ketevan Silagadze²² · Satti Sitanggang¹²² · Oleg Skugarevsky¹²³ · Anna Spikina¹²⁴ · Sridevi Sira Mahalingappa¹²⁵ · Maria Stoyanova⁶⁷ · Anna Szczegieliński¹²⁶ · Simona Claudia Tamasan¹⁰⁰ · Giuseppe Tavormina^{52,127,128} · Maurilio Giuseppe Maria Tavormina⁵² · Pavlos N. Theodorakis¹²⁹ · Mauricio Tohen¹³⁰ · Eva-Maria Tsapakis^{131,132} · Dina Tukhvatullina¹³³ · Irfan Ullah¹³⁴ · Ratnaraj Vaidya¹³⁵ · Johann M. Vega-Dienstmaier¹³⁶ · Jelena Vrublevska^{109,110,140} · Olivera Vukovic^{103,137} · Olga Vysotska¹³⁸ · Natalia Widiasih⁴³ · Anna Yashikhina^{82,139} · Panagiotis E. Prezerakos¹⁴⁰ · Michael Berk^{12,13} · Sarah Levaj¹⁴ · Daria Smirnova^{82,141}

Received: 11 September 2022 / Accepted: 2 February 2023

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany 2023

Abstract

Introduction The current study aimed to investigate the rates of anxiety, clinical depression, and suicidality and their changes in health professionals during the COVID-19 outbreak.

Extended author information available on the last page of the article

Materials and methods The data came from the larger COMET-G study. The study sample includes 12,792 health professionals from 40 countries (62.40% women aged 39.76 ± 11.70 ; 36.81% men aged 35.91 ± 11.00 and 0.78% non-binary gender aged 35.15 ± 13.03). Distress and clinical depression were identified with the use of a previously developed cut-off and algorithm, respectively.

Statistical analysis Descriptive statistics were calculated. Chi-square tests, multiple forward stepwise linear regression analyses, and Factorial Analysis of Variance (ANOVA) tested relations among variables.

Results Clinical depression was detected in 13.16% with male doctors and ‘non-binary genders’ having the lowest rates (7.89 and 5.88% respectively) and ‘non-binary gender’ nurses and administrative staff had the highest (37.50%); distress was present in 15.19%. A significant percentage reported a deterioration in mental state, family dynamics, and everyday lifestyle. Persons with a history of mental disorders had higher rates of current depression (24.64% vs. 9.62%; $p < 0.0001$). Suicidal tendencies were at least doubled in terms of RASS scores. Approximately one-third of participants were accepting (at least to a moderate degree) a non-bizarre conspiracy. The highest Relative Risk (RR) to develop clinical depression was associated with a history of Bipolar disorder (RR = 4.23).

Conclusions The current study reported findings in health care professionals similar in magnitude and quality to those reported earlier in the general population although rates of clinical depression, suicidal tendencies, and adherence to conspiracy theories were much lower. However, the general model of factors interplay seems to be the same and this could be of practical utility since many of these factors are modifiable.

Keywords COVID-19 · Health professionals · Depression · Suicidality · Mental health · Conspiracy theories · Mental disorders · Psychiatry · Anxiety

Introduction

There are many reports in the literature suggesting that health professionals are at particular risk to experience a deterioration of their mental health during the COVID-19 pandemic [23]. Clinical depression, sleep disorders, and post-traumatic stress disorder (PTSD) was also reported both in the general population as well as in health care professionals (HCP) [55, 97]. Thus, while the COVID-19 pandemic started as an epidemic of an infectious agent, it soon gained a wider content and included all effects on all aspects of human life by this condition, even the overwhelming burst of information of questionable reliability and validity (‘infodemic’) [10]. The abuse of the terms ‘trauma’ and ‘PTSD’ is such an example. The vast majority of studies reported a ‘tsunami’-scale impact on mental health. It is highly possible that this could be an exaggeration [96]. In addition, changes to social behavior, as well as working conditions, daily habits, and routines have imposed secondary stress. Higher levels of anxiety, stress, and depressive feelings have been reported, but it seems that this depends on the temporal situation and the specific events; the response is by no means homogenous [37, 85, 96, 115], [75, 104]. Apart from the effect of the virus itself, in addition, changes in social behavior, as well as in working conditions, daily habits, and routine are expected to impose further stress, especially with the expectation of an upcoming economic crisis and possible unemployment. In this frame, mental health has gained a central position as an area that is expected to be affected by

the pandemic because of its threatening nature as well as because of the profound impact on the everyday life of people. Especially concerning the later, it has been suggested that lockdowns triggered feelings of loneliness, irritableness, restlessness, and nervousness in the general population [90]. Especially the expectation of an upcoming economic crisis and possible unemployment were stressful factors. Conspiracy theories and maladaptive behaviors were also prevalent, compromising the public defense against the outbreak. The issue of increased suicidality as a consequence of extreme stress and depression has been raised again [25, 84].

At the end of the day, although there are several empirical data papers, their methodology varies, it is very difficult to make comparisons among countries and it is also difficult to arrive at universally valid conclusions [55, 97]. Additionally, the literature is full of opinion papers, viewpoints, perspectives, guidelines, and narrations of activities to cope with the pandemic. These borrow from previous experiences with different pandemics and utilize common sense, but, as a result, they often obscure rather than clarify the landscape. The role of mass and social media has been discussed but remains poorly understood in empirical terms.

An early meta-analysis reported high rates of anxiety (25%) and depression (28%) in the general population [86] while a second one reported that 29.6% of people experienced stress, 31.9% anxiety and 33.7% depression [92]. Not only do we need more reliable and valid data, but we also need to identify risk and protective factors to be able to recommend measures that will eventually improve public

health by preventing the adverse impact on mental health and simultaneously improve health-related behaviors [4, 47, 73, 75].

The **aim** of the current study was to investigate the rates of anxiety, clinical depression, and suicidality and their changes in health professionals aged 18–69 internationally, during the COVID-19 pandemic. The secondary aims were to investigate their relations with several personal, interpersonal/social, and lifestyle variables. The aim also included the investigation of the spreading of conspiracy theories concerning the COVID-19 outbreak and their relationship with mental health in this specific population group.

Materials and methods

Methods

The protocol used is available in the webappendix; each question was given an ID code; these ID codes were used throughout the results for increased accuracy.

According to a previously developed method, [39, 40, 42] the cut-off score of 23/24 for the Center for Epidemiological Studies-Depression (CES-D) scale and a derived algorithm were used to identify cases of clinical depression. This algorithm utilized the weighted scores of selected CES-D items to arrive at the diagnosis of clinical depression, and has already been validated. Cases identified by only either method were considered cases of distress (false positive cases in terms of depression), while cases identified by both the cut-off and the algorithm were considered as clinical depression. The State-Trait Anxiety Inventory-State form (STAI-S) [100] and the Risk for Assessment of Suicidality Scale (RASS) [42] were used to assess anxiety and suicidality respectively.

The data were collected online and anonymously from April 2020 through March 2021, covering periods of full implementation of lockdowns as well as of relaxations of measures in countries around the world. Announcements and advertisements were done on the social media and through news sites, but no other organized effort had been undertaken. The first page included a declaration of consent which everybody accepted by continuing with the participation.

Approval was initially given by the Ethics Committee of the Faculty of Medicine, Aristotle University of Thessaloniki, Greece, and locally concerning each participating country.

Materials

The data came from the larger COMET-G study [41].

The study population was self-selected. It was not possible to apply post-stratification on the sample as it was done

in a previous study [40], because this would mean that we would utilize a similar methodology across many different countries and the population data needed were not available for all.

Statistical analysis

- Chi-square tests were used for the comparison of frequencies when categorical variables were present and for the post hoc analysis of the results a Bonferroni-corrected method of pair-wise comparisons was utilized [69].
- Factorial Analysis of Variance (ANOVA) was used to test for the main effect as well as the interaction among grouping variables concerning continuous variables. The Scheffe test was used as the post-hoc test.
- Multiple forward stepwise linear regression analysis (MFSLRA) was performed to investigate which variables could function as predictors and contribute to the development of others (e.g. clinical depression).

To correct for multiple comparisons, the level of $p < 0.001$ was accepted as the level of significance for ANOVA and MFSLRA results (but not for post-hoc tests)

Results

Demographics

From the 55,589 responses from 40 countries (Table 1) of the COMET study, 23.01% reported they were working in the health field. Thus, the study sample of the current paper includes 12,792 health professionals ($N = 7983$ –62.40% women aged 39.76 ± 11.70 ; $N = 4709$ –36.81% men aged 35.91 ± 11.00 and $N = 100$ –0.78% non-binary gender aged 35.15 ± 13.03). The contribution of each country and the gender and age composition are shown in Table 1. The sex-by-specific occupation composition is shown in Table 2. The sociodemographic characteristics are shown in webtables 1,2,3,4,5,6,7 of the appendix. Details concerning various sociodemographic variables (marital status, education, work, etc. are shown in the webappendix, in webtables 1,2,3,4,5,6,7,8,9).

History of health

Moderate or bad somatic health was reported by 16.70% and the presence of a chronic medical somatic condition was reported by 21.29%. Detailed results are shown in webtable 8. Being either relatives or caretakers of vulnerable persons was reported by 46.88% (webtable 9).

Table 1 List of participating countries by sex, with number of subjects and mean age

Country	Men				Women				Non-binary gender					
	Age				Age				Age					
	<i>N</i>	%	Mean	SD	<i>N</i>	%	Mean	SD	<i>N</i>	%	Mean	SD	<i>N</i>	%
Argentina	439	20.14	44.53	14.39	1725	79.13	40.60	14.49	16	0.73	37.44	17.29	2180	3.92
Australia	21	30.43	33.67	8.05	48	69.57	32.63	7.89		0.00			69	0.12
Azerbaijan	70	19.89	36.20	10.33	280	79.55	37.71	11.46	2	0.57	26.00	0.00	352	0.63
Bangladesh	1681	55.42	24.09	5.24	1333	43.95	23.98	5.48	19	0.63	27.42	8.88	3033	5.46
Belarus	200	18.30	38.62	12.46	893	81.70	39.15	11.11		0.00			1093	1.97
Brazil	86	40.19	31.36	13.06	127	59.35	28.80	9.97	1	0.47	31.00		214	0.38
Bulgaria	202	26.47			558	73.13			3	0.39			763	1.37
Canada	142	27.73	42.24	15.49	367	71.68	42.57	14.00	3	0.59	46.33	17.79	512	0.92
Chile	86	26.71	40.76	15.43	234	72.67	39.57	15.08	2	0.62	42.50	16.26	322	0.58
Croatia	1041	35.91	41.73	11.70	1835	63.30	42.32	11.84	23	0.79	44.26	13.75	2899	5.22
Egypt	24	14.55	37.38	14.18	141	85.45	39.66	11.82		0.00			165	0.30
France	64	24.33	38.98	14.70	197	74.90	37.89	15.53	2	0.76	27.50	10.61	263	0.47
Georgia	48	11.59	30.77	6.82	364	87.92	32.06	9.04	2	0.48	33.50	6.36	414	0.74
Germany	15	25.00	48.93	18.58	45	75.00	34.87	13.98		0.00			60	0.11
Greece	624	18.26	36.55	10.58	2772	81.10	34.00	9.87	22	0.64	29.59	6.68	3418	6.15
Honduras	74	33.48	28.19	7.17	147	66.52	32.05	11.09		0.00			221	0.40
Hungary	146	19.13	44.60	11.95	617	80.87	41.36	11.95		0.00			763	1.37
India	3044	61.01	33.51	8.94	1917	38.42	31.59	11.97	28	0.56	28.36	7.86	4989	8.97
Indonesia	909	27.68	33.64	12.06	2358	71.80	30.49	11.42	17	0.52	28.00	11.62	3284	5.91
Israel	28	19.44	48.79	18.24	116	80.56	38.97	13.56		0.00			144	0.26
Italy	257	26.22	43.10	16.17	717	73.16	41.22	14.17	6	0.61	42.17	21.14	980	1.76
Japan	182	70.00	45.31	11.61	78	30.00	41.71	11.10		0.00			260	0.47
Kyrgyz Republic	614	27.76	36.38	14.16	1561	70.57	38.87	14.58	37	1.67	33.57	12.60	2212	3.98
Latvia	1036	39.72	48.18	12.38	1570	60.20	45.26	14.64	2	0.08	48.00	18.38	2608	4.69
Lithuania	271	21.54	39.34	13.62	983	78.14	40.16	12.75	4	0.32	40.75	12.89	1258	2.26
Malaysia	311	32.29	41.95	12.08	578	60.02	39.24	11.71	74	7.68	39.03	12.66	963	1.73
Mexico	447	25.03	36.84	16.13	1332	74.58	38.18	14.74	7	0.39	22.86	4.78	1786	3.21
Nigeria	752	65.22	30.30	7.46	397	34.43	25.83	7.55	4	0.35	31.75	7.97	1153	2.07
Pakistan	575	28.24	25.46	6.35	1445	70.97	23.45	4.42	16	0.79	24.75	10.93	2036	3.66
Peru	56	36.13	43.80	15.80	99	63.87	38.72	14.03		0.00			155	0.28
Poland	286	18.58	33.46	11.54	1239	80.51	33.65	11.24	14	0.91	31.21	14.67	1539	2.77
Portugal	16	18.82	43.31	18.22	68	80.00	42.34	13.77	1	1.18	38.00		85	0.15
Romania	293	20.22	47.54	14.45	1144	78.95	46.77	14.21	12	0.83	51.58	15.45	1449	2.61
Russia	3825	38.50	30.34	12.03	5847	58.85	31.74	12.25	264	2.66	27.64	10.87	9936	17.87
Serbia	152	25.08	39.16	11.94	453	74.75	41.84	11.77	1	0.17	58.00		606	1.09
Spain	330	31.82	51.49	14.85	703	67.79	48.52	13.53	4	0.39	50.00	13.11	1037	1.87
Turkey	95	27.38	25.03	6.26	249	71.76	25.05	7.36	3	0.86	21.33	0.58	347	0.62
Ukraine	306	21.07	38.42	15.38	1132	77.96	39.09	13.13	14	0.96	35.93	17.88	1452	2.61
UK	55	34.38	43.53	11.12	105	65.63	44.56	11.95		0.00			160	0.29
USA	124	30.32	37.50	15.47	273	66.75	37.78	14.51	12	2.93	28.00	9.78	409	0.74
Total	18,927	34.05	34.90	13.29	36,047	64.85	35.80	13.61	615	1.11	31.64	13.15	55,589	100.00

In terms of mental health history and self-harm, 8.59% had a prior history of an anxiety disorder, 10.93% of depression, 0.71% of Bipolar disorder, 0.42% of psychosis, and 2.90% of other mental disorder. Any mental disorder history

was present in 23.58%. At least once, 17.20% had hurt themselves in the past and 8.20% had attempted at least once in the past. The detailed rates by sex and country are shown in webtable 10.

Table 2 Sex-by-occupation composition and rates of clinical depression and distress

	%	Age		Distress %	Clinical depression %	Distress plus clinical depression (%)
		Mean	SD			
Administrative staff in hospital (4.10%)						
Women	64.35	40.29	11.14	12.28	17.37	29.64
Men	34.10	38.01	10.83	13.56	15.82	29.38
Non-binary gender	1.54	39.00	12.42	37.50	37.50	75.00
Total				13.10	17.15	30.25
Doctor (42.66%)						
Women	70.70	39.88	11.84	15.36	15.23	30.60
Men	28.67	40.21	13.42	14.03	7.89	21.91
Non-binary gender	0.63	37.35	15.25	35.29	5.88	41.18
Total				15.11	13.07	28.17
Nurse (10.89%)						
Women	87.22	41.11	11.90	14.49	16.65	31.14
Men	12.20	34.99	11.25	16.67	11.31	27.98
Non-binary gender	0.58	32.00	12.74	0.00	37.50	37.50
Total				14.67	16.12	30.79
Other healthcare profession (36.21%)						
Women	44.97	38.77	11.31	15.69	14.81	30.50
Men	54.36	33.34	7.87	14.54	9.60	24.15
Non-binary gender	0.68	36.35	11.57	29.03	16.13	45.16
Total				15.16	11.99	27.15
Other hospital staff (6.14%)						
Women	60.05	39.35	11.78	17.38	15.45	32.83
Men	37.63	34.40	12.43	18.84	14.38	33.22
Non-binary gender	2.32	28.61	9.58	33.33	22.22	55.56
Total				18.30	15.21	33.51
Total study sample						
Women	62.40	39.76	11.70	15.30	15.44	30.75
Men	36.81	35.91	11.00	14.68	9.63	24.31
Non-binary gender	0.78	35.15	13.03	30.30	17.17	47.47
Total	100.00	38.31	11.61	15.19	13.31	28.50

Family

In terms of family status, 57.86% were married, 60.02% had at least one child and only 11.42% were living alone. The responses suggested an increased need for communication with family members in 41.79%, an increased need for emotional support in 30.26%, fewer conflicts in 37.36% and increased conflicts within families for 17.77%, an improvement of the quality of relationships in 25.62%, while in most cases (90.18%) there was maintenance of basic daily routine at least somehow (webtable 11). During lockdowns 80.81% continued to work, while 47.65% expected their economic situation to worsen because of the COVID-19 outbreak (webtable 12).

Present mental health

Concerning mental health, data 47.15% reported an increase in anxiety, and 39.95% reported a worsening in depressive affect. Suicidal thoughts were increased in 10.48%. Overall, current clinical depression was present in 13.16% of the study sample (unweighted average) with male doctors and 'non-binary genders' having the lowest rates (7.89 and 5.88%, respectively) and 'non-binary gender' nurses and administrative staff having the highest (37.50%). In detail, the results are shown in Table 2. However, after taking into consideration the expected rates of current clinical depression in the population, men with positive history had the highest Relative Risk (RR = 6.47) while the lowest

was observed in women without a history of mental disorder (RR = 1.81). Additionally, distress was present in 15.19%, with the highest rates being for 'non-binary genders' (> 30%) and the lowest for female administrative staff (12.28%). The complete rates by sex and occupation are shown in webtable 17.

Suicidal tendencies doubled according to RASS subscales scores (webtable 17) and in comparison to what is expected [42].

Persons with a history of mental disorders had higher rates of current clinical depression (24.64% vs. 9.62%, chi-square test = 454.90; df = 1; $p < 0.0001$) (webtable 14). In persons without mental health history, the RR ranged from 2.14 to 2.90. In persons with a mental health history, the RR was highest in those with a psychotic history (Bipolar disorder RR = 9.04; Psychosis RR = 8.08) and ranged between 3.82 and 6.40 for non-psychotic history, with the lowest RR in persons with a history of 'other mental disorder', in comparison to the expected prevalence of depression (3% for men and 6% for women). Of women with clinical depression, half were new cases (without any past history of mental disorder) while this was true for two thirds of men. Taking into consideration that the pandemic increased the risk by definition, the risk to develop clinical depression during the pandemic when having a previous history was highest for bipolar disorder (RR = 4.23), while the previous history of self-harm or suicidal attempts did not increase the risk (Table 3).

The mean scale scores were 43.52 ± 11.99 for the STAI-S, 19.36 ± 8.17 for the CES-D, and 70.23 ± 134.74 for the Intention subscale of the RASS. The complete results by sex and country are shown in webtable 17.

Table 3 Relative Risk (RR) to develop clinical depression vs. participants with no mental health history and no history of self-harm or suicidal acts

History	Risk to develop clinical depression			
	When alone		When history of self-harm/attempt is also present	
	%	RR	%	RR
No previous history at all	9.62	1.00		1.0
Any mental disorder	31.81	3.31	29.87	3.11
Anxiety	19.93	2.07	29.49	3.07
Clinical depression	28.81	2.99	30.46	3.17
Bipolar disorder	40.66	4.23	30.78	3.20
Psychosis	36.36	3.78	30.93	3.22
Other	17.20	1.79	30.52	3.17
Only history of self-harm/attempt	17.89	1.86		

From the total sample, 5.17% reported that they often thought much or very much about committing suicide if they had the chance. Men and women had similar rates (5.76% vs. 4.69%) but those self-identified as 'non-binary gender' had much higher rates (17.00%). In subjects with a history of psychotic disorder or self-harm/attempt the rate was 15.45% while in those with a history of non-psychotic disorder, it was 9.76%. In persons free of any mental disorder or self-harm/attempt history the rate was as low as 2.09%. This means that the RR for the manifestation of at least moderate suicidal thoughts was equal to 7.4 for psychotic history and 3.7 for non-psychotic history. In those identified as 'non-binary gender', the RR was approximately equal to 3.5.

Beliefs in conspiracy theories


Approximately one third of responders accepted at least a moderate degree some non-bizarre conspiracy theory. The acceptance of inflated death rates was 44.24% while that of the 5G antenna theory was 20.81%. Doctors had the lowest rates, but impressively, 37.6% of doctors and 50.86% of nurses reported they were believing in the deliberate inflation of death rates by governments and 14.75% and 27.22% respectively were accepting the 5G theory. In detail, the responses by sex and country are shown in webtable 21.

Modeling of mental health changes during the pandemic

The presence of any mental health history acted as a risk factor for the development of current clinical depression with all chi-square tests being significant at $p < 0.001$. Interestingly a history of self-harm or suicidality emerged as a risk factor even for persons without reporting mental health history. In persons with only a history of self-harm or suicidality, 17.89% developed clinical depression. The combination of both self-harm and suicidal attempts history with specific mental health history revealed that subjects without any such history at all had the lowest rate or current clinical depression (9.62%), while the presence of previous self-harm/attempts increased the risk in subjects with past anxiety (29.49%) and other mental disorders (30.52%), but not clinical depression (30.46%), Bipolar disorder (30.78%) and psychoses (30.93%). The highest relative risk (RR) was calculated for history of Bipolar disorder but history of self-harm/attempt played no role (RR = 4.23). All RR values are shown in Table 2 and webtable 22. After taking into consideration that the annual incidence of depression is 0.3% [66], the calculated risk because of the pandemic for the health professionals population to develop clinical depression is RR = 30 Fig. 1.

The presence of a chronic somatic condition acted as a significant but weak risk factor for the development of

Fig. 1 Map of the 40 participating countries



clinical depression (Chi-square = 14.61, $df = 1$, $p < 0.001$; In terms of rates, 15.35% of those with a chronic somatic condition manifested clinical depression vs. 12.56% of those without (RR = 1.22).

The results of the MFSLRA suggested that a significant number of variables acted either as risk or as protective factors (Table 4, Fig. 2, webtable 23). These factors explained 16.1% of the change in anxiety, 11.6% of change in depressive affect, 19.1% of the development of distress or clinical depression, and 5.1% of change in suicidal thoughts. The individual contribution of each predictor separately was very small (many b coefficients were very close to zero).

If we consider a more or less linear continuum from fear to anxiety to depressive emotions to clinical depression and eventually to suicidality, the model which can be derived suggests there is a core of variables (Fig. 2, webfigure 1) that exert a stable either adverse or protective effect throughout the course of the development of the mental state.

Factorial ANOVA with the scores of STAI-S, CES-D and RASS as continuous variables and sex and being a doctor or a nurse as grouping variables was always significant for sex ($p < 0.0001$). For doctors only the interaction with sex is significant (Wilks = 0.996, $F = 4.86$, $df = 10$, error $df = 25,562$, $p < 0.0001$), with ‘non-binary genders’ having higher psychopathology. For nurses it was significant both independently (Wilks = 0.999, $F = 7.768$, $df = 10$, error $df = 25,562$, $p = 0.009$) as well as in interaction with sex (Wilks = 0.998, $F = 2.618$, $df = 10$, error $df = 25,562$, $p = 0.004$). Overall nurses had lower psychopathological scores than the rest. The Scheffe post-hoc tests (at $p < 0.05$) revealed that most groups defined by sex and occupation differed from each other in a complex and difficult-to-explain matrix.

Conspiracy theories manifest a complex behavior with some of them exerting a protective effect at certain phases

(Fig. 2), but their overall impact was lower in comparison to the general population. The mean scores of responses to questions pertaining to different conspiracy beliefs by history of any mental disorder and current clinical depression are shown in Table 5 and webtable 24. Factorial ANOVA suggested that history of any mental disorder and current clinical depression as well as their interaction were significant factors concerning the belief in conspiracy theories (Table 5). The results of post-hoc tests are shown in webtable 25. They suggest that persons with a history of mental disorder have lower overall tendency in believing in both the threatening and the reassuring conspiracy theories. Not believing in any conspiracy theory had a different composition with history of having any mental disorder being the determining factor leading to lower adaption of conspiracy theories and with depression acting at a second level and further decreasing this tendency. These findings were consistent across disorders and conspiracy theories.

Discussion

This large international study in a convenient sample of 12,792 health professionals from 40 countries detected clinical depression in 13.31% (unweighted average) with men and ‘non-binary genders’ doctors having the lowest rates (7.89% and 5.88% respectively) and ‘non-binary gender’ nurses and administrative staff having the highest (37.50%). Distress was present in 15.19%, with the highest rates being for the ‘non-binary gender’ (> 30%) and the lowest for female administrative staff (12.28%). A significant percentage reported a deterioration in mental state, family dynamics and everyday lifestyle. Persons with a history of mental disorders had higher rates of current clinical depression

Table 4 Results of four separate Multiple Forward Stepwise Linear Regression Analysis (MFLSRA) with change in anxiety (F21), change in depressive affect (G21), change in suicidal thoughts (O11) and the development of distress or clinical depression as dependent variables


	Change in anxiety (F21)			Change in depressive affect (G21)			Development of distress or clinical depression (O11)			Change in suicidal thoughts (O11)						
	b	SE	t	b	SE	t	b	SE	t	b	SE	t	p			
Intercept	-0.87	0.06	-14.49	<0.001	-0.75	0.06	-11.95	<0.001	0.59	0.04	14.31	<0.001	0.44	0.06	7.24	<0.001
Demographics																
Sex (A1)- 'non-binary gender' was not included	0.06	0.02	3.88	<0.001				-0.04	0.01	-3.02	0.003	0.06	0.02	3.98	<0.001	
Age (A2)	0.00	0.00	3.23	0.001	0.00	0.00	3.67	<0.001	0.00	-4.00	<0.001	0.00	0.00	-3.79	<0.001	
Number of persons in household (A5)																
Education level (A7)	-0.02	0.01	-2.23	0.026	-0.02	0.01	-2.42	0.016								
Work and finance																
Continue to work during lockdown (A11)																
Change in economic situation (E7)	0.10	0.01	14.02	<0.001	0.11	0.01	15.11	<0.001	-0.02	0.01	-3.15	0.002	-0.02	0.01	-2.58	0.010
Health																
Condition of general health (B1)	0.10	0.01	13.90	<0.001	0.08	0.01	10.55	<0.001	-0.09	0.01	-15.30	<0.001	-0.02	0.01	-3.29	0.001
Presence of a chronic medical condition (B2)																
Family/social																
Being a carer of a person belonging to a vulnerable group (B4)	-0.04	0.02	-2.60	0.009				-0.04	0.01	-2.98	0.003	0.04	0.01	2.57	0.010	
Conflicts within family (E3)	-0.03	0.01	-4.53	<0.001	-0.04	0.01	-5.21	<0.001	0.04	0.01	7.37	<0.001	0.06	0.01	8.30	<0.001
Change in quality of relationships within family (E4)	0.12	0.01	12.27	<0.001	0.15	0.01	15.33	<0.001	-0.02	0.01	-3.24	0.001	-0.06	0.01	-6.67	<0.001
Keeping a basic routine during lockdown (E5)	0.11	0.01	12.78	<0.001	0.10	0.01	11.46	<0.001	-0.10	0.01	-14.79	<0.001	-0.05	0.01	-5.81	<0.001
Changes in religiousness/spirituality (P1)	0.02	0.01	2.63	0.009	0.02	0.01	2.22	0.026	0.03	0.01	4.99	<0.001	-0.05	0.01	-6.22	<0.001
Mental health history																
History of anxiety (B5)	-0.14	0.03	-5.37	<0.001	-0.12	0.03	-4.39	<0.001	0.20	0.02	9.37	<0.001	0.05	0.03	2.03	0.042
History of depression (B5)	-0.10	0.02	-4.14	<0.001	-0.13	0.02	-5.45	<0.001	0.36	0.02	18.61	<0.001	0.07	0.02	2.95	0.003
History of psychosis (B5)								0.37	0.09	4.26	<0.001					
History of bipolar disorder (B5)								0.44	0.07	6.36	<0.001	-0.19	0.08	-2.28	0.023	
History of other mental disorder (B5)	-0.20	0.04	-4.50	<0.001				0.15	0.03	4.46	<0.001					
History of self-harm (O12)								0.008	0.17	0.02	9.63	<0.001	0.05	0.02	2.48	0.013
History of suicidal attempt (O13)								0.14	0.02	5.65	<0.001	0.11	0.03	3.87	<0.001	
The effect of the pandemic																
Fears of getting COVID-19 (C1)	-0.10	0.01	-12.09	<0.001	-0.05	0.01	-6.41	<0.001	0.06	0.01	8.69	<0.001	0.02	0.01	2.08	0.037
Fears that a member of the family will get COVID-19 and die (C3)	-0.08	0.01	-11.22	<0.001	-0.04	0.01	-6.35	<0.001	0.05	0.01	9.92	<0.001	-0.01	0.01	-2.24	0.025
Time spent outside of house during lockdown (D1)	0.02	0.01	3.89	<0.001	0.02	0.01	3.21	0.001								

Table 4 (continued)

	Change in anxiety (F21) R ² = 0.161; F (25, 12,620) = 97.286 p < <0.00001; SE of est: 0.809			Change in depressive affect (G21) R ² = 0.116; F (25, 12,620) = 66.742 p < <0.00001; SE of est: 0.827			Development of distress or clinical depression R ² = 0.191; F (31,12,614) = 93,466 p < <0.00001; SE of est: 0.640			Change in suicidal thoughts (O11) R ² = 0.051; F (31,12,614) = 22,282 p < <0.00001; SE of est: 0.770						
	b	SE	t	b	SE	t	b	SE	t	b	SE	t	p			
Currently locked up in the house (D2)				-0.03	0.01	-4.03	<0.0001			0.02	0.01	3.05	0.002			
Satisfaction by availability of information (D4)	0.06	0.01	8.09	<0.0001	0.06	0.01	7.88	<0.0001		-0.06	0.01	-8.12	<0.0001			
Beliefs in conspiracy theories																
The vaccine was ready before the virus broke out and they conceal it (J1)	0.04	0.01	5.15	<0.0001	0.03	0.01	4.41	<0.0001	0.03	0.01	4.38	<0.0001	0.003			
COVID-19 was created in a laboratory as a biochemical weapon (J2)									-0.03	0.01	-4.09	<0.0001	0.004			
COVID-19 is the result of 5G technology antenna (J3)				-0.03	0.01	-3.19	0.001	0.05	0.01	6.71	<0.0001					
COVID-19 appeared accidentally from human contact with animals (J4)									0.01	0.01	2.42	0.016	-0.02	0.01	-3.75	<0.0001
COVID-19 has much lower mortality rate but there is terror-inducing propaganda (J5)				0.01	0.01	2.03	0.042									
COVID-19 is a creation of the world's powerful leaders to create a global economic crisis (J6)									0.02	0.01	2.43	0.015	-0.02	0.01	-2.15	0.031
COVID-19 is a sign of divine power to destroy our planet (J7)	0.04	0.01	4.43	<0.0001					0.04	0.01	6.12	<0.0001	0.02	0.01	2.99	0.003
Occupation (A10)																
Doctor									-0.04	0.01	-3.08	0.002				
Nurse																
Other Hospital staff	0.07	0.02	2.89	0.004												
Other healthcare professional																
Administrative staff																

The predictors are shown in the left column

Fig. 2 The model which was previously developed in the general population and was proven valid also in the population of health professionals. It includes multiple vulnerabilities representing the mechanism through which the COVID-19 outbreak in combination a great number of factors could lead to clinical depression through stress, and eventually to suicidality. A number of variables act as risk factors (red) or as protective factors (green), while some of them change direction of action depending on the phase (green/red). Three core clusters emerge (delineated with the dotted lines)



(24.64%) while persons without any such history had the lowest rate of current clinical depression (9.62%). The highest rate was for the history of Bipolar disorder (40.66%; RR = 4.23). In those with a chronic somatic condition, the rate of clinical depression was 15.35% vs. 12.56% in those without (RR = 1.22). Believing in conspiracy theories was significant with at least one-third of cases accepting at least to a moderate degree a non-bizarre conspiracy.

The model developed suggested that a significant number of variables acted either as risk or as protective factors, explaining 19.1% of the development of distress or clinical depression, but their individual contribution was very small. Conspiracy theories manifested a complex behavior with some of them exerting a protective effect at certain phases. Current clinical depression acted as a risk factor and past history acted as a protective for the development of such beliefs.

The overall levels of clinical depression were lower than the rates reported in the literature, probably because of the stringent criteria of the algorithm in the current study. The large heterogeneity among countries probably reflects different phases of the pandemic in each country during the data collection. Rates of depression and mental health deterioration, in general, are probably higher in those that actually suffered from COVID-19 [29]. Other studies reported that half or more of health care professionals might suffer from depression. [7, 24, 32, 48, 76, 81, 82, 103, 117], Mira et al. 2020; [111, 112, 118]. Our results are identical to two reports [26, 52]. Meta-analyses suggested that depression rates range from 27% to 36% [45, 102, 109, 116] which is two to three times higher in comparison to our findings. In comparison, it has been reported that more than two-thirds of the general population experienced at least severe distress

[18, 31, 50, 62, 80, 83, 110], and high levels of suicidality [19]. Furthermore, our findings are in accord with a recently published meta-analysis that reported much lower depression rates in the general population [21].

An important observation is that while the rate of clinical depression was much higher in persons with a history of a mental disorder the proportion of depressed persons without such a history is much higher than expected, taking into consideration that the annual incidence of depression is 0.3% [66]. This might mean that the pandemic posed a RR = 30 on the population of health professionals to develop clinical depression.

The multivariable analysis of the data allowed the current paper to confirm a staged model previously proposed concerning the effect of the pandemic on mental health (Fig. 2). This model had been developed concerning the general population and it seems that in principle it is valid for health professionals, although with some differences, especially concerning the attenuating effect of conspiracy theories and religiousness/spirituality.

According to it, with the onset of the pandemic, its psychological impact and the development of severe anxiety and distress were determined by several sociodemographic and interpersonal variables including age, fears specific to the pandemic, the quality of relationships within the family, keeping a basic daily routine, change in the economic situation, history of any mental disorder and being afraid that him/herself or a family member will get COVID-19 and die. Similar findings concerning the effects of these factors have been reported in the literature [7, 17, 32, 35, 36, 46, 56, 57, 65, 67, 70, 80, 95, 98, 107, 111–113], Garre-Olmo et al. 2021; [89], but until now their detailed contribution had not been identified and no comprehensive model had been

Table 5 Means of responses (from -2 to +2) to all conspiracy theories by current clinical depression and history of any mental disorder and ANOVA results

Current clinical depression	History of any mental dis		Reassuring conspiracy theories						Threatening conspiracy theories						No believing in conspiracy theories					
			J1		J5		J7		J2		J3		J6		J4					
	Yes	No	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD				
Means of response scores by clinical depression and history																				
No	Yes		0.61	1.02	1.20	1.29	0.41	0.89	0.97	1.15	0.39	0.85	0.92	1.19	2.01	1.21				
No	No		0.81	1.10	1.32	1.25	0.57	0.99	1.28	1.24	0.56	0.95	1.20	1.24	1.69	1.19				
Yes	Yes		0.94	1.23	1.28	1.28	0.58	1.05	1.31	1.29	0.63	1.08	1.22	1.31	1.87	1.23				
Yes	No		1.06	1.21	1.41	1.25	0.95	1.28	1.43	1.28	0.89	1.16	1.44	1.30	1.73	1.21				
All Grps	0.80		1.11	1.31	1.26	0.57	1.01	1.24	1.24	0.55	0.96	1.17	1.25	1.76	1.21					
ANOVA results																				
			Wilks						F						Effect df		Error df		p	
Clinical depression			0.986						26.54						7		12,780		<0.001	
History of any mental disorder			0.988						23.06						7		12,780		<0.001	
Clinical depression* history of any mental disorder			0.997						5.51						7		12,780		<0.001	

developed. On the other hand, several factors not assessed by the current study, including the level of training, whether the person worked in the frontline against COVID or in an ICU, etc. [8, 54, 67, 111, 112], Mira et al. 2020; [7, 20, 24, 48, 51, 52, 71, 82, 95, 101, 103, 107] were reported as contributing in the development of clinical depression. The current paper suggests that from all health occupations, nurses might be at a higher risk to develop severe stress and clinical depression, and this is in accord with the literature [33, 48, 51, 58, 118]. Previous reports on the role of temperament are in accord with this [74]

At the pandemic onset, we might not have imagined the important role and the impact of conspiracy theories, which are largely social media driven. They are currently widely accepted as being important since the literature strongly supports their relationship with anxiety and depression [22, 28]. According to the results of the current study, approximately one-third of responders accepted at least to a moderate degree a non-bizarre conspiracy theory, and this was true both for ‘threatening’ as well as for ‘reassuring’ theories. The acceptance of inflated death rates was 44.24% while that of the 5G antenna theory was 20.81%. Doctors had the lowest rates, but impressively, 37.6% of doctors and 50.86% of nurses reported they were believing in the deliberate inflation of death rates by governments, and 14.75% and 27.22% respectively were accepting the 5G theory. Interestingly, believing in conspiracy theories pertaining to COVID-19 was lower in comparison to the general population and played an attenuated role in the development of anxiety and depression, however, these beliefs seem to be an important factor even among doctors. The high rates of believing in conspiracy theories are in accord with findings from various countries [1, 64, 91, 108] and are a worrying manifestation. Conspiracy beliefs – especially those regarding science, medicine, and health-related topics – are widespread [78], are widely distributed in social media [1, 11] and they challenge the capacity of the average person to distill and assess the content [30, 34]. They exert a well-documented adverse effect on health behaviors, especially vaccination [2, 3, 14–16, 43, 49, 59, 63, 72, 88, 91, 93, 99, 105]. There seems to be some relationship between believing in bizarre conspiracy theories and psychotic tendencies or a history of psychotic disorders [60].

As was found in the general population, current clinical depression and past history of mental disorders are both critical factors related to believing in conspiracy theories. Our results could mean that the critical factor which increases belief is the presence of current clinical depression, while the past history acts at a second level. As correlation does not imply causation, conspiracy theories could be either the cause of clinical depression, a copying mechanism against clinical depression, or a marker of maladaptive psychological patterns of cognitive appraisal.

After taking into consideration the complete model, and especially the relationship to past mental health history, the authors propose that the beliefs in conspiracy theories are a copying mechanism against stress. The finding of the relationship between current clinical depression and believing in conspiracies is in accord with the literature [28, 44, 106]. One explanation could be found in the theory concerning ‘Depressive Realism’ [5, 6], Alloy et al. 1981; [12, 68, 77] which suggests that depressive persons are more able than others to realistically interpret the world, however, this higher ability leads to pessimism.

At the most extreme end, when the emergence of suicidal thinking is possible, the family environment and family responsibilities and care act either as risk or protective factors, depending on their quality, while religiosity/spirituality and all beliefs in conspiracy theories act as protective factors, except for one which includes religious content. These results are in accord with the reports in the literature [9, 56, 57, 61, 65, 79, 113].

A difficult-to-answer question is how many of the cases detected by questionnaires and sophisticated algorithms correspond to real major clinical depression. The underlying neurobiology is opaque and maybe much diagnosed clinical depression might simply be an extreme form of a normal adjustment reaction [53]. However, there is no better way to psychometrically achieve higher validity and the algorithm we utilized is the best available method. The impressive increase in new cases of clinical depression (9.62% of persons without any history of mental disorders developed depression) which was found in our sample is in accord with the literature [87]. However, a large part of clinical depressions emerged from a previous mental health history. Of the 13.15% with current clinical depression, 7.35% were new cases while 5.80% had previous history. This suggests that almost beyond doubt true clinical depression increased by 30% (in the extreme scenario that none of cases without previous history was a case of true clinical depression. This extremely positive scenario also suggests that maybe relapses expected to occur in the next several years occurred earlier.

Concerning those without a previous history of mental disorder, it is expected that much of the adverse effects on mental health will rapidly attenuate with the end of the pandemic [27] but enduring effects will impact some vulnerable populations. So far studies investigating the long-term outcome and the long-term impact of the pandemic on mental health display equivocal findings [13, 114]. Especially sociability and the sense of belonging could be important factors determining mental health and health-related behaviors [15], and these factors seem to correspond to specific vulnerabilities seen especially in western cultures.

Conclusion

The current paper reports high rates of clinical depression, distress, and suicidal thoughts among the population of health workers during the pandemic, with a high prevalence of beliefs in conspiracy theories. For the development of clinical depression, general health status, previous mental health history, self-harm and suicidal attempts, family responsibility, economic change, and age acted as risk factors while keeping a daily routine, religiousness/spirituality, and belief in conspiracy theories were acting mostly as protective factors. These findings, although they should be closely monitored longitudinally, support previous suggestions by other authors concerning the need for a proactive intervention to protect the mental health of the general population but more specifically of vulnerable groups [38, 94]

Strengths and limitations

The strengths of the current paper include the large number of persons who filled out the questionnaire and the large bulk of information obtained, as well as the detailed way of post-stratification of the study sample.

The major limitation was that the data were obtained anonymously online through the self-selection of the responders. Additionally, the assessment included only the cross-sectional application of self-report scales, although the advanced algorithm used for the diagnosis of clinical depression corrected the problem to a certain degree. However, what is included under the umbrella of ‘clinical depression’ in the stressful times of the pandemic remains a matter of debate. Also, the lack of baseline data concerning the mental health of a similar study sample before the pandemic is also a problem.

Finally, a limitation would be that data from different countries were pooled together and with a rather large difference in numbers among countries. So the interpretation of findings should bear in mind the possible bias induced by different cultural backgrounds. However, one should have also in mind that backgrounds are not so different as one might think since the online survey of a specific professional population poses requirements that lead to similarities rather than differences among the populations from different countries. Each of these countries was also undergoing a different phase of the pandemic at each time point and phases were also different across individuals. This means that the results should be interpreted.

Author contributions All authors contributed equally to the paper. KNF and DS conceived and designed the study. The other authors participated formulating the final protocol, designing and supervising the data collection and creating the final dataset. KNF and DS did the data analysis and wrote the first draft of the paper. All authors

participated in interpreting the data and developing further stages and the final version of the paper.

Funding None.

Data availability statement Raw data are available upon request to the principal investigator.

Declarations

Conflict of interest None pertaining to the current paper.

References

- Ahmed W, Vidal-Alaball J, Downing J, Lopez Segui F (2020) COVID-19 and the 5G conspiracy theory: social network analysis of Twitter Data. *J Med Internet Res* 22(5):e19458. <https://doi.org/10.2196/19458>
- Allington D, Duffy B, Wessely S, Dhavan N, Rubin J (2021) Health-protective behaviour, social media usage and conspiracy belief during the COVID-19 public health emergency. *Psychol Med* 51(10):1763–1769. <https://doi.org/10.1017/S003329172000224X>
- Allington D, McAndrew S, Moxham-Hall V, Duffy B (2021) Coronavirus conspiracy suspicions, general vaccine attitudes, trust and coronavirus information source as predictors of vaccine hesitancy among UK residents during the COVID-19 pandemic. *Psychol Med*. <https://doi.org/10.1017/S0033291721001434>
- Alloy LB, Abramson LY, Viscusi D (1981) Induced mood and the illusion of control. *J Personal Soc Psychol* 41(6):1129
- Alloy LB, Abramson LY (1979) Judgment of contingency in depressed and nondepressed students: sadder but wiser? *J Exp Psychol Gen* 108(4):441–485. <https://doi.org/10.1037//0096-3445.108.4.441>
- Alloy LB, Abramson LY (1988) Depressive realism: Four theoretical perspectives. Cognitive processes in depression. The Guilford Press, New York, NY, US, pp 223–265
- Amin F, Sharif S, Saeed R, Durrani N, Jilani D (2020) COVID-19 pandemic- knowledge, perception, anxiety and depression among frontline doctors of Pakistan. *BMC Psychiatry* 20(1):459. <https://doi.org/10.1186/s12888-020-02864-x>
- Antonijevic J, Binic I, Zikic O, Manojlovic S, Tosic-Golubovic S, Popovic N (2020) Mental health of medical personnel during the COVID-19 pandemic. *Brain Behav* 10(12):e01881. <https://doi.org/10.1002/brb3.1881>
- Arslan G, Yildirim M (2021) Meaning-based coping and spirituality during the COVID-19 pandemic: mediating effects on subjective well-being. *Front Psychol* 12:646572. <https://doi.org/10.3389/fpsyg.2021.646572>
- Asmundson GJG, Taylor S (2021) Garbage in, garbage out: the tenuous state of research on PTSD in the context of the COVID-19 pandemic and infodemic. *J Anxiety Disord* 78:102368. <https://doi.org/10.1016/j.janxdis.2021.102368>
- Banerjee D, Meena KS (2021) COVID-19 as an “Infodemic” in public health: critical role of the social media. *Front Public Health* 9:610623. <https://doi.org/10.3389/fpubh.2021.610623>
- Beck AT, Brown G, Steer RA, Eidelson JI, Riskind JH (1987) Differentiating anxiety and depression: a test of the cognitive content-specificity hypothesis. *J Abnorm Psychol* 96(3):179–183. <https://doi.org/10.1037//0021-843x.96.3.179>
- Bendau A, Plag J, Kunas S, Wyka S, Strohle A, Petzold MB (2021) Longitudinal changes in anxiety and psychological distress, and associated risk and protective factors during the first

- three months of the COVID-19 pandemic in Germany. *Brain Behav* 11(2):e01964. <https://doi.org/10.1002/brb3.1964>
14. Bertin P, Nera K, Delouvee S (2020) Conspiracy beliefs, rejection of vaccination, and support for hydroxychloroquine: a conceptual replication-extension in the COVID-19 pandemic context. *Front Psychol* 11:565128. <https://doi.org/10.3389/fpsyg.2020.565128>
 15. Biddlestone M, Green R, Douglas KM (2020) Cultural orientation, power, belief in conspiracy theories, and intentions to reduce the spread of COVID-19. *Br J Soc Psychol* 59(3):663–673. <https://doi.org/10.1111/bjso.12397>
 16. Bogart LM, Wagner G, Galvan FH, Banks D (2010) Conspiracy beliefs about HIV are related to antiretroviral treatment nonadherence among African American men with HIV. *J Acquir Immune Defic Syndr* 53(5):648–655. <https://doi.org/10.1097/QAI.0b013e3181c57dbc>
 17. Bruffaerts R, Voorspoels W, Jansen L, Kessler RC, Mortier P, Vilagut G, De Vocht J, Alonso J (2021) Suicidality among healthcare professionals during the first COVID-19 wave. *J Affect Disord* 283:66–70. <https://doi.org/10.1016/j.jad.2021.01.013>
 18. Busch IM, Moretti F, Mazzi M, Wu AW, Rimondini M (2021) What we have learned from two decades of epidemics and pandemics: a systematic review and meta-analysis of the psychological burden of frontline healthcare workers. *Psychother Psychosom* 90(3):178–190. <https://doi.org/10.1159/000513733>
 19. Caballero-Dominguez CC, Jimenez-Villamizar MP, Campo-Arias A (2020) Suicide risk during the lockdown due to coronavirus disease (COVID-19) in Colombia. *Death Stud* <https://doi.org/10.1080/07481187.2020.1784312>
 20. Celmece N, Menekay M (2020) The effect of stress, anxiety and burnout levels of healthcare professionals caring for COVID-19 patients on their quality of life. *Front Psychol* 11:597624. <https://doi.org/10.3389/fpsyg.2020.597624>
 21. Cenat JM, Blais-Rochette C, Kokou-Kpolou CK, Noorishad PG, Mukunzi JN, McIntee SE, Dalexis RD, Goulet MA, Labelle PR (2021) Prevalence of symptoms of depression, anxiety, insomnia, posttraumatic stress disorder, and psychological distress among populations affected by the COVID-19 pandemic: a systematic review and meta-analysis. *Psychiatry Res* 295:113599. <https://doi.org/10.1016/j.psychres.2020.113599>
 22. Chen X, Zhang SX, Jahanshahi AA, Alvarez-Risco A, Dai H, Li J, Ibarra VG (2020) Belief in a COVID-19 conspiracy theory as a predictor of mental health and well-being of health care workers in Ecuador: cross-sectional survey study. *JMIR Public Health Surveill* 6(3):e20737. <https://doi.org/10.2196/20737>
 23. Claponea RM, Pop LM, Iorga M, Iurcov R (2022) Symptoms of burnout syndrome among physicians during the outbreak of COVID-19 pandemic—a systematic literature review. *Healthcare (Basel)*. <https://doi.org/10.3390/healthcare10060979>
 24. Conti C, Fontanesi L, Lanzara R, Rosa I, Doyle RL, Porcelli P (2021) Burnout status of Italian healthcare workers during the first COVID-19 pandemic peak period. *Healthcare (Basel)*. <https://doi.org/10.3390/healthcare9050510>
 25. Courtet P, Olie E (2021) Suicide in the COVID-19 pandemic: what we learnt and great expectations. *Eur Neuropsychopharmacol* 50:118–120. <https://doi.org/10.1016/j.euroneuro.2021.06.005>
 26. da Silva FCT, Neto MLR (2021) Psychological effects caused by the COVID-19 pandemic in health professionals: a systematic review with meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry* 104:110062. <https://doi.org/10.1016/j.pnpbp.2020.110062>
 27. Daly M, Robinson E (2021) Anxiety reported by US adults in 2019 and during the 2020 COVID-19 pandemic: population-based evidence from two nationally representative samples. *J Affect Disord* 286:296–300. <https://doi.org/10.1016/j.jad.2021.02.054>
 28. De Coninck D, Frissen T, Matthijs K, d’Haenens L, Lits G, Champagne-Poirier O, Carignan ME, David MD, Pignard-Cheynel N, Salerno S, Genereux M (2021) Beliefs in conspiracy theories and misinformation about COVID-19: comparative perspectives on the role of anxiety, depression and exposure to and trust in information sources. *Front Psychol* 12:646394. <https://doi.org/10.3389/fpsyg.2021.646394>
 29. Deng J, Zhou F, Hou W, Silver Z, Wong CY, Chang O, Huang E, Zuo QK (2021) The prevalence of depression, anxiety, and sleep disturbances in COVID-19 patients: a meta-analysis. *Ann N Y Acad Sci* 1486(1):90–111. <https://doi.org/10.1111/nyas.14506>
 30. Desta TT, Mulugeta T (2020) Living with COVID-19-triggered pseudoscience and conspiracies. *Int J Public Health* 65(6):713–714. <https://doi.org/10.1007/s00038-020-01412-4>
 31. Dominguez-Salas S, Gomez-Salgado J, Andres-Villas M, Diaz-Milanes D, Romero-Martin M, Ruiz-Frutos C (2020) Psycho-emotional approach to the psychological distress related to the COVID-19 pandemic in Spain: a cross-sectional observational study. *Healthcare (Basel)*. <https://doi.org/10.3390/healthcare8030190>
 32. Dong ZQ, Ma J, Hao YN, Shen XL, Liu F, Gao Y, Zhang L (2020) The social psychological impact of the COVID-19 pandemic on medical staff in China: a cross-sectional study. *Eur Psychiatry* 63(1):e65. <https://doi.org/10.1192/j.eurpsy.2020.59>
 33. Donkers MA, Gilissen V, Candel M, van Dijk NM, Kling H, Heijnen-Panis R, Pragt E, van der Horst I, Pronk SA, van Mook W (2021) Moral distress and ethical climate in intensive care medicine during COVID-19: a nationwide study. *BMC Med Ethics* 22(1):73. <https://doi.org/10.1186/s12910-021-00641-3>
 34. Duplaga M (2020) The determinants of conspiracy beliefs related to the COVID-19 pandemic in a nationally representative sample of internet users. *Int J Environ Res Public Health*. <https://doi.org/10.3390/ijerph17217818>
 35. Elbogen EB, Lanier M, Blakey SM, Wagner HR, Tsai J (2021) Suicidal ideation and thoughts of self-harm during the COVID-19 pandemic: the role of COVID-19-related stress, social isolation, and financial strain. *Depress Anxiety*. <https://doi.org/10.1002/da.23162>
 36. Elhai JD, Yang H, McKay D, Asmundson GJG, Montag C (2021) Modeling anxiety and fear of COVID-19 using machine learning in a sample of Chinese adults: associations with psychopathology, sociodemographic, and exposure variables. *Anxiety Stress Coping* 34(2):130–144. <https://doi.org/10.1080/10615806.2021.1878158>
 37. Fancourt D, Steptoe A, Bu F (2021) Trajectories of anxiety and depressive symptoms during enforced isolation due to COVID-19 in England: a longitudinal observational study. *Lancet Psychiatry* 8(2):141–149. [https://doi.org/10.1016/S2215-0366\(20\)30482-X](https://doi.org/10.1016/S2215-0366(20)30482-X)
 38. Fiorillo A, Gorwood P (2020) The consequences of the COVID-19 pandemic on mental health and implications for clinical practice. *Eur Psychiatry* 63(1):e32. <https://doi.org/10.1192/j.eurpsy.2020.35>
 39. Fountoulakis K, Iacovides A, Kleanthous S, Samolis S, Kaprinis SG, Sitzoglou K, St Kaprinis G, Bech P (2001) Reliability, validity and psychometric properties of the Greek translation of the Center for Epidemiological Studies-Depression (CES-D) Scale. *BMC Psychiatry* 1:3. <https://doi.org/10.1186/1471-244x-1-3>
 40. Fountoulakis KN, Apostolidou MK, Atsiouva MB, Filippidou AK, Florou AK, Gousiou DS, Katsara AR, Mantzari SN, Padouva-Markoulaki M, Papatriantafyllou EI, Sacharidi PI, Tonia AI, Tsagalidou EG, Zymara VP, Prezerakos PE, Koupidis SA, Fountoulakis NK, Chrousos GP (2021) Self-reported changes in anxiety, depression and suicidality during the

- COVID-19 lockdown in Greece. *J Affect Disord* 279:624–629. <https://doi.org/10.1016/j.jad.2020.10.061>
41. Fountoulakis KN, Karakatsoulis G, Abraham S, Adorjan K, Ahmed HU, Alarcón RD, Arai K, Auwal SS, Berk M, Bjedov S, Bobes J, Bobes-Bascaran T, Bourgin-Duchesnay J, Bredicean CA, Bukelskis L, Burkadze A, Abud IIC, Castilla-Puentes R, Cetkovich M, Colon-Rivera H, Corral R, Cortez-Vergara C, Crepin P, De Berardis D, Zamora Delgado S, De Lucena D, De Sousa A, Stefano RD, Dodd S, Elek LP, Elissa A, Erdelyi-Hamza B, Erzin G, Etchevers MJ, Falkai P, Farcas A, Fedotov I, Filatova V, Fountoulakis NK, Frankova I, Franza F, Frias P, Galako T, Garay CJ, Garcia-Álvarez L, García-Portilla MP, Gonda X, Gondek TM, González DM, Gould H, Grandinetti P, Grau A, Groudeva V, Hagin M, Harada T, Hasan TM, Hashim NA, Hilbig J, Hossain S, Iakimova R, Ibrahim M, Iftene F, Ignatenko Y, Irrarrazaval M, Ismail Z, Ismayilova J, Jakobs A, Jakovljević M, Jakšić N, Javed A, Kafali HY, Karia S, Kazakova O, Khalifa D, Khaustova O, Koh S, Kopishinskaja S, Kosenko K, Koupidis SA, Kovacs I, Kulig B, Lalljee A, Liewig J, Majid A, Malashonkova E, Malik K, Malik NI, Mammadzada G, Mandalia B, Marazziti D, Marčinko D, Martínez S, Matiekus E, Mejia G, Memon RS, Martínez XEM, Mickevičiūtė D, Milev R, Mohammed M, Molina-López A, Morozov P, Muhammad NS, Mustač F, Naor MS, Nassieb A, Navickas A, Okasha T, Pandova M, Panfil A-L, Panteleeva L, Papava I, Patsali ME, Pavlichenko A, Pejuskovic B, Da Costa MP, Popkov M, Popovic D, Raduan NJN, Ramírez FV, Rancans E, Razali S, Rebok F, Rewekant A, Flores ENR, Rivera-Encinas MT, Saiz P, de Carmona MS, Martínez DS, Saw JA, Saygili G, Schneiderreit P, Shah B, Shirasaka T, Silagadze K, Sitanggang S, Skugarevsky O, Spikina A, Mahalingappa SS, Stoyanova M, Szczegieliński A, Tamasan SC, Tavormina G, Tavormina MGM, Theodorakis PN, Tohen M, Tsapakis EM, Tukhvatullina D, Ullah I, Vaidya R, Vega-Dienstmaier JM, Vrublevska J, Vukovic O, Vysotska O, Widiastih N, Yashikhina A, Prezerakos PE, Smirnova D (2021) Results of the COVID-19 MEntal health inTernational for the General population (COMET-G) study. *Eur Neuropsychopharmacol*. <https://doi.org/10.1016/j.euroneuro.2021.10.004>
 42. Fountoulakis KN, Pantoula E, Siamouli M, Moutou K, Gonda X, Rihmer Z, Iacovides A, Akiskal H (2012) Development of the Risk Assessment Suicidality Scale (RASS): a population-based study. *J Affect Disord* 138(3):449–457. <https://doi.org/10.1016/j.jad.2011.12.045>
 43. Freeman D, Loe BS, Chadwick A, Vaccari C, Waite F, Rosebrock L, Jenner L, Petit A, Lewandowsky S, Vanderslott S, Innocenti S, Larkin M, Giubilini A, Yu LM, McShane H, Pollard AJ, Lambe S (2020) COVID-19 vaccine hesitancy in the UK: the Oxford coronavirus explanations, attitudes, and narratives survey (Oceans) II. *Psychol Med*. <https://doi.org/10.1017/S0033291720005188>
 44. Freyler A, Simor P, Szemerszky R, Szabolcs Z, Koteles F (2019) Modern health worries in patients with affective disorders. A pilot study. *Ideggyogy Sz* 72(9–10):337–341. <https://doi.org/10.18071/isz.72.0337>
 45. Galli F, Pozzi G, Ruggiero F, Mameli F, Cavicchioli M, Barbieri S, Canevini MP, Priori A, Pravettoni G, Sani G, Ferrucci R (2020) A systematic review and provisional metanalysis on psychopathologic burden on health care workers of coronavirus outbreaks. *Front Psychiatry* 11:568664. <https://doi.org/10.3389/fpsy.2020.568664>
 46. Gambin M, Sekowski M, Wozniak-Prus M, Wnuk A, Oleksy T, Cudo A, Hansen K, Huflejt-Lukasik M, Kubicka K, Lys AE, Gorgol J, Holas P, Kmita G, Lojek E, Maison D (2021) Generalized anxiety and depressive symptoms in various age groups during the COVID-19 lockdown in Poland. Specific predictors and differences in symptoms severity. *Compr Psychiatry* 105:152222
 47. Garre-Olmo J, Turro-Garriga O, Marti-Lluch R, Zacarias-Pons L, Alves-Cabrata L, Serrano-Sarboza D, Vilalta-Franch J, Ramos R, Girona Health Region Study G (2021) Changes in lifestyle resulting from confinement due to COVID-19 and depressive symptomatology: a cross-sectional a population-based study. *Compr Psychiatry* 104:152214. <https://doi.org/10.1016/j.comppsych.2020.152214>
 48. Giusti EM, Pedroli E, D'Aniello GE, Stramba Badiale C, Pietrabissa G, Manna C, Stramba Badiale M, Riva G, Castelnuovo G, Molinari E (2020) The psychological impact of the COVID-19 outbreak on health professionals: a cross-sectional study. *Front Psychol* 11:1684. <https://doi.org/10.3389/fpsyg.2020.01684>
 49. Gu F, Wu Y, Hu X, Guo J, Yang X, Zhao X (2021) The role of conspiracy theories in the spread of COVID-19 across the United States. *Int J Environ Res Public Health*. <https://doi.org/10.3390/ijerph18073843>
 50. Gualano MR, Lo Moro G, Voglino G, Bert F, Siliquini R (2020) Effects of COVID-19 lockdown on mental health and sleep disturbances in Italy. *Int J Environ Res Public Health*. <https://doi.org/10.3390/ijerph17134779>
 51. Hacimusalar Y, Kahve AC, Yasar AB, Aydin MS (2020) Anxiety and hopelessness levels in COVID-19 pandemic: a comparative study of healthcare professionals and other community sample in Turkey. *J Psychiatr Res* 129:181–188. <https://doi.org/10.1016/j.jpsychires.2020.07.024>
 52. Han S, Choi S, Cho SH, Lee J, Yun JY (2021) Associations between the working experiences at frontline of COVID-19 pandemic and mental health of Korean public health doctors. *BMC Psychiatry* 21(1):298. <https://doi.org/10.1186/s12888-021-03291-2>
 53. He L, Wei D, Yang F, Zhang J, Cheng W, Feng J, Yang W, Zhuang K, Chen Q, Ren Z, Li Y, Wang X, Mao Y, Chen Z, Liao M, Cui H, Li C, He Q, Lei X, Feng T, Chen H, Xie P, Rolls ET, Su L, Li L, Qiu J (2021) Functional connectome prediction of anxiety related to the COVID-19 pandemic. *Am J Psychiatry* 178(6):530–540. <https://doi.org/10.1176/appi.ajp.2020.20070979>
 54. Hesselink G, Straten L, Gallee L, Brants A, Holkenborg J, Barten DG, Schoon Y (2021) Holding the frontline: a cross-sectional survey of emergency department staff well-being and psychological distress in the course of the COVID-19 outbreak. *BMC Health Serv Res* 21(1):525. <https://doi.org/10.1186/s12913-021-06555-5>
 55. Hill JE, Harris C, Danielle LC, Boland P, Doherty AJ, Benedetto V, Gita BE, Clegg AJ (2022) The prevalence of mental health conditions in healthcare workers during and after a pandemic: Systematic review and meta-analysis. *J Adv Nurs* 78(6):1551–1573. <https://doi.org/10.1111/jan.15175>
 56. Huang Y, Zhao N (2020) Chinese mental health burden during the COVID-19 pandemic. *Asian J Psychiatr* 51:102052. <https://doi.org/10.1016/j.ajp.2020.102052>
 57. Huang Y, Zhao N (2021) Mental health burden for the public affected by the COVID-19 outbreak in China: who will be the high-risk group? *Psychol Health Med* 26(1):23–34. <https://doi.org/10.1080/13548506.2020.1754438>
 58. Jo SH, Koo BH, Seo WS, Yun SH, Kim HG (2020) The psychological impact of the coronavirus disease pandemic on hospital workers in Daegu, South Korea. *Compr Psychiatry* 103:152213. <https://doi.org/10.1016/j.comppsych.2020.152213>
 59. Jolley D, Douglas KM (2014) The effects of anti-vaccine conspiracy theories on vaccination intentions. *PLoS ONE* 9(2):e89177. <https://doi.org/10.1371/journal.pone.0089177>
 60. Jolley D, Paterson JL (2020) Pylons ablaze: Examining the role of 5G COVID-19 conspiracy beliefs and support for violence. *Br J Soc Psychol* 59(3):628–640. <https://doi.org/10.1111/bjso.12394>
 61. Jovancevic A, Milicevic N (2020) Optimism-pessimism, conspiracy theories and general trust as factors contributing to

- COVID-19 related behavior - A cross-cultural study. *Pers Individ Dif* 167:110216. <https://doi.org/10.1016/j.paid.2020.110216>
62. Knolle F, Ronan L, Murray GK (2021) The impact of the COVID-19 pandemic on mental health in the general population: a comparison between Germany and the UK. *BMC Psychol* 9(1):60. <https://doi.org/10.1186/s40359-021-00565-y>
 63. Lazarevic LB, Puric D, Teovanovic P, Lukic P, Zupan Z, Knezevic G (2021) What drives us to be (ir)responsible for our health during the COVID-19 pandemic? The role of personality, thinking styles, and conspiracy mentality. *Pers Individ Dif* 176:110771. <https://doi.org/10.1016/j.paid.2021.110771>
 64. Leibovitz T, Shamblaw AL, Rumas R, Best MW (2021) COVID-19 conspiracy beliefs: relations with anxiety, quality of life, and schemas. *Pers Individ Dif* 175:110704. <https://doi.org/10.1016/j.paid.2021.110704>
 65. Li J, Yang Z, Qiu H, Wang Y, Jian L, Ji J, Li K (2020) Anxiety and depression among general population in China at the peak of the COVID-19 epidemic. *World Psychiatry* 19(2):249–250. <https://doi.org/10.1002/wps.20758>
 66. Liu Q, He H, Yang J, Feng X, Zhao F, Lyu J (2020) Changes in the global burden of depression from 1990 to 2017: findings from the Global Burden of Disease study. *J Psychiatr Res* 126:134–140. <https://doi.org/10.1016/j.jpsychires.2019.08.002>
 67. Liu Y, Long Y, Cheng Y, Guo Q, Yang L, Lin Y, Cao Y, Ye L, Jiang Y, Li K, Tian K, Sun C, Zhang F, Song X, Liao G, Huang J, Du L (2020) Psychological impact of the COVID-19 outbreak on nurses in China: a nationwide survey during the outbreak. *Front Psychiatry* 11:598712. <https://doi.org/10.3389/fpsy.2020.598712>
 68. Lobitz WC, Post RD (1979) Parameters of self-reinforcement and depression. *J Abnorm Psychol* 88(1):33–41. <https://doi.org/10.1037//0021-843x.88.1.33>
 69. MacDonald PL, Gardner RC (2016) Type I error rate comparisons of post hoc procedures for I j Chi-Square tables. *Educ Psychol Measur* 60(5):735–754. <https://doi.org/10.1177/00131640021970871>
 70. Mahgoub IM, Abdelrahman A, Abdallah TA, Mohamed Ahmed KAH, Omer MEA, Abdelrahman E, Salih ZMA (2021) Psychological effects of the COVID-19 pandemic: Perceived stress, anxiety, work-family imbalance, and coping strategies among healthcare professionals in Khartoum state hospitals, Sudan, 2021. *Brain and behavior* 11(8):e2318
 71. Malik S, Ullah I, Irfan M, Ahorsu DK, Lin CY, Pakpour AH, Griffiths MD, Rehman IU, Minhas R (2021) Fear of COVID-19 and workplace phobia among Pakistani doctors: a survey study. *BMC Public Health* 21(1):833. <https://doi.org/10.1186/s12889-021-10873-y>
 72. Marinthe G, Brown G, Delouvee S, Jolley D (2020) Looking out for myself: Exploring the relationship between conspiracy mentality, perceived personal risk, and COVID-19 prevention measures. *Br J Health Psychol* 25(4):957–980. <https://doi.org/10.1111/bjhp.12449>
 73. Mira JJ, Carrillo I, Guilabert M, Mula A, Martin-Delgado J, Perez-Jover MV, Vicente MA, Fernandez C, Group SA-C-SVS (2020) Acute stress of the healthcare workforce during the COVID-19 pandemic evolution: a cross-sectional study in Spain. *BMJ Open* 10(11):e042555. <https://doi.org/10.1136/bmjopen-2020-042555>
 74. Moccia L, Janiri D, Pepe M, Dattoli L, Molinaro M, De Martin V, Chieffo D, Janiri L, Fiorillo A, Sani G, Di Nicola M (2020) Affective temperament, attachment style, and the psychological impact of the COVID-19 outbreak: an early report on the Italian general population. *Brain Behav Immun* 87:75–79. <https://doi.org/10.1016/j.bbi.2020.04.048>
 75. Mortier P, Vilagut G, Ferrer M, Serra C, Molina JD, Lopez-Fresnena N, Puig T, Pelayo-Teran JM, Pijoan JJ, Emparanza JJ, Espuga M, Plana N, Gonzalez-Pinto A, Orti-Lucas RM, de Salazar AM, Rius C, Aragones E, Del Cura-Gonzalez I, Aragon-Pena A, Campos M, Parellada M, Perez-Zapata A, Forjaz MJ, Sanz F, Haro JM, Vieta E, Perez-Sola V, Kessler RC, Bruffaerts R, Alonso J, Group MW (2021) Thirty-day suicidal thoughts and behaviors among hospital workers during the first wave of the Spain COVID-19 outbreak. *Depress Anxiety* 38(5):528–544. <https://doi.org/10.1002/da.23129>
 76. Naser AY, Dahmash EZ, Al-Rousan R, Alwafi H, Alrawashdeh HM, Ghoul I, Abidine A, Bokhary MA, Al-Hadithi HT, Ali D, Abuthawabeh R, Abdelwahab GM, Alhartani YJ, Al Muhaisen H, Dagash A, Alyami HS (2020) Mental health status of the general population, healthcare professionals, and university students during 2019 coronavirus disease outbreak in Jordan: a cross-sectional study. *Brain and behavior* 10(8):e01730. <https://doi.org/10.1002/brb3.1730>
 77. Nelson RE, Craighead WE (1977) Selective recall of positive and negative feedback, self-control behaviors, and depression. *J Abnorm Psychol* 86(4):379–388. <https://doi.org/10.1037//0021-843x.86.4.379>
 78. Oliver JE, Wood T (2014) Medical conspiracy theories and health behaviors in the United States. *JAMA Intern Med* 174(5):817–818. <https://doi.org/10.1001/jamainternmed.2014.190>
 79. Ozdin S, Bayrak Ozdin S (2020) Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: The importance of gender. *Int J Soc Psychiatry*. <https://doi.org/10.1177/0020764020927051>
 80. Ozdin S, Bayrak Ozdin S (2020) Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: The importance of gender. *Int J Soc Psychiatry* 66(5):504–511. <https://doi.org/10.1177/0020764020927051>
 81. Pearman A, Hughes ML, Smith EL, Neupert SD (2020) Mental Health Challenges of United States Healthcare Professionals During COVID-19. *Front Psychol* 11:2065. <https://doi.org/10.3389/fpsyg.2020.02065>
 82. Perera B, Wickramarachchi B, Samanmalie C, Hettiarachchi M (2021) Psychological experiences of healthcare professionals in Sri Lanka during COVID-19. *BMC Psychol* 9(1):49. <https://doi.org/10.1186/s40359-021-00526-5>
 83. Petzold MB, Bendau A, Plag J, Pyrkosch L, Mascarell Maricic L, Betzler F, Rogoll J, Grosse J, Strohle A (2020) Risk, resilience, psychological distress, and anxiety at the beginning of the COVID-19 pandemic in Germany. *Brain and behavior* 10(9):e01745. <https://doi.org/10.1002/brb3.1745>
 84. Pompili M (2021) Can we expect a rise in suicide rates after the Covid-19 pandemic outbreak? *Eur Neuropsychopharmacol* 52:1–2. <https://doi.org/10.1016/j.euroneuro.2021.05.011>
 85. Racine N, Hetherington E, McArthur BA, McDonald S, Edwards S, Tough S, Madigan S (2021) Maternal depressive and anxiety symptoms before and during the COVID-19 pandemic in Canada: a longitudinal analysis. *Lancet Psychiatry* 8(5):405–415. [https://doi.org/10.1016/S2215-0366\(21\)00074-2](https://doi.org/10.1016/S2215-0366(21)00074-2)
 86. Ren X, Huang W, Pan H, Huang T, Wang X, Ma Y (2020) Mental health during the Covid-19 outbreak in China: a meta-analysis. *Psychiatr Q* 91(4):1033–1045. <https://doi.org/10.1007/s11126-020-09796-5>
 87. Robillard R, Daros AR, Phillips JL, Porteous M, Saad M, Pennestri MH, Kendzerska T, Edwards JD, Solomonova E, Bhatla R, Godbout R, Kaminsky Z, Boafu A, Quilty LC (2021) Emerging New Psychiatric Symptoms and the Worsening of Pre-existing Mental Disorders during the COVID-19 Pandemic: A Canadian Multisite Study: Nouveaux symptômes psychiatriques émergents et détérioration des troubles mentaux préexistants durant la pandémie de la COVID-19: une étude canadienne multisite. *Can J Psychiatry*. <https://doi.org/10.1177/0706743720986786>

88. Romer D, Jamieson KH (2020) Conspiracy theories as barriers to controlling the spread of COVID-19 in the U.S. *Soc Sci Med* 263:113356. <https://doi.org/10.1016/j.socscimed.2020.113356>
89. Rossi R, Jannini TB, Succi V, Pacitti F, Lorenzo GD (2021) Stressful life events and resilience during the COVID-19 lockdown measures in Italy: association with mental health outcomes and age. *Front Psychiatry* 12:635832. <https://doi.org/10.3389/fpsy.2021.635832>
90. Saladino V, Algeri D, Auriemma V (2020) The psychological and social impact of Covid-19: new perspectives of well-Being. *Front Psychol*. <https://doi.org/10.3389/fpsyg.2020.577684>
91. Salali GD, Uysal MS (2020) COVID-19 vaccine hesitancy is associated with beliefs on the origin of the novel coronavirus in the UK and Turkey. *Psychol Med*. <https://doi.org/10.1017/S0033291720004067>
92. Salari N, Hosseini-Far A, Jalali R, Vaisi-Raygani A, Rasoulpoor S, Mohammadi M, Rasoulpoor S, Khaledi-Paveh B (2020) Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Global Health* 16(1):57. <https://doi.org/10.1186/s12992-020-00589-w>
93. Sallam M, Dababseh D, Yaseen A, Al-Haidar A, Taim D, Eid H, Ababneh NA, Bakri FG, Mahafzah A (2020) COVID-19 misinformation: Mere harmless delusions or much more? A knowledge and attitude cross-sectional study among the general public residing in Jordan. *PLoS ONE* 15(12):e0243264. <https://doi.org/10.1371/journal.pone.0243264>
94. Sani G, Janiri D, Di Nicola M, Janiri L, Ferretti S, Chieffo D (2020) Mental health during and after the COVID-19 emergency in Italy. *Psychiatry Clin Neurosci* 74(6):372. <https://doi.org/10.1111/pcn.13004>
95. Schneider J, Talamonti D, Gibson B, Forshaw M (2021) Factors mediating the psychological well-being of healthcare workers responding to global pandemics: a systematic review. *J Health Psychol*. <https://doi.org/10.1177/13591053211012759>
96. Shevlin M, Butter S, McBride O, Murphy J, Gibson-Miller J, Hartman TK, Levita L, Mason L, Martinez AP, McKay R, Stocks TVA, Bennett K, Hyland P, Bental RP (2021) Refuting the myth of a “tsunami” of mental ill-health in populations affected by COVID-19: evidence that response to the pandemic is heterogeneous, not homogeneous. *Psychol Med*. <https://doi.org/10.1017/S0033291721001665>
97. Slusarska B, Nowicki GJ, Niedorys-Karczmarczyk B, Chrzan-Rodak A (2022) Prevalence of depression and anxiety in nurses during the first eleven months of the COVID-19 pandemic: a systematic review and meta-analysis. *Int J Environ Res Public Health*. <https://doi.org/10.3390/ijerph19031154>
98. Solomou I, Constantinidou F (2020) Prevalence and predictors of anxiety and depression symptoms during the COVID-19 pandemic and compliance with precautionary measures: age and sex matter. *Int J Environ Res Public Health*. <https://doi.org/10.3390/ijerph17144924>
99. Soveri A, Karlsson LC, Antfolk J, Lindfelt M, Lewandowsky S (2021) Unwillingness to engage in behaviors that protect against COVID-19: the role of conspiracy beliefs, trust, and endorsement of complementary and alternative medicine. *BMC Public Health* 21(1):684. <https://doi.org/10.1186/s12889-021-10643-w>
100. Spielberger CD (2005) State-trait anxiety inventory for adults. Mind Garden, Redwood City California
101. Stuijzand S, Deforges C, Sandoz V, Sajin CT, Jaques C, Elmers J, Horsch A (2020) Psychological impact of an epidemic/pandemic on the mental health of healthcare professionals: a rapid review. *BMC Public Health* 20(1):1230. <https://doi.org/10.1186/s12889-020-09322-z>
102. Sun P, Wang M, Song T, Wu Y, Luo J, Chen L, Yan L (2021) The psychological impact of COVID-19 pandemic on health care workers: a systematic review and meta-analysis. *Front Psychol* 12:626547. <https://doi.org/10.3389/fpsyg.2021.626547>
103. Suryavanshi N, Kadam A, Dhumal G, Nimkar S, Mave V, Gupta A, Cox SR, Gupte N (2020) Mental health and quality of life among healthcare professionals during the COVID-19 pandemic in India. *Brain and behavior* 10(11):e01837. <https://doi.org/10.1002/brb3.1837>
104. Taquet M, Geddes JR, Husain M, Luciano S, Harrison PJ (2021) 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. *Lancet Psychiatry* 8(5):416–427. [https://doi.org/10.1016/S2215-0366\(21\)00084-5](https://doi.org/10.1016/S2215-0366(21)00084-5)
105. Teovanovic P, Lukic P, Zupan Z, Lazic A, Ninkovic M, Zezelj I (2020) Irrational beliefs differentially predict adherence to guidelines and pseudoscientific practices during the COVID-19 pandemic. *Appl Cogn Psychol*. <https://doi.org/10.1002/acp.3770>
106. Tomljenovic H, Bubic A, Erceg N (2020) It just doesn't feel right - the relevance of emotions and intuition for parental vaccine conspiracy beliefs and vaccination uptake. *Psychol Health* 35(5):538–554. <https://doi.org/10.1080/08870446.2019.1673894>
107. Torrente M, Sousa PA, Sanchez-Ramos A, Pimentao J, Royuela A, Franco F, Collazo-Lorduy A, Menasalvas E, Provencio M (2021) To burn-out or not to burn-out: a cross-sectional study in healthcare professionals in Spain during COVID-19 pandemic. *BMJ Open* 11(2):e044945. <https://doi.org/10.1136/bmjopen-2020-044945>
108. Uscinski J, Enders A, Klofstad C, Seelig M, Funchion J, Everett C, Wuchty S, Premaratne K, Murthi M (2020) Why do people believe COVID-19 conspiracy theories? The Harvard Kennedy School Misinformation Review 1 (Special Issue on COVID-19 and Misinformation)
109. Varghese A, George G, Kondaguli SV, Naser AY, Khakha DC, Chatterji R (2021) Decline in the mental health of nurses across the globe during COVID-19: a systematic review and meta-analysis. *J Glob Health* 11:05009. <https://doi.org/10.7189/jogh.11.05009>
110. Verma S, Mishra A (2020) Depression, anxiety, and stress and socio-demographic correlates among general Indian public during COVID-19. *Int J Soc Psychiatry* 66(8):756–762. <https://doi.org/10.1177/0020764020934508>
111. Vlah Tomicevic S, Lang VB (2021) Psychological outcomes amongst family medicine healthcare professionals during COVID-19 outbreak: A cross-sectional study in Croatia. *Eur J Gen Pract* 27(1):184–190. <https://doi.org/10.1080/13814788.2021.1954154>
112. Voorspoels W, Jansen L, Mortier P, Vilagut G, Vocht J, Kessler RC, Alonso J, Bruffaerts R (2021) Positive screens for mental disorders among healthcare professionals during the first covid19 wave in Belgium. *J Psychiatr Res* 140:329–336. <https://doi.org/10.1016/j.jpsychires.2021.05.024>
113. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, Ho RC (2020) Immediate psychological responses and associated factors during the initial stage of the 2019 Coronavirus Disease (COVID-19) epidemic among the general population in China. *Int J Environ Res Public Health*. <https://doi.org/10.3390/ijerph17051729>
114. Wang C, Pan R, Wan X, Tan Y, Xu L, McIntyre RS, Choo FN, Tran B, Ho R, Sharma VK, Ho C (2020) A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun* 87:40–48. <https://doi.org/10.1016/j.bbi.2020.04.028>
115. Wong SYS, Zhang D, Sit RWS, Yip BHK, Chung RY, Wong CKM, Chan DCC, Sun W, Kwok KO, Mercer SW (2020) Impact of COVID-19 on loneliness, mental health, and health service utilisation: a prospective cohort study of older adults with multimorbidity in primary care. *Br J Gen Pract* 70(700):e817–e824. <https://doi.org/10.3399/bjgp20X713021>

116. Yan H, Ding Y, Guo W (2021) Mental health of medical staff during the coronavirus disease 2019 pandemic: a systematic review and meta-analysis. *Psychosom Med* 83(4):387–396. <https://doi.org/10.1097/PSY.0000000000000922>
117. Yanez JA, Afshar Jahanshahi A, Alvarez-Risco A, Li J, Zhang SX (2020) Anxiety, distress, and turnover intention of health-care workers in Peru by their distance to the epicenter during the COVID-19 crisis. *Am J Trop Med Hyg* 103(4):1614–1620. <https://doi.org/10.4269/ajtmh.20-0800>
118. Zerbini G, Ebigbo A, Reicherts P, Kunz M, Messman H (2020) Psychosocial burden of healthcare professionals in times of COVID-19 - a survey conducted at the University Hospital Augsburg. *Ger Med Sci*. <https://doi.org/10.3205/000281>

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Konstantinos N. Fountoulakis¹  · Grigorios N. Karakatsoulis¹  · Seri Abraham^{2,3,4} · Kristina Adorjan⁵ · Helal Uddin Ahmed⁶ · Renato D. Alarcón^{7,8} · Kiyomi Arai⁹ · Sani Salihu Auwal^{10,11} · Julio Bobes^{15,16} · Teresa Bobes-Bascaran^{17,18} · Julie Bourgin-Duchesnay¹⁹ · Cristina Ana Bredicean²⁰ · Laurynas Bukelskis²¹ · Akaki Burkadze^{22,23} · Indira Indiana Cabrera Abud²⁴ · Ruby Castilla-Puentes²⁵  · Marcelo Cetkovich^{26,27} · Hector Colon-Rivera²⁸ · Ricardo Corral^{29,30} · Carla Cortez-Vergara³¹ · Piirika Crepin³² · Domenico de Berardis^{33,34,35} · Sergio Zamora Delgado³⁶ · David de Lucena³⁷ · Avinash de Sousa^{38,39} · Ramona di Stefano⁴⁰ · Seetal Dodd^{12,13,41} · Livia Priyanka Elek⁴² · Anna Elissa⁴³ · Berta Erdelyi-Hamza⁴² · Gamze Erzin⁴⁴ · Martin J. Etchevers⁴⁵ · Peter Falkai⁵ · Adriana Farcas⁴⁶ · Ilya Fedotov⁴⁷ · Viktoriia Filatova⁴⁸ · Nikolaos K. Fountoulakis⁴⁹  · Iryna Frankova⁵⁰ · Francesco Franza^{51,52} · Pedro Frias⁵³ · Tatiana Galako⁵⁴ · Cristian J. Garay⁴⁵ · Leticia Garcia-Álvarez¹⁸ · Paz García-Portilla^{15,55} · Xenia Gonda⁴² · Tomasz M. Gondek⁵⁶ · Daniela Morera González⁵⁷ · Hilary Gould⁵⁸ · Paolo Grandinetti³³ · Arturo Grau^{36,59} · Violeta Groudeva⁶⁰ · Michal Hagin⁶¹ · Takayuki Harada⁶² · Tasdik M. Hasan^{63,64} · Nurul Azreen Hashim⁶⁵ · Jan Hilbig²¹ · Sahadat Hossain⁶⁶ · Rossitza Iakimova⁶⁷ · Mona Ibrahim⁶⁸ · Felicia Iftene⁶⁹ · Yulia Ignatenko⁷⁰ · Matias Irrarazaval⁷¹ · Zaliha Ismail⁷² · Jamila Ismayilova⁷³ · Asaf Jacobs^{74,75} · Miro Jakovljević⁷⁶ · Nenad Jakšić¹⁴ · Afzal Javed^{77,78,79} · Helin Yilmaz Kafali⁸⁰ · Sagar Karia³⁸ · Olga Kazakova⁸¹ · Doaa Khalifa⁶⁸ · Olena Khaustova⁵⁰ · Steve Koh⁵⁸ · Svetlana Kopishinskaia^{82,83} · Korneliia Kosenko⁸⁴ · Sotirios A. Koupidis⁸⁵ · Illes Kovacs⁴² · Barbara Kulig⁴² · Alisha Lalljee³⁹ · Justine Liewig¹⁹ · Abdul Majid⁸⁶ · Evgeniia Malashonkova¹⁹ · Khamelia Malik⁴³ · Najma Iqbal Malik⁸⁷ · Gulay Mammadzada⁸⁸ · Bilvesh Mandalia³⁹ · Donatella Marazziti^{89,90,91} · Darko Marčinko^{14,76} · Stephanie Martinez⁵⁸ · Eimantas Matiekus²¹ · Gabriela Mejia⁵⁸ · Roha Saeed Memon⁹² · Xarah Elenne Meza Martínez⁹³ · Dalia Mickevičiūtė⁹⁴ · Roumen Milev⁶⁹ · Muftau Mohammed⁹⁵ · Alejandro Molina-López⁹⁶ · Petr Morozov⁹⁷ · Nuru Suleiman Muhammad⁹⁸ · Filip Mustač¹⁴ · Mika S. Naor⁹⁹ · Amira Nassieb⁶⁸ · Alvydas Navickas²¹ · Tarek Okasha⁶⁸ · Milena Pandova⁶⁷ · Anca-Livia Panfil¹⁰⁰ · Liliya Panteleeva¹⁰¹ · Ion Papava²⁰ · Mikaela E. Patsali^{102,103} · Alexey Pavlichenko⁷¹ · Bojana Pejuskovic^{104,105} · Mariana Pinto da Costa¹⁰⁶ · Mikhail Popkov¹⁰⁷ · Dina Popovic¹⁰⁸ · Nor Jannah Nasution Raduan⁶⁵ · Francisca Vargas Ramírez^{36,59} · Elmars Rancans^{109,110} · Salmi Razali⁶⁵ · Federico Rebok^{111,112} · Anna Rewekant¹¹³ · Elena Ninoska Reyes Flores¹¹⁴ · María Teresa Rivera-Encinas¹¹⁵ · Pilar A. Saiz¹⁵ · Manuel Sánchez de Carmona¹¹⁶ · David Saucedo Martínez¹¹⁷ · Jo Anne Saw⁶⁵ · Görkem Saygili¹¹⁸ · Patricia Schneiderit¹¹⁹ · Bhumika Shah¹²⁰ · Tomohiro Shirasaka¹²¹ · Ketevan Silagadze²² · Satti Sitanggang¹²² · Oleg Skugarevsky¹²³ · Anna Spikina¹²⁴ · Sridevi Sira Mahalingappa¹²⁵ · Maria Stoyanova⁶⁷ · Anna Szczegieliak¹²⁶ · Simona Claudia Tamasan¹⁰⁰ · Giuseppe Tavormina^{52,127,128} · Maurilio Giuseppe Maria Tavormina⁵² · Pavlos N. Theodorakis¹²⁹ · Mauricio Tohen¹³⁰ · Eva-Maria Tsapakis^{131,132} · Dina Tukhvatullina¹³³ · Irfan Ullah¹³⁴ · Ratnaraj Vaidya¹³⁵ · Johann M. Vega-Dienstmaier¹³⁶ · Jelena Vrublevska^{109,110,140} · Olivera Vukovic^{103,137} · Olga Vysotska¹³⁸ · Natalia Widiasih⁴³ · Anna Yashikhina^{82,139} · Panagiotis E. Prezerakos¹⁴⁰ · Michael Berk^{12,13} · Sarah Levaj¹⁴ · Daria Smirnova^{82,141} 

✉ Grigorios N. Karakatsoulis
gregkarakatsoulis@gmail.com

Konstantinos N. Fountoulakis
Kostasfountoulakis@gmail.com

Seri Abraham
seri.abraham@nhs.net

Kristina Adorjan
Kristina.Adorjan@med.uni-muenchen.de

Helal Uddin Ahmed
soton73@gmail.com

Renato D. Alarcón
renato.alarcon@upch.pe; alarcon.renato@mayo.edu

Kiyomi Arai
k_arai@shinshu-u.ac.jp

Sani Salihu Auwal
auwal01@yahoo.com

Julio Bobes
bobes@uniovi.es

Teresa Bobes-Bascaran
bobesmaria@uniovi.es

Julie Bourgin-Duchesnay
julie.bourgin@gmail.com

Cristina Ana Bredicean
brediceancristina@gmail.com

Laurynas Bukelskis
bukelskis@gmail.com

Akaki Burkadze
dr.burkadze@gmail.com

Indira Indiana Cabrera Abud
indira_ica@hotmail.com

Ruby Castilla-Puentes
rcastil4@its.jnj.com

Marcelo Cetkovich
mccetkovich@ineco.org.ar

Hector Colon-Rivera
hectorcolonriveramd@gmail.com

Ricardo Corral
rcorral33@gmail.com

Carla Cortez-Vergara
carla.cortez.v@upch.pe

Piirika Crepin
piirika.crepin@gmail.com

Domenico de Berardis
domenico.deberardis@aslteramo.it

Sergio Zamora Delgado
szamora@calvomackenna.cl

David de Lucena
alienista@ufc.br; dvdLucena@gmail.com

Avinash de Sousa
avinashdes888@gmail.com

Ramona di Stefano
ramonadist@gmail.com

Seetal Dodd
seetaldodd@gmail.com

Livia Priyanka Elek
elek.livia.priyanka@gmail.com

Anna Elissa
annaelissa.md@gmail.com

Berta Erdelyi-Hamza
hamzaberta@gmail.com

Gamze Erzin
gamze.erzin@gmail.com

Martin J. Etchevers
martinjetchevers@gmail.com

Peter Falkai
Peter.Falkai@med.uni-muenchen.de

Adriana Farcas
6amf@queensu.ca

Ilya Fedotov
ilyafdtv11@gmail.com

Viktoriia Filatova
filatovaviktorija@mail.ru

Nikolaos K. Fountoulakis
nikolasfountoulakis@gmail.com

Iryna Frankova
iryna.frankova@gmail.com

Francesco Franza
franza.francesco@virgilio.it

Pedro Frias
friaspn@gmail.com

Tatiana Galako
tatiana-galako@yandex.ru

Cristian J. Garay
cristiangaray@psi.uba.ar

Leticia Garcia-Álvarez
lettti@gmail.com

Paz García-Portilla
albert@uniovi.es

Xenia Gonda
kendermagos@yahoo.com

Tomasz M. Gondek
tomaszmgondek@gmail.com

Daniela Morera González
danielamorera@gmail.com

Hilary Gould
hgould@health.ucsd.edu

Paolo Grandinetti
grandinetti.paolo@gmail.com

Arturo Grau
doctorgrau@yahoo.com

Violeta Groudeva
violetagroudeva@gmail.com

Michal Hagin
michal.hagin@gmail.com

Takayuki Harada
tkharada77@gmail.com

Tasdik M. Hasan
tasdikhdip@yahoo.com

Nurul Azreen Hashim
azreen@uitm.edu.my

Jan Hilbig
hilbig.jan@gmail.com

Sahadat Hossain
sahadat.hossain@juniv.edu

Rossitza Iakimova
rosica.iakimova@abv.bg

Mona Ibrahim
monaawaad99@gmail.com

Felicia Iftene
iftenef@providencecare.ca

Yulia Ignatenko
hvala_korolevna@rambler.ru

Matias Irarrazaval
matias.irarrazaval@minsai.cl

Zaliha Ismail
zaliha78@uitm.edu.my

Jamila Ismayilova
ismayilova.d@gmail.com

Asaf Jacobs
asafjacobs@gmail.com

Miro Jakovljević
jakovljevic.miro@yahoo.com

Nenad Jakšić
nenad_jaksic@yahoo.com

Afzal Javed
afzalj@gmail.com

Helin Yilmaz Kafali
helinyilmaz136@gmail.com

Sagar Karia
kariabhai117@gmail.com

Olga Kazakova
olga.kazakova.md@gmail.com

Doaa Khalifa
doaaakhalifa72@gmail.com

Olena Khaustova
7974247@gmail.com

Steve Koh
shkoh@ucsd.edu

Svetlana Kopishinskaia
kopishinskaya@gmail.com

Korneliia Kosenko
sun2003@ukr.net

Sotirios A. Koupidis
sotirioskoupidis@yahoo.gr

Illes Kovacs
kovilles@gmail.com

Barbara Kulig
kulig.barbara@hotmail.com

Alisha Lalljee
alishalalljee@gmail.com

Justine Liewig
j.liewig@gmail.com

Abdul Majid
maajid72@gmail.com

Evgeniia Malashonkova
e.malashonkova@gh-nord-essonne.fr

Khamelia Malik
khameliapsi@gmail.com

Najma Iqbal Malik
Najma.iqbal@uos.edu.pk; najmamalik@gmail.com

Gulay Mammadzada
gulay.mammadzada@gmail.com

Bilvesh Mandalia
bilveshmandalia@gmail.com

Donatella Marazziti
dmarazzi@psico.med.unipi.it

Darko Marčinko
predstojnik.psi@kbc-zagreb.hr

Stephanie Martinez
stm032@health.ucsd.edu

Eimantas Matiekus
eimantasmatiekus@yahoo.com

Gabriela Mejia
ggmejia@health.ucsd.edu

Roha Saeed Memon
memon.roha@gmail.com

Xarah Elenne Meza Martínez
xarahmeza22@gmail.com

Dalia Mickeviciūtė
dalia.mickeviciute@gmail.com

Roumen Milev
roumen.milev@queensu.ca

Muftau Mohammed
miftahmuhammad101@gmail.com

Alejandro Molina-López
doctor.alex.psiquiatria@gmail.com

Petr Morozov
prof.morozov@gmail.com

Nuru Suleiman Muhammad
nurusulemuhammad@gmail.com

Filip Mustač
filip.mustac@gmail.com

Mika S. Naor
Mikanaor@mail.tau.ac.il

Amira Nassieb
amiraelbatrawy@gmail.com

Alvydas Navickas
alvydas.navickas@mf.vu.lt

Tarek Okasha
tarek.okasha@gmail.com

Milena Pandova
milena.pandova@gmail.com

Anca-Livia Panfil
anca.livia.panfil@gmail.com

Liliya Panteleeva
p.lilya12@gmail.com

Ion Papava
papava.ion@umft.ro

Mikaella E. Patsali
mikaellapatsali@gmail.com

Alexey Pavlichenko
apavlichenko76@gmail.com

Bojana Pejuskovic
bpejuskovic@hotmail.com

Mariana Pinto da Costa
mariana.pintodacosta@gmail.com

Mikhail Popkov
mihailpopkovanat@gmail.com

Dina Popovic
popovic.dina@gmail.com

Nor Jannah Nasution Raduan
norjannah@uitm.edu.my

Francisca Vargas Ramírez
fvargas.ra@gmail.com

Elmars Rancans
erancans@latnet.lv

Salmi Razali
drsalmi@uitm.edu.my

Federico Rebok
federicorebok@gmail.com

Anna Rewekant
reweana@gmail.com

Elena Ninoska Reyes Flores
elena.reyes@unah.edu.hn

María Teresa Rivera-Encinas
mriverae@usmp.pe

Pilar A. Saiz
frank@uniovi.es

Manuel Sánchez de Carmona
msanchezdecarmona@mac.com

David Saucedo Martínez
davidsaucedomartinez1@gmail.com

Jo Anne Saw
annejosaw@uitm.edu.my

Görkem Saygili
gsaygili@gmail.com

Patricia Schneiderei
p.schneiderei@klinikum-weissenhof.de

Bhumika Shah
bhumikarshah47@gmail.com

Tomohiro Shirasaka
shirasaka.t@gmail.com

Ketevan Silagadze
ketevani26@gmail.com

Satti Sitanggang
sattiradja_96@yahoo.co.id

Oleg Skugarevsky
skugarevsky@gmail.com

Anna Spikina
a-spikina@yandex.ru

Sridevi Sira Mahalingappa
sridevi.siramahalingappa@nhs.net

Maria Stoyanova
mb_milenkova@yahoo.com

Anna Szczegielniak
anna.szczegielniak@gmail.com

Simona Claudia Tamasan
simona_tamasan@yahoo.com

Giuseppe Tavormina
dr.tavormina.g@libero.it

Maurilio Giuseppe Maria Tavormina
mtavormina@virgilio.it

Pavlos N. Theodorakis
theodorakis@who.int

Mauricio Tohen
MTohen@salud.unm.edu

Eva-Maria Tsapakis
emtsapakis@doctors.org.uk

Dina Tukhvatullina
d.tukhvatullina@smd20.qmul.ac.uk

Irfan Ullah
Irfanullahecp2@gmail.com

Ratnaraj Vaidya
r.vaidya@newcastle.ac.uk

Johann M. Vega-Dienstmaier
johann.vega.d@upch.pe; johannvega@yahoo.com

Jelena Vrublevska
vrublevska@inbox.lv

Olivera Vukovic
olivukovic@gmail.com

Olga Vysotska
uafmed@gmail.com

Natalia Widiasih
widiasih_1973@yahoo.com

Anna Yashikhina
akvaraul@mail.ru

Panagiotis E. Prezerakos
prezerpot@gmail.com

Michael Berk
michael.berk@deakin.edu.au

Sarah Levaj
sarahbjedov@gmail.com

Daria Smirnova
daria.smirnova.md.phd@gmail.com

¹ 3rd Department of Psychiatry, School of Medicine, Aristotle University of Thessaloniki Greece, Thessaloniki, Greece

² Pennine Care NHS Foundation Trust, Ashton-under-Lyne, UK

³ Manchester Metropolitan University, Manchester, UK

⁴ Core Psychiatry Training, Health Education England North West, Manchester, UK

⁵ Department of Psychiatry, Ludwig-Maximilians-University, Munich, Germany

- 6 Child Adolescent and Family Psychiatry, National Institute of Mental Health, Dhaka, Bangladesh
- 7 Section of Psychiatry and Mental Health, Universidad Peruana Cayetano Heredia, Facultad de Medicina Alberto Hurtado, Lima, Peru
- 8 Department of Psychiatry and Psychology, Mayo Clinic School of Medicine, Rochester, MN, USA
- 9 School of Medicine and Health Science, Institute of Health Science Shinshu University, Matsumoto, Japan
- 10 Department of Psychiatry, Bayero University, Kano, Nigeria
- 11 Aminu Kano Teaching Hospital, Kano, Nigeria
- 12 IMPACT—the Institute for Mental and Physical Health and Clinical Translation, Deakin University, School of Medicine, Barwon Health, Geelong, Australia
- 13 Orygen The National Centre of Excellence in Youth Mental Health, Centre for Youth Mental Health, Florey Institute for Neuroscience and Mental Health and the Department of Psychiatry, The University of Melbourne, Melbourne, Australia
- 14 Department of Psychiatry and Psychological Medicine, University Hospital Centre Zagreb, Zagreb, Croatia
- 15 Psychiatry Area, Department of Medicine, University of Oviedo, ISPA, INEUROPA. CIBERSAM, Oviedo, Spain
- 16 Department of Psychiatry, Hospital Universitario Central de Asturias, ISPA, INEUROPA. CIBERSAM, Oviedo, Spain
- 17 Mental Health Center of La Corredoria, ISPA, INEUROPA. CIBERSAM, Oviedo, Spain
- 18 Department of Psychology, University of Oviedo, ISPA, INEUROPA. CIBERSAM, Oviedo, Spain
- 19 Division of Child and Adolescent Psychiatry, Department of Psychiatry, Groupe Hospitalier Nord Essonne, Orsay, France
- 20 Department of Neuroscience, Discipline of Psychiatry, “Victor Babes” University of Medicine and Pharmacy, Timisoara, Romania
- 21 Clinic of Psychiatry, Institute of Clinical Medicine, Medical Faculty, Vilnius University, Vilnius, Lithuania
- 22 Mental Hub, Tbilisi, Georgia
- 23 NGO Healthcare Research and Quality Agency, Tbilisi, Georgia
- 24 Hospital San Juan de Dios Hospital, Guadalajara, Mexico
- 25 Janssen Research and Development, Johnson and Johnson, American Society of Hispanic Psychiatry and WARMI Women Mental Health, Cincinnati, OH, USA
- 26 Institute of Translational and Cognitive Neuroscience (INCYT), INECO Foundation, Favaloro University, Buenos Aires, Argentina
- 27 National Scientific and Technical Research Council (CONICET), Buenos Aires, Argentina
- 28 APM Board Certified in General Psychiatry and Neurology, Addiction Psychiatry, and Addiction Medicine, UPMC, DDAP, Philadelphia, USA
- 29 Department of Teaching and Research, Hospital Borda, Buenos Aires, Argentina
- 30 University of Buenos Aires, Buenos Aires, Argentina
- 31 Universidad Peruana Cayetano Heredia, Clínica AngloAmericana, Lima, Perú
- 32 Sanitaire and Social Union for Accompaniment and Prevention, Center of Ambulatory Psychiatry of Narbonne and Lezigan, Narbonne, France
- 33 Department of Mental Health, Psychiatric Service of Diagnosis and Treatment, Hospital “G. Mazzini”, ASL Teramo, Teramo, Italy
- 34 School of Nursing, University of L’Aquila, L’Aquila, Italy
- 35 Department of Neuroscience and Imaging, School of Psychiatry, University of Chieti, Chieti, Italy
- 36 Child and Adolescent Psychiatry Department, Hospital Luis Calvo Mackenna, Santiago, Chile
- 37 Departamento de Fisiología E Farmacología, Universidade Federal Do Ceará, Fortaleza, Ceará, Brazil
- 38 Department of Psychiatry, Lokmanya Tilak Municipal Medical College, Mumbai, India
- 39 Desousa Foundation, Mumbai, India
- 40 Department of Biotechnological and Applied Clinical Sciences, Section of Psychiatry, University of L’Aquila, L’Aquila, Italy
- 41 University Hospital Geelong, Barwon Health, Geelong, VIC, Australia
- 42 Department of Psychiatry and Psychotherapy, Semmelweis University, Budapest, Hungary
- 43 Department of Psychiatry, Faculty of Medicine, Universitas Indonesia, Cipto Mangunkusumo National Referral Hospital, Jakarta, Indonesia
- 44 Psychiatry Department, Ankara Dışkapı Training and Research Hospital, Ankara, Turkey
- 45 Faculty of Psychology, University of Buenos Aires (UBA), Buenos Aires, Argentina
- 46 Centre of Neuroscience, Queen’s University, Kingston, ON, Canada
- 47 Department of Psychiatry and Narcology, Ryazan State Medical University N.a. Academician I.P. Pavlov, Ryazan, Russia
- 48 State Budgetary Institution of the Rostov Region “Psychoneurological Dispensary”, Rostov-On-Don, Russia
- 49 Faculty of Medicine, Medical University of Sofia, Sofia Center, Bulgaria
- 50 Medical Psychology, Psychosomatic Medicine and Psychotherapy Department, Bogomolets National Medical University, Kiev, Ukraine
- 51 Villa Dei Pini Psychiatric Rehabilitation Center, Avellino, Italy
- 52 Psychiatric Studies Centre, Provaglio d’Iseo, Italy
- 53 Hospital Magalhães Lemos, Porto, Portugal
- 54 Department of Psychiatry, Medical Psychology and Drug Abuse, Kyrgyz State Medical Academy, Bishkek, Kyrgyz Republic

- 55 Mental Health Center of La Ería, ISPA, INEUROPA. CIBERSAM, Oviedo, Spain
- 56 Specialty Training Section, Polish Psychiatric Association, Wrocław, Poland
- 57 Instituto Nacional de Psiquiatría Ramón De La Fuente Muñiz, Mexico City, Mexico
- 58 Department of Psychiatry, University of California San Diego, San Diego, USA
- 59 Universidad Diego Portales, Santiago, Chile
- 60 Department of Diagnostic Imaging, University Hospital Saint Ekaterina, Sofia, Bulgaria
- 61 Forensic Psychiatry Unit, Abarbanel Mental Health Center, Bat Yam, Israel
- 62 Faculty of Human Sciences, Education Bureau of the Laboratory Schools, University of Tsukuba, Tokyo, Japan
- 63 Department of Primary Care and Mental Health, University of Liverpool, Liverpool, UK
- 64 Public Health Foundation, Dhaka, Bangladesh
- 65 Department of Psychiatry, Faculty of Medicine, Universiti Teknologi MARA, Sungai Buloh, Selangor, Malaysia
- 66 Department of Public Health and Informatics, Jahangirnagar University, Dhaka, Bangladesh
- 67 Second Psychiatric Clinic, University Hospital for Active Treatment in Neurology and Psychiatry “Saint Naum”, Sofia, Bulgaria
- 68 Faculty of Medicine, Okasha Institute of Psychiatry, Ain Shams University, Cairo, Egypt
- 69 Department of Psychiatry, Queens University, Kingston, ON, Canada
- 70 Mental Health Clinic No 1 N.a. N.A. Alexeev of Moscow Healthcare Department, Education Center, Moscow, Russia
- 71 Ministry of Health, Millenium Institute for Research in Depression and Personality, Santiago, Chile
- 72 Department of Public Health Medicine, Faculty of Medicine, Universiti Teknologi MARA, Sungai Buloh, Selangor, Malaysia
- 73 National Mental Health Center of the Ministry of Health of the Republic of Azerbaijan, Baku, Azerbaijan
- 74 Department of Psychiatry, Westchester Medical Center Health System, Valhalla, NY, USA
- 75 New York Medical College, Valhalla, NY, USA
- 76 School of Medicine, University of Zagreb, Zagreb, Croatia
- 77 Institute of Applied Health Research, University of Birmingham, Birmingham, UK
- 78 Warwick Medical School, University of Warwick, Coventry, UK
- 79 Psychiatric Research Centre, Fountain House, Lahore, Pakistan
- 80 Child Psychiatry Department, Ankara City Hospital, Ankara, Turkey
- 81 Faculty of Medicine, Lund University, Malmö, Sweden
- 82 International Centre for Education and Research in Neuropsychiatry (ICERN), Samara State Medical University, Samara, Russia
- 83 Kirov State Medical University, Kirov, Russia
- 84 Drug Abuse and Psychology Department, Odessa National Medical University, Odessa, Ukraine
- 85 Occupational and Environmental Health Sector, Public Health Policy Department, School of Public Health, University of West Attica, Athens, Greece
- 86 Department of Psychiatry, SKIMS Medical College, Srinagar, India
- 87 Department of Psychology, University of Sargodha, Sargodha, Pakistan
- 88 Department of Psychiatry, Azerbaijan Medical University, Baku, Azerbaijan
- 89 Department of Clinical and Experimental Medicine, Section of Psychiatry, University of Pisa, Pisa, Italy
- 90 Unicamillus, Saint Camillus International University of Health Sciences, Rome, Italy
- 91 Brain Research Foundation Onus, Lucca, Italy
- 92 Dow Medical College, Dow University of Health Sciences, Karachi, Pakistan
- 93 Postgraduate Program in Psychiatry, National Autonomous University of Honduras, Tegucigalpa, Honduras
- 94 Private Outpatient Clinics “JSC InMedica Klinika”, Vilnius, Lithuania
- 95 Department of Clinical Services, Federal Neuropsychiatric Hospital, Kaduna, Nigeria
- 96 General Office for the Psychiatric Services of the Ministry of Health, Mexico City, Mexico
- 97 Department of Postgraduate Education, Russian National Research Medical University N.a. N.I. Pirogov, Moscow, Russia
- 98 Department of Community Medicine, Ahmadu Bello University Teaching Hospital, Zaria, Nigeria
- 99 Sackler School of Medicine New York State American Program, Tel Aviv University, Tel Aviv-Yafo, Israel
- 100 Compartment of Liaison Psychiatry, “Pius Brinzeu” County Emergency Clinical Hospital, Timisoara, Romania
- 101 Department of Medical Psychology, Psychiatry and Psychotherapy, Kyrgyz-Russian Slavic University, Bishkek, Kyrgyz Republic
- 102 School of Social Sciences, Hellenic Open University, Patras, Greece
- 103 Department of Internal Medicine, Nicosia General Hospital, Nicosia, Cyprus
- 104 Faculty of Medicine, University of Belgrade, Belgrade, Serbia
- 105 Clinical Department for Crisis and Affective Disorders, Institute of Mental Health, Belgrade, Serbia
- 106 Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, UK

- ¹⁰⁷ Department of the Introduction to Internal Medicine and Family Medicine, International Higher School of Medicine, Bishkek, Kyrgyz Republic
- ¹⁰⁸ Abarbanel Mental Health Center, Bat-Yam, Israel
- ¹⁰⁹ Department of Psychiatry and Narcology, Riga Stradins University, Riga, Latvia
- ¹¹⁰ Riga Centre of Psychiatry and Narcology, Riga, Latvia
- ¹¹¹ Servicio de Emergencia, Acute Inpatient Unit, Hospital Moyano, Buenos Aires, Argentina
- ¹¹² Argentine Institute of Clinical Psychiatry (IAPC), Buenos Aires, Argentina
- ¹¹³ General Psychiatry Unit I, Greater Poland Neuropsychiatric Center, Kościan, Poland
- ¹¹⁴ Department of Psychiatry, National Autonomous University of Honduras, Tegucigalpa, Honduras
- ¹¹⁵ Centro de Investigación en Salud Pública, Facultad de Medicina, Universidad de San Martín de Porres, Instituto Nacional de Salud Mental "Honorio Delgado – Hideyo Noguchi", Lima, Perú
- ¹¹⁶ Faculty of Health Sciences, Anahuac University, Mexico City, Mexico
- ¹¹⁷ Department of Psychiatry, Escuela Nacional de Medicina, TEC de Monterrey. Servicio de Geriátría. Hospital Universitario "José Eleuterio González" UANL, Monterrey, Nuevo León, México
- ¹¹⁸ Assistant Professor at Cognitive Science and Artificial Intelligence Department, Tilburg University, Tilburg, The Netherlands
- ¹¹⁹ Klinik Für Allgemeine Psychiatrie Und Psychotherapie Ost, Psychiatrische Institutsambulanz, Klinikum Am Weissenhof, Weissenhof, Germany
- ¹²⁰ DY Patil Medical College, Navi Mumbai, India
- ¹²¹ Department of Psychiatry, Teine Keijinkai Medical Center, Sapporo, Japan
- ¹²² Psychiatric Unit, Pambalah Batung General Hospital, South Kalimantan, Amuntai, Indonesia
- ¹²³ Department of Psychiatry and Medical Psychology, Belarusian State Medical University, Minsk, Belarus
- ¹²⁴ Saint Petersburg Psychoneurological Dispensary No2, Saint Petersburg, Russia
- ¹²⁵ Derbyshire Healthcare NHS Foundation Trust, The Liasion Team, Royal Derby Hospital, Derby, Derbyshire, UK
- ¹²⁶ Department of Psychiatric Rehabilitation, Department of Psychiatry and Psychotherapy, Faculty of Medical Sciences in Katowice, Medical University of Silesia, Katowice, Poland
- ¹²⁷ European Depression Association and Italian Association on Depression, Brussels, Belgium
- ¹²⁸ Bedfordshire Center for Mental Health Research, in association with the University of Cambridge, Cambridge, UK
- ¹²⁹ Health Policy, WHO Regional Office for Europe, Copenhagen, Denmark
- ¹³⁰ Department of Psychiatry and Behavioral Sciences, School of Medicine, University of New Mexico, Albuquerque, NM, USA
- ¹³¹ Agios Charalambos Mental Health Clinic, Heraklion, Crete, Greece
- ¹³² 1st Department of Academic Psychiatry, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece
- ¹³³ Centre for Global Public Health, Institute of Population Health Sciences, Queen Mary University of London, London, UK
- ¹³⁴ Kabir Medical College, Gandhara University, Peshawar, Pakistan
- ¹³⁵ Faculty of Medical Sciences, Newcastle University, Newcastle Upon Tyne, UK
- ¹³⁶ Facultad de Medicina Alberto Hurtado, Universidad Peruana Cayetano Heredia, Lima, Perú
- ¹³⁷ Institute of Public Health, Riga Stradins University, Riga, Latvia
- ¹³⁸ Department for Research and Education, Institute of Mental Health, Belgrade, Serbia
- ¹³⁹ Educational and Research Center–Ukrainian Family Medicine Training Center, Bogomolets National Medical University, Kiev, Ukraine
- ¹⁴⁰ Department of Psychiatry, Narcology, Psychotherapy and Clinical Psychology, Samara State Medical University, Samara, Russia
- ¹⁴¹ Department of Nursing, University of Peloponnese, Laboratory of Integrated Health Care, Tripoli, Greece