


ORIGINAL ARTICLE

OPEN

Interleukin-18 signaling promotes activation of hepatic stellate cells in mouse liver fibrosis

Jana Knorr^{1,2} | Benedikt Kaufmann^{3,4} | Maria Eugenia Inzaugarat⁵ |
 Theresa Maria Holtmann¹ | Lukas Geisler¹ | Jana Hundertmark¹ |
 Marlene Sophia Kohlhepp¹ | Laela M. Boosheri³ | Daisy R. Chilin-Fuentes⁶ |
 Amanda Birmingham⁶ | Kathleen M. Fisch⁶ | Joel D. Schilling⁷ |
 Sven H. Loosen⁸ | Christian Trautwein⁵ | Christoph Roderburg⁸ |
 Münevver Demir¹ | Frank Tacke¹ | Hal M. Hoffman³ | Ariel E. Feldstein³ |
 Alexander Wree^{1,3} 

¹Department of Hepatology and Gastroenterology, Charité - Universitätsmedizin Berlin, Campus Virchow Klinikum and Charité Campus Mitte, Berlin, Germany

²Department of Biology, Humboldt-Universität zu Berlin, Berlin, Germany

³Department of Pediatrics, University of California, San Diego, Rady Children's Hospital San Diego California, San Diego, California, USA

⁴Department of Surgery, TUM School of Medicine, Klinikum rechts der Isar, Technical University of Munich, Munich, Germany

⁵Department of Internal Medicine III, RWTH University Hospital Aachen, Aachen, Germany

⁶Center for Computational Biology & Bioinformatics, University of California, San Diego, La Jolla, California, USA

⁷Division of Cardiovascular Diseases, Department of Internal Medicine, Washington University School of Medicine, St. Louis, Missouri, USA

⁸Clinic for Gastroenterology, Hepatology and Infectious Diseases, Medical Faculty of Heinrich Heine University Düsseldorf, University Hospital Düsseldorf, Düsseldorf, Germany

Correspondence

Alexander Wree, Department of Hepatology and Gastroenterology, Charité - Universitätsmedizin Berlin, Augustenburger Platz 1, 13353 Berlin, Germany.
 Email: alexander.wree@charite.de

Funding information

Deutsche Forschungsgemeinschaft, Grant/Award Number: KA 5089/1-1, SFB/ TRR57, TR285/10-2 and WR173/3-1; Deutsche Krebshilfe, Grant/Award Number: 70113000; Foundation for the National Institutes of Health, Grant/Award Number: R01 DK113592, U01 AA024206 and UL1TR001442

Abstract

Background and Aims: Nucleotide-binding oligomerization domain-like receptor-family pyrin domain-containing 3 (NLRP3) inflammasome activation has been shown to result in liver fibrosis. Mechanisms and downstream signaling remain incompletely understood. Here, we studied the role of IL-18 in hepatic stellate cells (HSCs), and its impact on liver fibrosis.

Approach and Results: We observed significantly increased serum levels of IL-18 (128.4 pg/ml vs. 74.9 pg/ml) and IL-18 binding protein (BP; 46.50 ng/ml vs. 15.35 ng/ml) in patients with liver cirrhosis compared with healthy controls. Single cell RNA sequencing data showed that an immunoregulatory subset of

Abbreviations: α SMA, alpha smooth muscle actin; Acta2, actin alpha 2; ALT, alanine aminotransferase; ATP, adenosine triphosphate; BP, binding protein; CCl₄, carbon tetrachloride; CD, control diet; CDAA-HFD, choline-deficient, L-amino acid-defined high fat diet; Col1a1, collagen type I alpha 1; Ctgf, connective tissue growth factor; CreL, cre recombinase under control of an endogenous lysozyme 2 promoter; ECM, extracellular matrix; F4/80, murine macrophage marker; HSCs, hepatic stellate cells; LPS, lipopolysaccharide; Lrat, lecithin retinol acyltransferase; MFB, myofibroblast; Mmp, matrix metalloproteinase; MPO, myeloperoxidase; NLRP3, nucleotide-binding oligomerization domain leucine rich repeat containing receptor-family pyrin domain-containing 3; rm, recombinant mouse; Timp1, tissue inhibitor of matrix metalloproteinase 1; WT, wild type.

Supplemental Digital Content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website, www.hepjournal.com.

Jana Knorr and Benedikt Kaufmann contributed equally.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Copyright © 2023 The Author(s). Published by Wolters Kluwer Health, Inc.

murine HSCs highly expresses *Il18* and *Il18r1*. Treatment of cultured primary murine HSC with recombinant mouse IL-18 accelerated their transdifferentiation into myofibroblasts. *In vivo*, IL-18 receptor-deficient mice had reduced liver fibrosis in a model of fibrosis induced by HSC-specific NLRP3 overactivation. Whole liver RNA sequencing analysis from a murine model of severe NASH-induced fibrosis by feeding a choline-deficient, L-amino acid-defined, high fat diet showed that genes related to IL-18 and its downstream signaling were significantly upregulated, and *Il18*^{-/-} mice receiving this diet for 10 weeks showed protection from fibrotic changes with decreased number of alpha smooth muscle actin-positive cells and collagen deposition. HSC activation triggered by NLRP3 inflammasome activation was abrogated when IL-18 signaling was blocked by its naturally occurring antagonist IL-18BP. Accordingly, we observed that the severe inflammatory phenotype associated with myeloid cell-specific NLRP3 gain-of-function was rescued by IL-18BP.

Conclusions: Our study highlights the role of IL-18 in the development of liver fibrosis by its direct effect on HSC activation identifying IL-18 as a target to treat liver fibrosis.

INTRODUCTION

The cytokine IL-18 was initially described as an interferon gamma-inducing proinflammatory factor in a septic shock model, which stimulated T cell proliferation.^[1] Like IL-1 β , IL-18 belongs to the IL-1 superfamily and is synthesized as biologically inactive 24-kDa precursor protein (pro-IL-18) lacking a signal peptide required for secretion. An intracellular cysteine protease called caspase-1 that is also known as IL-1 β -converting enzyme, cleaves pro-IL-18 into its mature and active form.^[2] Activation of caspase-1 itself is mediated by multiprotein complexes called inflammasomes. The most studied member of the nucleotide-binding oligomerization domain-like receptor (NLR) family is the NLR family pyrin domain-containing 3 (NLRP3) inflammasome, which detects a broad range of endogenous or exogenous danger signals including pathogens and sterile triggers.^[3] IL-18 precursors were shown to be constitutively expressed in peripheral blood mononuclear cells, macrophages, monocytes, keratinocytes, intestinal epithelial cells, Kupffer cells, and in spleen cells. The transcription of IL-18 was not induced by inflammatory stimuli in these cells, demonstrating that IL-18 plays a role in homeostasis and is not regulated by inflammation like IL-1 β .^[4–9] IL-18 signaling through its receptor is mediated by a unique extracellular receptor complex that consists of the ligand-binding chain IL-18R α and the nonbinding signaling-transducing chain IL-18R β .^[10,11]

Besides transcription and caspase-1 processing, IL-18 is also regulated by IL-18 binding protein (BP), which acts as an endogenous antagonist with extremely high affinity.^[12] The 40 kDa glycoprotein lacks a

transmembrane domain and impedes the binding of IL-18 to its receptor, blocking the initiation of signal transduction. In healthy humans, about 2–3 ng/ml of IL-18BP are found in blood circulation, showing a 20-fold molar excess in comparison with IL-18.^[13]

Hepatic fibrosis is a progressive form of chronic liver injury characterized by an excessive deposition of extracellular matrix (ECM) like type I collagen that has been shown to contribute to the development of cirrhosis and hepatocellular carcinoma.^[14] Activation of hepatic stellate cells (HSCs) by other liver cells, predominantly liver macrophages, has been identified to have a pivotal role in fibrogenesis. Under physiological conditions, HSCs maintain in a quiescent state located in the space of Disse and participate in the homeostasis of ECM, regeneration, repair, as well as control of retinoid storage and metabolism. Upon liver damage, quiescent HSCs transdifferentiate into an activated and proliferating state characterized by a myofibroblast (MFB)-like appearance, the loss of retinoid droplets, the expression of alpha smooth muscle actin (α SMA), and the secretion of ECM and cytokines.^[15] However, HSCs can also contribute to the regression of liver fibrosis by undergoing apoptosis or reverting to an inactive HSC state with the capacity of rapid reactivation in the case of profibrotic stimuli.^[16,17]

Recently, our research group demonstrated that NLRP3 activation and pyroptosis are important contributors to liver inflammation and fibrosis.^[18–20] Specifically, we discovered that hyperactivation of NLRP3 in murine HSCs provoked transdifferentiation into an activated MFB phenotype. Moreover, using a genetic gain-of-function

mouse model, we found that persistent NLRP3 activation in HSCs resulted in increased expression of profibrotic genes and proteins and liver fibrosis by 24 weeks of age. These results imply a direct role of NLRP3 inflammasome activation in HSCs and hepatic fibrogenesis.^[21]

Several inflammatory diseases are associated with significantly increased IL-18 expression in serum such as type 2 diabetes mellitus,^[22] rheumatoid arthritis,^[23] Crohn's disease,^[24] and multiple sclerosis.^[25] Growing evidence suggests an important role of IL-18 in liver diseases. Elevated amounts of IL-18 in serum have been observed for patients with primary biliary cholangitis,^[26] biliary atresia,^[27] and chronic liver disease of different etiologies,^[12] and are correlated with disease severity. Further studies demonstrated that IL-18 inhibition by exogenous IL-18BP, neutralizing antibodies, or caspase-1 deficiency protected mice from endotoxin-induced liver injury or nonalcoholic fatty liver disease, whereas IL-1 β blockade does not rescue the phenotype.^[1,28–30] Therefore, we initiated this study to elucidate the role of the proinflammatory cytokine IL-18 on HSC activation and the development of liver fibrosis.

MATERIALS AND METHODS

Patient cohort

Patients were recruited from the Department of Hepatology and Gastroenterology, Charité - Universitätsmedizin Berlin. The study protocol was approved by the ethics committee of Charité - Universitätsmedizin Berlin, Germany (ethical approval number EA2/091/19) and was in accordance with the 1975 Declaration of Helsinki. Diagnosis of liver fibrosis/cirrhosis was based on liver imaging consistent with liver cirrhosis (e.g., nodular hepatic contour, changes in volume distribution indicating portal hypertension in the absence of portal vein thrombosis, secondary phenomena of portal hypertension such as splenomegaly, enlarged caudate lobe, and left lobe lateral segment, regenerative nodules) together with clinical and laboratory signs of portal hypertension/cirrhosis (e.g., low platelets, albumin and prothrombin time, esophageal varices). For all patients with cirrhosis, individual Model for End-Stage Liver Disease scores were calculated. Individuals without evidence of liver disease were recruited as control group. Written informed consent was obtained from all patients.

Single cell RNA sequencing

Gene expression analysis of *Il18* and *Il18r* in HSCs was performed by analyzing recently published data with permission of the authors.^[31] Raw data are available at <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE132662>. In this previously published dataset, activated HSCs were isolated from wild type (WT) mice

(C57BL6/J) receiving carbon tetrachloride (CCl₄) for 3 weeks. Quiescent HSCs were isolated from untreated animals and processed as described in the previous manuscript. Feature plots were generated by using the package “Seurat” (version 2.3.2) for R (version 3.5) (<https://www.r-project.org/>).^[31,32]

Mouse strains

Different mutant *Nlrp3* gain-of-function mouse models of increasing severity were used, including: *Nlrp3*^{D301N^{NeoR}} mice (B6N.129-Nlrp3^{tm3^{Hhf}/J}) carrying the aspartic acid 301 to asparagine (D301N), *Nlrp3*^{A350V^{NeoR}} mice (B6N.129-Nlrp3^{tm1^{Hhf}/J}) carrying the alanine 350 to valine (A350V) substitution, and *Nlrp3*^{L351P^{NeoR}} mice (B6N.129-Nlrp3^{tm2^{Hhf}/J}) carrying the leucine 351 to proline (L351P) substitution.^[33] *Nlrp3*^{A350V^{NeoR}} mice were crossed with mice expressing Cre recombinase under control of the endogenous lysozyme 2 promoter (CreL; B6.129P2-Lyz2^{tm1[cre]lfo/J}), so that mutant *Nlrp3* expression is only restricted to myeloid lineage cells (*Nlrp3*^{A350V CreL}). Furthermore, *Nlrp3*^{A350V CreL} mice were bred to *Il18*^{-/-} (B6.129P2-*Il18*^{tm1Aki/J}) or *Il1r*^{-/-} (B6.129S7-*Il1r1*^{tm1Imx/J}) and *Il18r*^{-/-} (B6.129P2-*Il18r1*^{tm1Aki/J}) mice. *Nlrp3*^{L351P^{NeoR}} mice were crossed with transgenic mice expressing lecithin retinol acyltransferase (*Lrat*)-driven Cre kindly provided by the laboratory of R.F. Schwabe (Columbia University), in which expression of mutant *Nlrp3*^{L351P} is restricted to HSCs (*Nlrp3*^{L351P/+ CreLrat}). *Nlrp3*^{L351P/+ CreLrat} mice were additionally bred with *Il18r*^{-/-} mice to inhibit the IL-18 signaling cascade. The group size ($n = 3–5$) was based on a former publication, the 3R principle, and governmental regulations.^[21] The all-male mice were maintained on a standard rodent chow for 24 weeks. At this time point, mice were sacrificed, and liver tissue and serum were harvested. Depending on the experiment, Cre-negative littermates, WT or PBS-treated mice were used as control group. WT mice had a C57BL/6 background. Genetically modified mice were purchased from The Jackson Laboratory and have been bred with mice with C57BL/6 background over several generations. All experiments were approved by German local governmental authorities (Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen, LANUV NRW) and the Institutional Animal Care and Use Committee at the University of California, San Diego. Mice were housed in filter top cages on a 12-h light/dark cycle with wood shaving bedding, ad libitum access to rodent chow and water unless otherwise noted.

Dietary-related fibrosis induction

8-week-old WT ($n = 7–8$) or *Il18*^{-/-} ($n = 5–6$) male mice were either fed with choline-deficient, L-amino acid-defined high fat diet (CDAA-HFD; 60%; #A06071302;

Research Diets) or with choline-supplemented, L-amino acid-defined control diet (CD) over a period of 10 weeks. Mice were randomly assigned to CDAA-HFD or CD groups so that all groups included mice from several litters.

IL-18BP treatment in mice

Recombinant human IL-18BP (rhIL-18BP; Tadekinig alfa) was provided by AB2 Bio LTD (Lausanne, Switzerland) because of the ability of human IL-18BP to also neutralize murine IL-18.^[34] In this study, *Nlrp3^{D301NneoR+/-}* (B6N.129-Nlrp3^{tm3Hhf/J}) bred to *Lyz2^{tm1[cre]lfo+/-}* mice (*Nlrp3^{D301N} CreL*) to generate myeloid-specific NLRP3 gain-of-function mutants (male and female) were injected beginning at the age of 2 days subcutaneously with 6 µg/g body weight rhIL-18BP ($n = 6$) or PBS ($n = 5$) six times per week. After 14 days of treatment, mice were euthanized, and livers and sera were collected. Mice were randomly assigned to control and treatment groups from several litters.

HSC isolation and *in vitro* stimulation

For isolation of HSCs, 16–24-week-old WT (C57BL/6J) female or male mice were used. Briefly, mice were anesthetized by ketamine/xylazine injection and perfused *in situ* through the inferior vena cava with sequential Pronase E (0.4 mg/ml) and Collagenase D (0.8 mg/ml) solutions. Liver was removed and digested *in vitro* with Collagenase D (0.5 mg/ml), Pronase E (0.5 mg/ml), and DNase I (0.02 mg/ml). After 20 minutes, tissue was filtered through a 70 µm mesh. Cells were separated using a Nycodenz gradient centrifugation. The isolated HSCs were cultured in Dulbecco's Modified Eagle's Medium supplemented with 10% sera plus (PAN Biotech) and 100 U/ml penicillin/streptomycin. After cells attached overnight, HSCs were washed with PBS and stimulated with recombinant mouse (rm)IL-18 (100 ng/ml and 400 ng/ml; R&D, 9139-IL) in sera-free medium for 24 h. In an independent experiment, inflammasome activation in HSCs was induced by lipopolysaccharide (LPS; 1 µg/ml) for 3 h and adenosine triphosphate (ATP; 5 mM) for 1 h. In addition, HSCs were incubated with 400 ng/ml rmIL-18BPd (R&D, 122-BP) to inhibit binding of IL-18 to its receptor. As all substances were either dissolved in PBS or water, culture medium without serum was used as vehicle. In all experiments, cells were incubated at 37°C and 5% CO₂.

Statistical analysis

Statistical analysis was performed with GraphPad Prism 8.0 (GraphPad Software Inc.). Unpaired two-tailed

Student *t* test was used for two-group comparison, whereas analysis of variance followed by Tukey's multiple comparison was used in order to analyze three or more groups. The significance level was set at $\alpha = 5\%$ for all comparisons. Unless otherwise stated, data are presented as means \pm standard deviation.

Other materials and methods are described in the Supporting Materials and Methods.

RESULTS

IL-18 signaling is a key regulator of liver fibrosis/cirrhosis in humans and mice

To investigate the potential role of IL-18 and IL-18BP as serum-based biomarkers in patients with NASH-induced liver cirrhosis, we measured the serum levels of IL-18 and IL-18BP in 38 patients and 50 volunteers. Patient and control characteristics are summarized in Table S1. Interestingly, the level of circulating IL-18 was elevated in patients with liver cirrhosis (128.4 pg/ml; 76.48–209.8 pg/ml) in comparison with controls (74.9 pg/ml; 47.73–103.1 pg/ml) (Figure 1A). A significant increase was also observed for IL-18BP in sera of patients with cirrhosis (46.50 ng/ml; 34.89–64.07 ng/ml) when compared with individuals in the control (15.36 ng/ml; 13.50–17.84 ng/ml) (Figure 1B).

To delineate the expression pattern of IL-18 in quiescent and active HSCs during liver fibrosis, we leveraged single cell RNA sequencing data from a recent publication by Krenkel et al.^[31] In this study, activated HSCs were isolated from mice treated with CCl₄ to induce liver fibrosis. Based on the 50 most significant principal components they could be divided into four subpopulations (myofibroblasts; MFB I to MFB IV). T-distributed stochastic neighbor embedding analysis shows that the expression of *Il18* and *Il18r1* were found to be highly expressed in a subset of activated HSCs (named MFB II) but not in quiescent HSCs (Figure 1C,D). Interestingly, this specific cluster of activated HSCs was characterized as an immunoregulatory subset that expressed various collagens as well as inflammatory mediators, suggesting a potential role in modulating the inflammatory environment in their surroundings.^[31]

Stimulation with IL-18 activated HSCs, whereas IL-18BP mitigated LPS-induced HSC activation

To characterize the direct effect of IL-18 on HSCs, we stimulated primary mouse HSCs from WT livers with two different concentrations of rmIL-18 (100 ng/ml and 400 ng/ml) for 24 h. Immunofluorescence staining demonstrated an increase in α SMA expression but not for collagen I in primary HSCs treated with rmIL-18. These




FIGURE 1 IL-18 signaling impacts liver fibrosis/cirrhosis in humans and mice. (A) Circulating IL-18 levels in patients with NASH-induced liver cirrhosis (128.4 pg/ml; 76.48–209.8 pg/ml) compared with healthy controls (74.9 pg/ml; 47.73–103.1 pg/ml). (B) Serum IL-18BP of patients with liver cirrhosis (46.50 ng/ml; 34.89–64.07 ng/ml) in comparison with the control group (15.36 ng/ml; 13.50–17.84 ng/ml). Data shown: median; IQR. (Control, $n = 50$; Liver cirrhosis, $n = 38$; **** $p < 0.0001$; Mann–Whitney U test). T-SNE plots mapping distribution of HSCs expressing (C) *Il18* and (D) *Il18r1*. Activated HSCs from CCl_4 -treated mice that induce liver fibrosis were differentiated into four subpopulations (MFB I to MFB IV) according to their different gene expression patterns.^[31] T-SNE plots show that especially HSCs from the immunoregulatory MFB II cluster express *Il18* and *Il18r1*. BP, binding protein; CCl_4 , carbon tetrachloride; HSC, hepatic stellate cell; IQR, interquartile range; MFB, myofibroblast; t-SNE, T-distributed stochastic neighbor embedding.

findings were observed especially for the higher dosage (Figure 2A,B) because some cells already show αSMA -positive stress fibers that were not observed for the lower concentration. Correspondingly, IL-18 treatment induced the expression of profibrotic genes including *connective tissue growth factor (Ctgf)*, *collagen type I alpha 1 (Col1a1)*, and *actin alpha 2 (Acta2)* the gene of αSMA (Figure 2C).

Next, we investigated the impact of inhibiting IL-18 signaling using its naturally occurring antagonist IL-

18BP on primary HSCs from mouse livers. We found that HSC activation induced by LPS/ATP treatment was attenuated by IL-18BP, which was evidenced by decreased expression of αSMA protein (Figure 2D) and significantly downregulated levels of *Ctgf*, *Col1a1*, and *Acta2* close to basal conditions (Figure 2E). Taken together, these *in vitro* results suggest that IL-18 signaling plays a role in HSC activation, and that IL-18BP inhibits self-amplifying activation, which might have an impact on fibrogenesis.

IL-18 signaling deficiency prevented liver fibrosis induced by persistent NLRP3 expression in HSCs

As described previously by our group, the enzyme *Lrat* is uniquely expressed in liver by HSCs and breeding of *Lrat* Cre expressing mice to *Nlrp3^{L351P/+}* (*Nlrp3^{L351P/+}CreLrat*) generates a model of NLRP3 hyperactivation in HSCs that is associated with “spontaneous” liver fibrosis.^[21] We used this model to examine IL-18 signaling deficiency in mice with early fibrosis by crossing these mice with IL-18 receptor knockout animals (*Il18r^{-/-}*). In *Il18r^{-/-}* mice, mRNA levels of *Acta2* and *Col1a1* were significantly reduced compared with *Nlrp3^{L351P/+}CreLrat Il18r^{+/+}* littermates (Figure 3A). Sirius Red liver staining displayed increased collagen deposition in *Nlrp3^{L351P/+}CreLrat Il18r^{+/+}* compared with WT control, which was reduced to baseline conditions in *Nlrp3^{L351P/+}CreLrat Il18r^{-/-}* (Figure 3B,D). Interestingly, mice with IL-18 signaling deficiency revealed a trend toward reduced macrophage infiltration represented by quantification of murine macrophage marker (F4/80) positive cells in liver sections (Figure 3B and E). Importantly, serum levels of alanine aminotransferase (ALT)/aspartate aminotransferase and hematoxylin–eosin staining of liver tissue indicated that liver injury was similar between the genotypes (Figure 3B,C).

IL-18 signaling and downstream regulatory network are activated in CDAA-HFD-induced fibrotic NASH

To assess the relevance of IL-18 signaling in a model of diet-induced fibrosis, we used CDAA-HFD because this diet recapitulates many features of human NASH with fibrosis.^[35] Mice were put on CDAA-HFD or CD for 10 weeks, and the gene expression profile of whole liver tissue was analyzed by RNA sequencing (Figure 4A). IL-18 signaling was activated in mice fed a CDAA-HFD compared with mice fed a CD (Figure 4B). Genes including *Il18r1*, *Il18 receptor accessory protein*, and *Il18bp* were upregulated in mice fed a CDAA-HFD, associated with increased expression of genes downstream of IL-18 signaling such as *Cxcl10*, *Il16*, and *Fth1*. In addition, inflammasome components that act upstream of IL-18 activation were also upregulated including *Nlrp3* and *Casp1*. Fibrosis development is a major characteristic of NASH. Mice fed a CDAA-HFD also showed a strong activation of fibrosis-related genes including *Col1a1*, *Col3a1*, *Vimentin*, *tissue inhibitor of matrix metalloproteinase 1* (*Timp1*), *Acta2*, and transforming growth factor beta (Figure 4C). These data highlight that many molecular players involved in IL-18 pathway are transcriptionally induced in fibrotic NASH.

IL-18 knockout mice are protected from CDAA-HFD-induced fibrosis

Based on the RNA sequencing data, we next determined the role of ablation of IL-18 in NASH associated liver fibrosis. CDAA-HFD-fed animals showed prominent macrovesicular steatosis with incipient NASH in both WT and *Il18r^{-/-}* mice (Figure S1A). In addition, CDAA-HFD mice had increased liver weight and lower body weight (Figure S1B) compared with those mice fed with CD. Liver sections from CDAA-HFD-fed *Il18r^{-/-}* mice displayed less intrahepatic collagen accumulation and bridging fibrosis (Figure 5A,B) compared with WT mice on CDAA-HFD that was supported by a trend toward reduced amount of hydroxyproline in the *Il18r^{-/-}* liver tissue (Figure 5C). In line with the fibrosis data, the positive area of the HSC activation marker α SMA (Figure 5D,E) was significantly reduced in *Il18r^{-/-}* liver tissue. The gene expression levels of *Ctgf*, *Col3a*, *Timp1*, and *Vimentin* increased in response to CDAA-HFD in both groups compared with CD, but to a reduced extent in *Il18r^{-/-}* mice (Figure 5F). To analyze the impact of ductular reaction, which is a common feature in CDAA-HFD-fed mice,^[36] the amount of cytokeratin 19 (CK19)-positive cells (apart from mature bile ducts, reactive cholangiocytes and hepatic progenitor cells) were determined. Both, WT and *Il18r^{-/-}* mice show ductular reaction, but without significant difference (Figure S1D,E).

IL-18BP treatment alleviated liver inflammation and fibrosis in myeloid cell-specific *Nlrp3* mutant mice

In a previous study we demonstrated that disruption of IL-1 β signaling partially rescued the hepatic and systemic inflammatory phenotype concomitant with growth impairment and shortened lifespan of myeloid cell-specific *Nlrp3* mutant mice (Figure S2).^[18] Interestingly, IL-18 ablation combined with IL-1R deficiency by generating a double knockout (*Nlrp3^{A350V} CreL Il1r^{-/-} Il18r^{-/-}*) displayed significantly mitigated liver fibrosis (Figure S2).

Based on the critical role of endogenous IL-18 for HSC activation and fibrosis, we hypothesized that IL-18 antagonism, by its natural inhibitor IL-18BP, could be a strategy to attenuate liver fibrosis. To investigate therapeutic effects of rhIL-18BP, we injected *Nlrp3^{D301N} CreL* mice with IL-18BP or PBS (vehicle) subcutaneously for 14 days. The characteristic growth retardation and the urticarial-like rash of *Nlrp3^{D301N} CreL* mice were ameliorated by rhIL-18BP treatment (Figure 6A). In the *Nlrp3^{D301N} CreL* mice treated with vehicle, liver histology demonstrated severe inflammation with numerous inflammatory foci and infiltrating cells. However, the mice treated with rhIL-18BP had a substantial attenuation of the liver inflammatory response (Figure 6B). In particular, infiltrating neutrophils




FIGURE 2 Stimulation with recombinant IL-18 accelerates HSC activation *in vitro*, whereas blockage of IL-18 signaling inhibits HSC activation. (A) Representative immunofluorescence images and (B) their quantification by the corrected total cell fluorescence showing α SMA and Collagen I of HSCs after treatment with either 100 ng/ml or 400 ng/ml mrlL-18 after 24 h. (C) mRNA levels of *Ctgf*, *Col1a1*, and *Acta2* of primary murine HSCs after mrlL-18 stimulation. (D) Immunofluorescence images and (E) mRNA levels of *Ctgf*, *Col1a1*, and *Acta2* of primary HSCs treated with LPS/ATP and additional pretreatment with mIL-18BP. α SMA, alpha smooth muscle actin; ATP, adenosine triphosphate; BP, binding protein; *Col1a1*, collagen type I alpha 1; *Ctgf*, connective tissue growth factor; HSC, hepatic stellate cell; LPS, lipopolysaccharide; rm, recombinant mouse. 3–4 independent replicates were included for all measured values; upper scale bars represent 50 μ m. * $p < 0.05$; ** $p < 0.01$; **** $p < 0.0001$.




FIGURE 3 IL-18 receptor deficiency significantly reduced expression of the HSC activation marker in 24-week-old mice with HSC-specific NLRP3 hyperactivation. (A) mRNA levels of *Ctgf*, *Col1a1*, and *Acta2* in control, *Nlrp3^{L351P/+}CreLrat Il18^{+/+}*, and *Nlrp3^{L351P/+}CreLrat Il18^{-/-}* mice. (B) Representative pictures of H&E-, Sirius Red-, and F4/80-stained liver sections. (C) Liver aminotransferases and quantification of (D) Sirius Red and (E) F4/80-positive cells in mice with IL-18R deficiency compared with *Nlrp3^{L351P/+}CreLrat Il18^{+/+}* and control mice. *Acta2*, actin alpha 2; *Col1a1*, collagen type I alpha 1; *Ctgf*, connective tissue growth factor; F4/80, murine macrophage marker; H&E, hematoxylin–eosin; HSC, hepatic stellate cell; Lrat, lecithin retinol acyltransferase; NLRP3, nucleotide-binding oligomerization domain leucine rich repeat containing receptor-family pyrin domain-containing 3. $n = 3$ –5 mice per group for all measured values; scale bars represent 100 μm . * $p < 0.05$; ** $p < 0.01$.

were prevalent in liver of nontreated mice visualized by myeloperoxidase (MPO) staining, but not in rhIL-18BP-treated ones (Figure S3A,B). Consistently, liver mRNA levels of immune cell markers and chemoattractants (*Tnfa*, *Cxcl1*, *Ly6C*, *Ly6G*, *Mpo*) as well as matrix metalloproteinase 8 (*Mmp8*, neutrophil collagenase) were significantly downregulated for *Nlrp3^{D301N} CreL* mice injected with IL-18BP, which are abundantly expressed by macrophages, granulocytes, and/or neutrophils (Figure S3C). Furthermore, we observed a reduction of collagen deposition in treated mice compared with vehicle visualized by Sirius Red staining (Figure 6C) as well as significantly decreased gene expression level of the ECM regulating enzyme *Timp1*

(Figure 6D). In contrast, *Mmp10* and *Mmp13* were increased in treated mice compared with vehicle group (Figure 6D). Interestingly, immunofluorescence staining for αSMA and platelet-derived growth factor receptor (PDGFR) show that IL-18BP-treated mice show fewer activated, αSMA -expressing HSCs, suggesting that IL-18 is a mediator of HSC activation (Figure 6E,F).

DISCUSSION

In the present study, we found a significant elevation of circulating IL-18 and IL-18BP in patients with liver

cirrhosis in comparison with healthy controls. Subsequently, we provided strong evidence that IL-18 signaling and the corresponding downstream signaling cascade play a pivotal role in experimental NASH-induced liver fibrogenesis, which was confirmed by *Il18*^{-/-} mice that showed protection from the fibrotic changes in a NASH model. Finally, we identified IL-18BP as a therapeutic approach that has potential for the treatment of liver fibrosis.

IL-18 is a pleiotropic cytokine with proinflammatory properties, whose biological activity is controlled by IL-18BP. It has been reported that IL-18 is involved in the activation and differentiation of T helper cells in concert with IL-12 or IL-15, as well as in activation of natural killer cells.^[37] In line with our results, Ludwiczek and colleagues indicated a parallel elevation of IL-18 and IL-18BP in patients' sera with different stages of liver cirrhosis induced by different etiologies. Interestingly, patients with end-stage liver disease (Child-Pugh stage C) presented a decreased level of IL-18BP and therefore significantly elevated amounts of unbound IL-18, suggesting a causative role of IL-18 in the development and progression of liver diseases.^[12] Of importance, in patients with cirrhosis who undergo liver transplantation, the serum level of IL-18 normalized, indicating a key role of the liver in IL-18 production.^[38] Furthermore, IL-18 has been associated previously with progression of cardiac,^[39] pulmonary,^[40] and renal fibrosis.^[41,42] The activation of HSCs is a well-studied and essential event in fibrogenesis by their transdifferentiation into a proliferative and migratory MFB-like cell type. Current data indicate that stimulation with IL-18 *in vitro* directly induced HSC activation by upregulation of *Ctgf*, *Col1a1*, and *Acta2*, as well as protein expression of α SMA in a concentration-dependent manner (Figure 2). These observations were supported *in vivo* by using an HSC-specific NLRP3 overactivation fibrosis model that is independent of inflammation and liver injury.^[21] We demonstrated that collagen deposition (Sirius Red) and key fibrosis-associated genes (*Acta2*, *Col1a1*) are decreased in the absence of IL-18 signaling in *Nlrp3*^{L351P/+} *CreLrat* *Il18*^{-/-} mice (Figure 3).

In line with our results that IL-18 induces transition of quiescent HSCs to α SMA-expressing MFB, inhibition of IL-18 signaling showed a protective effect on renal as well as pulmonary fibrosis due to inhibition of cell transdifferentiation into MFB.^[40,42] This leads to the assumption that IL-18 might drive cell differentiation. It has been reported that hepatic IL-18 expression and its release mainly originated from Kupffer cells and macrophages.^[1,43] Inflammasome activation has previously been implicated in the development of fibrosis.^[18,20,44] To this end, fibrogenesis in different experimental mouse models was suppressed by inflammasome-related (*Nlrp3*, *caspase-1*, *Il1b*) knockout mice

in which the production and gene expression of *Col1a* and *Acta2* was diminished.^[20,45,46] Correspondingly, persistent global expression of NLRP3 resulted in severe liver inflammation and advanced stages of liver fibrosis by increased HSC activation and remarkable collagen deposition. Blocking IL-1 β signaling by an IL-1 receptor antagonist (anakinra) decreased liver inflammation but did not change HSC activation level or the degree of liver fibrosis.^[18] Further studies have observed bivalent results regarding the role of IL-1 β in liver fibrosis depending on the mouse model. In ATP-binding cassette transporter b4 knockout animals, fibrosis was associated with increased gene expression of *Il1b*. *In vitro* experiments with primary HSCs revealed decreased proliferation when treated with anakinra, but anakinra failed to improve liver injury or fibrosis *in vivo*.^[47] In another study, IL-1 receptor antagonist knockout mice exhibited more fibrosis in the bile duct ligation model, but less fibrosis in the CCl₄ model. In contrast, when WT mice were treated with anakinra, fibrosis was improved after bile duct ligation but aggravated after CCl₄ injections.^[48]

However, the role of IL-18 was not investigated in these prior studies. Our findings indicate the profibrotic capability of IL-18 in liver disease. Sequencing data demonstrated that the gene regulatory network both up and downstream of IL-18 including compartments of inflammasomes and IL-18-related ligands are significantly increased in our murine fibrotic NASH model. Further experiments elucidated that IL-18 deficiency was protective against the fibrotic changes when fed with CDAA-HFD. Especially the collagen- and α SMA-positive area as well as profibrotic genes were significantly reduced in the *Il18*^{-/-} mice compared with WT, which highlights again the impact of IL-18 signaling on HSC activation and fibrogenesis. Hohenester et al. showed that liver injury in nonalcoholic fatty liver disease, which increases the risk for fibrosis, was dependent on IL-18 but not IL-1 β signaling.^[30] *Il18*^{-/-} mice were protected from hepatocellular injury assessed by reduced ALT levels in a sedentary lifestyle-induced obesity syndrome-diet mouse model, but it was not further supported by histological stainings or other methods.^[30] In contrast, in our study serum ALT levels showed no difference when comparing WT and *Il18*^{-/-} mice on CDAA-HFD, which might be caused by the strong inflammatory phenotype related to this diet.

Using mice that have persistent hyperactivation of NLRP3 in myeloid cells (*Nlrp3*^{A350V} *CreL*), a double knockout of *Il1r* and *Il18* prevented liver fibrosis whereas the *Il1r* single knockout had no protection. Brydges et al. demonstrated that 9-day-old *Nlrp3*^{A350V} mice on an *Il18*^{-/-} background feature almost no neutrophilic inflammation compared with *Nlrp3*^{A350V} mice on an *Il1r*^{-/-} background, which might drive the development of fibrosis in these mice when aging.^[49] These findings suggest that IL-18 signaling may have a more prominent




FIGURE 4 IL-18 signaling and downstream regulatory network are activated in CDAA-HFD-induced fibrotic NASH. Bulk RNA sequencing analysis of liver tissue after 10 weeks of CDAA-HFD or CD. (A) An overview of experimental design. (B) Clustering of IL-18 signaling-related gene transcripts. (C) Clustering of fibrosis-related gene transcripts. CD, control diet; CDAA-HFD, choline-deficient, L-amino acid-defined high fat diet. $n = 4$ mice per group.

role in fibrogenesis than IL-1 β . In our experiments, administration of IL-18BP not only alleviated inflammation and liver fibrosis in the highly inflammatory *Nlrp3*^{D301N} *CreL* mouse model (Figure 6B,C), but also significantly reduced neutrophil infiltration assessed by

MPO staining (Figure S3). Interestingly, treatment with IL-18BP also resulted in remarkably lower number of α SMA-positive HSCs in the liver (Figure 6E), which points out the correlation of IL-18BP on HSC activation. Accordingly, *in vitro* experiments with IL-18BP-treated




FIGURE 5 IL-18 deficiency reduced CDAA-HFD-induced fibrosis. (A, B) Collagen deposition assessed by Sirius Red staining and (C) hydroxyproline, (D, E) αSMA-positive cells, and (F) mRNA expression levels of the profibrotic genes *Ctgf*, *Col3a*, *Timp1*, and *Vimentin* in WT and *Il18*^{-/-} mice fed with CDAA-HFD compared with mice on CD. αSMA, alpha smooth muscle actin; CD, control diet; CDAA-HFD, choline-deficient, L-amino acid-defined high fat diet; *Ctgf*, connective tissue growth factor; *Col3a*, collagen type III alpha 1 chain; *Timp1*, tissue inhibitor of matrix metalloproteinase 1; WT, wild type. $n = 5-8$ mice per group for all measured values; scale bars represent 250 μm . * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

primary HSCs inhibited their activation suggesting that IL-18BP binds IL-18 released in response to NLRP3 inflammasome activation by LPS/ATP treatment in order to interrupt their self-amplifying activation (Figure 2). Consistent with our results, other studies have shown that IL-18 neutralization resulted in reduced neutrophil

accumulation in experimental hepatic ischemia/reperfusion injury and lethal endotoxemia.^[29,50] IL-18BP prevents mature IL-18 from binding to the IL-18 receptor by virtue of its exceptionally high binding affinity that is greater than IL-18R α .^[10,11] Recently, rhIL-18BP (Tadekinig Alfa) was tested in clinical trial phase II for




FIGURE 6 Treatment of *Nlrp3*^{D301N} *CreL* mice with IL-18BP rescued highly inflammatory phenotype. *Nlrp3*^{D301N} *CreL* mice were treated with 6 μ g/g rhIL-18BP (Tadekinig Alfa) or PBS from day 2 to 14. (A) Growth impairment, rash, and reduced body weight, consistent with the phenotype of *Nlrp3*^{D301N} *CreL* mice, were clearly normalized by IL-18BP treatment, whereas serum alanine aminotransferase (ALT) did not significantly change. (B) H&E-stained liver sections and quantification showing less inflammation in IL-18BP-treated mice. (C) Sirius Red staining. (D) mRNA expression of ECM regulating enzymes (*Timp1*, *Mmp10*, *Mmp13*) were measured from total liver lysate. (E) Immunofluorescence staining and (F) quantification of PDGFR and α SMA showing more activated HSCs in the nontreated mice. α SMA, alpha smooth muscle actin; BP, binding protein; ECM, extracellular matrix; H&E, hematoxylin–eosin; HSCs, hepatic stellate cells; Mmp, matrix metalloproteinase; Nlrp3, nucleotide-binding oligomerization domain leucine rich repeat containing receptor-family pyrin domain-containing 3; PDGFR, platelet-derived growth factor receptor; Timp1, tissue inhibitor of matrix metalloproteinase 1; rh, recombinant human. $n = 5$ –6 mice per group for all measured values; black scale bars represent 250 μ m; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

the autoinflammatory disorder Adult-onset Still's disease.^[51] In summary, our results highlight the pivotal role of IL-18 signaling in liver fibrogenesis through the activation of HSCs *in vitro* and *in vivo* in three different mouse models. Its blockage by the use of IL-18BP, as well as potential other approaches such as neutralizing antibodies, may provide a therapeutic strategy for the treatment of liver fibrosis.

AUTHOR CONTRIBUTIONS

Study concept, design and supervision: Alexander Wree, Ariel E. Feldstein, Maria Eugenia Inzaugarat. Acquisition of data: Jana Knorr, Benedikt Kaufmann, Maria Eugenia Inzaugarat, Laela M. Boosheri, Jana Hundertmark, Lukas Geisler, Marlene Sophia Kohlhepp. Analysis and interpretation of data: Jana Knorr, Benedikt Kaufmann, Theresa Maria Holtmann. Drafting of the manuscript: Jana Knorr. Critical revision of the manuscript for important intellectual content: Benedikt Kaufmann, Alexander Wree, Frank Tacke, Ariel E. Feldstein, Hal M. Hoffman, Maria Eugenia Inzaugarat, Münevver Demir, Christian Trautwein, **Joel D. Schilling**. Provided samples and study materials: Münevver Demir, Christoph Roderburg, Hal M. Hoffman, Sven H. Loosen. Obtained funding: Alexander Wree, Ariel E. Feldstein, Hal M. Hoffman, Benedikt Kaufmann, Christian Trautwein.

ACKNOWLEDGMENTS

The authors wish to thank AB2 Bio LTD for providing recombinant human rhIL-18BP (Tadekinig alfa) and discussions related to its use, and Dr. Cem Gabay for advice on the experimental conditions.

FUNDING INFORMATION

This work was funded by National Institutes of Health (NIH) grants R01 DK113592 to Hal M. Hoffman and Ariel E. Feldstein, U01 AA024206 to Ariel E. Feldstein, German Research Foundation (WR173/3-1 TR285/10-2 and SFB/TRR57) to Alexander Wree and Christian Trautwein, German Cancer Aid (Deutsche Krebshilfe 70113000) to Alexander Wree and Christian Trautwein, German Research Foundation (DFG-Grant KA 5089/1-1) to Benedikt Kaufmann. In addition, the project described was partially supported by the NIH grant UL1TR001442 of the Clinical and Translational Science Awards Program. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

CONFLICTS OF INTEREST

Frank Tacke advises and received grants from Gilead. He consults for Novo Nordisk and Pfizer. He is on the speakers' bureau for Falk and AbbVie. He received grants from Inventiva, Bristol-Myers Squibb, and Allergan. Hal M. Hoffman consults for Novartis.

DATA AVAILABILITY STATEMENT

Single cell sequencing data are deposited in Gene Expression Omnibus with accession number GSE132662 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE132662>). Other RNA sequencing data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Alexander Wree  <https://orcid.org/0000-0002-2968-1335>

REFERENCES

- Okamura H, Tsutsi H, Komatsu T, Yutsudo M, Hakura A, Tanimoto T, et al. Cloning of a new cytokine that induces IFN-gamma production by T cells. *Nature*. 1995;378(6552):88–91.
- Ghayur T, Banerjee S, Hugunin M, Butler D, Herzog L, Carter A, et al. Caspase-1 processes IFN-gamma-inducing factor and regulates LPS-induced IFN-gamma production. *Nature*. 1997;386(6625):619–23.
- Guo H, Callaway JB, Ting JPY. Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nat Med*. 2015;21(7):677–87.
- Puren AJ, Fantuzzi G, Dinarello CA. Gene expression, synthesis, and secretion of interleukin 18 and interleukin 1beta are differentially regulated in human blood mononuclear cells and mouse spleen cells. *Proc Natl Acad Sci USA*. 1999;96(5):2256–61.
- Mehta VB, Hart J, Wewers MD. ATP-stimulated release of interleukin (IL)-1beta and IL-18 requires priming by lipopolysaccharide and is independent of caspase-1 cleavage. *J Biol Chem*. 2001;276(6):3820–6.
- Gulinelli S, Salaro E, Vuerich M, Bozzato D, Pizzirani C, Bolognesi G, et al. IL-18 associates to microvesicles shed from human macrophages by a LPS/TLR-4 independent mechanism in response to P2X receptor stimulation. *Eur J Immunol*. 2012;42(12):3334–45.
- Naik SM, Cannon G, Burbach GJ, Singh SR, Swerlick RA, Wilcox JN, et al. Human keratinocytes constitutively express interleukin-18 and secrete biologically active interleukin-18 after treatment with pro-inflammatory mediators and dinitrochlorobenzene. *J Invest Dermatol*. 1999;113(5):766–72.
- Pizarro TT, Michie MH, Bentz M, Woraratanadharm J, Smith MFJ, Foley E, et al. IL-18, a novel immunoregulatory cytokine, is up-regulated in crohn's disease: expression and localization in intestinal mucosal cells. *J Immunol*. 1999;162(11):6829–35.
- Matsui K, Yoshimoto T, Tsutsui H, Hyodo Y, Hayashi N, Hiroishi K, et al. Propionibacterium acnes treatment diminishes CD4+ NK1.1+ T cells but induces type I T cells in the liver by induction of IL-12 and IL-18 production from Kupffer cells. *J Immunol*. 1997;159(1):97–106.
- Born TL, Thomassen E, Bird TA, Sims JE. Cloning of a novel receptor subunit, AcPL, required for interleukin-18 signaling. *J Biol Chem*. 1998;273(45):29445–50.
- Tsutsumi N, Kimura T, Arita K, Ariyoshi M, Ohnishi H, Yamamoto T, et al. The structural basis for receptor recognition of human interleukin-18. *Nat Commun*. 2014;5(1):5340.
- Ludwiczek O, Kaser A, Novick D, Dinarello CA, Rubinstein M, Vogel W, et al. Plasma levels of interleukin-18 and interleukin-18 binding protein are elevated in patients with chronic liver disease. *J Clin Immunol*. 2002;22(6):331–7.
- Novick D, Schwartsburd B, Pinkus R, Suissa D, Belzer I, Sthoeger Z, et al. A novel IL-18BP ELISA shows elevated serum

- IL-18BP in sepsis and extensive decrease of free IL-18. *Cytokine*. 2001;14(6):334–42.
14. Yin C, Evason KJ, Asahina K, Stainier DYR. Hepatic stellate cells in liver development, regeneration, and cancer. *J Clin Invest*. 2013;123(5):1902–10.
 15. Tsuchida T, Friedman SL. Mechanisms of hepatic stellate cell activation. *Nat Rev Gastroenterol Hepatol*. 2017;14(7):397–411.
 16. Kisseleva T, Cong M, Paik Y, Scholten D, Jiang C, Benner C, et al. Myofibroblasts revert to an inactive phenotype during regression of liver fibrosis. *Proc Natl Acad Sci USA*. 2012;109(24):9448–53.
 17. Troeger JS, Mederacke I, Gwak GY, Dapito DH, Mu X, Hsu CC, et al. Deactivation of hepatic stellate cells during liver fibrosis resolution in mice. *Gastroenterology*. 2012;143(4):1073–83.e22.
 18. Wree A, Eguchi A, McGeough MD, Pena CA, Johnson CD, Canbay A, et al. NLRP3 inflammasome activation results in hepatocyte pyroptosis, liver inflammation, and fibrosis in mice. *Hepatology*. 2014;59(3):898–910.
 19. Wree A, McGeough MD, Inzaugarat ME, Eguchi A, Schuster S, Johnson CD, et al. NLRP3 inflammasome driven liver injury and fibrosis: roles of IL-17 and TNF in mice. *Hepatology*. 2018;67(2):736–49.
 20. Wree A, McGeough MD, Peña CA, Schlattjan M, Li H, Inzaugarat ME, et al. NLRP3 inflammasome activation is required for fibrosis development in NAFLD. *J Mol Med (Berl)*. 2014;92(10):1069–82.
 21. Inzaugarat ME, Johnson CD, Holtmann TM, McGeough MD, Trautwein C, Papouchado BG, et al. NLR family pyrin domain-containing 3 inflammasome activation in hepatic stellate cells induces liver fibrosis in mice. *Hepatology*. 2019;69(2):845–59.
 22. Moriwaki Y, Yamamoto T, Shibutani Y, Aoki E, Tsutsumi Z, Takahashi S, et al. Elevated levels of interleukin-18 and tumor necrosis factor- α in serum of patients with type 2 diabetes mellitus: relationship with diabetic nephropathy. *Metabolism*. 2003;52(5):605–8.
 23. Yamamura M, Kawashima M, Taniai M, Yamauchi H, Tanimoto T, Kurimoto M, et al. Interferon- γ -inducing activity of interleukin-18 in the joint with rheumatoid arthritis. *Arthritis Rheum*. 2001;44(2):275–85.
 24. Ludwiczek O, Kaser A, Novick D, Dinarello CA, Rubinstein M, Tilg H. Elevated systemic levels of free interleukin-18 (IL-18) in patients with Crohn's disease. *Eur Cytokine Netw*. 2005;16(1):27–33.
 25. Losy J, Niezgodna A. IL-18 in patients with multiple sclerosis. *Acta Neurol Scand*. 2001;104(3):171–3.
 26. Yamano T, Higashi T, Nouso K, Nakatsukasa H, Kariyama K, Yumoto E, et al. Serum interferon- γ -inducing factor/IL-18 levels in primary biliary cirrhosis. *Clin Exp Immunol*. 2000;122(2):227–31.
 27. Urushihara N, Iwagaki H, Yagi T, Kohka H, Kobashi K, Morimoto Y, et al. Elevation of serum interleukin-18 levels and activation of kupffer cells in biliary atresia. *J Pediatr Surg*. 2000;35(3):446–9.
 28. Faggioni R, Cattley RC, Guo J, Flores S, Brown H, Qi M, et al. IL-18-binding protein protects against lipopolysaccharide-induced lethality and prevents the development of Fas/Fas ligand-mediated models of liver disease in mice. *J Immunol*. 2001;167(10):5913–20.
 29. Netea MG, Fantuzzi G, Kullberg BJ, Stuyt RJL, Pulido EJ, McIntyre RC, et al. Neutralization of IL-18 reduces neutrophil tissue accumulation and protects mice against lethal *Escherichia coli* and *Salmonella typhimurium* endotoxemia. *J Immunol*. 2000;164(5):2644–9.
 30. Hohenester S, Kanitz V, Schiergens T, Einer C, Nagel J, Wimmer R, et al. IL-18 but not IL-1 signaling is pivotal for the initiation of liver injury in murine non-alcoholic fatty liver disease. *Int J Mol Sci*. 2020;21(22):8602.
 31. Krenkel O, Hundertmark J, Ritz TP, Weiskirchen R, Tacke F. Single cell RNA sequencing identifies subsets of hepatic stellate cells and myofibroblasts in liver fibrosis. *Cells*. 2019;8(5):503.
 32. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell gene expression data. *Nat Biotechnol*. 2015;33(5):495–502.
 33. Brydges SD, Mueller JL, McGeough MD, Pena CA, Misaghi A, Gandhi C, et al. Inflammasome-mediated disease animal models reveal roles for innate but not adaptive immunity. *Immunity*. 2009;30(6):875–87.
 34. Kim SH, Eisenstein M, Reznikov L, Fantuzzi G, Novick D, Rubinstein M, et al. Structural requirements of six naturally occurring isoforms of the IL-18 binding protein to inhibit IL-18. *Proc Natl Acad Sci USA*. 2000;97(3):1190–5.
 35. Matsumoto M, Hada N, Sakamaki Y, Uno A, Shiga T, Tanaka C, et al. An improved mouse model that rapidly develops fibrosis in non-alcoholic steatohepatitis. *Int J Exp Pathol*. 2013;94(2):93–103.
 36. Wei G, An P, Vaid KA, Nasser I, Huang P, Tan L, et al. Comparison of murine steatohepatitis models identifies a dietary intervention with robust fibrosis, ductular reaction, and rapid progression to cirrhosis and cancer. *Am J Physiol Gastrointest Liver Physiol*. 2020;318(1):G174–88.
 37. Kaplanski G. Interleukin-18: biological properties and role in disease pathogenesis. *Immunol Rev*. 2018;281(1):138–53.
 38. Yasudo H, Ando T, Maehara A, Ando T, Izawa K, Tanabe A, et al. A possible association between a nucleotide-binding domain LRR-containing protein family PYD-containing protein 1 mutation and an autoinflammatory disease involving liver cirrhosis. *Hepatology*. 2021;74:2296–9.
 39. Xiao H, Li H, Wang JJ, Zhang JS, Shen J, An XB, et al. IL-18 cleavage triggers cardiac inflammation and fibrosis upon β -adrenergic insult. *Eur Heart J*. 2018;39(1):60–9.
 40. Zhang LM, Zhang Y, Fei C, Zhang J, Wang L, Yi ZW, et al. Neutralization of IL-18 by IL-18 binding protein ameliorates bleomycin-induced pulmonary fibrosis via inhibition of epithelial-mesenchymal transition. *Biochem Biophys Res Commun*. 2019;508(2):660–6.
 41. Liang H, Xu F, Zhang T, Huang J, Guan Q, Wang H, et al. Inhibition of IL-18 reduces renal fibrosis after ischemia-reperfusion. *Biomed Pharmacother*. 2018;106:879–89.
 42. Bani-Hani AH, Leslie JA, Asanuma H, Dinarello CA, Campbell MT, Meldrum DR, et al. IL-18 neutralization ameliorates obstruction-induced epithelial-mesenchymal transition and renal fibrosis. *Kidney Int*. 2009;76(5):500–11.
 43. Seki E, Tsutsui H, Nakano H, Tsuji NM, Hoshino K, Adachi O, et al. Lipopolysaccharide-induced IL-18 secretion from murine Kupffer cells independently of myeloid differentiation factor 88 that is critically involved in induction of production of IL-12 and IL-1 β . *J Immunol*. 2001;166(4):2651–7.
 44. Mridha AR, Wree A, Robertson AAB, Yeh MM, Johnson CD, Van Rooyen DM, et al. NLRP3 inflammasome blockade reduces liver inflammation and fibrosis in experimental NASH in mice. *J Hepatol*. 2017;66(5):1037–46.
 45. Dixon LJ, Berk M, Thapaliya S, Papouchado BG, Feldstein AE. Caspase-1-mediated regulation of fibrogenesis in diet-induced steatohepatitis. *Lab Invest*. 2012;92(5):713–23.
 46. Gieling RG, Wallace K, Han YP. Interleukin-1 participates in the progression from liver injury to fibrosis. *Am J Physiol Gastrointest Liver Physiol*. 2009;296(6):G1324–31.
 47. Reiter FP, Wimmer R, Wottke L, Artmann R, Nagel JM, Carranza MO, et al. Role of interleukin-1 and its antagonism of hepatic stellate cell proliferation and liver fibrosis in the *Abcb4*($-/-$) mouse model. *World J Hepatol*. 2016;8(8):401–10.
 48. Meier RPH, Meyer J, Montanari E, Lacotte S, Balaphas A, Muller YD, et al. Interleukin-1 receptor antagonist modulates liver inflammation and fibrosis in mice in a model-dependent manner. *Int J Mol Sci*. 2019;20(6):1295.
 49. Brydges SD, Broderick L, McGeough MD, Pena CA, Mueller JL, Hoffman HM. Divergence of IL-1, IL-18, and cell death in NLRP3 inflammasomopathies. *J Clin Invest*. 2013;123(11):4695–705.

50. Takeuchi D, Yoshidome H, Kato A, Ito H, Kimura F, Shimizu H, et al. Interleukin 18 causes hepatic ischemia/reperfusion injury by suppressing anti-inflammatory cytokine expression in mice. *Hepatology*. 2004;39(3):699–710.
51. Gabay C, Fautrel B, Rech J, Spertini F, Feist E, Kötter I, et al. Open-label, multicentre, dose-escalating phase II clinical trial on the safety and efficacy of tadekinig alfa (IL-18BP) in adult-onset Still's disease. *Ann Rheum Dis*. 2018;77(6):840–7.

How to cite this article: Knorr J, Kaufmann B, Inzaugarat ME, Holtmann TM, Geisler L, Hundertmark J, et al. Interleukin-18 signaling promotes activation of hepatic stellate cells in mouse liver fibrosis. *Hepatology*. 2023;77:1968–1982. <https://doi.org/10.1002/hep.32776>