

Research



**Cite this article:** Nguyen T, Zimmer L, Hoehl S. 2023 Your turn, my turn. Neural synchrony in mother–infant proto-conversation. *Phil. Trans. R. Soc. B* **378**: 20210488. <https://doi.org/10.1098/rstb.2021.0488>

Received: 5 May 2022  
Accepted: 11 October 2022

One contribution of 15 to a discussion meeting issue ‘Face2face: advancing the science of social interaction’.

**Subject Areas:**  
behaviour, neuroscience

**Keywords:**  
mother–infant communication, functional near-infrared spectroscopy, hyperscanning, synchrony, turn-taking, vocalizations

**Author for correspondence:**  
Trinh Nguyen  
e-mail: [trinh.nguyen@iit.it](mailto:trinh.nguyen@iit.it)

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6412288>.

# Your turn, my turn. Neural synchrony in mother–infant proto-conversation

Trinh Nguyen<sup>1,2</sup>, Lucie Zimmer<sup>3</sup> and Stefanie Hoehl<sup>1</sup>

<sup>1</sup>Department of Developmental and Educational Psychology, University of Vienna, Liebiggasse 5, 1010 Vienna, Austria

<sup>2</sup>Neuroscience of Perception and Action Lab, Italian Institute of Technology (IIT), Center for Life Nano and Neuro Science, Viale Regina Elena 291, 00161 Rome, Italy

<sup>3</sup>Department of Psychology, Ludwig-Maximilians University Munich, Leopoldstrasse 13, 80802 Munich, Germany

**id** TN, 0000-0003-0420-2147; LZ, 0000-0002-5766-2991; SH, 0000-0003-0472-0374

Even before infants utter their first words, they engage in highly coordinated vocal exchanges with their caregivers. During these so-called proto-conversations, caregiver–infant dyads use a presumably universal communication structure—turn-taking, which has been linked to favourable developmental outcomes. However, little is known about potential mechanisms involved in early turn-taking. Previous research pointed to interpersonal synchronization of brain activity between adults and preschool-aged children during turn-taking. Here, we assessed caregivers and infants at 4–6 months of age ( $N = 55$ ) during a face-to-face interaction. We used functional-near infrared spectroscopy hyperscanning to measure dyads’ brain activity and microcoded their turn-taking. We also measured infants’ inter-hemispheric connectivity as an index for brain maturity and later vocabulary size and attachment security as developmental outcomes potentially linked to turn-taking. The results showed that more frequent turn-taking was related to interpersonal neural synchrony, but the strength of the relation decreased over the course of the proto-conversation. Importantly, turn-taking was positively associated with infant brain maturity and later vocabulary size, but not with later attachment security. Taken together, these findings shed light on mechanisms facilitating preverbal turn-taking and stress the importance of emerging turn-taking for child brain and language development.

This article is part of a discussion meeting issue ‘Face2face: advancing the science of social interaction’.

## 1. Introduction

Long before acquiring verbal language, human infants engage with their caregivers in reciprocal vocal exchanges, so-called proto-conversations [1–6], which have a strikingly similar temporal structure to later verbal conversations [7]. Both types of exchanges are characterized by a rapid back-and-forth (i.e. turn-taking) between the interaction partners, with short lags between the turns and limited overlaps. This temporal structure is documented across many languages and cultures, suggesting that the turn-taking system might be a universal feature of human communication [8]. In the current study, we set out to explore the early origins of turn-taking, rooted in the infant–caregiver interaction. What are the mechanisms of early turn-taking, and what are its functions in preverbal human infants?

In human adults, turn-taking enables highly efficient verbal conversations [9]. Across languages, response latencies between speakers are remarkably short (around 200 ms on average), implying that speakers must predict the speech act of their interlocutor while still listening, and they must plan their response before their partner’s speech offset [10]. Combined with rare overlaps of partners speaking simultaneously, these cognitive processes ensure a fluid conversation and highly efficient exchange of information. Turn-taking is a

fundamentally cooperative endeavour as partners must pay close attention to each other to enable such fine-grained mutual coordination. Accordingly, shorter response latencies have been associated with greater feelings of connection between interaction partners [11].

In early human ontogeny, infants as young as eight weeks of age actively engage in turn-taking in vocal interactions with their caregivers [1]. These early exchanges already display a remarkable level of temporal coordination [12]. In the first six months after birth, infants' response latencies are quite similar to latencies observed in adult conversations [2]. The early ontogenetic onset of turn-taking in human social exchanges and the structural similarity of preverbal proto-conversations with later verbal conversations beg the question of whether similar mechanisms are involved in early and later turn-taking. The objective of the current study is twofold: (i) we aim for a better understanding of the mechanisms of early turn-taking exchanges between young infants and their mothers, and (ii) we want to assess potential links between early dyadic turn-taking and later developmental outcomes, in particular infant attachment quality and vocabulary size.

### (a) Mechanisms of early turn-taking

As turn-taking depends on precise temporal coordination, one key candidate underlying mechanism is the mutual entrainment and synchronization of endogenous neural oscillators in the brains of the conversation partners [13]. According to this view, a listener's neural oscillations become entrained to the rhythms of their partner's speech and vice versa, leading to mutual synchronization of brain rhythms [14] which enables fine-tuned conversational turn-taking. Indeed, ample empirical evidence shows that listeners' neural oscillations entrain to perceived speech rhythms in adults [15,16] and infants [17,18]. Using hyperscanning, meaning simultaneous measurements of brain activity of at least two participants, interpersonal synchronization of brain rhythms has been documented in verbal conversations between adults [19,20] and between caregivers and their preschool-aged children [21]. The growing body of research underscores the involvement of temporal and especially frontal brain regions in neural synchrony during a conversation [22]. Synchrony in these brain regions has been related to precise mutual predictions, which are integral cognitive processes to turn-taking. Indeed, neural synchronization during mother-child conversation increased more steeply when dyads took more turns in their conversation [21].

Importantly, we know little about when the relation between interpersonal neural synchrony and turn-taking emerges during development. Gaining insights into the functional role of interpersonal neural synchrony is important as it could pave the way for interventions promoting neural synchrony and its related benefits [23,24]. Furthermore, if turn-taking is linked to neural synchrony already in infant-caregiver interactions this might point to neural synchrony as a potential shared mechanism underlying turn-taking across ontogeny. Given that brain rhythms undergo protracted development in humans [25], mutual adaptation of rhythmic brain activities, resulting in neural synchrony during turn-taking, could be an important factor in shaping the development of infant brain dynamics [26]. In the present study, we therefore tested whether interpersonal neural synchrony relates to turn-taking already in mother-infant dyads. We hypothesized

that turn-taking is linked to higher mother-infant neural synchrony, which should increase over the course of a free proto-conversation. Specifically, we predicted that higher interpersonal neural synchrony relates to more frequent and faster turns, indicating higher turn-taking quality [7] (Hypothesis 1).

### (b) Links between early turn-taking and developmental outcomes

As infants continue to partake in early proto-conversations, including turn-taking, they become more competent at turn-taking and increasingly initiate turn-taking sequences [1]. We suggest that individual differences in becoming more mature participants in proto-conversations might be reflected in variability in infants' functional brain network connectivity [27]. Especially functional connectivity in the homologous inter-hemispheric network, which is considered a marker for neural maturity [27,28], could be linked to infants' early turn-taking abilities. According to previous literature, these abilities should include more frequent turn-taking initiated by the infant as well as faster infant responses to turns initiated by the mother [1,2,7]. In the present study, we measured infant maturity at a neural level using inter-hemispheric connectivity and linked it to turn-taking behaviour between mothers and their infants. We hypothesized that more frequent infant-initiated turn-taking and, potentially, turns with shorter response latencies are associated with higher inter-hemispheric connectivity (Hypothesis 2).

Early turn-taking has been associated with developmental benefits in language processing and acquisition [10,29]. For instance, higher rates of turn-taking have been related to advanced language skills [30] and larger vocabulary in 2-year-olds [31]. By contrast, weaker vocal coordination of preverbal infants has been linked to poorer cognitive outcomes [12]. In line with these findings, we hypothesized that more frequent turns between mothers and their infants and turns with shorter response latencies are linked to larger vocabulary at two years of age (Hypothesis 3).

Next to infants' individual development, turn-taking in mother-infant proto-conversations is assumed to be one fundamental feature of interpersonal connectedness [11,22] and the basis for parent-child bonding [32]. More specifically, insecurely attached children, a measure of parent-child relationship quality [33], show longer response latencies during preverbal turn-taking than securely attached children. Still, there is little evidence on the role of preverbal turn-taking for attachment development, even though there is ample evidence for the role of behaviourally coordinated interactions for attachment security. In the present study, we thus examined whether turn-taking patterns in mother-infant proto-conversation can inform us about infants' attachment development. Accordingly, we assumed that more frequent turns and faster turns, i.e. shorter response latencies, are related to higher attachment security (Hypothesis 4).

## 2. Methods

### (a) Participants

Fifty-five mother-infant dyads (25 female infants; M age = 145.36 days, s.d. = 16.93 days; range = 121–180 days) were included in the final sample. Infants were all born healthy and full-term (after the 37th gestation week and with a birth weight of more

than 2.500 g). Twenty-six additional dyads of the original sample ( $N=81$ ) were excluded from the neural synchrony analysis due to early termination of the experiment ( $n=15$ ), not codable video/audio ( $n=5$ ), and bad functional near-infrared spectroscopy (fNIRS) signal quality ( $n=6$ ). Subsequent sample sizes were further decreased because parents either did not return the language development questionnaire ( $n=27$ ) or did not participate in the home visits to assess attachment ( $n=23$ ). Mothers were, on average, 33.35 years old (s.d. = 4.80 years; range = 24–44 years), of European White origin, and highly educated (76% of mothers graduated with a university degree). Participants were recruited from the database of the Department of Developmental Psychology (Wiener Kinderstudien). Most of the registered families were recruited at Vienna General Hospital, and their children were added to the database after their parents gave their written consent. Mothers and infants received a small present for their participation. The local Ethics Committee approved the study (reference no. 00352).

Previous publications, including data from the same sample, comprised analyses on the role of proximity and touch in neural and physiological synchrony [34] and intrapersonal coupling between prefrontal brain activity and respiratory sinus arrhythmia in infants and adults [35].

## (b) Procedure

Mothers and infants participated in three conditions: distal joint watching, proximal joint watching and free play. In the joint watching conditions, the infant was seated in a highchair (distal joint watching) or in the mother's lap (proximal joint watching) and watched a calm aquarium video for 90 s. The order of distal and proximal joint watching conditions was counterbalanced. In the free play condition, mother and infant were instructed to freely play face-to-face, as they would at home, but without toys and without singing for five minutes. The free play condition always followed the two watching conditions. Only interpersonal synchrony data from the free play condition are reported here (see [36] for a comparison of experimental conditions). Mothers' and infants' brain activity was measured using fNIRS. Their behaviour was micro-coded using ELAN (v. 5.9; Max Planck Institute for Psycholinguistics, The Language Archive, Nijmegen). The dyads' electrocardiography was also measured but analysed in separate papers [34,35]. Mothers were sent questionnaires on affect, postnatal depression, attachment style, and infant temperament (not reported here) before the first visit to the laboratory.

At 12 months of age, we visited families at their homes, observed the mother–infant dyad for at least 90 min, and rated them on the Attachment Q-Sort [37]. Parents filled out a questionnaire on infants' expressive language skills (ELFRA-2 [38]) when infants were 24 months old.

## (c) Measures

### (i) fNIRS

We used two NIRSport 8-8 (NIRx Medizintechnik GmbH, Germany) devices in the tandem setting and simultaneously recorded concentration changes in oxy-haemoglobin (HbO) and deoxy-haemoglobin (HbR) in mother and infant. The  $8 \times 2$  probe sets were attached to an electroencephalography cap with a 10–20 configuration (electronic supplementary material, figure S1). The probe sets over the left and right inferior frontal gyrus (IFG) surrounded F7 and F8, whereas the probes on the medial prefrontal area (mPFC) surrounded FP1 and FP2. These regions of interest were based on previous work involving adult–child interactions [24,39]. In each probe set, eight sources and eight detectors were positioned, which resulted in 22 measurement channels with equal distances of approximately 2.3 cm between the infants'

optodes and 3 cm between the mothers' optodes. The absorption of near-infrared light was measured at the wavelengths 760 and 850 nm, and the sampling frequency was 7.81 Hz.

fNIRS measurements were processed using MATLAB-based functions derived from Homer 2 [40]. Raw data were converted into optical density. Next, optical density data were motion-corrected with a wavelet-based algorithm with an interquartile range of 0.5. Motion-corrected time series were further visually inspected during a quality check procedure (see [41] for further information). Before continuing, we removed 22.87% of the channels from both mother and child from further analyses due to bad signal-to-noise ratio and motion artefacts. Then, slow drifts and physiological noise were removed from the signals using a band-pass second-order Butterworth filter with 0.01 and 0.5 Hz cut-offs. Based on the modified Beer-Lambert law, we converted the filtered data to changes ( $\mu\text{Mol}$ ) in HbO and HbR. HbR analyses are reported in the supplements.

We assessed the relation between the fNIRS time series in each caregiver and infant using (Morlet) wavelet transform coherence (WTC) as a function of frequency and time [42]. WTC is more suitable than correlational approaches, as it is invariant to inter-regional differences in the hemodynamic response function (HRF) [43]. On the other hand, correlations are sensitive to the shape of the HRF, which is assumed to be different between individuals (especially of different ages) and different brain areas. A high correlation may be observed among regions with no blood flow fluctuations. Based on previous studies [21], visual inspection, and spectral analyses, the frequency band of 0.063–0.167 Hz (corresponding to 6–16 s) was identified as the frequency of interest. In the present study, coherence was calculated for each channel combination in 60-second intervals of the interactive free play condition, resulting in 5 (intervals)  $\times$  22 (channels) coherence values for each dyad.

Using WTC, we assessed infants' intra-personal inter-hemispheric functional connectivity between homologous channels in the passive distal watching condition (as an age-appropriate resting phase). The frequency band of interest was determined to be 0.063 Hz–0.167 Hz (corresponding to 6–16 s), according to the potential infant HRF function [44], by visual inspection and spectral analyses. The channels were assigned to the right and left hemispheres, resulting in ten channel combinations connecting both hemispheres. The two channels placed in the middle were excluded for this analysis, as no calculation of inter-hemispheric functional connectivity was possible due to their spatial proximity. Coherence values were calculated per channel combination and then averaged, resulting in one WTC value for each infant.

### (ii) Turn-taking coding

To assess vocal turn-taking in the free play condition (adapted from [1,45]), trained graduate students coded audio recordings of the free-play sessions using ELAN (v. 5.9; Max Planck Institute for Psycholinguistics, The Language Archive, Nijmegen). The experimental sessions were recorded at 25 frames per second. Vocalizations of mother and infant (excluding crying) were micro-coded frame-by-frame. Subsequently, we analysed vocalizations in terms of turn-taking, which is defined as turns that are taken up by others [1]. A turn-taking sequence included two related turns separated by 0–3 s. We assessed a) the number of turn-taking sequences from mother to infant (mother vocalization followed by infant vocalization), b) the number of turn-taking sequences from infant to mother (infant vocalization followed by mother vocalization), c) the number of bi-directional turn-taking sequences (sum of all turn-taking sequences), d) mother response latency, e) infant response latency, f) overall response latencies, g) the number of mother-to-infant overlaps (when infant's vocalization overlaps the mother's preceding vocalization), h) mother-to-infant overlap duration, i) infant-to-mother overlaps, and j) infant-to-mother

overlap duration. Two trained research assistants coded 25% of randomly chosen videos to establish inter-rater reliability. Inter-rater reliability for vocalizations was high (mother vocalizations:  $\kappa = 0.98$ , infant vocalizations:  $\kappa = 0.78$ ).

### (iii) Language outcome questionnaire

The ELFRA-2 [38] is a parent-report questionnaire to assess infants' expressive language skills at two years of age. The questionnaire comprises a list of 260 words, 25 questions on syntax and 11 questions on morphology. In the present study, we only used the word list to assess infants' word production at 24 months. Based on the Communicative Development Inventories [46], the list includes 20 categories, such as animals, verbs and food. The reliability of the items was high, Cronbach's alpha = 0.99.

### (iv) Attachment assessment

We used the Attachment Q-Sort (AQS [37]) to observe the attachment security of the mother–infant dyads at 12 months of age. Two trained research assistants observed the dyad for at least 90 min during each home visit and independently evaluated the attachment security. The interrater reliability between raters averaged at  $r = 0.75$ , thus, an averaged AQS score was used in subsequent analyses. These AQS scores ranged between  $-1.0$  and  $+1.0$ , with higher scores representing higher attachment security of the observed dyad.

### (v) Statistical analysis

All statistical analyses were calculated in RStudio (RStudio Team, 2020). We used generalized linear mixed models (GLMM), generalized linear models (GLM), and linear models (LM) to analyse the hypothesized associations between turn-taking, interpersonal neural synchrony, inter-hemispheric functional connectivity, infant's language development and attachment.

Hypothesis 1: Turn-taking and interpersonal neural synchrony over time. WTC values were entered as the response variable (assuming a beta distribution and logit link) of a GLMM with interval (1–5), the number of bidirectional turn-taking, response latencies, and region of interest (ROI: IFG versus IPFC versus mPFC) as fixed factors. We included the interaction between intervals, number of turns, and ROI. ROI was inserted as random slopes and dyads as random intercepts. Random slopes for interval and turn-taking did not lead to model convergence and were thus excluded. To further examine significant main and interaction effects, *post-hoc* analyses (emmeans) were applied to contrast factors. Multiple comparisons were corrected by using Tukey's honest significant difference. All continuous predictor variables were z-standardized, and distributions of residuals were visually inspected for each model. Models were estimated using maximum likelihood. Model fit was compared using a Chi-square test (likelihood ratio test [47]).

Hypothesis 2: Turn-taking and infants' inter-hemispheric connectivity. We calculated two separate GLM including inter-hemispheric connectivity values as the response variable (assuming a Gaussian distribution and logit link). The first GLM tested infant response latencies, and number of infant-to-mother turns as the main effects. The second GLM tested the effects of mother-to-infant and infant-to-mother overlaps. Further turn-taking variables are considered in analyses reported in the supplements.

Hypothesis 3: Infants' expressive language abilities and turn-taking. Infants' vocabulary at 24 months assessed by the ELFRA questionnaire was entered as the response variable (assuming a Gaussian distribution) of three LM. One included the main effects of dyadic turn-taking (number of turns, response latency). Another one included the main effects of infant turn-taking variables (number of turns, response latency, number of overlaps and

overlap duration). The third LM included mother turn-taking variables (number of turns and response latency).

Hypothesis 4: Infants' attachment and turn-taking. Infants' attachment security score at 12 months of age assessed by the AQS was entered as the response variable (Fisher's z-transformed, assuming a Gaussian distribution) of three LM. One included the main effects of dyadic turn-taking (number of turns, response latency). Another one included the main effects of infant turn-taking variables (number of turns, response latency, number of overlaps and overlap duration). The third LM included mother turn-taking variables (number of turns and response latency).

The *p*-values of the multiple GLMMTMB/GLM/LM were corrected for multiple comparisons when necessary (false discovery rate [36]). We also conducted a power analysis (using a bespoke web app [48]) in relation to the statistical analysis on the sample size of 55. According to previous research [21], we assumed a medium effect size ( $d = 0.5$ ), which yielded  $1 - \beta = 0.856$ . We, thus, deemed the sample size to be appropriate for this set of analyses.


## 3. Results

### (a) Turn-taking patterns

Descriptive statistics for the turn-taking patterns of mother–child dyads ( $N = 55$ ) throughout the free play condition are included in electronic supplementary material, table S1. Number of bidirectional turn-taking and response latencies were highly correlated (turn-taking:  $\beta = 0.869$ , s.e. = 0.068,  $t_{53} = 12.78$ ,  $p < 0.001$ ; response latencies:  $\beta = 0.541$ , s.e. = 0.116,  $t_{53} = 4.685$ ,  $p < 0.001$ ). Neither frequency of turn-taking nor overlaps from mother to infant or infant to mother were significantly related,  $p > 0.189$ .

### (b) Turn-taking and interpersonal neural synchrony over time

Firstly, we analysed whether more frequent turn-taking during a proto-conversation was related to mother–infant neural synchrony in frontal regions. The GLMM ( $n = 55$ ) provided a significantly better fit to the data than the null model ( $\chi^2 = 28.472$ , d.f. = 11,  $p = 0.002$ ). The comparison of the full model with the reduced models revealed that the three-way interaction turns\*intervals\*ROI ( $\chi^2 = 11.914$ , d.f. = 2,  $p = 0.003$ ) and the main effect ROI ( $\chi^2 = 11.920$ , d.f. = 2,  $p = 0.003$ ) were statistically significant. These effects indicate that interpersonal neural synchrony was different between brain regions and that this difference was modulated by the frequency of turn-taking and that their slopes were different over time. Neither the two-way interactions turn-taking\*intervals, intervals\*ROI, turn-taking\*ROI nor the simple effects of turn-taking, intervals and response latencies reached significance ( $p > 0.139$ ). To understand the three-way interaction, we conducted three separate GLMM to test the interaction between turns and intervals in relation to interpersonal neural synchrony in each ROI. We corrected *p*-values (false discovery rate) for multiple comparisons. The GLMM for the inferior frontal gyrus and lateral prefrontal cortex revealed no significant main effects and interaction effects ( $p > 0.270$ ). Only the GLMM for the medial prefrontal cortex revealed a significant main effect of turns (estimate = 0.005, s.e. = 0.002, 95% CI = [0.002 0.010],  $z = 2.742$ ,  $p = 0.006$ , *p*-corrected = 0.009), a significant main effect of interval (estimate = 0.049, s.e. = 0.019, 95% CI = [0.011 0.086],  $z = 2.564$ ,  $p = 0.010$ , *p*-corrected = 0.010), and a significant interaction




**Figure 1.** The graph depicts the interaction effect between interval ( $x$  axis) and the number of turns (traces, divided in  $-1$  s.d. = red, estimated mean = blue,  $+1$  s.d. = green) on interpersonal neural synchrony (measured in WTC;  $y$  axis). A higher turn-taking frequency is associated with higher neural synchrony between mother–infant at the beginning of the proto-conversation. Neural synchrony in later intervals is not associated with the number of turns. The shaded area depicts the 95% confidence interval of the predicted values.

between turns and intervals (estimate =  $-0.002$ , s.e. =  $0.001$ , 95% CI =  $[-0.003 -0.0005]$ ,  $z = -2.748$ ,  $p = 0.006$ ,  $p$ -corrected =  $0.009$ ). This significant set of effects shows that more frequent turns were related to higher neural synchrony, specifically in the medial prefrontal cortex. Neural synchrony between mother and infant also increased over time. But more specifically, more frequent turns were associated with higher neural synchrony in the medial prefrontal cortex in the first intervals (up to the second minute; figure 1). The relation between turns and neural synchrony is then attenuated over later intervals (third to fifth minute). In addition, we examined whether interpersonal neural synchrony in HbR was related to turn-taking but found no significant effects (see supplements for further details). Next, we explored whether response latencies or overlap frequency were associated with mother–infant neural synchrony. The relations were, however, not significant ( $p > 0.114$ ).

### (c) Turn-taking and infants' inter-hemispheric connectivity

Next, we tested infants' inter-hemispheric connectivity as a neural correlate to infant-initiated (infant-to-mother) turns and infants' (mother-to-infant) response latencies. The first model ( $n = 54$ ) output displayed that the frequency of infant-to-mother turns was significantly associated with infants' inter-hemispheric connectivity (estimate =  $0.078$ , s.e. =  $0.038$ ,  $z = 2.031$ ,  $p = 0.048$ ), as depicted in figure 2. We were able to replicate these results using infants' inter-hemispheric connectivity in HbR (detailed in supplements). Mother-to-infant response latencies were however not related to infants' interhemispheric connectivity ( $p = 0.701$ ). Moreover, infants' and mother's frequency of overlaps were not significantly associated with inter-hemispheric connectivity ( $p > 0.255$ ). Taken together, more infant-initiated turns in




**Figure 2.** Infant-to-mother turn-taking ( $x$  axis) is positively associated with infants' inter-hemispheric connectivity (assessed using WTC;  $y$  axis). The predicted values are plotted in black, and the shaded area depicts the 95% confidence interval.

the proto-conversation were associated with higher inter-hemispheric connectivity in infants' brains.

### (d) Infants' expressive language abilities and turn-taking

We tested the relation between infants' vocabulary at 24 months and dyadic as well as individual turn-taking patterns during the mother–infant interaction at 4–6 months ( $n = 28$ ). The first LM assessed the relation between infants' vocabulary and dyadic turn-taking patterns and revealed a significant correlation between the number of turns of the dyad and infant



**Figure 3.** (a) The graph depicts the relation between dyadic turn-taking frequency (x axis) at 4–6 months of age and infants' vocabulary at 24 months of age (y axis). (b) The graph depicts the relation between infant overlaps (x axis) at 4–6 months and infants' vocabulary at 24 months of age (y axis). Predicted values are plotted in black, and the shaded area depicts the 95% confidence interval.

vocabulary (estimate = 0.137, s.e. = 0.021,  $z = 6.509$ ,  $p < 0.001$ ). More frequent turn-taking was related to infants' larger vocabulary at 24 months (figure 3a). Next, we split the dyadic turns into uni-directional turns and ran an additional regression analysis. The result showed that the number of mother-to-infant turn-takings explained more variance in infants' vocabulary than infant-to-mother turn-taking (estimate = 0.164, s.e. = 0.030,  $z = 5.414$ ,  $p < 0.001$ ). The third LM assessed the relation between infants' vocabulary and mother-to-infant overlap frequency and found a significant negative relation (estimate =  $-0.066$ , s.e. = 0.021,  $z = -3.151$ ,  $p < 0.001$ ). Fewer mother-to-infant overlaps were associated with infants' larger vocabulary (figure 3b). Infants' vocabulary was not significantly related to infant-to-mother overlaps ( $p = 0.794$ ).

### (e) Infants' attachment and turn-taking

Next, we tested the relation between mother–infant turn-taking patterns at 4–6 months and infants' attachment security at 12 months. Attachment security was not related to mother–infant turn-taking patterns,  $p$ -corrected  $> 0.122$ . In an exploratory analysis, we dichotomized infants' attachment scores into securely and insecurely attached infants (split at  $r = 0.3$  as per [37]). The tested relation between secure versus insecure attachment and turn-taking patterns was not significant,  $p$ -corrected  $> 0.140$ .

## 4. Discussion

In the present study, we examined turn-taking patterns in mother–infant proto-conversations in relation to interpersonal neural synchrony and developmental outcome measures: infants' inter-hemispheric connectivity, expressive language abilities, and attachment. The more turn-taking mother and infant showed the more they displayed interpersonal neural synchrony during the proto-conversation. The relation was stronger in the initial phase of the interaction and decreased over time. Next, infants' number of turns was related to higher inter-hemispheric connectivity in infants' frontal brain regions. In addition, the number of turns in general, but especially infants' turns, was related to a larger vocabulary at 24 months of age. However, mother–infant turn-taking was not related to the infant's attachment security. The results

underscore the role of interpersonal neural synchrony in early turn-taking as well as brain maturation and language development as potential outcomes related to coordinated proto-conversation in early caregiver–infant interactions.

As predicted, we observed a positive relation between the frequency of turns and neural synchronization between mothers and their infants in a proto-conversation. This association was limited to the medial prefrontal cortex among our regions of interest. Medial prefrontal regions have been implicated in caregiver–infant synchrony, especially in face-to-face exchanges [49]. This is consistent with the role of the medial prefrontal cortex in processing communicative signals directed to the self and, more generally, in social cognition and mentalizing processes [39,50,51]. Synchronization of rhythmic brain activities in this region could be linked to high levels of mutual engagement, thus facilitating mutual prediction and fluid turn-taking between interaction partners. Also, in line with our predictions, neural synchrony in the mother–infant dyads generally increased over the course of the free play exchange. However, the positive association between frequency of turns and neural synchrony was unexpectedly limited to the early phases of the mother–infant interaction.

Our hypothesis that neural synchrony might be a shared underlying mechanism for turn-taking across earlier and later human development can therefore neither be fully confirmed nor dismissed. By contrast to the current results, neural synchrony increased over time in mothers and their preschool-aged children in association with more frequent turn-taking in a previous study [21]. There are several possible interpretations for the partial discrepancy between the current findings and these earlier results with older children. While neural synchrony might help mother–infant dyads getting attuned to each other early on during a proto-conversation, upholding a preverbal exchange with frequent turns might not rely on persistent neural synchrony in the same way as might be the case in later (more complex) verbal conversations. Alternatively, the oscillator model of turn-taking [13] might indeed be a universal mechanism for precisely timed turn-taking, but we were not able to capture this over the course of the proto-conversations here, e.g. due to our focus on slower haemodynamic brain responses instead of temporally better resolved electroencephalography. Future research could elucidate the functional link between neural synchrony and turn-taking in different

age groups further by experimentally manipulating turn-taking and measuring resulting effects on neural synchrony and vice versa, by manipulating neural synchrony through neurofeedback or transcranial stimulation methods [52].

Next, we examined the relationship between mother–infant turn-taking and infant maturity at the neural level, measured by inter-hemispheric connectivity. Consistent with our hypothesis, we found that more frequent infant-to-mother turn-taking, but not mother-to-infant response latencies, was positively related to brain maturation. These results support previous findings that infants' individual differences in inter-hemispheric connectivity can be identified and related to individual emotional and sensorimotor differences [27,28]. Interestingly, at 4–6 months, more mature infants might more likely engage in early proto-conversations and express themselves vocally, independent of how well coordinated their own vocalizations might be [30,53]. In sum, our results show that the frequency of proto-conversational patterns (i.e. turn-taking), independent of their quality and thus response latency, is connected to infant brain maturation. The direction of this link is, however, unclear as we assessed turn-taking and inter-hemispheric connectivity at the same time point.

Even though early social interactions are maintained to be a core component of children's language development [10], few studies have explicitly evidenced a relation between early turn-taking and infants' later language abilities. Here, we find that more frequent turn-taking between mothers and infants as young as 4–6 months of age, especially mother-to-infant turns, is related to their expressive vocabulary skills at 24 months. Our results extend recent findings showing that turn-taking in children between 2 and 24 months of age and their parents is related to their vocabulary growth [31] and emerging communicative capacity [29]. Not only do the number of overall turns matter, but the results again point to the integral and contingent response of the caregiver as a factor relating to infants' language skills. Importantly, frequent (mother-to-)infant overlaps during the proto-conversation were negatively related to infants' later vocabulary. We suspect that the negative relation between overlaps and infants' language development might reflect infants' drastic reduction of overlap frequency and duration during proto-conversations as they grow [1]. Even though call overlapping is discussed as an important communicative signal in several species [54], mostly, overlap avoidance is discussed in relation to optimized human communication (e.g. [7]). Vocal matching and vocal imitation in human mother–infant interactions occur less often [55] and seem to be more relevant for mother–infant musical interactions [56]. Overlap avoidance becomes even more important when infants start to speak their first words, implicating infants' inhibitory and predictive abilities to coordinate with others as relevant for infants' language development [10]. Consistent with previous results, response latencies were again not significantly related to infants' expressive vocabulary at 24 months. This finding indicates that infants accept a range of latencies, especially by their caregivers, when engaging in a proto-conversation [57]. Taken together, the results highlight the importance of emerging turn-taking in frequency but not latency in caregiver–infant proto-conversations when preparing preverbal infants for their first words. While we focussed exclusively on vocabulary size, future studies might consider including other potentially relevant variables, such as syntax development or pragmatics, to gain a more comprehensive view on the relation between turn-taking and language development.

Next to supporting the development of infants' language skills, conversational turn-taking is suggested to help build connections between people [11,22]. Previous findings indicate that specifically insecurely attached children showed longer response latencies during caregiver–child proto-conversations [32]. However, we did not detect a significant relation between early preverbal turn-taking patterns and infant attachment security. The lack of a significant association could be due to several reasons: firstly, we assessed very early forms of turn-taking and attachment at a much younger age, while the previous study assessed older infants and toddlers. The relation between turn-taking and attachment could thus develop as infants grow beyond the preverbal phase, with infants' attachment representations maturing [58]. Secondly, our sample might have been too homogeneous for infants' turn-taking abilities to differ significantly between groups of securely and insecurely attached children. Overall, our results indicate that future studies might need to include a larger sample of insecurely attached infants to precisely examine early turn-taking as a way to facilitate the connection between caregiver and child.

Our results highlight the importance of preverbal turn-taking between caregivers and infants in early development. Turn-taking arises together with interpersonal neural synchronization in the medial prefrontal cortex, a potential biomarker of mutual engagement and prediction. Moreover, turn-taking in proto-conversations is related to infants' brain maturation and language development. Our results also indicate that the range of turn-taking qualities in the present sample might comprise 'good enough' communication patterns to facilitate the bond between healthy and (mostly) securely attached infants and their caregivers [12]. Therefore, a future research avenue could include investigating samples that show altered interaction patterns, such as insecurely attached infants or anxious caregivers [7,59].

Taken together, our results show that mother–infant turn-taking, similar to later in development [21], relies on a high level of mutual engagement. More mature infants initiate more turns and more responsive mothers have infants with better language outcomes. In addition, bidirectional turn-taking frequency is related to higher levels of neural synchrony, pointing to the intricate link between both partners being mutually engaged with one another (to establish neural synchrony) and a fluid turn-taking exchange. Yet, the directionality of that link and thus mechanistic underpinnings to turn-taking remain to be tested.

**Ethics.** The local Ethics Committee approved the study (reference no. 00352).

**Data accessibility.** Processed fNIRS and vocalization data, MATLAB processing, and RStudio analysis code and files are made accessible on OSF: <https://osf.io/fx4ud/>. The conditions of our ethics approval do not permit public archiving of raw fNIRS and video data. Readers seeking access to the data should contact the lead author T.N. Access will be granted to named individuals in accordance with ethical procedures governing the reuse of sensitive data. Specifically, requestors must meet the following conditions to obtain the data: completion of a formal data sharing agreement.

**Authors' contributions.** T.N.: conceptualization, formal analysis, investigation, methodology, project administration, software, validation, visualization, writing—original draft, writing—review and editing; L.Z.: data curation, formal analysis, investigation, writing—original draft, writing—review and editing; S.H.: conceptualization, funding acquisition, methodology, resources, supervision, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

**Conflict of interest declaration.** We declare we have no competing interests.

**Funding.** This research was funded by the University of Vienna and by a doctoral stipend from the Studienstiftung des Deutschen Volkes granted to T.N.

**Acknowledgements.** We are grateful to Liesbeth Forsthuber, our student assistants Anja Lueger, Melanie Huber and Hannah Meller for their support in data acquisition, video and attachment coding. Additionally, we thank all families who participated in the study and the Department of Obstetrics and Gynaecology of the Vienna General Hospital for supporting our participant recruitment.

## References

- Gratier M, Devouche E, Guellai B, Infanti R, Yilmaz E, Parlato-Oliveira E. 2015 Early development of turn-taking in vocal interaction between mothers and infants. *Front. Psychol.* **6**, 1167. (doi:10.3389/fpsyg.2015.01167)
- Hilbrink EE, Gattis M, Levinson SC. 2015 Early developmental changes in the timing of turn-taking: a longitudinal study of mother–infant interaction. *Front. Psychol.* **6**, 1492. (doi:10.3389/fpsyg.2015.01492)
- Rohlfing KJ, Fritsch J, Wrede B, Jungmann T. 2006 How can multimodal cues from child-directed interaction reduce learning complexity in robots? *Adv. Robot.* **20**, 1183–1199. (doi:10.1163/156855306778522532)
- Trevarthen C, Aitken KJ. 2001 Infant intersubjectivity: research, theory, and clinical applications. *J. Child Psychol. Psychiatry* **42**, 3–48. (doi:10.1111/1469-7610.00701)
- Murray L, Trevarthen C. 1986 The infant's role in mother–infant communications. *J. Child Lang.* **13**, 15–29. (doi:10.1017/S0305000900000271)
- Cohn JF, Tronick EZ. 1987 Mother–infant face-to-face interaction: the sequence of dyadic states at 3, 6, and 9 months. *Dev. Psychol.* **23**, 68–77. (doi:10.1037/0012-1649.23.1.68)
- Nguyen V, Versyp O, Cox C, Fusaroli R. 2022 A systematic review and Bayesian meta-analysis of the development of turn taking in adult-child vocal interactions. *Child Dev.* **93**, 1181–1200. (doi:10.1111/cdev.13754)
- Stivers T *et al.* 2009 Universals and cultural variation in turn-taking in conversation. *Proc. Natl Acad. Sci. USA* **106**, 10 587–10 592. (doi:10.1073/pnas.0903616106)
- Sacks H, Schegloff EA, Jefferson G. 1974 A simplest systematics for the organization of turn taking for conversation. *Language* **50**, 696–735. (doi:10.2307/412243)
- Levinson SC. 2016 Turn-taking in human communication – origins and implications for language processing. *Trends Cogn. Sci.* **20**, 6–14. (doi:10.1016/j.tics.2015.10.010)
- Templeton EM, Chang LJ, Reynolds EA, Cone LeBeaumont MD, Wheatley T. 2022 Fast response times signal social connection in conversation. *Proc. Natl Acad. Sci. USA* **119**, e2116915119. (doi:10.1073/pnas.2116915119)
- Jaffe J, Beebe B, Feldstein S, Crown CL, Jasnow MD, Rochat P, Stern DN. 2001 Rhythms of dialogue in infancy: coordinated timing in development. *Monogr. Soc. Res. Child Dev.* **66**, i–149.
- Wilson M, Wilson TP. 2005 An oscillator model of the timing of turn-taking. *Psychon. Bull. Rev.* **12**, 957–968. (doi:10.3758/BF03206432)
- Hasson U, Ghazanfar AA, Galantucci B, Garrod S, Keysers C. 2012 Brain-to-brain coupling: a mechanism for creating and sharing a social world. *Trends Cogn. Sci.* **16**, 114–121. (doi:10.1016/j.tics.2011.12.007)
- Zion Golumbic EM *et al.* 2013 Mechanisms underlying selective neuronal tracking of attended speech at a 'cocktail party'. *Neuron* **77**, 980–991. (doi:10.1016/j.neuron.2012.12.037)
- Luo H, Poeppel D. 2007 Phase patterns of neuronal responses reliably discriminate speech in human auditory cortex. *Neuron* **54**, 1001–1010. (doi:10.1016/j.neuron.2007.06.004)
- Attaheri A *et al.* 2022 Delta- and theta-band cortical tracking and phase-amplitude coupling to sung speech by infants. *Neuroimage* **247**, 118698. (doi:10.1016/j.neuroimage.2021.118698)
- Menn KH, Michel C, Meyer L, Hoehl S, Männel C. 2022 Natural infant-directed speech facilitates neural tracking of prosody. *Neuroimage* **251**, 118991. (doi:10.1016/j.neuroimage.2022.118991)
- Jiang J, Dai B, Peng D, Zhu C, Liu L, Lu C. 2012 Neural synchronization during face-to-face communication. *J. Neurosci.* **32**, 16 064–16 069. (doi:10.1523/JNEUROSCI.2926-12.2012)
- Pérez A, Dumas G, Karadag M, Duñabeitia JA. 2019 Differential brain-to-brain entrainment while speaking and listening in native and foreign languages. *Cortex* **111**, 303–315. (doi:10.1016/j.cortex.2018.11.026)
- Nguyen T, Schleihau H, Kayhan E, Matthes D, Vrtička P, Höhl S. 2021 Neural synchrony during mother-child conversation. *Soc. Cogn. Affect. Neurosci.* **16**, 93–102. (doi:10.1093/scan/nsaa079)
- Jiang J, Zheng L, Lu C. 2021 A hierarchical model for interpersonal verbal communication. *Soc. Cogn. Affect. Neurosci.* **16**, 246–255. (doi:10.1093/scan/nsaa151)
- Kostorz K, Nguyen T, Pan Y, Steyrl D, Melinscak F, Hu Y, Sorger B, Hoehl S, Scharnowski F. 2022 Syncing brains: real-time fNIRS-neurofeedback of inter-brain synchrony. In *Human Brain Mapping Conf., Glasgow, UK, 19-23 June 2022*.
- Dikker S, Montgomery S, Tunca S. 2019 Using synchrony-based neurofeedback in search of human connectedness. In *Brain art* (ed. A Nijholt), pp. 161–206. Cham, Switzerland: Springer International Publishing.
- Saby JN, Marshall PJ. 2012 The utility of EEG band power analysis in the study of infancy and early childhood. *Dev. Neuropsychol.* **37**, 253–273. (doi:10.1080/87565641.2011.614663)
- Reindl V, Gerloff C, Scharke W, Konrad K. 2018 Brain-to-brain synchrony in parent-child dyads and the relationship with emotion regulation revealed by fNIRS-based hyperscanning. *Neuroimage* **178**, 493–502. (doi:10.1016/j.neuroimage.2018.05.060)
- Kelsey CM, Farris K, Grossmann T. 2021 Variability in infants' functional brain network connectivity is associated with differences in affect and behavior. *Front. Psychiatry* **12**, 685754. (doi:10.3389/fpsyg.2021.685754)
- Allievi AG, Arichi T, Tumor N, Kimpton J, Arulkumaran S, Counsell SJ, Edwards AD, Burdet E. 2016 Maturation of sensori-motor functional responses in the preterm brain. *Cereb. Cortex* **26**, 402–413. (doi:10.1093/cercor/bhv203)
- Zimmerman FJ, Gilkerson J, Richards JA, Christakis DA, Xu D, Gray S, Yapanel U. 2009 Teaching by listening: the importance of adult-child conversations to language development. *Pediatrics* **124**, 342–349. (doi:10.1542/peds.2008-2267)
- Ginsburg GP, Kilbourne BK. 1988 Emergence of vocal alternation in mother-infant interchanges. *J. Child Lang.* **15**, 221–235. (doi:10.1017/S0305000900012344)
- Donnelly S, Kidd E. 2021 The longitudinal relationship between conversational turn-taking and vocabulary growth in early language development. *Child Dev.* **92**, 609–625. (doi:10.1111/cdev.13511)
- Klann-Delius G, Hofmeister C. 1997 The development of communicative competence of securely and insecurely attached children in interactions with their mothers. *J. Psycholinguist. Res.* **26**, 69–88. (doi:10.1023/A:1025012221407)
- Ainsworth MS. 1979 Infant–mother attachment. *Am. Psychol.* **34**, 932.
- Nguyen T, Abney DH, Salamander D, Bertenthal BI, Hoehl S. 2021 Proximity and touch are associated with neural but not physiological synchrony in naturalistic mother-infant interactions. *Neuroimage* **244**, 118599. (doi:10.1016/j.neuroimage.2021.118599)
- Nguyen T, Hoehl S, Bertenthal BI, Abney DH. 2022 Coupling between prefrontal brain activity and respiratory sinus arrhythmia in infants and adults. *Dev. Cogn. Neurosci.* **58**, 101184. (doi:10.1016/j.dcn.2022.101184)
- Benjamini Y, Hochberg Y. 1995 Controlling the false discovery rate: a practical and powerful approach to



- multiple testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300. (doi:10.1111/j.2517-6161.1995.tb02031.x)
37. Waters E. 1995 Caregiving, cultural, and cognitive perspectives on secure-base behavior and working models: new growing points of attachment theory and research. *Monogr. Soc. Res. Child Dev.* **60**, 234–246.
38. Grimm H, Doil H. 2000 *ELFRA 2: Elternfragebogen für zweijährige kinder*. Göttingen, Germany: Hogrefe.
39. Kampe KKW, Frith CD, Frith U. 2003 'Hey John': signals conveying communicative intention toward the self activate brain regions associated with 'mentalizing', regardless of modality. *J. Neurosci.* **23**, 5258–5263. (doi:10.1523/JNEUROSCI.23-12-05258.2003)
40. Huppert TJ, Diamond SG, Franceschini MA, Boas DA. 2009 HomER: a review of time-series analysis methods for near-infrared spectroscopy of the brain. *Appl. Opt.* **48**, D280–D298. (doi:10.1364/ao.48.00d280)
41. Nguyen T, Hoehl S, Vrtička P. 2021 A guide to parent-child fNIRS hyperscanning data processing and analysis. *Sensors* **21**, 4075. (doi:10.3390/s21124075)
42. Grinsted A, Moore JC, Jevrejeva S. 2004 Application of the cross wavelet transform and wavelet coherence to geophysical time series. *Nonlinear Process. Geophys.* **11**, 561–566. (doi:10.5194/np-11-561-2004)
43. Sun FT, Miller LM, D'Esposito M. 2004 Measuring interregional functional connectivity using coherence and partial coherence analyses of fMRI data. *Neuroimage* **21**, 647–658. (doi:10.1016/j.neuroimage.2003.09.056)
44. Siddiqui MF, Lloyd-Fox S, Kaynezhad P, Tachtsidis I, Johnson MH, Elwell CE. 2017 Non-invasive measurement of a metabolic marker of infant brain function. *Sci. Rep.* **7**, 1330. (doi:10.1038/s41598-017-01394-z)
45. Oller DK, Buder EH, Ramsdell HL, Warlaumont AS, Chorna L, Bakeman R. 2013 Functional flexibility of infant vocalization and the emergence of language. *Proc. Natl Acad. Sci. USA* **110**, 6318–6323. (doi:10.1073/pnas.1300337110)
46. Fenson L, Dale P, Reznick JS, Thal D, Bates E, Hartung J, Pethick S, Reilly JS 1993 *The MacArthur communicative development inventories: user's guide and technical manual*. San Diego, CA: Singular Publishing Group.
47. Dobson AJ. 2002 *An introduction to generalized linear models*. London, UK: Chapman and Hall.
48. Judd CM, Westfall J, Kenny DA. 2017 Experiments with more than one random factor: designs, analytic models, and statistical power. *Annu. Rev. Psychol.* **68**, 601–625. (doi:10.1146/annurev-psych-122414-033702)
49. Piazza E, Hasenfratz L, Hasson U, Lew-Williams C. 2020 Infant and adult brains are coupled to the dynamics of natural communication. *Psychol. Sci.* **31**, 6–17. (doi:10.1177/0956797619878698)
50. Amodio DM, Frith CD. 2006 Meeting of minds: the medial frontal cortex and social cognition. *Nat. Rev. Neurosci.* **7**, 268–277. (doi:10.1038/nrn1884)
51. Schilbach L, Wohlschlaeger AM, Kraemer NC, Newen A, Shah NJ, Fink GR, Vogeley K. 2006 Being with virtual others: neural correlates of social interaction. *Neuropsychologia* **44**, 718–730. (doi:10.1016/j.neuropsychologia.2005.07.017)
52. Novembre G, Iannetti GD. 2021 Hyperscanning alone cannot prove causality. Multibrain stimulation can. *Trends Cogn. Sci.* **25**, 96–99. (doi:10.1016/j.tics.2020.11.003)
53. Feldman R, Eidelman AI. 2007 Maternal postpartum behavior and the emergence of infant-mother and infant-father synchrony in preterm and full-term infants: the role of neonatal vagal tone. *Dev. Psychobiol.* **49**, 290–302. (doi:10.1002/dev.20220)
54. Pika S, Wilkinson R, Kendrick KH, Vernes SC. 2018 Taking turns: bridging the gap between human and animal communication. *Proc. R. Soc. B* **285**, 20180598. (doi:10.1098/rspb.2018.0598)
55. Athari P, Dey R, Rvachew S. 2021 Vocal imitation between mothers and infants. *Infant Behav. Dev.* **63**, 101531. (doi:10.1016/j.infbeh.2021.101531)
56. Carvalho ME, Justo JM, Gratiier M, Ferreira Rodrigues H. 2021 Infants' overlapping vocalizations during maternal humming: contributions to the synchronization of preterm dyads. *Psychol. Music* **49**, 1654–1670. (doi:10.1177/0305735620968920)
57. Bigelow AE, Rochat P. 2006 Two-month-old infants' sensitivity to social contingency in mother–infant and stranger–infant interaction. *Infancy* **9**, 313–325. (doi:10.1207/s15327078in0903\_3)
58. Waters E, Merrick S, Treboux D, Crowell J, Albersheim L. 2000 Attachment security in infancy and early adulthood: a twenty-year longitudinal study. *Child Dev.* **71**, 684–689. (doi:10.1111/1467-8624.00176)
59. Smith CG, Jones EJH, Charman T, Clackson K, Mirza FU, Wass SV. 2021 Anxious parents show higher physiological synchrony with their infants. *Psychol. Med.* **14**, 3040–3050. (doi:10.1017/S0033291720005085)