

RESEARCH ARTICLE

Tree species composition along environmental and disturbance gradients in tropical sub-montane forests, Tanzania

Nandera Juma Lolila^{1*}, Deo D. Shirima^{2,3}, Ernest William Mauya¹

1 Department of Forest Engineering and Wood Sciences, College of Forestry, Wildlife and Tourism, Sokoine University of Agriculture, Morogoro, Tanzania, **2** Department of Ecosystems and Conservation, College of Forestry, Wildlife and Tourism, Sokoine University of Agriculture, Morogoro, Tanzania, **3** National Carbon Monitoring Centre, Sokoine University of Agriculture, Morogoro, Tanzania

* njlolila@gmail.com

Abstract

Understanding the environmental and disturbance determinants of tree species dominance and community composition in an ecosystem, is important for informing management and conservation decisions, through maintaining or improving the existing forest composition and structure. This study was carried out to quantify the relationship between forest tree composition structure and environmental and disturbance gradients, in a tropical sub-montane forest of Eastern Usambara. Vegetation, environmental, and anthropogenic disturbance data for 58 plots across Amani and Nilo nature forest reserves were obtained. Agglomerative hierarchical cluster analysis and canonical correspondence analysis (CCA) were used to identify plant communities and analyze the influence of environmental variables and anthropogenic disturbances on tree species and community composition respectively. Four communities were identified and CCA results showed that the variation was significantly related to elevation, pH, Annual mean temperature, temperature seasonality, phosphorus nutrients and pressures from adjacent villages and roads. Likewise, environmental factors (climate, soil and topography) explained the most variation (14.5%) of tree and community composition in relation to disturbance pressure (2.5%). The large and significant variation in tree species and community patterns explained by environmental factors suggests a need for site-specific assessment of environmental properties for biodiversity conservation plans. Similarly, the intensification of human activities and associated impacts on natural environment should be minimized to maintain forest species composition patterns and communities. The findings are useful in guiding in policy interventions that focus on minimizing human disturbances in the forests and could aid in preserving and restoring the functional organization and tree species composition of the sub-tropical montane forests.

OPEN ACCESS

Citation: Lolila NJ, Shirima DD, Mauya EW (2023) Tree species composition along environmental and disturbance gradients in tropical sub-montane forests, Tanzania. PLoS ONE 18(3): e0282528. <https://doi.org/10.1371/journal.pone.0282528>

Editor: Sehrish Sadia, University of Veterinary and Animal Sciences, PAKISTAN

Received: August 8, 2022

Accepted: February 17, 2023

Published: March 8, 2023

Copyright: © 2023 Lolila et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Tropical montane forests are characterized by complex ecological systems resulting from dramatic changes of physical and geographic properties [1] such as topography [2–5], including: slope [6], aspect [1,5], climate [7,8], soil [9] and their interactions [10–12] from the base to the

summit of mountains [13] creating a multitude of microhabitats [3]. Tropical montane forests are the most unique, with high diversity of vegetation [14,15], but they are also among the most threatened ecosystems by anthropogenic activities [5,16,17]. Forest disturbances, particularly selective removal of trees [18], have impact on the species composition and dominance on mountainous landscape [11,19], particularly on the lower elevations [20]. The dynamics of interaction among individual species and their environmental and disturbances influence species composition through coexistence or exclusion [8]. Hence, plant species that respond equally to the same factors, will usually coexist giving rise to distinct patterns of plant species communities across the landscape [21,22] while plant species with restricted habitat requirements are most susceptible to extinction in the face of anthropogenic pressures [23,24].

Species composition and structure of the tropical montane forests vary across a wide range of ecological gradients [1,25]. African mountainous areas have large human population densities encircling and depending on forest resources for their livelihoods [26,27]. This results to high human impacts on plant species composition including resource utilization, road construction, and residential development [20], increasing the rates of deforestation and forest degradation due to over exploitation of resources [28]. The intensity of these disturbances is likely to decrease as the distance of the roads and villages from the boundary of the forest increases [29]. Tanzania government acknowledges deforestation as a major threat to biodiversity, and has committed to the establishment and management of protected areas (PAs), significantly reducing the rate of forest degradation and ecosystem fragmentation [30]. However, the effectiveness of PAs management and their ability to withstand anthropogenic pressures varies, depending on the nature of the anthropogenic activities that pose a threat to the PAs [31].

Several studies have demonstrated that disturbances, physical and geographic factors [25,32], shape the structure of the forest [20,33,34] reported that, the relationship between tree species determinants and community composition is complicated because it involves a wide range of factors and considerations that provide a variety of clues, which partially explain the relationship, and whose findings cannot be generalized to other ecosystems. Despite being classified as protected areas (PAs) [35], the scenic beauty and unique biodiversity of the Amani and Nilo nature forest reserves, along with the high degree of forest accessibility [36,37], makes them vulnerable to mass tourism pressure [38,39]. On the other hand, there hasn't been a thorough analysis of how the combined environment and anthropogenic pressures are affecting the tree species composition patterns in both nature reserves. Meanwhile, being able to predict how tropical forests will react to environmental change requires an understanding of how quickly the structure and composition of these forests recover and which factors govern these processes [1,40]. Hence, understanding the influence of environmental and disturbance factors on plant species composition [41] in an ecosystem, is a prerequisite in developing management plans [42], important for prioritizing conservation activities at local, regional and global scales [43]. Thus, this study analyzed forest tree species composition structure along varying environmental and disturbance gradients in Tropical Sub-montane forests, exploring (i) How does environmental (climate, topography and soil) and potential disturbance pressures explain the tree species and community composition in tropical sub-montane forests? (ii) How influential are environmental factors relative to disturbance pressures in explaining observed species composition?

Materials and methods

Study area

This study was conducted in Amani and Nilo nature forest reserve (NFR), located in Muheza, Mkinga and Korogwe Districts in Tanga Region of Tanzania. Amani nature Forest reserve




Fig 1. Map of the study area showing Eastern Arc mountains, Amani and Nilo nature forest reserve sites. The Eastern Arc layer source is from [35].

<https://doi.org/10.1371/journal.pone.0282528.g001>

(NFR) is located between $5^{\circ}04'30''$ - $5^{\circ}14'10''$ S and $38^{\circ}30'34''$ - $38^{\circ}40'06''$ E with an area of about 8360 ha. The elevation of ANFR is ranging from 190m-1130 m above the sea level [36,39]. NNFR is located between $4^{\circ}50'$ - $4^{\circ}59'S$ and $38^{\circ}37'$ - $38^{\circ}41'06''$ E, covering about 6025 ha (Fig 1). The location of NNFR is at an elevation range of 400 to 1506 m above the sea level. Both Amani NFR and Nilo NFR are confined to the East Usambara Mountains forests (EUMFs) located in eastern Tanzania and they are part of the Eastern Arc Mountains (EAM), which is a chain of mountains that stretches down the coast of East Africa from southern Kenya to southern Tanzania [36].

The rainfall distribution in EUMFs is bi-modal, peaking between March and May and between October and December [39]. Rainfall is greatest at higher altitudes and in the south-east of the mountains, increasing from 1,200 mm annually in the foothills to over 2,300 mm at higher altitudes [36]. The dry seasons are from June to August and January to March. The vegetation of these forests' ranges includes lowland forest at 300 m on Eastern side, sub-montane forests and montane forests. Tree species composition varies considerably, but species such as *Khaya anthotheca*, *Milicia excelsa* are found in the lowlands and others such as *Myrianthus holstii*, *Albizia gummifera*, *Allanblackia stuhlmannii* and *Newtonia buchananii* are dominant at high altitudes [37,44].

Sampling design

We utilized the existing systematic 450 by 900 m grids of east-west and north-south transect method in 1999–2000 by Frontier Tanzania in each of the two nature forest reserves [44]. The grids were intended for surveying the flora and fauna where 182 rectangular plots were established in Amani NFR and 122 rectangular plots of 20 x 50 were established in Nilo NFR.

In this study, we measured a sub-sample of 30 circular plots of 18 m radius within the Frontier grids in Amani NFR in 2017/18 as described in [45]. Sixteen (16) plots were on the exact position as in the frontier plots, with difference in shape and sizes only, 14 plots were established along the grid lines of Frontier plots and located exactly midway between two existing Frontier plots. Thus, the inter plot distance between our plots was 225 rather than 450m. The 30 plots covered an elevation range from 200 to 1000 m above sea level so that both the lowland forests (<800 m above sea level) and the sub-mountain forests (>800 m above sea level) were covered [37].

In Nilo NFR 28, circular plots of 18 m radius were established exactly on the same location with the existing Frontier plots. Same grids were used only with difference in plot size and shapes, thus the inter plot distance was 450m. We decided to use circular plots, because they are easier to establish in the field, and because they have one dimension (i.e., radius) that defines the plot boundary. Therefore, in total we had 58 field plots measured in the two nature forest reserves.

Data collection

Field data. Existing plot coordinate's location established by [45] were used to navigate to the center of field plots in Amani NFR using handheld GPS. Similarly, in Nilo NFR the plot coordinate's locations as reported by [46] for the Frontier were used to locate the center position of the plots. Field data were collected in Dec 2017 and May 2018 in Amani NFR and Nilo NFR respectively. As stated above, circular plots with a radius of 18 m were established on each of the two-nature forest reserves. On each plot, all trees with diameter at breast height (*dbh*) greater or equal to 5cm were measured and recorded for *dbh*, scientific and local name. Plant materials were collected for further identification or confirmation at the Tanzania National Herbarium. A caliper was used to measure *dbh*, however for larger and trunked trees diameter tape was used.

Forest tree species composition and dominance. To determine the species composition of the forests, we summarized the species datasets into list of all species recorded and their abundance values.

The dominance of tree species was determined from the estimation of their species importance value index (SIVI). The Importance Values Index (IVI), indicates the ecological importance of a tree species and hence the dominance [20]. The Importance Value Index (IVI) for each species was computed as the sum of relative density, relative dominance and relative frequency of the species in each plot following [47].

Environmental data. To estimate the influence of environmental factors on tree species, we used information on topography, climate and soil factors.

Topographic factors including; elevation, slope and aspect were extracted from the raster layer derived from the SRTM 30 m-based DEM-USGS Earth Explorer (<https://earthexplorer.usgs.gov/>). A total of 58 points consisting of plot locations were imported into QGIS 3.22, where a spatial analyst tool of the QGIS was employed to obtain values of aspect, elevation and slope as well as the Topographic Position Index (TPI). Climatic data was downloaded from WorldClim site (<https://www.worldclim.org>) with a 30 arc seconds resolution and soil data was extracted from the Re-gridded Harmonized World Soil Database: ISRIC Data Hub

Table 1. The environmental and disturbance factors (Mean \pm SD) within the tropical sub -montane forests of East Usambara, Tanzania. The bolded factors significantly explained species composition.

Factors	Variables	Codes	Mean	SD	Min	Max
Soil	pH	pH	5.398	0.17	5.00	6.00
	Organic carbon	OC	15.88	5.24	6.00	30.00
	Electrical conductivity	EC	0.67	0.30	0.13	1.49
	Cation exchange capacity	CEC	12.17	2.82	8.00	20.00
	Available water holding capacity	AWHC	26.83	1.39	24.00	30.00
	Bulk density	BD	1,364.48	54.55	1,260.00	1,490.00
	Potassium	K	153.83	45.12	72.00	265.00
	Total available Nitrogen	N	2,023.07	211.57	1,355.00	2,682.00
	Sodium	Na	217.71	42.79	179.00	381.00
	Phosphorous	P	849.74	204.95	428.00	1,299.00
Climatic	Mean Annual Temperature	MAT	21.82	1.42	20.14	24.48
	Temperature seasonality	TS	164.31	12.07	140.53	186.86
	Mean Annual precipitation	MAP	1,337.71	147.32	1,073.00	1,589.00
	Precipitation seasonality	PS	57.57	2.33	53.57	62.31
Topographic	Aspect	As	164.32	88.08	3.57	355.10
	Elevation	El	704.36	301.46	222.65	1,120.00
	Slope	Sl	29.33	16.52	3.58	70.57
	Topographical position Index	TPI	0.47	2.81	-4.74	9.44
Disturbance	Distance to the road	Rd	6,378.83	4,355.10	19.08	12,385.68
	Distance to the villages	Vl	1,561.55	634.05	461.22	3,384.29
	Distance to conservation offices	HQDist	4,538.13	3,014.47	117.93	9,871.30

<https://doi.org/10.1371/journal.pone.0282528.t001>

(International Soil Reference and Information Centre) <https://data.isric.org/geonetwork/srv/eng/catalog.search#/home>. For each plot, values for all spatially interpolated climate and soil predictors were extracted using “raster package” in R software sampled using the coordinates of the plot center and averaged across plots per site (Table 1).

Disturbance data. The degree of forest accessibility was included as proxy for anthropogenic disturbances [29,48]. Likewise, the presence of a conservation office building in the NFR was regarded as a factor for the decrease in intensity of human disturbances.

To quantify the influence of disturbance on tree species composition, we used distance from each field sampling plot to the nearest primary, secondary, or tertiary road, as well as the distance to the nearest city, village, or town and the distance from the plot to the conservators’ office. We obtained spatial data for roads and villages as shapefiles using road and settlement vector data from OpenStreetMap (<http://download.geofabrik.de/africa/tanzania.html>). The positions of the conservators’ offices were digitized from high resolution base-maps of the areas. The distances were obtained by calculating the Cartesian distance (in kilometers), using the gDistance function within the rgeos package of R statistical software. The function returns the minimum Cartesian distance between the two points (plots to the road or villages).

Data analysis

Dominant tree species composition. We used detrended correspondence analysis (DCA) to detect the magnitude of the ecological gradient in our species composition matrix and hence the constrained ordination to be used [49,50]. The DCA results showed an axis length greater than 4.0 in both study sites, suggesting that the data is heterogeneous. We therefore used Canonical Correspondence analysis (CCA) [51] to determine the influence of environmental and disturbance on dominant tree species and community types.

Prior to CCA, we checked for collinearity among variables on the basis of Pearson correlation (r) and Variance Inflation Factor (VIF). Predictor variables with a VIF value greater than ten (10) [52] and $r > 0.7$ [20] were considered to be high collinear [51] and were trimmed out from the list of predictors. Species abundance data were log-transformed to meet assumptions of multivariate normality [50]. Species that occurred in less than five plots were considered to have no major ecological significance (rare species) and were removed from the matrix [53]. Stepwise automatic forward selection was used to identify significant explanatory variables to be constrained in multivariate analysis [50]. The selected explanatory variables were then constrained against the tree species and community composition using CCA. The effect of the obtained explanatory variables on tree species composition in CCA model was determined using the Monte Carlo test with 999 unrestricted permutations [49] at 95% confidence interval.

Community composition. Similar communities were identified using cluster analysis. First, we computed the Bray–Curtis’s distance matrix using the abundance data and then, performed hierarchical Ward’s minimum variance clustering on the Bray–Curtis’s distance matrix. The decision on the number of clusters was based on Silhouette validation technique using ‘*Nbclust package*’ [54]. Indicator species analysis was performed to identify significant dominant species of the communities, using package *labdsv*. Then, community types were named after the two most dominant species [43] based on high synoptic cover abundance values (mean frequency multiplied by mean cover-abundance), following [49] (S1 Appendix).

Additionally, the community types obtained was subjected to ANOVA to describe environmental and disturbance differences among individual clusters. For the explanatory variables which were significant, pairwise differences tested among clusters; using Tukey HSD comparison was done.

Tree species composition variance partitioning. Explanatory variables that were significant in explaining variability in tree species composition were retained and considered for subsequent variance partitioning analysis. The significant explanatory variables were categorized into two main groups i.e., environmental and disturbance groups, then sub-grouped into four sets of climatic, topographic, edaphic and disturbance. We performed variance partitioning by partial CCA to determine independent and joint contributions of each group predictor variables to explain variation in species composition. The proportion of variation explained was given as the adjusted R^2 of the explanatory variable in the CCA, as unbiased estimates of the variation [41]. DCA and CCA were performed by the ‘vegan’ package and variation partitioning with CCA by the ‘varpart’ package in the R software (v. 4.1.2) [21].

Results

Trees species and community structure

Tree species composition structure. A total of 2910 individuals were recorded belonging to 174 species, 128 genera, and 46 families. The most dominant families with a high number of species were: Leguminosae (28), Rubiaceae (19), Moraceae (13), Malvaceae (11) and Sapotaceae (10). The most common genera with the highest number of trees were *Albizia* (5), *Ficus* (5) and *Celtis* (5). Also, the most dominant species, in terms of IVI were *Leptonychia usambarensis* K. Schum(151.8), *Cephalosphaera usambarensis* (Warb.) Warb. (114.6), *Antiaris toxicaria* Lesch (95.2) and *Maesopsis eminii* Engl. (85.7) (S1 Appendix).

Our results show that, 28% of variation of dominant tree species composition was explained by the environmental and disturbance factors. The overall CCA model ($\chi^2 = 0.58$, $P = 0.001$), as well as the first two ordination axes were statistically significant ($P = 0.001$), showing that the observed patterns differed from a random relationship. The first two ordinations

accounted for 15.9% of the total variation, having eigenvalues of 0.58 and 0.21 respectively. The first ordination axis was significantly correlated with elevation (El), mean Annual Precipitation (MAP) gradients, pH and Temperature seasonality (TS), while the second canonical axis showed strong significant correlations ($P < 0.05$) with distance to the nearest road (Rd) and villages (VI) (Fig 2).

The biplots indicated that, on axis 1, species like *Isoberlinia scheffleri* (Harms) Greenway, *Alsodeiopsis schumannii* (Engl.) Engl., *Parinari excelsa* Sabine have positive scores while species like *Pouteria alnifolia* (Baker) Roberty, *Celtis mildbraedii* Engl. and *Lecaniodiscus fraxinifolia* Baker have high negative scores and are highly significantly influenced by Elevation and Precipitation (MAP). On axis 2, species like; *Trichilia dregeana* Sond., *Alangium chinense* (Lour.) Harms and *Ficus sur* Forssk. had positive scores and are positively influenced by pressure of nearest roads (Fig 2).

Community composition. We identified five distinct significant tree species communities ($p < 0.001$) in the study area (Fig 3). Most of the species were shared across communities; however each community is characterized by a distinct set of dominant species (S1 Appendix).

The community composition structure in the two forests were influenced by elevation (El), Phosphorous (P), pH, mean annual precipitation (MAP), temperature seasonality (TS), nearest distance to the road (Rd) and villages (VI). The *Isoberlinia scheffleri*-*Sorindeia madagascariensis* forest community is significantly influenced by pressures from nearest roads ($P = 0.001$), *Tabernaemontana ventricose*-*Leptonychia usambarensis* and *Ficus sur*-*Myrianthus holstii* by pH ($P < 0.05$), while *Artocarpus heterophyllus*—*Lecaniodiscus fraxinifolia* forest community is influenced by mean annual precipitation, elevation and temperature seasonality ($P < 0.05$) (Fig 4).

Tree communities' compositions are influenced by elevation (El), Phosphorous (P), pH, mean annual precipitation (MAP), temperature seasonality (TS), disturbance pressure from nearest road (Rd) and villages (VI). The tree communities occupy significant different elevation gradients ($P < 0.05$) (Fig 5A). Likewise, there is significance difference of the mean annual precipitation received by the communities with an exception of *Artocarpus heterophyllus*—*Lecaniodiscus fraxinifolia* (Arhe.Lefr) and *Tabernaemontana ventricosa*-*Leptonychia usambarensis* (Tave.Leus) communities (Fig 5B). There is no significant difference in the influence of disturbance pressures from the nearest villages (VI) to tree communities ($P > 0.05$).

Tree species composition variance partitioning

Results showed that environmental factors (14.5%) explained variability in tree species composition five times more than the potential disturbance factors (2.5%). Overall, the climate factors accounted for 6.95% of the variation in tree species composition (Table 2). The variation in species composition is also a result of interaction of multitude of these factors. Example; the interaction between climate and topography (6.3%) and climate, topography and soil (6.5%) explained high variation in species composition (Fig 6).

Discussion

Tree species community structure

Divergent distribution patterns of species and community composition of mountain ecosystems are influenced by dramatic changes in environmental conditions over short altitudinal distance [33,49,55,56]. Nevertheless, human exploitation may be imposing constraints that result in environmental modification [57] causing a shift in species range [58] and thus influencing species composition [59]. Our study found that tree species and community patterns are determined by climate, topography, soil and disturbance pressures that exhibit heterogeneity over the sub-montane forest of East Usambara.




Fig 2. Canonical correspondence analysis diagram showing the ordination of dominant species represented and their correlation with environmental and disturbance variables in the sub-tropical montane forest plots. Canonical correspondence analysis biplot of Amani and Nilo FR species distribution (blue dots) based on tree species abundance. The gray axes vectors indicate the significant explanatory variables in their abbreviated form: El = Elevation, TS = Temperature seasonality, MAP = Mean Annual precipitation, P = Soil phosphorus, Rd = Distance to the road and VI = Distance to the villages. We also used abbreviated species scientific names of each tree by using first two letters each from genus and species; *Al.ch* = *Alangium chinense* (Lour.) Harms, *Al.gu* = *Albizia gummifera* (J.F.Gmel.) C.A.Sm., *Al.me* = *Allophylus melliodorus* Gilg ex Radlk., *Al.st* = *Allanblackia stuhlmannii* (Engl.) Engl., *Al.sc* = *Alsodeiopsis schumannii* (Engl.) Engl., *An.ku* = *Annickia kummeriae* (Engl. & Diels) Setten & Maas, *An.gr* = *Anthocleista grandiflora* Gilg, *An.me* = *Antidesma membranaceum* Müll.Arg, *An.to* = *Antiaris toxicaria* Lesch., *Ar.he* = *Artocarpus heterophyllus* Lam, *Bl.un* = *Blighia unijugata* Baker, *Ce.go* = *Celtis gomphophylla* Baker, *Ce.mi* = *Celtis mildbraedii* Engl., *Ce.us* = *Cephalosphaera usambarensis* (Warb.) Warb., *Ch.pe* = *Chrysophyllum perpulchrum* Mildbr. ex Hutch. & Dalziel, *Dr.ge* = *Drypetes gerrardii* Hutch., *En.us* = *Englerodendron usambarense* Harms, *Er.su* = *Erythrophleum suaveolens* (Guill. & Perr.) Brenan, *Fi.ex* = *Ficus exasperata* Vahl, *Fi.su* = *Ficus sur* Forssk., *Fu.af* = *Funtumia africana* (Benth.) Stapf, *Gr.su* = *Greenwayodendron suaveolens* (Engl. & Diels) Verdc., *Is.sc* = *Isobertinia scheffleri* (Harms) Greenway, *Le.fr* = *Lecaniodiscus fraxinifolia* Baker, *Le.us* = *Leptonychia usambarensis* K. Schum. *Ma.ca* = *Macaranga capensis* (Baill.) Sim, *Ma.*

em = *Maesopsis eminii* Engl., *Ma.lu* = *Markhamia lutea* (Benth.) K.Schum., *Me.in* = *Mesogyne insignis* Engl., *My.ho* = *Myrianthus holstii* Engl., *Ne.bu* = *Newtonia buchananii* (Baker) G.C.C.Gilbert & Boutiqu, *Pa.ex* = *Parinari excelsa* Sabine, *Po.al* = *Pouteria alnifolia* (Baker) Roberty, *Qu.un* = *Quassia undulata* (Guill. & Perr.) D.Dietr. *Ri.he* = *Ricinodendron heudelotii* (Baill.) Heckel, *Ro.ma* = *Rothmannia manganjae* (Hiern) Keay, *Sh.el* = *Shirakiopsis elliptica* (Hochst.) Esser, *So.ma* = *Sorindeia madagascariensis* Thouars ex DC., *St.sc* = *Strombosia scheffleri* Engl., *Sy.ce* = *Synsepalum cerasiferum* (Welw.) T.D.Penn., *Sy.ms* = *Synsepalum msolo* (Engl.) T.D.Penn., *Ta.pa* = *Tabernaemontana pachysiphon* Stapf, *Ta.ve* = *Tabernaemontana ventricosa* Hochst. ex A.DC., *Ta.ni* = *Tarennia nigrescens* (Hook.f.) Hiern, *Tr.dr* = *Trichilia dregeana* Sond., *Tr.ma* = *Trilepisium madagascariense* DC., *Tr.my* = *Tricalysia myrtifolia* S.Moore, *Xy.mo* = *Xymalos monospora* (Harv.) Baill.

<https://doi.org/10.1371/journal.pone.0282528.g002>

Our results showed that elevation significantly influences differences in species communities, similar to mean annual precipitation and temperature seasonality as shown in the CCA (Fig 5), suggesting correlations among topographic, elevation and precipitation [60] perhaps due to adiabatic decrease of temperature with elevation [61] and elevation-dependent wetting (EDWE) [62] phenomenon. Nevertheless, elevation is linked to variation in humidity [59] and soil moisture [9], that may indirectly affect tree nutrient availability [43] and other factors, which affecting tree species establishment, and their community composition [6,63]. Several studies have shown that elevation-temperature relationship significantly influenced the physiological attributes of plant species [59].

Our study shows similar biogeographical patterns of communities where, *Artocarpus heterophyllus*-*Lecaniodiscus franxinifolia* and *Isoberlinia schefflera*-*Sorindeia madagascariensis* communities, occupying lower to mid- elevation have higher species richness in comparison to other communities occupying high elevations. Studies suggest that, middle to low elevation zones have relatively mild climate and favorable nutritional conditions [64], meaning which means high chance for any species' growth and survival. Many of the species in the forests are limited by extremities in temperatures [24,59] as well as precipitation [65]. Seasonal variation in temperature across most tropical forests, maybe minor [66], but recent studies suggest that small changes in temperature are likely to affect species distribution patterns according to their species physiological drought tolerance [65,67,68]. Climatic conditions can thus exert selective pressure on composition species [8], given that some tree species may not prefer areas of high elevation because of low temperatures [55,69]. Thus, elevation can be used as a proxy for understanding tree species adaptation to climate change [56].

Furthermore, we found that species and community composition are significantly explained by the variation of pH and phosphorous (P) nutrients (Figs 2 and 4). Our study sites occupy areas of acidic to slightly acidic (i.e. pH between 5 and 6). Similar to [70], our results suggests that soil pH is a substantial, driver of species composition and distribution in East Usambara. Others studies, also show that soil pH has the ability to influence tree species nutrient uptake and their ecological behavior at different localities. [70–72]. Moreover, we found that soil phosphorus significantly ($P < 0.05$) influences the variability of the tree species and communities. Regardless of phosphorus being important for regulating plant primary productivity, and other biological processes such root allocation and growth [73–75], it is limited to plants due to its low solubility and the sorption processes in soils [72]. Nevertheless, soil P availability can be affected by direct and indirect climatic processes through sorption and desorption [76]. The relationship between soil properties and climate can thus never be inseparable [9,22] as well as topography, as there's evidently reports of elevation-climate dependency [43,59]. This suggests that any huge changes to the ecosystem may have knock- on effects on the whole forest system.

We found that disturbance had a significant influence on species composition in sub-tropical forest of East Usambara. *Isoberlinia schefflera*-*Sorindeia madagascariensis* communities forest community was influenced by disturbance factors and is the richest (59 species), while *Tabernaemontana ventricose*-*Leptonychia usambarensis* forest community is less affected by

Agglomerative Hierarchical classification(Wards' Linkage)




Fig 3. Dendrogram of hierarchical clustering using similarity ratio showing four community types in different sites (plot 1–59) of the study area (Ward’s method, Bray-Curtis distance) across Amani and Nilo nature forest reserve in sub-tropical East Usambara montane forest of Eastern Arc. The communities are named by using number of communities denoted as (Cn) and they are named after by two dominant species: C1 = *Tabernaemontana ventricosa*-*Leptonychia usambarensis*, C2 = *Ficus sur*- *Myrianthus holstii*, C3 = *Artocarpus heterophyllus*-*Lecaniodiscus fraxinifolia* and C4 = *Isobertlinia schefflera*- *Sorindeia madagascariensis* communities (S1 Appendix).

<https://doi.org/10.1371/journal.pone.0282528.g003>

disturbances and had the least tree species richness (24 species). The patterns of species richness, corresponds to the intermediate disturbance hypothesis, which suggests that local species richness is maximized when ecological disturbance is neither too rare nor too frequent and not too intense [77]. We also found that a dominant species like *Maesopsis eminii* Engl, a light




Fig 4. Canonical correspondence analysis diagram displaying the ordination of community types represented and their correlation with environmental and disturbance variables, indicating abundance relationships in subtropical East Usambara montane forest of Eastern Arc mountains. Communities: Tave.Leus = *Tabernaemontana ventricosa*-*Leptonychia usambarensis*, Arhe.Lefr = *Artocarpus heterophyllus*-*Lecaniodiscus fraxinifolia*, Fisu.Myho = *Ficus sur*-*Myrianthus holstii* and *Isoberlinia schefflera*-*Sorindeia madagascariensis* communities (S1 Appendix).

<https://doi.org/10.1371/journal.pone.0282528.g004>

-demanding pioneer tree species, was influenced by disturbance i.e., adjacent to villages (Fig 2). It is evidently supported that disturbance pressures to the forests may exert effects on the natural environment, thus changing the site condition and species habitat, enabling conditions for new species to emerge and/or diminish the existing species [33]. Light demanding species like *M. eminii* Engl have mostly been in high proportion in disturbances gaps, due to available maximum light necessary for triggering their establishment and growth [78], but responses to disturbance pressures vary among tree species, depending on their ability to colonize and to compete [79].

Tree species composition variance partitioning

The results of partial CCA (pCCA) revealed that species composition was defined best by climate and disturbance factors, with most of the variation being explained by the climate




Fig 5. Boxplot showing difference among individual communities with respect to environmental and disturbance factors. The horizontal line crossing the box in the center represents the median; the box represents the 25th and the 75th percentiles. The vertical line outside the box represents the minimum and maximum values. Difference in letters (a, b, c, d) signifies that the communities are different in the specified factor. Communities: 1 = *Tabernaemontana ventricosa*-*Leptonychia usambarensis*, 2 = *Ficus sur-* *Myrianthus holstii*, 3 = *Artocarpus heterophyllus*-*Lecaniodiscus fraxinifolia*, 4 = *Isoberlinia schefflera*- *Sorindeia madagascariensis*.

<https://doi.org/10.1371/journal.pone.0282528.g005>

variable similar to [79]. Our results support the argument that mountain tree communities are sensitive to climate and, thus, able to reveal its effects sooner than others [56]. Variation of species is also explained by interaction of factors, including: Climate, topography and soil (6.5%), as well as climate and topography (6.3%) (Fig 6). This confirms that species composition is a

Table 2. CCA variation partitioning (%) on overall plant species composition of Tropical sub- montane forests of East Usambara, Northern Tanzania.

Effect and main variable	Covariables	Variation explained (%)	F-value	P-value
Total effect (Environmental and disturbance factors)		28.90	8.18	0.001
Partial effect				
Environmental	Anthropogenic pressure	14.50	1.93	0.001
Disturbance	Environmental	2.50	2.75	0.001
Soil	Climate, Topography and Anthropogenic pressure	4.80	1.63	0.009
Topography	Soil, Climate and Anthropogenic pressures	2.30	1.59	0.030
Climate	Soil, Topography and Anthropogenic pressure	6.95	2.01	0.001
Anthropogenic pressure	Soil, Topography and Climate	5.80	1.80	0.001

<https://doi.org/10.1371/journal.pone.0282528.t002>




Fig 6. The Venn diagram showing variation partitioning results (partial CCA model) and the contribution of the four studied environmental and disturbance variable groups (i.e., climatic, soil, topography, and disturbance) that drive the plant species distribution. The proportion of variance in tree species composition explained by environmental and disturbance factors. The values represent adjusted R^2 (%) of independent and shared effects by 4 the factors.

<https://doi.org/10.1371/journal.pone.0282528.g006>

result of combination of multiple factors [20]. Although the influences among the environmental and disturbance variables covaried, neither the effect of one variable was entirely nested within the other (% variation of each individual variable >1). This indicates that the individual contribution of each factor has important and independent effects [21] on community structure in a wide variety of tropical forest communities [74].

Next to climate (mean annual precipitation and temperature seasonality), disturbance explained the most variability of species composition. We expect that the protected reserves are supposed to be free from outside human intrusions, but the protective efficiency of the protected forest is still in question, as anthropogenic activities continue to pose threats to the forest reserves [30,31]. Several studies [19,33,79] have recognized the importance of assessing human activities in shaping the floristic composition of the forests. Our study used proxy values of nearest distance to the roads and villages, which indicated that human-induced threats such as selective logging, [26,38], had significant influence on the variability in species and community composition. Chronic disturbance generated by anthropogenic activities normally gives rise to alterations in habitat conditions through the modification of physical structure of the forest [18].

Conclusion

Understanding the influence of environmental and disturbance factors on tree species dominance and community composition in an ecosystem, is important to inform conservation of existing forest structures. Our findings suggest that there is a strong relationship among forest structure, environmental and disturbances factors. The large and significant variation in tree species and community patterns explained by environmental factors suggests a need for site-specific assessment of environmental properties for biodiversity conservation plans. We also showed that, human induced disturbances from easy road access and villages near to forests may influence tree species composition. The effects of site condition modification as a result of human induced disturbance vary with location and existence of flora composition depending on the ability of tree species to adapt either at small scale or massive changes. Human induced disturbances should be discouraged in order to maintain species composition and encourage continuous monitoring and sustainable use of forest community structure. Likewise, understanding tree species composition and community structure patterns and their underlying environmental factors is worthwhile when considering conservation of montane sub-tropical forests. Thus, our findings may be useful for formulation of forest restoration policy interventions to sustain the tree species composition and the functioning in the subtropical montane forests of Eastern Usambara Mountain and similar ecosystems elsewhere.

Supporting information

S1 Appendix. List of dominant species in Sub-tropical forest of Eastern Arc montane forests, Tanzania. The bolded species are indicator species with highest indicator values which were used for naming the communities in Sub-tropical East Usambara montane forest of Eastern Arc mountains, Tanzania. IVI = Importance value index and CCA1 and CCA2 = Canonical correspondence analysis showing values for species along axis 1 and 2. (PDF)

Acknowledgments

We acknowledge the project titled “Meta-Analysis of Forest Carbon Information Using Existing Data and Remote Sensing Techniques”, under the Eastern Arc Mountain Conservation

programme, for providing field inventory data. Sincere acknowledgments are also extended to Prof. Tuyeni Heita Mwampamba for her constructive ideas and suggestions which helped to improve this study.

Author Contributions

Conceptualization: Nandera Juma Lolila, Deo D. Shirima, Ernest William Mauya.

Data curation: Ernest William Mauya.

Formal analysis: Nandera Juma Lolila.

Methodology: Nandera Juma Lolila, Deo D. Shirima.

Software: Nandera Juma Lolila.

Supervision: Deo D. Shirima, Ernest William Mauya.

Validation: Nandera Juma Lolila, Deo D. Shirima, Ernest William Mauya.

Visualization: Nandera Juma Lolila.

Writing – original draft: Nandera Juma Lolila.

Writing – review & editing: Nandera Juma Lolila, Deo D. Shirima, Ernest William Mauya.

References

1. Jucker T, Bongalov B, Burslem DFRP, Nilus R, Dalponte M, Lewis SL, et al. Topography shapes the structure, composition and function of tropical forest landscapes. *Ecol Lett*. 2018; 21: 989–1000. <https://doi.org/10.1111/ele.12964> PMID: 29659115
2. Wang S, Qi G, Knapp BO. Topography affects tree species distribution and biomass variation in awarm temperate, secondary forest. *Forests*. 2019; 10: 1–16. <https://doi.org/10.3390/f10100895>
3. Muscarella R, Kolyaie S, Morton DC, Zimmerman JK, Uriarte M. Effects of topography on tropical forest structure depend on climate context. *J Ecol*. 2020; 108: 145–159. <https://doi.org/10.1111/1365-2745.13261>
4. Tiwari OP, Sharma CM, Rana YS. Influence of altitude and slope-aspect on diversity, regeneration and structure of some moist temperate forests of Garhwal Himalaya. *Trop Ecol*. 2020; 61: 278–289. <https://doi.org/10.1007/s42965-020-00088-4>
5. Báez S, Fadrique B, Feeley K, Homeier J. Changes in tree functional composition across topographic gradients and through time in a tropical montane forest. *PLoS One*. 2022; 17: 1–20. <https://doi.org/10.1371/journal.pone.0263508> PMID: 35442987
6. Sharma CM, Baduni NP, Gairola S, Ghildiyal SK, Suyal S. Effects of slope aspects on forest compositions, community structures and soil properties in natural temperate forests of Garhwal Himalaya. *J For Res*. 2010; 21: 331–337. <https://doi.org/10.1007/s11676-010-0079-y>
7. Prager CM, Jing X, Henning JA, Read QD, Meidl P, Lavorel S, et al. Climate and multiple dimensions of plant diversity regulate ecosystem carbon exchange along an elevational gradient. *Ecosphere*. 2021; 12. <https://doi.org/10.1002/ecs2.3472>
8. Couralet C, Sterck FJ, Sass-Klaassen U, Van Acker J, Beeckman H. Species-specific growth responses to climate variations in understory trees of a central african rain forest. *Biotropica*. 2010; 42: 503–511. <https://doi.org/10.1111/j.1744-7429.2009.00613.x>
9. Zhao L, Xiang W, Li J, Lei P, Deng X, Fang X, et al. Effects of topographic and soil factors on woody species assembly in a Chinese subtropical evergreen broadleaved forest. *Forests*. 2015; 6: 650–669. <https://doi.org/10.3390/f6030650>
10. Das A, Nagendra H, Anand M, Bunyan M. Topographic and bioclimatic determinants of the occurrence of forest and grassland in tropical montane forest-grassland mosaics of the Western Ghats, India. *PLoS One*. 2015; 10: 1–19. <https://doi.org/10.1371/journal.pone.0130566> PMID: 26121353
11. Prada CM, Stevenson PR. Plant composition associated with environmental gradients in tropical montane forests (Cueva de Los Guacharos National Park, Huila, Colombia). *Biotropica*. 2016; 48: 568–576. <https://doi.org/10.1111/btp.12331>

12. Omijeh JE. Variations in species composition and forest structure among fragments along altitudinal gradients of Bagale forest reserve, Adamawa state, Nigeria. *Int J For Ecol Environ*. 2020; 1: 35–47. <https://doi.org/10.18801/ijfee.010220.05>
13. Hernández L, Dezzeo N, Sanoja E, Salazar L, Castellanos H. Changes in structure and composition of evergreen forests on an altitudinal gradient in the Venezuelan Guayana Shield. *Rev Biol Trop*. 2012; 60: 11–33. <https://doi.org/10.15517/rbt.v60i1.2360> PMID: 22458207
14. Christmann T, Menor IO. A synthesis and future research directions for tropical mountain ecosystem restoration. *Sci Rep*. 2021; 11: 1–17. <https://doi.org/10.1038/s41598-021-03205-y> PMID: 34907235
15. Willcock S, Phillips OL, Platts PJ, Balmford A, Burgess ND, Lovett JC, et al. Quantifying and understanding carbon storage and sequestration within the Eastern Arc Mountains of Tanzania, a tropical biodiversity hotspot. *Carbon Balance Manag*. 2014; 9: 1–17. <https://doi.org/10.1186/1750-0680-9-2> PMID: 24891875
16. Jürgen H, Siegmar WB, Sven G, Rütger TR, Christoph L. Tree Diversity, Forest Structure and Productivity along Altitudinal and Topographical Gradients in a Species-Rich Ecuadorian Montane Rain Forest. *Biotropica*. 2010; 42: 140–148. Available: <https://doi.org/http%3A//dx.doi.org/10.1111/j.1744-7429.2009.00547.x>
17. Homeier J, Breckle SW, Günter S, Rollenbeck RT, Leuschner C. Tree diversity, forest structure and productivity along altitudinal and topographical gradients in a species-rich Ecuadorian montane rain forest. *Biotropica*. 2010; 42: 140–148. <https://doi.org/10.1111/j.1744-7429.2009.00547.x>
18. Jara-Guerrero A, González-Sánchez D, Escudero A, Espinosa CI. Chronic Disturbance in a Tropical Dry Forest: Disentangling Direct and Indirect Pathways Behind the Loss of Plant Richness. *Front For Glob Chang*. 2021;4. <https://doi.org/10.3389/ffgc.2021.723985>
19. Sahu PK, Sagar R, Singh JS. Tropical forest structure and diversity in relation to altitude and disturbance in a Biosphere Reserve in central India. *Appl Veg Sci*. 2008; 11: 461–470. <https://doi.org/10.3170/2008-7-18537>
20. Thammanu S, Marod D, Han H, Bhusal N, Asanok L, Ketdee P, et al. The influence of environmental factors on species composition and distribution in a community forest in Northern Thailand. *J For Res*. 2021; 32: 649–662. <https://doi.org/10.1007/s11676-020-01239-y>
21. Baldeck CA, Harms KE, Yavitt JB, John R, Turner BL, Valencia R, et al. Soil resources and topography shape local tree community structure in tropical forests. *Proc R Soc B Biol Sci*. 2013;280. <https://doi.org/10.1098/rspb.2012.2532> PMID: 23256196
22. Zhang C, Li X, Chen L, Xie G, Liu C, Pei S. Effects of topographical and edaphic factors on tree community structure and diversity of subtropical mountain forests in the Lower Lancang River Basin. *Forests*. 2016;7. <https://doi.org/10.3390/f7100222>
23. Sheil D. Disturbance and distributions: Avoiding exclusion in a warming world. *Ecol Soc*. 2016;21. <https://doi.org/10.5751/ES-07920-210110>
24. Lee B, Park J, Lee H, Kim TK, Cho S, Yoon J, et al. Changes of tree species composition and distribution patterns in Mts. Jiri and Baegun, s. *Forests*. 2020; 11: 1–14. <https://doi.org/10.3390/f11020186>
25. Vacek Z, Vacek S, Bílek L, Remeš J, Štefančík I. Changes in horizontal structure of natural beech forests on an altitudinal gradient in the Sudetes. *Dendrobiology*. 2015; 73: 33–45. <https://doi.org/10.12657/denbio.073.004>
26. Kacholi DS. Analysis of Structure and Diversity of the Kilengwe Forest in the Morogoro Region, Tanzania. *Int J Biodivers*. 2014; 2014: 1–8. <https://doi.org/10.1155/2014/516840>
27. Mansourian S, Sumbi P, Bonifasi E, Meshack C, Malugu I, Vallauri D. Lessons Learnt from 10 Years of Restoration of Coastal and Sub-montane Tropical Forests: The East Usambara Landscape (Tanzania). *Wwf*. 2019.
28. Cronin DT, Libalah MB, Bergl RA, Hearn GW. Biodiversity and conservation of tropical montane ecosystems in the Gulf of Guinea, West Africa. *Arctic, Antarct Alp Res*. 2014; 46: 891–904. <https://doi.org/10.1657/1938-4246-46.4.891>
29. Muposhi VK, Chademana TC, Gandiwa E, Muboko N. Edge effects: impact of anthropogenic activities on vegetation structure and diversity in western Umfuduzi Park, Zimbabwe. *Afr J Ecol*. 2016; 54: 450–459. <https://doi.org/10.1111/aje.12300>
30. Gizachew B, Rizzi J, Shirima DD, Zahabu E. Deforestation and connectivity among protected areas of Tanzania. *Forests*. 2020; 11: 1–16. <https://doi.org/10.3390/f11020170>
31. Mwakosya J, Mligo C. The impacts of anthropogenic activities on the vegetation communities and structure in the western part of Rungwe forest reserve, Tanzania. *Tanzania J Sci*. 2014; 40: 60–78. Available: <https://www.ajol.info/index.php/tjs/article/view/148968/138470>.
32. Munishi PKT, Shear TH, Wentworth T, Temu RAPC. Compositional gradients of plant communities in submontane rainforests of Eastern Tanzania. *J Trop For Sci*. 2007; 19: 35–45.

33. Hailemariam MB, Temam TD. Pattern of plant community distribution along the elevational gradient and anthropogenic disturbance in Gole forest, Ethiopia. *Int J Ecol.* 2020;2020. <https://doi.org/10.1155/2020/6536374>
34. Estrada-Villegas S, Bailón M, Hall JS, Schnitzer SA, Turner BL, Caughlin T, et al. Edaphic factors and initial conditions influence successional trajectories of early regenerating tropical dry forests. *J Ecol.* 2020; 108: 160–174. <https://doi.org/10.1111/1365-2745.13263>
35. Platts PJ, Burgess ND, Gereau RE, Lovett JC, Marshall AR, McClean CJ, et al. Delimiting tropical mountain ecoregions for conservation. *Environ Conserv.* 2011; 38: 312–324. <https://doi.org/10.1017/S0376892911000191>
36. Burgess ND, Butynski TM, Cordeiro NJ, Doggart NH, Fjeldså J, Howell KM, et al. The biological importance of the Eastern Arc Mountains of Tanzania and Kenya. *Biol Conserv.* 2007; 134: 209–231. <https://doi.org/10.1016/j.biocon.2006.08.015>
37. Doggart N, Burgess N, Njana M, Sumbi P. Tanzania 's Nature Reserves Profiles of Tanzania 's 12 Nature Reserves. Tanzania For Conserv Group Tanzania For Conserv Gr. 2017; 1–32.
38. Milledge S, Elibariki R. Green gold: Ongoing efforts towards preventing illegal harvesting and exports of Tanzania's most valuable hardwoods. *Arc J.* 2004;17.
39. Kacholi DS. Floristic Similarity and Diversity Gradients in the Eastern Arc and Coastal Forests of Tanzania. *J Educ Humanit Sci.* 2018; 7: 93–104. Available: <https://www.researchgate.net/publication/329866671>.
40. Powers JS, Becknell JM, Irving J, Pérez-Aviles D. Diversity and structure of regenerating tropical dry forests in Costa Rica: Geographic patterns and environmental drivers. *For Ecol Manage.* 2009; 258: 959–970. <https://doi.org/10.1016/j.foreco.2008.10.036>
41. Legendre P. Studying beta diversity: ecological variation partitioning by multiple regression and canonical analysis. *J Plant Ecol.* 2008; 1: 3–8. <https://doi.org/10.1093/jpe/rtm001>
42. Bhutia Y, Gudasalamani R, Ganesan R, Saha S. Assessing forest structure and composition along the altitudinal gradient in the state of Sikkim, Eastern Himalayas, India. *Forests.* 2019; 10: 1–17. <https://doi.org/10.3390/f10080633>
43. Teshome M, Asfaw Z, Dalle G. Effect of environmental gradients on diversity and plant community distribution in remnant dry Afromontane forest of Mount Duro, Nagelle Arsi, Ethiopia. *Biodivers Res Conserv.* 2020; 58: 21–31. <https://doi.org/10.2478/biocr-2020-0004>
44. Doody K, Howell K FEA. Amani Nature Reserve, A Biodiversity Survey. East Usambara Conservation Area Management Programme, Tech Paper 52. In: Ministry of Natural Resources and Tourism Tanzania and Frontier-Tanzania. Tanga. 2001.
45. Mauya EW, Hansen EH, Gobakken T, Bollandsås OM, Malimbwi RE, Næsset E. Effects of field plot size on prediction accuracy of aboveground biomass in airborne laser scanning-assisted inventories in tropical rain forests of Tanzania. *Carbon Balance Manag.* 2015; 10: 1–14. <https://doi.org/10.1186/s13021-015-0021-x> PMID: 25983857
46. Beharrell NK, Hall SM, Ntemi SA. Vegetation in Nilo Forest Reserve: A biodiversity survey. East Usambara Conserv Area Manag Program Tech Pap 53 Front Tanzania; For Beekeep Div Metsahall Consult Dar es Salaam, Tanzania Vantaa, Finl. 2002.
47. Beals EW. Bray-curtis ordination: An effective strategy for analysis of multivariate ecological data. *Adv Ecol Res.* 1984; 14: 1–55. [https://doi.org/10.1016/S0065-2504\(08\)60168-3](https://doi.org/10.1016/S0065-2504(08)60168-3)
48. Requena Suarez D, Rozendaal DMA, De Sy V, Gibbs DA, Harris NL, Sexton JO, et al. Variation in aboveground biomass in forests and woodlands in Tanzania along gradients in environmental conditions and human use. *Environ Res Lett.* 2021;16. <https://doi.org/10.1088/1748-9326/abe960>
49. Hejzmanova-Nežerková P, Hejzman M. A canonical correspondence analysis (CCA) of the vegetation-environment relationships in Sudanese savannah, Senegal. *South African J Bot.* 2006; 72: 256–262. <https://doi.org/10.1016/j.sajb.2005.09.002>
50. Andrew SM, Moe SR, Totland Ø, Munishi PKT. Species composition and functional structure of herbaceous vegetation in a tropical wetland system. *Biodivers Conserv.* 2012; 21: 2865–2885. <https://doi.org/10.1007/s10531-012-0342-y>
51. Quinn GP, Keough MJ. Introduction experimental design and data analysis for biologists. Cambridge Univ Press. 2012; 1–13.
52. Zare Chahouki M a, Zare Chahouki A. Predicting the distribution of plant species using logistic regression (Case study: Garizat rangelands of Yazd province). *Desert.* 2010; 15: 151–158.
53. Kyaw Myo K, Thwin S, Khaing N. Floristic Composition, Structure and Soil Properties of Mixed Deciduous Forest and Deciduous Dipterocarp Forest: Case Study in Madan Watershed, Myanmar. *Am J Plant Sci.* 2016; 07: 279–287. <https://doi.org/10.4236/ajps.2016.72027>

54. Thinsungnoen T, Kaoungku N, Durongdumronchai P. The Clustering Validity with Silhouette and Sum of Squared Errors. 2015; 3: 4–11. <https://doi.org/10.12792/iciae2015.012>
55. Zhang W, Huang D, Wang R, Liu J, Du N. Altitudinal patterns of species diversity and phylogenetic diversity across temperate mountain forests of northern China. *PLoS One*. 2016; 11: 1–13. <https://doi.org/10.1371/journal.pone.0159995> PMID: 27454311
56. Rahman IU, Hart RE, Ijaz F, Afzal A, Iqbal Z, Calixto ES, et al. Environmental variables drive plant species composition and distribution in the moist temperate forests of Northwestern Himalaya, Pakistan. *PLoS One*. 2022; 17: 1–27. <https://doi.org/10.1371/journal.pone.0260687> PMID: 35202409
57. Bentsi-Enchill F, Dampney FG, Pappoe ANM, Ekumah B, Akotoye HK. Impact of anthropogenic disturbance on tree species diversity, vegetation structure and carbon storage potential in an upland evergreen forest of Ghana, West Africa. *Trees, For People*. 2022; 8: 100238. <https://doi.org/10.1016/j.tfp.2022.100238>
58. Tilman D, Lehman C. Human-caused environmental change: Impacts on plant diversity and evolution Environmental Constraints in Plant Communities. *Proc Natl Acad Sci U S A*. 2001; 98: 5433–5440.
59. Sinha S, Badola HK, Chhetri B, Gaira KS, Lepcha J, Dhyani PP. Effect of altitude and climate in shaping the forest compositions of Singalila National Park in Khangchendzonga Landscape, Eastern Himalaya, India. *J Asia-Pacific Biodivers*. 2018; 11: 267–275. <https://doi.org/10.1016/j.japb.2018.01.012>
60. Hu W, Yao J, He Q, Chen J. Elevation-dependent trends in precipitation observed over and around the tibetan plateau from 1971 to 2017. *Water (Switzerland)*. 2021; 13: 1–17. <https://doi.org/10.3390/w13202848>
61. Iqbal Z, Zeb A, Abd_Allah EF, Rahman IU, Khan SM, Ali N, et al. Ecological assessment of plant communities along the edaphic and topographic gradients of biha valley, District Swat, Pakistan. *Appl Ecol Environ Res*. 2018; 16: 5611–5631. https://doi.org/10.15666/aeer/1605_56115631
62. Napoli A, Crespi A, Ragone F, Maugeri M, Pasquero C. Variability of orographic enhancement of precipitation in the Alpine region. *Sci Rep*. 2019; 9: 1–8. <https://doi.org/10.1038/s41598-019-49974-5> PMID: 31527776
63. Brown R, Fredericksen T. Topographic Factors Affecting the Tree Species Composition of Forests in the Upper Piedmont of Virginia. *Va J Sci*. 2008; 59: 1.
64. Cirimwami L, Doumenge C, Kahindo JM, Amani C. The effect of elevation on species richness in tropical forests depends on the considered lifeform: results from an East African mountain forest. *Trop Ecol*. 2019; 60: 473–484. <https://doi.org/10.1007/s42965-019-00050-z>
65. Duchesne L, Ouimet R. Relationships between Structure, Composition, and Dynamics of the Pristine Northern Boreal Forest and Air Temperature, Precipitation, and Soil Texture in Quebec (Canada). *Int J For Res*. 2009; 2009: 1–13. <https://doi.org/10.1155/2009/398389>
66. Amisshah L, Mohren GMJ, Bongers F, Hawthorne WD, Poorter L. Rainfall and temperature affect tree species distribution in Ghana. *J Trop Ecol*. 2014; 30: 435–446. <https://doi.org/10.1017/S026646741400025X>
67. Krishnadas M, Kumar A, Comita LS. Environmental gradients structure tropical tree assemblages at the regional scale. *J Veg Sci*. 2016; 27: 1117–1128. <https://doi.org/10.1111/jvs.12438>
68. Méndez-Toribio M, Ibarra-Manríquez G, Navarrete-Segueda A, Paz H. Topographic position, but not slope aspect, drives the dominance of functional strategies of tropical dry forest trees. *Environ Res Lett*. 2017; 12. <https://doi.org/10.1088/1748-9326/aa717b>
69. Iqbal M, Khan SM, Azim Khan M, Ahmad Z, Ahmad H. A novel approach to phytosociological classification of weeds flora of an agro-ecological system through Cluster, Two Way Cluster and Indicator Species Analyses. *Ecol Indic*. 2018; 84: 590–606. <https://doi.org/10.1016/j.ecolind.2017.09.023>
70. Chitiki AK. Edaphic correlates of tree species diversity, composition, and distribution in an eastern arc biodiversity hotspot, Tanzania. *Cent Asian J Environ Sci*. . . . 2020; 4: 206–218. Available: https://www.cas-press.com/article_111290_f65676500931133451dbd6ff697e0f84.pdf.
71. Riesch F, Stroh HG, Tonn B, Isselstein J, Riesch F, Stroh HG, et al. Soil pH and phosphorus drive species composition and richness in semi-natural heathlands and grasslands unaffected by twentieth-century agricultural intensification. *Plant Ecol Divers*. 2018; 00: 1–15. <https://doi.org/10.1080/17550874.2018.1471627>
72. Bueis T, Bravo F, Pando V, Kissi YA, Turrión MB. Phosphorus availability in relation to soil properties and forest productivity in *Pinus sylvestris* L. plantations. *Ann For Sci*. 2019; 76. <https://doi.org/10.1007/s13595-019-0882-3>
73. Baribault TW, Kobe RK, Finley AO. Tropical tree growth is correlated with soil phosphorus, potassium, and calcium, though not for legumes. *Ecol Monogr*. 2012; 82: 189–203. <https://doi.org/10.1890/11-1013.1>

74. Birhanu L, Bekele T, Tesfaw B, Demissew S. Relationships between topographic factors, soil and plant communities in a dry afro-montane forest patches of Northwestern Ethiopia. *PLoS One*. 2021; 16: 1–18. <https://doi.org/10.1371/journal.pone.0247966> PMID: 33711027
75. Long A, Long C, Yang X, Long W, Li D. Soil Nutrients Influence Plant Community Assembly in Two Tropical Coastal Secondary Forests. *Soil Nutrients Influence Plant Community Assembly in Two Tropical Coastal Secondary Forests*. 2022;11. <https://doi.org/10.1177/1940082918817956>
76. Hou E, Chen C, Luo Y, Zhou G, Yuanwen K, Zhang Y, et al. Effects of climate on soil phosphorus cycle and availability in natural terrestrial ecosystems. *Global*. 2018; 1–13. <https://doi.org/10.1111/gcb.14093> PMID: 29450947
77. Dornelas M. Disturbance and change in biodiversity. *Philos Trans R Soc B Biol Sci*. 2010; 365: 3719–3727. <https://doi.org/10.1098/rstb.2010.0295> PMID: 20980319
78. Epila J, Verbeeck H, Otim-Epila T, Okullo P, Kearsley E, Steppe K. The ecology of *Maesopsis eminii* Engl. in tropical Africa. *Afr J Ecol*. 2017; 55: 679–692. <https://doi.org/10.1111/aje.12408>
79. Huebner CD, McGill DW. The importance of disturbance versus physiography in defining vegetation composition and predicting possible successional trajectories. *Castanea*. 2018; 83: 54–76. <https://doi.org/10.2179/17-139>