

Supplementary Table 5 . Age-standardised 10-year net survival (%) with 95% confidence interval (CI), 2000-2004. Children (0-14 years) diagnosed with low-grade astrocytoma (WHO grade I and II) or medulloblastoma, by country.

Country	Period of diagnosis	Astrocytoma, WHO grade I and II		Astrocytoma, WHO grade III and IV		Medulloblastoma	
		NS	95% CI	NS	95% CI	NS	95% CI
AMERICA (CENTRAL AND SOUTH)							
Argentina*	2000-2004	79.6	71.2 - 87.9	4.6	0.0 - 9.5	44.1	37.6 - 50.6
Brazil	2000-2004	92.9	78.9 - 100.0	-	-	50.3	28.0 - 72.6
Colombia	2000-2004	60.9 §	32.0 - 89.7	0.0 §	0.0 - 0.0	-	-
Ecuador	2000-2004	-	-	-	-	46.4 §	21.3 - 71.5
Puerto Rico*	2000-2004	94.9	87.9 - 100.0	-	-	58.4	32.0 - 84.8
AMERICA (NORTH)							
Canada	2000-2004	92.4	87.2 - 97.6	33.4	11.0 - 55.8	65.8	50.4 - 81.3
United States	2000-2004	93.9	92.9 - 94.8	28.0	24.1 - 31.9	71.4	68.2 - 74.5
ASIA							
Israel*	2000-2004	93.5	88.3 - 98.6	13.3	0.0 - 28.4	63.9	53.1 - 74.8
Japan	2000-2004	80.9	61.9 - 99.8	13.1	0.5 - 25.7	64.4	48.5 - 80.4
Jordan*	2000-2004	74.3 §	45.1 - 100.0	61.7 §	42.5 - 80.9	75.5 §	61.6 - 89.3
Korea*	2000-2004	93.8	82.3 - 100.0	20.7	11.0 - 30.5	56.5	49.4 - 63.5
Singapore*	2000-2004	-	-	-	-	40.0	12.2 - 67.9
Taiwan*	2000-2004	88.9	80.5 - 97.3	31.9	18.4 - 45.3	46.2	35.2 - 57.3
Thailand	2000-2004	-	-	13.4 §	0.0 - 28.4	31.8 §	11.9 - 51.7
EUROPE							
Austria*	2000-2004	-	-	17.4	3.0 - 31.8	66.7	47.1 - 86.3
Belarus*	2000-2004	75.0	64.9 - 85.1	8.4	0.0 - 20.7	57.4	43.6 - 71.1
Belgium*	2000-2004	85.8	71.2 - 100.0	-	-	-	-
Croatia*	2000-2004	72.8	47.8 - 97.8	-	-	56.1	37.2 - 75.1
Czech Republic*	2000-2004	84.7	73.5 - 95.9	20.0	3.7 - 36.4	42.9	18.5 - 67.4
Denmark*	2000-2004	93.1	85.6 - 100.0	-	-	62.2	44.9 - 79.4
Estonia*	2000-2004	100.0	100.0 - 100.0	-	-	-	-

Supplementary Table 5 . Age-standardised 10-year net survival (%) with 95% confidence interval (CI), 2000-2004. Children (0-14 years) diagnosed with low-grade astrocytoma (WHO grade I and II) or medulloblastoma, by country.

Country	Period of diagnosis	Astrocytoma, WHO grade I and II		Astrocytoma, WHO grade III and IV		Medulloblastoma	
		NS	95% CI	NS	95% CI	NS	95% CI
Finland*	2000-2004	97.1	93.0 - 100.0	25.0	2.9 - 47.1	68.8	47.0 - 90.7
France*	2000-2004	87.3	84.3 - 90.2	10.0	1.3 - 18.7	55.6	50.1 - 61.1
Germany	2000-2004	82.5	69.9 - 95.1	7.7	0.0 - 18.7	64.6	48.1 - 81.1
Ireland*	2000-2004	87.1	77.5 - 96.8	-	-	-	-
Italy	2000-2004	94.8	90.6 - 98.9	17.7	5.6 - 29.7	59.8	49.6 - 69.9
Lithuania*	2000-2004	-	-	-	-	35.9	12.3 - 59.5
Netherlands*	2000-2004	90.1	85.2 - 95.1	25.0	12.1 - 38.0	48.5	37.9 - 59.1
Norway*	2000-2004	92.3	85.2 - 99.4	-	-	66.8	45.7 - 87.8
Poland*	2000-2004	84.3	76.9 - 91.8	27.6	16.7 - 38.5	53.7	46.0 - 61.3
Portugal*	2000-2004	84.2	73.6 - 94.9	-	-	54.0	38.6 - 69.3
Slovakia*	2000-2004	84.3	75.2 - 93.5	-	-	34.9	16.2 - 53.7
Slovenia*	2000-2004	81.8	60.2 - 100.0	-	-	-	-
Spain	2000-2004	90.0	82.6 - 97.4	-	-	50.1	35.2 - 65.0
Sweden*	2000-2004	95.6	90.6 - 100.0	20.5	0.0 - 42.3	74.1	57.9 - 90.3
Switzerland*	2000-2004	95.1	89.6 - 100.0	9.1	0.0 - 22.4	60.4	44.0 - 76.9
United Kingdom*	2000-2004	92.9	90.4 - 95.4	11.3	5.1 - 17.4	59.3	53.4 - 65.2
OCEANIA							
Australia*	2000-2004	84.0	75.0 - 92.9	16.3	7.1 - 25.5	68.6	58.8 - 78.4
New Zealand*	2000-2004	80.1	63.0 - 97.1	-	-	70.7	49.8 - 91.7

* Countries with 100% coverage of the national population. § Survival estimates considered less reliable because the proportion of patients lost to follow-up or censored alive prior to five years, and/or the proportion of diagnoses based on a death certificate or autopsy, and/or the proportion of patients registered with incomplete dates, was 15% or more. Survival estimates in bold are age-standardised; survival estimates in italics are not age-standardised.