

Supplement 2. The demographic characteristics of the emergency clinicians and their reported acceptability of the palliative dyspnea protocol.

Clinician demographic	Frequency	Neutral*	Somewhat Helpful*	Very Helpful*	P-value
	N=35 (%)	n=20 (%)	n=9 (%)	n=6 (%)	
Role					0.692
Attending physician	14 (40.0)	10 (50.0)	2 (22.2)	2 (33.3)	
Resident physician	11 (31.4)	5 (25.0)	4 (44.4)	2 (33.3)	
Physician assistant	10 (28.6)	5 (25.0)	3 (33.3)	2 (33.3)	
Year of clinical experience					0.501
1-5	19 (54.3)	9 (45.0)	7 (77.8)	3 (50.0)	
6-10	5 (14.3)	4 (20.0)	0	1 (16.7)	
>10	11 (31.4)	7 (35.0)	2 (22.2)	2 (33.3)	
Age (year)					0.647
20-29	6 (17.1)	2 (10.0)	3 (33.3)	1 (16.7)	
30-39	17 (48.6)	11 (55.0)	4 (44.4)	2 (33.3)	
40-49	8 (22.9)	4 (20.0)	2 (22.2)	2 (33.3)	
>49	4 (11.4)	3 (15.0)	0	1 (16.7)	
Gender					0.587
Male	14 (40.0)	7 (35.0)	5 (55.6)	2 (33.3)	
Female	19 (54.3)	12 (60.0)	4 (44.4)	3 (50.0)	
Decline to answer	2 (5.7)	1 (5.0)	0	1 (16.7)	

*No participant rated 'somewhat unhelpful' and 'very unhelpful'