

## Changes of lymphocyte membrane fluidity in rheumatoid arthritis: a fluorescence polarisation study

E BECCERICA,<sup>1</sup> G PIERGIACOMI,<sup>1</sup> G CURATOLA,<sup>2</sup> AND G FERRETTI<sup>2</sup>

*From the Institutes of <sup>1</sup>Rheumatology and <sup>2</sup>Biochemistry, Medical Faculty, University of Ancona, Via Ranieri, 60131 Ancona, Italy*

**SUMMARY** Fluorescence polarisation of 1,6-diphenyl-1,3,5-hexatriene was used to study the lymphocyte membrane in rheumatoid arthritis. The increase of polarisation value in the patients (n=27) compared with healthy controls (n=32) suggests a decrease of membrane fluidity. Moreover, erythrocyte sedimentation rate (ESR) and plasma fibrinogen concentrations were positively correlated with lymphocyte fluorescence polarisation values (r=0.66 and r=0.76 respectively). The results suggest that the changes in lymphocyte membrane fluidity could be involved in the pathogenetic mechanism of rheumatoid arthritis.

**Key words:** 1,6-diphenyl-1,3,5-hexatriene fluorescence polarisation.

Rheumatoid arthritis (RA) is characterised by complex abnormalities of the immunological system and by the presence of defects in the regulation of the immune response. The principal immunopathogenetic features are the production of rheumatoid factor, the presence of circulating immune complexes, and alterations in the functional activities of polymorphonuclear cells<sup>1,2</sup> and lymphocytes.<sup>3-5</sup> Immunoregulation is mediated by a complex network of cellular interactions involving both cell-cell contacts and soluble factors.<sup>6</sup> In these events a key role is performed by cell membranes, both plasmatic and intracellular, as they are involved in cellular recognition phenomena<sup>7</sup> and in signal transduction.<sup>8</sup> Although the aetiology of RA is still unknown, it has become evident that lymphocytes have an important role in the pathogenesis of RA,<sup>9</sup> as suggested by the observation that profound clinical improvements follow depletion of recirculating lymphocytes either by lymphopheresis or thoracic duct drainage,<sup>10,11</sup> and that lymphocyte reinfusion causes transient exacerbations.<sup>4</sup> Several studies have shown abnormal lymphoid cell morphology<sup>12</sup> and functions in the peripheral blood of patients with RA<sup>3-5,13</sup> and an increase of circulating activated lymphocytes.<sup>14,15</sup>

As cell membranes have been shown to be primarily involved in immune responses<sup>16</sup> in inflammatory and cytotoxic events,<sup>16,17</sup> the possibility of lymphocyte membrane alterations at the molecular level has to be considered. Alterations in the behaviour of lymphocytes from patients with RA may be due to abnormalities in membrane fluidity. Membrane fluidity has an important role in regulating cellular functions affecting the conformation of membrane-bound enzymes,<sup>18</sup> the diffusion<sup>19</sup> or the degree of receptor protein exposure, or both, and their recycling<sup>20</sup>; fluidity as measured by spectroscopic techniques is a complex physicochemical feature in which mobility and order of membrane components and membrane permeability properties are strictly interconnected.<sup>21</sup> Changes in composition and in molecular organisation are the principal determinants of alterations of membrane fluidity observed in many human diseases.<sup>22-27</sup>

In this study we used the fluorescence polarisation of 1,6-diphenyl-1,3,5-hexatriene (DPH), which has been widely used to analyse membrane fluidity of blood cells in normal and pathological conditions.<sup>23-27</sup> The lipophilic probe DPH, though primarily embedded in the plasma membrane, is not located exclusively there, but also penetrates the intracellular membranes, thus characterising all cell membrane apparatus.<sup>28</sup>

Our principal aim was to study the physicochemical state of the lymphocyte membrane in patients

Accepted for publication 25 November 1987.

Correspondence to Professor G Curatola, Istituto di Biochimica, Facoltà di Medicina e Chirurgia, Università degli Studi di Ancona, Via Ranieri, 60131 Ancona, Italia.

with RA and to try to correlate this with clinical and biochemical indices of RA activity. The results are discussed in the light of the possible involvement of lymphocyte membrane in the pathogenetic mechanism of RA.

## Patients and methods

### PATIENTS

Twenty seven patients (19 women, eight men; mean age 53.8 (SD 14.4) years, range 20–79) with classical or definite RA according to the criteria of the American Rheumatism Association were studied. None had taken disease modifying anti-rheumatic drugs for at least six months and non-steroidal anti-inflammatory drugs for at least three days. Disease activity was assessed clinically by the Ritchie index and by measurement of a series of objective indices, including erythrocyte sedimentation rate (ESR),<sup>29</sup>  $\alpha_2$  globulins,<sup>30</sup> C reactive protein (CRP),<sup>31</sup> plasma fibrinogen concentrations,<sup>32</sup> platelet count, plasma sulphhydryl groups,<sup>33</sup> and plasma viscosity.<sup>34</sup>

Thirty two healthy subjects (19 women, 13 men; mean age 45 (15.4) years, range 22–78) acted as controls.

### LYMPHOCYTES

Lymphocytes were obtained from freshly drawn heparinised blood by Ficoll-Isopaque (Lymphoprep; Nyegard and Co, A/S, Oslo, Norway) centrifugal sedimentation.<sup>35</sup> A population of cells containing 98% lymphocytes was obtained. Possible contamination by monocytes was tested by staining with  $\alpha$ -naphthol acetate esterase.<sup>36</sup> The isolated lymphocytes were washed twice, resuspended in phosphate buffered saline to a concentration of  $2 \times 10^6$  cells/ml, and immediately used for fluorescence measurements.

### FLUORESCENCE LABELLING OF LYMPHOCYTES

1,6-Diphenyl-1,3,5-hexatriene was used as the fluorescent probe for monitoring the degree of fluidity of the cell membrane. DPH was dissolved in tetrahydrofuran at a concentration of  $2 \times 10^{-3}$  mol/l. For cell labelling, the solution of DPH was diluted 1000-fold by injection into vigorously stirred phosphate buffered saline. The dispersion of DPH obtained ( $2 \times 10^{-6}$  mol/l) was clear and practically lacking in fluorescence. One millilitre of this fluorescent probe solution was added to 1 ml of cellular suspension ( $2 \times 10^6$  cells/ml) to give a final probe concentration of  $10^{-6}$  mol/l.<sup>27, 28</sup>

The incorporation of DPH into lymphocyte membranes was followed by a steep increase in fluor-

escence intensity. With our experimental conditions we did not observe significant changes in spectral characteristics and kinetic incorporations of DPH into lymphocyte membranes of patients with RA in comparison with control membranes. After 45 minutes' incubation at room temperature the labelled cells were used for fluorescence studies.

### FLUORESCENCE POLARISATION ANALYSIS

DPH fluorescence polarisation was measured with a Perkin Elmer spectrofluorimeter MPF 44A. The excitation and emission wavelengths were respectively 365 nm and 430 nm. The degree of DPH fluorescence polarisation (P) was obtained by the following equation:

$$P = \frac{I_{\parallel} - I_{\perp} g}{I_{\parallel} + I_{\perp} g}$$

where g is an instrumental correction factor,  $I_{\parallel}$  and  $I_{\perp}$  are respectively the emission intensities polarised vertically and horizontally to the direction of the polarised light. For DPH the fluorescence polarisation values depend on probe rotational mobility and on the degree of order of membrane molecules; changes in fluorescence polarisation values have been correlated with changes in membrane microviscosity. In general, an increase of fluorescence polarisation indicates an increase of microviscosity and therefore a decrease of fluidity, assuming, on physical grounds, that fluidity is the reciprocal of viscosity.<sup>21</sup>

### STATISTICAL METHODS

All results are expressed as mean (SD). Data were analysed by Student's *t* test and the Mann-Whitney U test, and linear regression analysis was used for the calculation of correlation coefficients.

## Results

Fig. 1 shows the individual values of DPH fluorescence polarisation in lymphocytes from healthy controls and patients with RA. Of the 27 patients with RA studied, 16 showed polarisation values clearly above the range of the controls, while the remaining cases were above the control mean value. Among the patients with RA there was a significant increase of DPH polarisation (Table 1) ( $p < 0.01$  Mann-Whitney test), indicating a decreased lymphocyte membrane fluidity. The applicability of this molecular approach in the study of RA pathogenetic mechanisms is confirmed by the lack of variability of polarisation values in relation to the sex and the age of the subjects. There were no significant differences in the mean polarisation values in relation to the sex and age either in the controls or in the

patients with RA (Tables 1 and 2). In healthy women and men we obtained mean (SD) fluorescence polarisation values of 0.215 (0.004) and 0.218 (0.006) respectively. These values were




Fig. 1 Fluorescence polarisation values of 1,6-diphenyl-1,3,5-hexatriene in lymphocyte membranes of patients with rheumatoid arthritis (■, □) and of control subjects (●, ○). The dashed line indicates the control mean fluorescence polarisation value, and the continuous lines the range of values for the controls.

Table 1 DPH fluorescence polarisation values related to sex of the subjects in lymphocytes from controls and patients with rheumatoid arthritis

	Controls		Patients with rheumatoid arthritis	
	Fluorescence polarisation values <sup>†</sup>	Number	Fluorescence polarisation values <sup>†</sup>	Number
Men	0.218 (0.006)	13	0.235 (0.008)*	8
Women	0.215 (0.004)	19	0.232 (0.009)*	19
Total	0.217 (0.005)	32	0.233 (0.009)*	27

\* $p < 0.01$  Mann-Whitney test.

<sup>†</sup>Values are mean (SD).

Table 2 DPH fluorescence polarisation values related to age in lymphocytes from controls and patients with rheumatoid arthritis

Subjects	Age (years)			
	20-35	35-50	50-65	65-80
Controls	0.215 (0.010) <sup>†</sup> (n=12)	0.217 (0.004) (n=8)	0.218 (0.004) (n=6)	0.218 (0.007) (n=6)
Patients with rheumatoid arthritis	0.228 (0.010)* (n=6)	0.231 (0.007)** (n=5)	0.231 (0.009)*** (n=8)	0.237 (0.012)**** (n=8)

\* $p = 0.008$ , \*\* $p = 0.001$ , \*\*\* $p = 0.005$ , \*\*\*\* $p = 0.016$  as compared with controls.

<sup>†</sup>Data are presented as mean (SD).

significantly different from those in women and men with RA (0.232 (0.009) and 0.235 (0.008) respectively;  $p < 0.01$  Mann-Whitney test) (Table 1). These results indicate an average increase of membrane fluorescence polarisation of 7-8% in the RA lymphocytes as compared with normal lymphocytes. Although this difference between groups appears to be small in absolute value, the standard deviation of DPH fluorescence polarisation measurements is extremely small so that the difference in the mean values for the two groups is highly significant ( $p < 0.01$ ). To determine possible differences in membrane fluidity in cells other than lymphocytes the fluidity of erythrocyte membrane isolated from patients with RA and from healthy controls was tested by fluorescence polarisation. No difference was detected in fluidity of erythrocyte membranes from patients with RA and from control subjects (data not shown). When 0.229 was taken as the fluorescence polarisation cut off point, this being the upper limit of lymphocyte membrane fluidity in the normals (Fig. 1), the patients with RA were divided into two classes, one having fluorescence polarisation values lower than 0.229 and the other having values higher than 0.229. The patients with decreased lymphocyte membrane fluidity had the

Table 3 Comparison of disease activity indices and DPH fluorescence polarisation values (P) in lymphocytes of patients with rheumatoid arthritis

	$P < 0.229$	$P > 0.229$
ESR (mm/h)	51 (19) <sup>†</sup>	71 (30)*
Fibrinogen (g/l)	3.68 (0.65)	6.16 (1.34)*
$\alpha_2$ Globulins (g/l)	0.78 (0.07)	0.79 (0.07)
Platelet count ( $\times 10^9/l$ )	300 (70)	313 (101)
Plasma viscosity (cP):	37°C	1.38 (0.16)
	18°C	2.11 (0.21)
C reactive protein (mg/l)	28 (18)	34 (12)
Plasma sulphhydryl groups (mg/l)	13.7 (1.5)	13.1 (0.9)
Ritchie index	17.6 (9)	20 (17)

\* $p < 0.01$ .

<sup>†</sup>Data are presented as mean (SD).

highest ESR, CRP concentrations, Ritchie index, and plasma fibrinogen concentrations, whereas the patients with the lowest ESR, CRP concentrations, Ritchie index, and plasma fibrinogen concentrations had lymphocyte membrane fluidity in the normal range (Table 3). There were no differences in  $\alpha_2$  globulin concentrations, platelet count, plasma sulphhydryl groups, and plasma viscosity in the two groups. Moreover, both ESR and plasma fibrinogen concentrations were positively correlated with lymphocyte fluorescence polarisation measurements ( $r=0.66$ ,  $p<0.01$ ;  $r=0.76$ ,  $p<0.01$  respectively (Figs 2 and 3).

**Discussion**

Using fluorescence polarisation, we have shown a decrease of fluidity in lymphocytes from patients with RA. Although apparently slight (average increase 7–8%), the observed changes in membrane physical state are noteworthy as much experimental evidence has shown that even small modifications of fluidity can affect membrane functions through the control exerted by phospholipid matrix on membrane protein activities. Moreover, changes similar to those found by us have been reported in cell membranes either in experimental or in human disease.<sup>24 37-39</sup>




Fig. 2 Correlation between erythrocyte sedimentation rate (ESR) and fluorescence polarisation of 1,6-diphenyl-1,3,5-hexatriene in lymphocyte membranes from patients with rheumatoid arthritis ( $r=0.66$ ,  $p<0.01$ ).




Fig. 3 Correlation between plasma fibrinogen concentrations and fluorescence polarisation of 1,6-diphenyl-1,3,5-hexatriene in lymphocyte membranes from patients with rheumatoid arthritis ( $r=0.76$ ,  $p<0.01$ ).

The study of cell membrane fluidity in various human diseases has received increasing attention in recent years and it has been used to investigate disease at the molecular level. The alterations we observed in lymphocyte membrane can be considered specifically linked to this cell as non-specific effects can be excluded by the finding that erythrocyte membrane fluidity was not modified. Further relevance to these results comes from the association of the highest polarisation values with the highest Ritchie index and CRP concentrations; fluorescence polarisation values were also positively correlated with ESR and plasma fibrinogen concentrations. As ESR and plasma fibrinogen concentrations are considered to be indicators of disease activity our results suggest that alterations in lymphocyte membrane fluidity could be related to inflammation; it is not possible to explain at present, however, why positive correlations have been found for only two indices and not for all the indices studied.

A hypothesis can be advanced about the molecular mechanisms involved in the changes of membrane physical state shown in RA lymphocytes. The membrane fluidity is related to the structural organisation of phospholipid molecules, in particular in the membrane hydrophobic core probed by DPH; a decrease of acyl chain unsaturation produces a decrease of fluidity. After the respiratory burst the release of free radicals and active oxygen species causes damage to membrane proteins<sup>40</sup> and induces the peroxidation of polyunsaturated fatty acids of membrane phospholipids.<sup>41-42</sup> In phospholipid liposomes lipid peroxidation has been shown to increase membrane rigidity.<sup>43</sup> This effect could be of importance in lymphocyte membranes, which have a high content of unsaturated fatty acid and a very high susceptibility to lipid peroxidation. Moreover, in RA there is evidence of increased levels of free radical oxidation products in serum and synovial fluid<sup>44-45</sup> and evidence of decreased levels of free radical scavengers and antioxidant enzyme systems, which normally protect the biological membranes from peroxidation. The increase of lipid peroxidation or the decrease of antioxidant systems could induce the observed changes in lymphocyte membrane fluidity, and these could be either a passive consequence of the disease condition or be directly involved in its pathogenetic mechanism. Although further studies are necessary to elucidate these aspects of membrane damage, we suggest that the lymphocyte membrane can be used to investigate molecular events of RA.

The authors are grateful to Professor Claudio Cervini for his helpful comments in the preparation of the manuscript.

## References

- Gale R, Bertouch J V, Gordon T P, Bradley J, Roberts-Thompson P J. Neutrophil activation by immune complexes and the role of rheumatoid factor. *Ann Rheum Dis* 1984; **43**: 34-9.
- King S L, Parker J, Cooper R, Sturrock R, Gemmel C G. Polymorphonuclear leucocyte function in rheumatoid arthritis. *Br J Rheumatol* 1986; **25**: 26-33.
- Van Boxel J A, Paget S A. Predominantly T-cell infiltrate in rheumatoid membranes. *N Engl J Med* 1975; **293**: 517-20.
- Cervini C, Numo R. Artrite reumatoide. In: Introzzi P, ed. *Trattato Italiano di medicina interna*. Part 10. Florence: USES, 1983: 233-320.
- Wooley P H, Panayi G S. Studies of lymphocytes in rheumatoid arthritis. *Ann Rheum Dis* 1978; **37**: 343-6.
- Waldmann T A, Broder S. Polyclonal B-cell activators in the study of the regulation of immunoglobulin synthesis in the human system. *Adv Immunol* 1982; **32**: 1-25.
- Zanders E. Lymphocyte receptors and transmembrane signalling. In: Cohen P, Houslay M D, eds. *Molecular mechanisms of transmembrane signalling* Amsterdam: Elsevier, 1985: 411-27.
- Weigandt H. Hormone reception and the cell surface—an overview. In: Bertoli E, Chapman D, Cambia A, Scapagnini U, eds. *Biomembrane and receptor mechanism*. Padova: Liviana Press, 1987: 269-77.
- Pearson C M, Paulus H E, Mackleder H. The role of the lymphocyte and its products in the propagation of joint disease. *Ann NY Acad Sci* 1975; **256**: 150-68.
- Paulus H E, Machleder H I, Levine S, Yu D T Y, MacDonald N S. Lymphocyte involvement in rheumatoid arthritis: in a longitudinal thoracic duct drainage. *Arthritis Rheum* 1977; **20**: 1249-62.
- Karsh J, Klippel J M, Plotz P H, Decker J L, Wright D G, Flye M W. Lymphapheresis in rheumatoid arthritis: a randomized trial. *Arthritis Rheum* 1981; **24**: 867-73.
- De Vries E, Van Buijssen A C, Van Der Weij J P, Haasnoot C J P, Meijer C J L M, Cats A. Morphometric analysis of peripheral blood and synovial fluid lymphocytes of patients with rheumatic disease. *J Rheumatol* 1983; **10**: 12-18.
- Corrigal V, Panayi G S, Laurent R. Lymphocyte studies in rheumatoid arthritis. *Scand J Rheumatol* 1979; **8**: 10-16.
- Papadimitrou G M, Bacon P A, Hall N D. Circulating activated lymphocytes in rheumatoid arthritis: a marker of synovial inflammation. *J Rheumatol* 1982; **9**: 217-23.
- Froebel K S, Lewis D, Dickson R, Sturrock R D. Spontaneous lymphocyte activity in rheumatoid arthritis in a longitudinal study in relation to gold therapy. *Clin Rheumatol* 1985; **4**: 301-7.
- Loor F. Cell surface-cell cortex transmembranous interactions with special reference to lymphocyte functions. In: Poste G, Nicholson G L, eds. *Cytoskeletal elements and plasma membrane organization*. Vol 6. Amsterdam: North Holland 1981: 253-335.
- Roozmond R C, Bonavida B. Effect of altered membrane fluidity on NK cell-mediated cytotoxicity. *J Immunol* 1985; **134**: 2209-14.
- Lenaz G, Curatola G, Fiorini R M, Parenti Castelli G. Membrane fluidity and its role in the regulation of cellular processes. *Biology of Cancer* 1983; **2**: 25-34.
- Edidin M. Rotational and translational diffusion in membranes. *Annual Reviews of Biophysics and Bioengineering* 1974; **3**: 179-201.
- Borochoy H, Shinitzky M. Vertical displacement of membrane proteins mediated by changes in microviscosity. *Proc Natl Acad Sci USA* 1976; **73**: 4526-30.
- Shinitzky M. Membrane fluidity and cellular functions. In: Shinitzky M, ed. *Physiology of membrane fluidity*. Boca Raton, Florida: CRC Press, 1984; 1-52.
- Cooper R A. Abnormality of cell membrane fluidity in the pathogenesis of disease. *N Engl J Med* 1977; **197**: 371-7.

- 23 Curatola G, Ferretti G, Bertoli E, Dotti M, Bartolotta E, Giorgi P L. Changes of membrane fluidity in erythrocytes of obese children: a spin label study. *Pediatr Res* 1987; **22**: 141-4.
- 24 Inbar M, Goldman R, Inbar I, et al. Fluidity differences of membrane lipids in human and leukemic lymphocytes as controlled by serum components. *Cancer Res* 1977; **37**: 3037-41.
- 25 Benedetti A, Birarelli A M, Brunelli E, et al. Effect of chronic ethanol abuse on the physico-chemical properties of erythrocyte membranes in man. *Pharmacol Res Commun* 1986; **18**: 1003-14.
- 26 Baba Y, Kai M, Kamadett T, Setayama S, Otsuji S. Higher levels of erythrocyte membrane microviscosity in diabetes. *Diabetes* 1979; **28**: 1138-40.
- 27 Shinitzky M, Inbar M. Difference in microviscosity induced by different cholesterol levels in the surface membrane lipid layer of normal lymphocytes and malignant lymphoma cells. *J Mol Biol* 1974; **85**: 603-15.
- 28 Shinitzky M, Barenholz Y. Fluidity parameters of lipid regions determined by fluorescence polarization. *Biochim Biophys Acta* 1978; **515**: 367-94.
- 29 Westergren A. The technique of the red cell sedimentation reaction. *American Review of Tuberculosis* 1926; **14**: 94-101.
- 30 Kohn J. Cellular acetate electrophoresis. In: Smith I, ed. *Chromatographic and electrophoretic techniques*. Vol 2. London: Heinemann, 1968: 84-9.
- 31 Singer J M, Plotz C M, Elster S K. The latex fixation test. III. Agglutination test for C reactive protein and comparison with the capillary precipitin method. *Am J Clin Pathol* 1957; **28**: 611-7.
- 32 Mancini G, Carbonara A O, Heremans J F. Immunochemical quantitation of antigens by single radioimmunoassay. *Immunochemistry* 1965; **2**: 235-8.
- 33 Simplicio P. An improved method of determination of SH group concentration and reactivity in plasma of patients with various clinical disorders and in rats after administration of indomethacin or carbon tetrachloride. *Pharmacol Res Commun* 1983; **15**: 805-14.
- 34 Harkness J. The viscosity of human blood plasma—its measurement in health and disease. *Biorheology* 1971; **8**: 171-93.
- 35 Spiegelberg H, Dainer P. Fc receptors for IgG, IgM and IgE on human leukemic lymphocytes. *Clin Exp Immunol* 1979; **35**: 286-95.
- 36 Fox R I, Thompson L F, Huddleston J R. Ty cells express T-cell associated antigens. *J Immunol* 1981; **126**: 2062-3.
- 37 Ferretti G, Falcioni A M, Curatola G, Valentini B, Saccucci F. Duchenne muscular dystrophy: alterations in lymphocyte fluidity. *Boll Soc Ital Biol Sper* 1983; **59**: 1035-40.
- 38 Zubenko G S, Cohen B M, Growdon J, Corkin S. Cell membrane abnormality in Alzheimer's disease. *Lancet* 1984; **ii**: 235.
- 39 Inbar M, Yuli I, Raz A. Contact-mediated changes in the fluidity of membrane lipids in normal and malignant transformed mammalian fibroblast. *Exp Cell Res* 1977; **105**: 325-35.
- 40 Fantone J C, Ward P A. Polymorphonuclear leukocyte-mediated cell and tissue injury: oxygen metabolites and their relations to human disease. *Hum Pathol* 1985; **16**: 973-8.
- 41 Clavel J P, Emerit J, Thuillier A. Lipidperoxydation et radicaux libres. *Pathol Biol (Paris)* 1985; **33**: 61-9.
- 42 Stossel T P, Mason R J, Smith A L. Lipid peroxidation by human blood phagocytes. *J Clin Invest* 1974; **54**: 638-45.
- 43 Dobretsov G E, Borschevskaya T A, Petrov V A, Vladimirov Y A. The increase of phospholipid bilayer rigidity after lipid peroxidation. *FEBS Lett* 1977; **84**: 125-8.
- 44 Lunec J, Halloran S P, White A G, Dormandy T L. Free-radical oxidation (peroxidation) products in serum and synovial fluid in rheumatoid arthritis. *J Rheumatol* 1981; **8**: 233-45.
- 45 Gutteridge J M C, Hill C, Blake D R. Copper stimulated phospholipid membrane peroxidation: antioxidant activity of serum and synovial fluid from patients with rheumatoid arthritis. *Clin Chim Acta* 1984; **139**: 85-90.