

Multifunctional Polyplex Micelles for Efficient MicroRNA Delivery and Accelerated Osteogenesis

Qian Li, Zhiai Hu, Xin Rong, and Xiaohua Liu*

*Department of Biomedical Sciences,
Texas A&M University College of Dentistry, Dallas, TX 75246, USA*

*Correspondence to:

Xiaohua Liu, PhD

Associate Professor

Department of Biomedical Sciences

Texas A&M University College of Dentistry

3302 Gaston Ave., Dallas, TX 75246

Phone: 214-370-7007

Fax: 214-874-4538

Email: xliu1@tamu.edu

Table S1. Primer sequences for qRT-PCR.

Gene	Primer sequence
miR-218	UUGUGCUUGAUCUAACCAUGU
ALP	F: ACTGCGCTCCTTAGGGCT R: GGCAGCGTCAGATGTTAATTG
RUNX2	F: AGTGCTCTAACCACAGTCCATGCA R: TACAAACCATACCCAAGTACCTGTTT
Col1	F: GATGGTGCATCACTCAATGG R: TATGCGCACTTCAAAAACCA
OCN	F: GGTAGTGAACAGACTCCGGC R: CAAGCAGGGTTAAGCTCACA
DKK2	F: TCAGTCAGCCAACCGATCTG R: TCTCTGTGGCATCGTTTCTTTT
SFRP2	F: CCCCTGTCTGTCTCGACGA R: CTTCACACACCTTGGGAGCTT
SOST	F: AGCCTTCAGGAATGATGCCAC R: CTTTGGCGTCATAGGGATGGT
PCNA	F: TTTCACAAAAGCCACTCCACTG R: CTTTAAGTGTCCCATGTCAGCAAT
GAPDH	F: TTGATGGCAACAATCTCCAC R: CGTCCCGTAGACAAAATGGT




Fig. S1 FT-IR spectra of synthesized PLGA copolymers and PLGA-PEI triblock copolymer.




Fig. S2 (a) miR-218 condensation ability of various complexes. (b) Agarose gel electrophoresis for the assessment of miRNA stabilization against RNase degradation after incubation for 4 h. Line 1: miR-218 only as negative control, Line 2: PLGA-PEI/miR-218, Line 3: PPP/miR-218, Line 4: PPP-RG11/miR-218, Line 5: PPP-RG12/miR-218.

Line 4: PPP-RGI1/miR-218, and Line 5: PPP-RGI2/miR-218. (c, d) Sizes and zeta potentials of micelles/miR-218 complexes at various w/w ratios.




Fig. S3 Mean fluorescence intensity of PPP-RGI2/miR-218 treated group after cultured for 4 h and 24 h.




Fig. S4 Relative cell viability of BMSCs at different miR-218 concentration after treated for three days. DNA concentration ranging from 0.4 to 4.5 µg/mL at w/w ratio of 5.