

Supplement Table 1. Univariable analyses for severe erectile dysfunction in patients undergoing hemodialysis

Variable	Factor	p-value	Odds ratio	95% confidence interval
Age	Continuous	0.095	1.054	0.991–1.120
Body mass index	Continuous	0.721	0.983	0.894–1.081
Hypertension	Positive	0.516	0.472	0.049–4.546
Dyslipidemia	Positive	0.843	0.888	0.274–2.879
Diabetes mellitus	Positive	0.703	0.802	0.258–2.490
Duration of dialysis	Continuous	0.098	1.010	0.998–1.023
β-blockers	Taking	0.706	0.808	0.267–2.445
Calcium-blockers	Taking	0.691	0.791	0.249–2.514
Serum iPTH level	Continuous	0.221	0.997	0.992–1.002
Serum testosterone level	Continuous	0.826	1.000	0.996–1.003
Smoking status	Current	0.636	1.422	0.331–6.118
Depression	Positive	>0.999	1.000	0.085–11.78
Cardio-ankle vascular index	Continuous	0.652	1.073	0.791–1.455
Ankle-brachial index	Continuous	0.098	0.068	0.003–1.641
Aortic calcification index	Continuous	0.050	1.019	1.000–1.039

iPTH: intact parathyroid hormone.