

1 **SUPPLEMENTARY INFORMATION**

2 **mRNA vaccines and hybrid immunity use different B cell germlines against Omicron BA.4 and BA.5**

3 Emanuele Andreano¹, Ida Paciello¹, Giulio Pierleoni², Giuseppe Maccari³, Giada Antonelli¹, Valentina
4 Abbiento¹, Piero Pileri¹, Linda Benincasa², Ginevra Giglioli², Giulia Piccini⁴, Concetta De Santi¹, Claudia Sala¹,
5 Duccio Medini³, Emanuele Montomoli^{2,4,5}, Piet Maes⁶, Rino Rappuoli^{7,8,*}

6

7 ¹Monoclonal Antibody Discovery (MAD) Lab, Fondazione Toscana Life Sciences, Siena, Italy

8 ²VisMederi Research S.r.l., Siena, Italy

9 ³Data Science for Health (DaScH) Lab, Fondazione Toscana Life Sciences, Siena, Italy

10 ⁴VisMederi S.r.l, Siena, Italy

11 ⁵Department of Molecular and Developmental Medicine, University of Siena, Siena, Italy


12 ⁶KU Leuven, Rega Institute, Department of Microbiology, Immunology and Transplantation, Laboratory of
13 Clinical and Epidemiological Virology, Leuven, Belgium

14 ⁷Department of Biotechnology, Chemistry and Pharmacy, University of Siena, Siena, Italy

15 ⁸Fondazione Biotecnopolo di Siena, Siena, Italy


16 *Corresponding author: Rino Rappuoli rino.rappuoli@biotecnopolo.it

17 **SUPPLEMENTARY FIGURES**



18

19 **Supplementary Fig. 1. Distribution of omicron mutations on SARS-CoV-2 S protein. a**, the heatmap shows
20 the SARS-CoV-2 S protein mutations in the NTD (cyan), RBD (dark gray) and S2 domain (light gray) of omicron
21 BA.1, BA.2, BA.4 and BA.5. **b**, Representation of the RBD with all omicron mutations highlighted in red.



22

23 **Supplementary Fig. 2. Distribution and neutralization potency of RBD and NTD-targeting nAbs against**
 24 **omicron BA.4 and BA.5. a-b**, dot charts compare the distribution of nAbs isolated from subjects belonging
 25 to the SN3 and SP2 cohorts against omicron BA.4 (a) and BA.5 (b) omicron variants. The number, percentage
 26 and GM-IC₁₀₀ (black lines and colored bars) of nAbs are denoted on each graph. **c-d**, table summarizing the
 27 GM-IC₁₀₀ of Class 1/2, Class 3, Class 4 and NTD nAbs against BA.4 (c) and BA.5 (d). Source data are provided
 28 as a Source Data file.

29 **SUPPLEMENTARY TABLES**30 **Supplementary Table 1. Numbers and frequency of nAbs in the SN3 cohort.**

Subject	Vaccine	Wuhan – Neutralizing antibodies	Omicron BA.1 – Neutralizing antibodies (%)	Omicron BA.2 – Neutralizing antibodies (%)	Omicron BA.4 – Neutralizing antibodies (%)	Omicron BA.5 – Neutralizing antibodies (%)
VAC-001	BNT162b2	106	22 (20.8)	24 (22.6)	11 (10.4)	9 (8.5)
VAC-002	BNT162b2	57	15 (26.3)	17 (29.8)	9 (15.8)	9 (15.8)
VAC-008	BNT162b2	36	7 (19.4)	9 (25.0)	4 (11.1)	5 (13.9)
VAC-010	mRNA-1273	7	1 (14.3)	2 (28.6)	0 (0.0)	0 (0.0)
Total		206	45 (21.8)	52 (25.2)	24 (11.6)	23 (11.2)

31

32 **Supplementary Table 2. Frequency of predominant germlines against omicron variants.**

Seronegative 2 nd doses (SN2)					
Germlines	Wuhan Virus (%)	Omicron (BA.1) (%)	Omicron (BA.2) (%)	Omicron (BA.4) (%)	Omicron (BA.5) (%)
IGHV1-69;IGHJ4-1	2 (4.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
IGHV3-30;IGHJ6-1	2 (4.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
IGHV3-53;IGHJ6-1	6 (13.0)	0 (0.0)	1 (25.0)	0 (0.0)	0 (0.0)
IGHV3-66;IGHJ4-1	3 (6.5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Tot. nAbs	46	1	4	0	0
Tot. germline specific nAbs	13 (28.3)	0 (0.0)	1 (25.0)	0 (0.0)	0 (0.0)
Seronegative 3 rd doses (SN3)					
Germlines	Wuhan Virus (%)	Omicron (BA.1) (%)	Omicron (BA.2) (%)	Omicron (BA.4) (%)	Omicron (BA.5) (%)
IGHV1-58;IGHJ3-1	12 (6.8)	5 (13.9)	4 (9.5)	1 (4.5)	0 (0.0)
IGHV1-69;IGHJ3-1	10 (5.7)	4 (11.1)	5 (11.9)	1 (4.5)	3 (13.6)
IGHV1-69;IGHJ4-1	16 (9.1)	9 (25.0)	9 (21.4)	4 (18.2)	4 (18.2)
IGHV3-66;IGHJ4-1	9 (5.1)	0 (0.0)	1 (2.4)	2 (9.1)	3 (13.6)
IGHV3-66;IGHJ6-1	10 (5.7)	3 (8.3)	4 (9.5)	4 (18.2)	3 (13.6)
Tot. nAbs	176	36	42	22	22
Tot. germline specific nAbs	57 (32.4)	21 (58.3)	23 (54.8)	12 (54.6)	13 (59.1)
Seropositive 2 nd doses (SP2)					
Germlines	Wuhan Virus (%)	Omicron (BA.1) (%)	Omicron (BA.2) (%)	Omicron (BA.4) (%)	Omicron (BA.5) (%)
IGHV1-24;IGHJ6-1	5 (2.4)	0 (0.0)	3 (6.1)	3 (10.0)	3 (10.3)
IGHV1-58;IGHJ3-1	10 (4.8)	7 (18.9)	5 (10.2)	0 (0.0)	0 (0.0)
IGHV2-5;IGHJ4-1	9 (4.3)	2 (5.4)	6 (12.2)	6 (20.0)	6 (20.7)
Tot. nAbs	208	37	49	30	29
Tot. germline specific nAbs	24 (11.5)	9 (24.3)	14 (28.6)	9 (30.0)	9 (31.0)

33

34 **Supplementary Table 3. Neutralization potency of predominant germlines against omicron variants.**

Seronegative 2 nd doses (SN2)					
Germlines	Wuhan Virus GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.1) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.2) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.4) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.5) GM-IC ₁₀₀ ng ml ⁻¹
IGHV1-69;IGHJ4-1	191.63	N/A	N/A	N/A	N/A
IGHV3-30;IGHJ6-1	382.71	N/A	N/A	N/A	N/A
IGHV3-53;IGHJ6-1	79.58	N/A	25.60	N/A	N/A
IGHV3-66;IGHJ4-1	66.42	N/A	N/A	N/A	N/A
Seronegative 3 rd doses (SN3)					
Germlines	Wuhan Virus GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.1) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.2) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.4) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.5) GM-IC ₁₀₀ ng ml ⁻¹
IGHV1-58;IGHJ3-1	93.79	157.36	192.01	4261.36	N/A
IGHV1-69;IGHJ3-1	50.21	155.79	124.13	39.67	616.76
IGHV1-69;IGHJ4-1	212.38	987.84	1069.88	1318.53	1927.94
IGHV3-66;IGHJ4-1	327.61	N/A	66.17	253.58	578.35
IGHV3-66;IGHJ6-1	272.30	295.43	223.63	494.97	453.70
Seropositive 2 nd doses (SP2)					
Germlines	Wuhan Virus GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.1) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.2) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.4) GM-IC ₁₀₀ ng ml ⁻¹	Omicron (BA.5) GM-IC ₁₀₀ ng ml ⁻¹
IGHV1-24;IGHJ6-1	314.04	N/A	130.20	413.36	463.99
IGHV1-58;IGHJ3-1	52.19	852.49	982.36	N/A	N/A
IGHV2-5;IGHJ4-1	28.10	379.68	197.85	264.09	166.37

35