

**Supplementary Table 1:** Comparison of patients' characteristics in the three cohorts (CHU Lille, CHU Liege, UZ Antwerp) included in the study

	<b>CHU Lille</b> (101 patients)		<b>CHU Liege</b> (68 patients)		<b>UZ Antwerp</b> (61 patients)	
	AEC < 220/ $\mu$ L	AEC $\geq$ 220/ $\mu$ L	AEC < 220/ $\mu$ L	AEC $\geq$ 220/ $\mu$ L	AEC < 220/ $\mu$ L	AEC $\geq$ 220/ $\mu$ L
	<b>83</b> (82.2%)	<b>18</b> (17.8%)	<b>46</b> (67.65%)	<b>22</b> (32.35%)	<b>40</b> (65.6%)	<b>21</b> (34.4%)
<b>Sex, n</b>						
Male	56	16	41	18	28	20
Female	27	2	5	4	12	2
<b>Histological subtype, n</b>						
Epithelioid	71	14	34	18	39	16
Non-epithelioid	10	4	8	4	1	5
Sarcomatoid	5	2	1	0	0	3
Biphasic	4	2	5	3	1	2
Desmoplastic	1	0	2	1	0	0
Unknown	2	0	4	0	0	0
<b>ECOG performance status, n</b>						
0	23	6	5	2	N/A	N/A
1	47	10	37	16	N/A	N/A
2	5	2	2	0	N/A	N/A
Unknown	0	0	1	4	41	20

n = number of patients