

Supplementary Table 1. Inclusion and Exclusion Criteria^a

Inclusion criteria	Exclusion criteria
<ul style="list-style-type: none">- Clinical diagnosis of PD according to the UK PD Society Brain Bank criteria¹⁷- Modified Hoehn & Yahr Stage I to IV in the ON state- Age between 40 and 90 years- Mini Mental State Examination (MMSE) score of ≥ 21 or > 16 on the 26-item MMSE screening- Stable PD medication during the previous month- Self-reported FOG severity of at least one FOG episode per day, irrespective of FOG occurring ON- or OFF-medication- Observed FOG during the provoking protocol in baseline testing	<ul style="list-style-type: none">- Participation in another clinical study or usage of cueing devices- A fall frequency of more than once a day- Acute musculoskeletal, cardiovascular, other neurological or any other medical condition affecting gait- Occurrence of any of the following within 3 months prior to informed consent: orthopedic surgery of the lower extremity, myocardial infarction, hospitalization for unstable angina, coronary artery bypass graft, percutaneous coronary intervention, implantation of a cardiac resynchronization therapy device, implantation of deep brain stimulation- Substance abuse, major depressive disorder or clinical apathy that affects daily walking activity or may interfere with the patient's compliance- Inability to walk without a rollator indoors- Use of a duodopa pump or apomorphine injections, jeopardizing OFF-medication assessments

^a FOG: Freezing of Gait; PD: Parkinson disease.

¹⁷Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992;**55(3)**:181–4.