

Supplementary material has been published as submitted. It has not been copyedited, typeset or checked for scientific content by Acta Dermato-Venereologica

**Table SI. Baseline screening recommendation and their periodicity.**

<b>Risk factor</b>	<b>Baseline screening</b>	<b>Periodicity</b>
<b>Tobacco use</b>	All adults should be inquired about tobacco consumption <sup>21-24</sup>	<ul style="list-style-type: none"> <li>At every healthcare visit <sup>23,24</sup></li> </ul>
<b>Obesity</b>	BMI calculation and waist circumference measurement <sup>21-24</sup>	<ul style="list-style-type: none"> <li>At least annually <sup>23</sup></li> </ul>
<b>HTN</b>	Blood pressure measurement in all adults (18 years or older) <sup>25,26</sup>	<ul style="list-style-type: none"> <li>If &lt;120/80, every 5 years <sup>25,26</sup></li> <li>If 120-130/80-85, every 3 years <sup>25,26</sup></li> <li>If &gt;130/85, at least annually <sup>25,26</sup></li> </ul>
<b>Dyslipidemia</b>	Fasting or nonfasting lipid profile <sup>23,24,29</sup> <ul style="list-style-type: none"> <li>Men &gt; 40 years old and women &gt; 50 years of age or post-menopausal <sup>27</sup></li> <li>Adults who are 20 years of age or older and not on lipid-lowering therapy <sup>28</sup></li> </ul>	<ul style="list-style-type: none"> <li>Interval for subsequent screening not provided <sup>24,27</sup></li> <li>Adults 20 to 39 years of age, at least every 4-6 years; For adults 40 to 75 years of age, assess traditional CVRFs routinely <sup>23</sup></li> <li>Annual screening in middle-aged adults without CVRFs (men aged 45-65 years, women aged 55-65 years) <sup>29</sup></li> </ul>
<b>DM</b>	Fasting plasma glucose, HbA1c <sup>22,30</sup> <ul style="list-style-type: none"> <li>Testing should begin at 45 years <sup>30</sup></li> </ul>	<ul style="list-style-type: none"> <li>If results are normal, testing should be repeated at least every 3 years <sup>30</sup></li> </ul>
<b>10-year risk of CVD calculation</b>	<ul style="list-style-type: none"> <li>SCORE2 in apparent healthy men &gt; 40 years old and in women &gt; 50 years of age or post-menopausal; SCORE2-OP in apparent healthy people <math>\geq</math> 70 years <sup>24</sup></li> <li>Pooled cohort equations for adults 40 to 75 years of age <sup>23</sup></li> </ul>	<ul style="list-style-type: none"> <li>Every 4 to 6 years <sup>23,24</sup></li> </ul>

The clinical practice guideline on which each recommendation is based is mentioned in superscript at the end of each statement. BMI: Body Mass Index; CVD: Cardiovascular disease; CVRF: Cardiovascular risk factors; DM: Diabetes mellitus; ESC: European Society of Cardiology; HTN: Hypertension; \*\* In the survey question, considering the questionnaire was distributed in a European country, the discrepancy in the age at screening beginning was resolved in favor of ESC guidelines.

**Table SII. Baseline characteristics of the dermatologists surveyed and AEDV members**

	<b>Dermatologists surveyed</b>	<b>AEDV members</b>
<b>Female, %</b>	63.1	48.3
<b>Fellows, %</b>	8.7	12.0
<b>Private activity*, %</b>	58.3	82.2

AEDV: Spanish Society of Dermatology and Venereology (Academia Española de Dermatología y Venereología). \*With or without public activity.

**Table SIII. Baseline characteristics of the rheumatologists surveyed and SER members**

	<b>Rheumatologists surveyed</b>	<b>Rheumatologists from the SER</b>
<b>Female, %</b>	67.0	61.0
<b>Fellows, %</b>	3.2	15.9
<b>Private activity*, %</b>	22.3	29.4

SER: Spanish Society of Rheumatology (Sociedad Española de Reumatología). \*With or without public activity.

**Table SIV. Statin prescription multivariable analysis**

	Univariable			Multivariable		
	OR	95% CI	p-value	OR	95% CI	p-value
<b>Specialty</b>						
Dermatologists vs Rheumatologists	0.24	0.12-0.50	<b>&lt;0.001</b>	0.04	0.01-0.16	<b>&lt;0.001</b>
<b>Sex</b>						
Men vs women	1.08	0.54-2.16	0.82	—	—	—
<b>Practice setting</b>			0.702			—
Private vs Public	0.81	0.30-2.20	0.678	—	—	—
Private + public vs only public	0.73	0.33-1.59	0.420	—	—	—
<b>Years since completing residency training</b>			0.012			0.003
1-10 years vs resident	2.20	0.25-19.24	0.476	1.82	0.11-29.90	0.674
10-20 years vs resident	3.88	0.45-33.34	0.216	1.98	0.12-31.89	0.629
20-30 years vs resident	2.48	0.28-21.70	0.413	1.49	0.08-26.46	0.785
>30 years vs resident	9.78	1.13-84.37	<b>0.038</b>	13.45	0.80-226.62	0.071
<b>Patients with psoriasis treated per month</b>			0.369			0.433
5-20 vs <5	1.10	0.21-5.57	0.920	0.44	0.04-4.88	0.500
>20 vs <5	1.75	0.36-8.59	0.489	0.80	0.07-9.21	0.859
<b>Specialized psoriasis outpatient clinic</b>				—	—	—
No	—	—		—	—	—
Yes	1.76	0.86-3.59	0.122	7.93	2.06-30.59	<b>0.003</b>
<b>Research projects related to psoriasis</b>						
No	—	—		—	—	
Yes	1.62	0.84-3.12	0.151	1.28	0.50-3.31	0.609
<b>University teaching including psoriasis lectures</b>						
No	—	—		—	—	
Yes	3.13	1.51-6.48	<b>0.002</b>	2.85	1.05-7.75	<b>0.041</b>

CI: Confidence interval; OR: Odds Ratio.