| Antibody | Clone | Supplier/Info | Catalogue # | Lot # | Validation |
|---|---|---|---|---|---|
| anti-CD4-APC-Cy7 | GK1.5 | Biolegend | 100414 | B326190 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Dialynas DP, et al. 1983. J. Immunol. 131:2445. (Block, IP)<br>• Dialynas DP, et al. 1983. Immunol. Rev. 74:29. (IP, Deplete)<br>• Wu L, et al. 1991. J. Exp. Med. 174:1617. (Costim) |
| anti-CD8-APC-R700 | 53-6.7 | BD Biosciences | 564983 | 0080396 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Bierer BE, Sleckman BP, Ratnofsky SE, Burakoff SJ. The biologic roles of CD2, CD4, and CD8 in T-cell activation. Annu Rev Immunol. 1989; 7:579-599. (Biology).<br>• Fujiura Y, Kawaguchi M, Kondo Y, et al. Development of CD8 alpha alpha+ intestinal intraepithelial T cells in beta 2-microglobulin- and/or TAP1-deficient mice. J Immunol. 1996; 156(8):2710-2715. (Clone-specific: Flow cytometry).<br>• Hathcock KS. T cell depletion by cytotoxic elimination. Curr Protoc Immunol. 1991; 1:3.4.1-3.4.3. (Clone-specific: Cell separation, Depletion, Flow cytometry). |
| anti-CD44-BV785 | IM7 | Biolegend | 103041/103059 | B346798 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Trowbridge IS, et al. 1982. Immunogenetics 15:299. (ICFC, IP, CMCD)<br>• Katoh S, et al. 1994. J. Immunol. 153:3440. (ELISA)<br>• Budd RC, et al. 1987. J. Immunol. 138:3120. (IP) |
| anti-CD44-FITC | IM7 | BD Pharmigen | 553133 | 3331648 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript. |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | • Bendelac A. Mouse NK1+ T cells. Curr Opin Immunol. 1995; 7(3):367-374. (Biology).<br>• Brocke S, Piercy C, Steinman L, Weissman IL, Veromaa T. Antibodies to CD44 and integrin alpha4, but not L-selectin, prevent central nervous system inflammation and experimental encephalomyelitis by blocking secondary leukocyte recruitment. Proc Natl Acad Sci U S A. 1999; 96(12):6896-6901. (Clone-specific).<br>• Budd RC, Cerottini JC, Horvath C, et al. Distinction of virgin and memory T lymphocytes. Stable acquisition of the Pgp-1 glycoprotein concomitant with antigenic stimulation. J Immunol. 1987; 138(10):3120-3129. (Biology). |
| anti-CD62L BV711 | MEL-14 | Biolegend | 104445 | B305757 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Zallatin WM, et al. 1983. Nature 304:30. (IP, Block)<br>• Siegelman MH, et al. 1990. Cell 61:611. (IP, Block)<br>• Reichert RA, *et al.* 1986. *J. Immunol.* 136:3535. (IHC, FC) |
| anti-PD-1-BV605 | J43 | BD Biosciences | 563059 | 1152563 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Agata Y, Kawasaki A, Nishimura H, et al. Expression of the PD-1 antigen on the surface of stimulated mouse T and B lymphocytes. Int Immunol. 1996 May; 8(5):765-772. (Immunogen: Flow cytometry, Immunoprecipitation).<br>• Ansari MJ, Salama AD, Chitnis T, et al. The programmed death-1 (PD-1) pathway regulates autoimmune diabetes in nonobese diabetic (NOD) mice. J Exp Med. 2003 July; 198(1):63-69. (Clone-specific: Blocking).<br>• Carreno BM, Collins M. The B7 family of ligands and its receptors: New pathways for costimulation and |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | inhibition of immune responses. Annu Rev Immunol. 2002; 20:29-53. (Biology). |
| anti-CXCR5-BV421 | L138D7 | Biolegend | 145511/145512 | B308119 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Wrzner K, et al. 2021. EBioMedicine. 63:103197.<br>• Wilfahrt D, et al. 2021. Elife. 10:.<br>• Hill DL, et al. 2021. Elife. 10:. |
| anti-GL-7-FITC | GL7 | BD Pharmigen | 562080 | 2270851 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Han S, Dillon SR, Zheng B, Shimoda M, Schlissel MS, Kelsoe G. V(D)J recombinase activity in a subset of germinal center B lymphocytes. Science. 1997; 278(5336):301-305. (Biology).<br>• Han S, Zheng B, Schatz DG, Spanopoulou E, Kelsoe G. Neoteny in lymphocytes: Rag1 and Rag2 expression in germinal center B cells. Science. 1996; 274(5295):2094-2097. (Biology).<br>• Han S, Zheng B, Takahashi Y, Kelsoe G. Distinctive characteristics of germinal center B cells. Semin Immunol. 1997; 9(4):255-260. (Biology). |
| anti-CD38-Pe-Cy7 | 90 | Biolegend | 102718 | B316990 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Hudson WH, et al. 2020. Immunity. 51(6):1043-1058.e4..<br>• Degn SE et al. 2017. Cell. 170(5):913-926<br>• Bauer KM, et al. 2022. JCI Insight. 7:. |

| | | | | | |
|---|---|---|---|---|---|
| anti-IgD-APC-Cy7 | 11-26c.2a | Biolegend | 405715/405716 | B310012 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Nitschke L, et al. 1993. P. Natl. Acad. Sci. USA 90:1887. (FC)<br>• Weih D, et al. 2001. J. Immunol. 167:1909. (IHC)<br>• Koni PA, et al. 2001. J. Exp. Med. 193:741. (IHC) |
| anti-CD19-APC-Red | 1D3 | BD Biosciences | 565473 | 104814 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Engel P, Zhou LJ, Ord DC, Sato S, Koller B, Tedder TF. Abnormal B lymphocyte development, activation, and differentiation in mice that lack or overexpress the CD19 signal transduction molecule. Immunity. 1995; 3(1):39-50. (Biology).<br>• Fearon DT. The CD19-CR2-TAPA-1 complex, CD45 and signaling by the antigen receptor of B lymphocytes. Curr Opin Immunol. 1993; 5(3):341-348. (Biology). Gommerman JL, Oh DY, Zhou X, et al. A role for CD21/CD35 and CD19 in responses to acute septic peritonitis: a potential mechanism for mast cell activation. J Immunol. 2000; 165(12):6915-6921. (Biology). |
| anti-IgM-BV786 | II/41 | BD Biosciences | 743328 | 2291206 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Hennet T, Chui D, Paulson JC, Marth JD. Immune regulation by the ST6Gal sialyltransferase.. Proc Natl Acad Sci USA. 1998; 95(8):4504-9. (Clone-specific: Flow cytometry).<br>• Laszlo G, Hathcock KS, Dickler HB, Hodes RJ. Characterization of a novel cell-surface molecule |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | expressed on subpopulations of activated T and B cells. J Immunol. 1993; 150(12):5252-5262. (Clone-specific). |
| anti-IgG1-BV510 | RMG1-1 | Biolegend | 406621 | B343134 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Rucci F, et al. 2010. P. Natl. Acad. Sci. USA 107:3024. (FC)<br>• Zhang JG, et al. 2011. Int. Immunopharmacol. 11:1685 |
| anti-IgG2a/b-BV510 | R2-40 | BD Biosciences | 744293 | 1148164 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Benson MJ, Dillon SR, Castigli E, et al. Cutting edge: the dependence of plasma cells and independence of memory B cells on BAFF and APRIL.. J Immunol. 2008; 180(6):3655-9. (Clone-specific: Flow cytometry).<br>• Chen CL, Lee CT, Liu YC, Wang JY, Lei HY, Yu CK. House dust mite Dermatophagoides farinae augments proinflammatory mediator productions and accessory function of alveolar macrophages: implications for allergic sensitization and inflammation.. J Immunol. 2003; 170(1):528-36. (Clone-specific: ELISA) |
| anti-IgG3-BV510 | R40-82 | BD Biosciences | 744134 | 1131187 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Duong BH, Ota T, Aït-Azzouzene D, et al. Peripheral B cell tolerance and function in transgenic mice expressing an IgD superantigen.. J Immunol. 2010; 184(8):4143-58. (Clone-specific: ELISA, Flow cytometry). |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | • Kuzin II, Snyder JE, Ugine GD, et al. Tetracyclines inhibit activated B cell function.. Int Immunol. 2001; 13(7):921-31. (Clone-specific: Flow cytometry) |
| anti-TACI-BV421 | 8F10 | BD Biosciences | 742840 | 2034081 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Ng LG, Sutherland AP, Newton R, et al. B cell-activating factor belonging to the TNF family (BAFF)-R is the principal BAFF receptor facilitating BAFF costimulation of circulating T and B cells.. J Immunol. 2004; 173(2):807-17. (Immunogen: Flow cytometry).<br>• Siegel RM, Lenardo MJ.. To B or not to B: TNF family signaling in lymphocytes. Nat Immunol. 2001; 2(7):577-578. (Biology).<br>• Yan M, Wang H, Chan B, et al. Activation and accumulation of B cells in TACI-deficient mice.. Nat Immunol. 2001; 2(7):638-43. (Biology). |
| anti-CD19-BV510 | 6D5 | Biolegend | 115546 | B264815 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Shoham T, et al. 2003. J. Immunol. 171:4062. (FC)<br>• Goodyear CS, et al. 2004. J. Immunol. 172:2870. (FC)<br>• Kamimura D, et al. 2006. J. Immunol. 177:306. (FC) |
| anti-CD45R/B220-BUV395 | RA3-6B2 | BD Biosciences | 563793 | 9028674 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Allman DM, Ferguson SE, Cancro MP. Peripheral B cell maturation. I. Immature peripheral B cells in adults are heat-stable antigenhi and exhibit unique signaling characteristics. J Immunol. 1992; 149(8):2533-2540. (Clone-specific: Flow cytometry). |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | • Asensi V, Kimeno K, Kawamura I, Sakumoto M, Nomoto K. Treatment of autoimmune MRL/lpr mice with anti-B220 monoclonal antibody reduces the level of anti-DNA antibodies and lymphadenopathies. Immunology. 1989; 68(2):204-208. (Clone-specific: Flow cytometry, In vivo exacerbation).<br>• Ballas ZK, Rasmussen W. Lymphokine-activated killer cells. VII. IL-4 induces an NK1.1+CD8 alpha+beta- TCR-alpha beta B220+ lymphokine-activated killer subset. J Immunol. 1993; 150(1):17-30. (Clone-specific: Flow cytometry, Fluorescence activated cell sorting, Immunofluorescence). |
| anti-GL7-eFluor450 | GL7 | eBiosience | 48-5902-82 | 2062737 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Khan AR, Hams E, Floudas A, Sparwasser T, Weaver CT, Fallon PG. PD-L1hi B cells are critical regulators of humoral immunity. Nat Commun. 2015 Jan 22;6:5997. doi: 10.1038/ncomms6997. PMID: 25609381.<br>• Khan AR, Hams E, Floudas A, Sparwasser T, Weaver CT, Fallon PG. PD-L1hi B cells are critical regulators of humoral immunity. Nat Commun. 2015 Jan 22;6:5997. doi: 10.1038/ncomms6997. PMID: 25609381. |
| anti-CD95-PE-Cy7 | Jo2 | BD Biosciences | 557653 | 9039631 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Hiromatsu K, Aoki Y, Makino M, et al. Increased Fas antigen expression in murine retrovirus-induced immunodeficiency syndrome, MAIDS. Eur J Immunol. 1994; 24(10):2446-2451. (Clone-specific: Flow cytometry, Functional assay). |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | • Kagi D, Vignaux F, Ledermann B, et al. Fas and perforin pathways as major mechanisms of T cell-mediated cytotoxicity. Science. 1994; 265(5171):528-530. (Clone-specific: Flow cytometry, Functional assay).<br>• Ogasawara J, Suda T, Nagata S. Selective apoptosis of CD4+CD8+ thymocytes by the anti-Fas antibody. J Exp Med. 1995; 181(2):485-491. (Clone-specific: Flow cytometry, Functional assay). |
| anti-CD8-APC-Cy7 | 53-6.7 | BD Bioscience | 557654 | 8319924 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Beavis AJ, Pennline KJ. Allo-7: a new fluorescent tandem dye for use in flow cytometry. Cytometry. 1996; 24(4):390-395. (Biology).<br>• Bierer BE, Sleckman BP, Ratnofsky SE, Burakoff SJ. The biologic roles of CD2, CD4, and CD8 in T-cell activation. Annu Rev Immunol. 1989; 7:579-599. (Biology).<br>• Fujiura Y, Kawaguchi M, Kondo Y, et al. Development of CD8 alpha alpha+ intestinal intraepithelial T cells in beta 2-microglobulin- and/or TAP1-deficient mice. J Immunol. 1996; 156(8):2710-2715. (Biology). |
| anti-CD4-APCCy7 | GK1.5 | BD Bioscience | 552051 | 8333906 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Beavis AJ, Pennline KJ. Allo-7: a new fluorescent tandem dye for use in flow cytometry. Cytometry. 1996; 24(4):390-395. (Methodology: Flow cytometry).<br>• Dialynas DP, Quan ZS, Wall KA, et al. Characterization of the murine T cell surface molecule, designated L3T4, identified by monoclonal antibody GK1.5: similarity of L3T4 to the human Leu-3/T4 molecule. J Immunol. |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | 1983; 131(5):2445-2451. (Immunogen: Blocking, Depletion, Immunoprecipitation).<br>• Dialynas DP, Wilde DB, Marrack P, et al. Characterization of the murine antigenic determinant, designated L3T4a, recognized by monoclonal antibody GK1.5: expression of L3T4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. Immunol Rev. 1983; 74:29-56. (Clone-specific: Blocking, Depletion, Immunoprecipitation).<br>• |
| anti-Ly-6G-APC-Cy7 | RB6-8C5 | BD Pharmingen | 557661 | 8318838 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Beavis AJ, Pennline KJ. Allo-7: a new fluorescent tandem dye for use in flow cytometry. Cytometry. 1996; 24(4):390-395. (Methodology).<br>• Conlan JW, North RJ. Neutrophils are essential for early anti-Listeria defense in the liver, but not in the spleen or peritoneal cavity, as revealed by a granulocyte-depleting monoclonal antibody. J Exp Med. 1994; 179(1):259-268. (Clone-specific: Depletion).<br>• Fleming TJ, Fleming ML, Malek TR. Selective expression of Ly-6G on myeloid lineage cells in mouse bone marrow. RB6-8C5 mAb to granulocyte-differentiation antigen (Gr-1) detects members of the Ly-6 family. J Immunol. 1993; 151(5):2399-2408. (Clone-specific: Immunoprecipitation, Inhibition). |
| anti-F4/80-APC-Cy7 | BM8 | Biolegend | 123118 | B274179 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Schaller E, et al. 2002. Mol. Cell. Biol. 22:8035. (IHC) |

| | | | | | |
|---|---|---|---|---|---|
| | | | | | • Stevceva L, et al. 2001. BMC Clin Pathol. 1:3. (IHC)<br>• Kobayashi M, et al.2008. J. Leukoc. Biol. 83:1354 |
| anti-CD11c-APC-Cy7 | HL3 | BD Pharmingen | 561241 | 8235909 | Broadly used by many laboratories and flow cytometric cell staining demonstrated in this manuscript.<br>• Barclay NA, Brown MH, Birkeland ML, et al, ed. The Leukocyte Antigen FactsBook. San Diego, CA: Academic Press; 1997.<br>• Gao JX, Liu X, Wen J, et al. Differentiation of monocytic cell clones into CD8 alpha+ dendritic cells (DC) suggests that monocytes can be direct precursors for both CD8 alpha+ and CD8 alpha- DC in the mouse. J Immunol. 2003; 170(12):5927-5935. (Biology).<br>• Huleatt JW, Lefrançois L. Antigen-driven induction of CD11c on intestinal intraepithelial lymphocytes and CD8+ T cells in vivo. J Immunol. 1995; 154(11):5684-93. (Immunogen). |
| CIS43 | NA | Human recombinant CIS43 (Kisalu et al., 2018) | NA | NA | validated in Kisalu et al, 2018 by ELISA, ITC and X-ray-crystallography, and in this paper by BLI in Figure S3. |
| 1210 | NA | Human recombinant 1210 (Imkeller et al., 2018) | NA | NA | validated in Imkeller et al, 2018 by ELISA, ITC and X-ray crystallography and in this paper by BLI in Figure S3. |
| 1710 | NA | Human recombinant 1710 (Scally et al., 2018) | NA | NA | validated in Scally et al, 2018 by ELISA, BLI and X-ray crystallography and in this paper by BLI in Figure S3. |

| | | | | | |
|---|---|---|---|---|---|
| 5D5 | NA | Humanized recombinant 5D5 (Espinosa et al., 2015; Thai et al., 2020) | NA | NA | validated in Thai et al, 2020 by BLI and X-ray crystallography and in this paper by BLI in Figure S3. |
| Anti-Mouse IgG-Alexa Fluor® 555 | Polyclonal | Molecular Probes | A-21422 | 1094410 | Secondary Antibody, NA |
| | | | | | |