Appendix, table E1. Assessed data.

Assessed data	Item list
Family constellation	
Biometric information, including sex at birth, body height, body weight, ethnicity, number of children and grand-children and profession	
Tests for SARS-CoV-2 in the last 4 weeks, inlcuding type, date and result	
Vaccination against SARS-CoV-2, including manufacturer, date(s) and symptoms after vaccination	
Temporal course of infection in the family	
Index case and assumed point of infection	
Information about household, including number of household members, pet animals and information about smoking (including smoked substances)	
Paediatric anamnesis, including biological parents and legal parents, gestational week at birth, problems during pregnancy, birth and perinatal period, childcare or daycare or school or training, previous RSV infections	
Medical conditions and therapy, including smoking (including substance and length), atopic disposition, allergies, asthma and previous hospitalizations	
Symptoms (including duration) as listed and symptom severity according to WHO scales	Fever or elevated temperature, exhaustion and fatigue, loss of appetite, dizziness, diarrhea, nausea, vomiting, impaired sense of taste or smell, rhinorrhea, congested nose, sneezing, unproductive cough, productive cough, shortness of breath, headache, abdominal pain, sore throat, muscle ache, chest pain, ear pain, joint pain, conjunctivitis, lymphadenitis, skin efflorescence, shivering, paresis, impaired consciousness
Hygiene measures to prevent spreading of disease	
Ability to smell, including parosmia	
Consequences of SARS-CoV-2 infection, including hospitalization, concomitant disease(s) and treatment	
Symptoms of Post-COVID or fatigue as listed, including their duration, whether dependent in occurence or intensity on physical or mental stress or posture	Pain, nausea, vomiting, lymphadenitis, loss of appetite, abdominal complaint(s), loss of smell or taste, palpitation, shortness of breath, coughing, fever, skin efflorescence, conjunctivitis, hair loss, reddening or swelling of hand or feet, reddening or swelling of tongue, lips or cheeks, exhaustion or fatigue, reduced physical and exercise capacity, sleep ineffectivity, disturbed sleeping rhythm, disturbed sleep duration, mood disturbance, cognitive impairment, behavioral changes, sadness or depression, restlessness or excitement, orthostatic dysregulation, increased sensibility to trigges, such as light, noise, smell, touch or others, impaired adaptability to cold, heat or others