

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Prenatal Exposure to Air Pollution and Pre-Labor Rupture of Membranes in a Prospective Cohort Study: The Role of Maternal Hemoglobin and Iron Supplementation

Lin Wu, Wan-jun Yin, Li-jun Yu, Yu-hong Wang, Xiao-min Jiang, Ying Zhang, Fang-biao Tao, Rui-xue Tao, and Peng Zhu

Table of Contents

Table S1. The general characteristics of the study population during 2015 to 2021 in Hefei [n (%)].

Table S2. Correlations of air pollution in the prenatal period.

Table S3. The PROM risk in association with week-specific prenatal air pollution exposure during pregnancy.

Table S4. Cumulative effects and average effects between air pollutant exposure and PROM risk Distributed Lag Model and logistic regression model.

Table S5. Cumulative effects and average effects between air pollutant exposure and PROM risk Distributed Lag Model and logistic regression model after adjusting the trimester.

Table S6. Association of prenatal air pollution exposure with the term PROM and preterm PROM.

Table S7. Association of air pollution exposure and PROM in co-pollutant models.

Table S8. The estimated change in hemoglobin levels in third trimester was calculated for each quartile and each unit increment in PM_{2.5}, PM₁₀, SO₂, and CO during the second and third trimesters using linear regression model.

Table S9. Association of air pollution exposure and hemoglobin levels in co-pollutant models.

Table S10. The relationship between hemoglobin levels and PROM risk in logistic models.

Table S11. Correlation matrix eigenvalues of multiple air pollutants model on second to third trimester and corresponding eigenvector.

Table S12. The association between iron supplementation and PROM risk stratified by air

Figure S1. Directed acyclic graph for the association between air pollution exposure and PROM
GDM: Gestational diabetes mellitus, PROM: pre-labor rupture of membranes.